

T. Ikeda · Y. Kanno · K. Ozaki · A. Shinada

Metabolic rates of epipelagic marine copepods as a function of body mass and temperature

Published online: 31 October 2001
© Springer-Verlag 2001

Mar Biol (2001) 139:587–596

In the third sentence of the Abstract, reference is made to “22 species and 32 data sets for phosphate excretion”. The correct number of species is 20.

In the section of the Discussion entitled “Metabolism–growth coupling”, owing to conversion problems a

wrong character appears in the equations and text relating to them. In each case, ϕ should be a prime sign (ϕ').

The online version of the original article can be found at <http://dx.doi.org/10.1007/s002270100608>

T. Ikeda (✉) · Y. Kanno · K. Ozaki · A. Shinada
Faculty of Fisheries Science, Hokkaido University,
3-1-1 Minato-cho, Hakodate 041-0821, Japan

E-mail: tom@pop.fish.hokudai.ac.jp
Tel.: + 81-138-405541
Fax: + 81-138-405542

Present address: K. Ozaki
Faculty of Applied Biological Science,
Hiroshima University, Saijo-cho,
Higashi-Hiroshima 739-8528, Japan

Present address: A. Shinada
Institute of Hydrospheric-Atmospheric Sciences,
Nagoya University, Furo-cho, Chikusa-ku,
Nagoya 464-8601, Japan