

Stefano Lisini, Antonio Marigonda

On a class of modified Wasserstein distances induced by concave mobility functions defined on bounded intervals

Received: 12 September 2009 / Revised: 8 April 2010

Published online: 27 June 2010

Abstract. We study a new class of distances between Radon measures similar to those studied in Dolbeault et al. (Calc Var Partial Differ Equ 34:193–231, 2009). These distances (more correctly pseudo-distances because can assume the value $+\infty$) are defined generalizing the dynamical formulation of the Wasserstein distance by means of a concave mobility function. We are mainly interested in the physical interesting case (not considered in Dolbeault et al. (Calc Var Partial Differ Equ 34:193–231, 2009)) of a concave mobility function defined in a bounded interval. We state the basic properties of the space of measures endowed with this pseudo-distance. Finally, we study in detail two cases: the set of measures defined in \mathbb{R}^d with finite moments and the set of measures defined in a bounded convex set. In the two cases we give sufficient conditions for the convergence of sequences with respect to the distance and we prove a property of boundedness.

1. Introduction

In [13], Dolbeault, Nazaret and Savaré introduce and study the basic properties of a new class of distances between non-negative Radon measures on \mathbb{R}^d . These distances are defined generalizing the dynamical characterization of the Wasserstein distance. We briefly recall that the Wasserstein distance between two non-negative measures with the same mass can be defined as a relaxed optimal transportation problem (see [3, 26, 27] for a reference on this interesting topic)

$$W_p(\mu_0, \mu_1) := \inf \left\{ \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} |y - x|^p d\Sigma \right)^{\frac{1}{p}} : \Sigma \in \Gamma(\mu_0, \mu_1) \right\} \quad (1)$$

where $\Gamma(\mu_0, \mu_1)$ is the set of all *transport plans* between μ_0 and μ_1 : they are non-negative measures Σ on $\mathbb{R}^d \times \mathbb{R}^d$ with the same mass of μ_0 and μ_1 whose

S. Lisini (✉): Dipartimento di Matematica, „F. Casorati“, Università degli Studi di Pavia, Via Ferrata 1, 27100 Pavia, Italy. e-mail: stefano.lisini@unipv.it

S. Lisini: LAMA (Laboratorio di matematica applicata), Fondazione per l'Università di Mantova, Mantua, Italy.

A. Marigonda: Dipartimento di Informatica, Università degli Studi di Verona, Strada Le Grazie 15, 37134 Verona, Italy. e-mail: antonio.marigonda@univr.it

Mathematics Subject Classification (2000): 49J27, 49J40.

first and second marginals are respectively μ_0 and μ_1 , i.e. $\Sigma(B \times \mathbb{R}^d) = \mu_0(B)$ and $\Sigma(\mathbb{R}^d \times B) = \mu_1(B)$ for all Borel set B of \mathbb{R}^d .

In [4], Benamou and Brenier prove that the Wasserstein distance defined in (1) can be characterized, for absolutely continuous measures with respect to the Lebesgue measure \mathcal{L}^d , with compactly supported smooth densities, as follows

$$W_p^p(\mu_0, \mu_1) = \inf \left\{ \int_0^1 \int_{\mathbb{R}^d} |\mathbf{v}_t(x)|^p \rho_t(x) dx dt : \partial_t \rho_t + \nabla \cdot (\rho_t \mathbf{v}_t) = 0 \text{ in } \mathbb{R}^d \times (0, 1), \mu_0 = \rho_{|t=0} \mathcal{L}^d, \mu_1 = \rho_{|t=1} \mathcal{L}^d \right\}. \quad (2)$$

The proof of the dynamical characterization for general non-negative Borel measures was given in [3] where the continuity equation in (2) was considered in distributional sense.

The generalization of (2) studied in [13], roughly speaking, replace the mobility coefficient ρ in (2) with a non-linear one $h(\rho)$, where $h : [0, +\infty) \rightarrow [0, +\infty)$ is a concave increasing function such that $h(0) = 0$ (particularly important examples are the functions $h(\rho) = \rho^\alpha, \alpha \geq 0$) and the new “distance” is defined modifying (2) as follows

$$W_{p,h}^p(\mu_0, \mu_1) = \inf \left\{ \int_0^1 \int_{\mathbb{R}^d} |\mathbf{v}_t(x)|^p h(\rho_t(x)) dx dt : \partial_t \rho_t + \nabla \cdot (h(\rho_t) \mathbf{v}_t) = 0 \text{ in } \mathbb{R}^d \times (0, 1), \mu_0 = \rho_{|t=0} \mathcal{L}^d, \mu_1 = \rho_{|t=1} \mathcal{L}^d \right\}. \quad (3)$$

This “definition” is not rightly stated because it is necessary to specify the spaces where ρ and \mathbf{v} has to belong, and the notion of solution of the modified continuity equation in (3). The right framework is that of Radon measures and distributional solutions.

The motivation for studying distances defined like in (3) arises from physical problems. Indeed many interesting models are described by partial differential equations whose solutions can be seen as trajectory of the gradient flow of a suitable energy functional with respect to this distance (see for instance the introduction of [11, 13]).

On the other hand, the concave mobility $h(\rho) \geq 0$ considered in [13] is defined on the unbounded interval $[0, +\infty)$ and has to be necessarily non-decreasing. If we want to consider non-monotone concave mobilities $h(\rho) \geq 0$, then the domain of h has to be a bounded interval. This case, not considered in [13], is physically interesting. Indeed, examples of equations that can be modeled as gradient flows with respect to this kind of distances are a version of Cahn-Hilliard equation [14], some equation modelling chemotaxis with prevention of overcrowding [7, 8, 12],

equations describing the relaxation of gas of fermions [9, 10, 15, 16, 20, 21], studies of phase segregation [18, 25], and studies of thin liquid films [5].

The principal example of mobility function in the papers cited above is

$$h(\rho) = \rho(1 - \rho), \quad \text{defined in } [0, 1],$$

or $h(\rho) = 1 - \rho^2$ defined in $[-1, 1]$, mainly for the Cahn-Hilliard equation, the relaxation of fermion gas and the chemotaxis with overcrowding prevention. A more general example is of the form $h(\rho) = (\rho - a)^\alpha(b - \rho)^\beta$ defined in $[a, b]$ for some $\alpha, \beta \in [0, 1]$. In the previous examples, if $a < 0$ then the density could be negative at some points and we have to consider signed measures instead of non-negative measures.

The first goal of the paper is to show that an application on the couple of measures like in (3) (see the precise definition (29)) is really a pseudo-distance (it can be equal to $+\infty$) also in the case of the mobility h defined on a bounded interval, without assuming monotonicity of the mobility and sign of the measure. We are able to prove that a large part of the properties studied in [13] can be extended to the new case. The second goal of the paper is to provide sufficient conditions for the finiteness of the new distance and for the convergence of sequences.

As previously observed, in order to give a precise meaning of the dynamical characterization (2) and to define in a rigorous way the modified distance (3), the right framework is that of time dependent families of Radon measures and distributional solutions of the continuity equation. Following the explanation given in the introduction of [13], we replace ρ_t by a continuous curve $t \in [0, 1] \mapsto \mu_t$ ($\mu_t = \rho_t \mathcal{L}^d$ in the absolutely continuous case) in the space $\mathcal{M}^+(\mathbb{R}^d)$ of nonnegative Radon measures in \mathbb{R}^d endowed with the usual weak* topology. We replace the vector field \mathbf{v}_t in (2) with a time dependent family of vector measures $\mathbf{v}_t := \mathbf{v}_t \mu_t \ll \mu_t$. The continuity equation in (2) can be written in terms of the couple (μ, \mathbf{v})

$$\partial_t \mu_t + \nabla \cdot \mathbf{v}_t = 0 \quad \text{in the sense of distributions in } \mathcal{D}'(\mathbb{R}^d \times (0, 1)), \quad (4)$$

and it is a linear equation. Since $\mathbf{v}_t = d\mathbf{v}_t/d\mu_t$ is the density of \mathbf{v}_t with respect to μ_t , the action functional which has to be minimized in (2) is

$$\int_0^1 \Phi(\mu_t, \mathbf{v}_t) dt, \quad \Phi(\mu, \mathbf{v}) := \int_{\mathbb{R}^d} \left| \frac{d\mathbf{v}}{d\mu} \right|^p d\mu. \quad (5)$$

In the case of absolutely continuous measures with respect to \mathcal{L}^d , i.e. $\mu = \rho \mathcal{L}^d$ and $\mathbf{v} = \mathbf{w} \mathcal{L}^d$, the functional Φ can be expressed as

$$\Phi(\mu, \mathbf{v}) := \int_{\mathbb{R}^d} \phi(\rho, \mathbf{w}) d\mathcal{L}^d(x), \quad \phi(\rho, \mathbf{w}) := \rho \left| \frac{\mathbf{w}}{\rho} \right|^p. \quad (6)$$

Denoting by $\mathcal{CE}(0, 1)$ the class of measure-valued distributional solutions (μ, \mathbf{v}) of the continuity equation (4), we can state the dynamical characterization of the Wasserstein distance as follows

$$W_p^p(\mu_0, \mu_1) := \inf \left\{ \int_0^1 \Phi(\mu_t, \mathbf{v}_t) dt : (\mu, \mathbf{v}) \in \mathcal{CE}(0, 1), \right. \\ \left. \mu|_{t=0} = \mu_0, \mu|_{t=1} = \mu_1 \right\} \tag{7}$$

(as already observed, the Benamou-Brenier characterization (7) for Borel non-negative measures was proven in [3]). We observe that the function ϕ defined in (6) is p -homogeneous w.r.t. \mathbf{w} , is convex with respect to (ρ, \mathbf{w}) , and positively 1-homogeneous with respect to (ρ, \mathbf{w}) . By the 1-homogeneity it is immediate to check that the functional Φ in (6) is independent on the Lebesgue measure, in the sense that if $\gamma \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$ is another reference measure such that $\text{supp}(\gamma) = \mathbb{R}^d$ and $\mu = \tilde{\rho}\gamma$ and $\mathbf{v} = \tilde{\mathbf{w}}\gamma$, then

$$\Phi(\mu, \mathbf{v}) = \int_{\mathbb{R}^d} \phi(\tilde{\rho}, \tilde{\mathbf{w}}) d\gamma. \tag{8}$$

We explain the main idea of [13] for state rigorously the intuitive “definition” (3). Given a concave mobility function $h : (a, b) \rightarrow (0, +\infty)$, we consider still the linear continuity equation (4) and modify the action density ϕ in the following way: $\phi : (a, b) \times \mathbb{R}^d \rightarrow [0, +\infty)$

$$\phi(\rho, \mathbf{w}) := h(\rho) \left| \frac{\mathbf{w}}{h(\rho)} \right|^p. \tag{9}$$

The concavity of h is a necessary and sufficient condition for the convexity of ϕ in (9) (see [24] and Theorem 7). We observe that ϕ still satisfies the p -homogeneity with respect to \mathbf{w} and is globally convex, but it is no longer positively 1-homogeneous with respect to (ρ, \mathbf{w}) . Hence, in order to consider the integral functional Φ like (8) it is necessary to precise the reference measure $\gamma \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$ for that ρ and \mathbf{w} are the densities of μ and \mathbf{v} respectively.

Defining

$$\Phi(\mu, \mathbf{v}|\gamma) := \begin{cases} \int_{\mathbb{R}^d} \phi(\rho, \mathbf{w}) d\gamma & \text{if } \mu = \rho\gamma \ll \gamma, \mathbf{v} = \mathbf{w}\gamma \ll \gamma \\ +\infty & \text{otherwise} \end{cases},$$

the definition of the generalized Wasserstein distance associated to (ϕ, γ) is therefore

$$\mathcal{W}_{\phi, \gamma}^p(\mu_0, \mu_1) := \inf \left\{ \int_0^1 \Phi(\mu_t, \mathbf{v}_t) dt : (\mu, \mathbf{v}) \in \mathcal{CE}(0, 1), \right. \\ \left. \mu|_{t=0} = \mu_0, \mu|_{t=1} = \mu_1 \right\}. \tag{10}$$

Particularly important for the applications are the following choices of γ :

- $\gamma := \mathcal{L}_{|\Omega}^d = \chi_{\Omega} \mathcal{L}^d$, with Ω an open subset of \mathbb{R}^d ;
- $\gamma := e^{-V} \mathcal{L}^d$ for some C^1 potential $V : \mathbb{R}^d \rightarrow \mathbb{R}$;
- $\gamma := \mathcal{H}_{\mathbb{M}}^k$, where \mathbb{M} is a smooth k -dimensional manifold embedded in \mathbb{R}^d with the Riemannian metric induced by the Euclidean distance and \mathcal{H}^k denotes the k -dimensional Hausdorff measure.

In the paper [11], the authors used this kind of distance in the case $\gamma = \mathcal{L}_{|\Omega}^d$ in order to study the problem of the convexity of integral functionals along geodesics induced by the distance. The forthcoming paper [22] will be devoted to the study of fourth orders equations (Cahn-Hilliard type with nonlinear mobility and thin-film like equations), with the proof of the existence of solutions by means of the minimizing movements approximation scheme (see [3]) for the distance like (10) and a first order integral functional.

We conclude this introduction stating the principal properties obtained in this paper for the distance like (10) with $h : (a, b) \rightarrow (0, +\infty)$, referring to Sect. 3 for the precise definitions and the complete statements. We recall that the choice of consider the mobility with bounded domain (a, b) allow to consider also the distance between signed measures.

- The space $\mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ endowed with the distance $\mathcal{W}_{\phi, \gamma}$ is a complete pseudo-metric space (the distance can assume the value $+\infty$), inducing as strong as, or stronger topology than the weak* one. Bounded sets with respect to $\mathcal{W}_{\phi, \gamma}$ are weakly* relatively compact. The distance $\mathcal{W}_{\phi, \gamma}$ is lower semi continuous with respect to the weak* convergence.
- In order to avoid that the distance could be $+\infty$ we consider the space $\mathcal{M}[\sigma] := \{\mu \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d) : \mathcal{W}_{\phi, \gamma}(\mu, \sigma) < +\infty\}$ for a given measure $\sigma \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$. The space $\mathcal{M}[\sigma]$ turns out to be a complete metric space.
- $\mathcal{M}[\sigma]$ is a geodesic space and the geodesic are unique if h is strictly concave.
- If $\tilde{m}_{-q}(\gamma) < +\infty$, where q is the conjugate exponent of p and the generalized momentum is defined in Definition 7, then $\mu(\mathbb{R}^d) = \sigma(\mathbb{R}^d)$ for every $\mu \in \mathcal{M}[\sigma]$.

We would like to point out that the key to generalize the work [13] to the case of mobility h defined on bounded intervals, is Proposition 4: a compactness result on solutions of the continuity equation with equibounded action.

Finally, in Sect. 4 we give sufficient conditions on the measures μ_0, μ_1 in order to have finiteness of the distance $\mathcal{W}_{\phi, \gamma}(\mu_0, \mu_1)$, and we prove two results: one for the all space \mathbb{R}^d with the Lebesgue measure as a reference, the other one for convex bounded domains in \mathbb{R}^d . In the two cases we study also the relation between the weak-* convergence of measures and the convergence with respect to the distance $\mathcal{W}_{\phi, \gamma}$.

2. Preliminaries

In this section we introduce the necessary tools in order to define in the next Section the modified Wasserstein distance and prove its basic properties. The contents are an adaptation of Sects. 2–4 of [13].

2.1. Action function

Definition 1. (*Admissible action density functions*) Given $a, b \in \mathbb{R}, a < b$, and $1 < p < \infty$, we say that $\phi \in \mathcal{A}_p(a, b)$ if the following properties holds:

- (ϕ_1) $\phi : \mathbb{R} \times \mathbb{R}^d \rightarrow [0, +\infty]$ is a l.s.c. nonnegative convex function,
- (ϕ_2) $\text{int}(\text{dom}(\phi)) = (a, b) \times \mathbb{R}^d$,
- (ϕ_3) $\mathbf{w} \mapsto \phi(\cdot, \mathbf{w})$ is p -homogeneous, i.e. for every $\rho \in \mathbb{R}$ such that $\{\rho\} \times \mathbb{R}^d \subset \text{dom}(\phi)$ we have $\phi(\rho, 0) = 0$ and for every $\lambda \neq 0, \mathbf{w} \in \mathbb{R}^d$ we have $\phi(\rho, \lambda \mathbf{w}) = |\lambda|^p \phi(\rho, \mathbf{w})$,
- (ϕ_4) there exists $\rho_0 \in \mathbb{R}$ such that $\{\rho_0\} \times \mathbb{R}^d \subseteq \text{dom}(\phi)$ and $\phi(\rho_0, \mathbf{w}) > 0$ for all $\mathbf{w} \neq 0$.

Let q be the conjugate exponent of p . We construct the *partial dual* $\mathcal{A}_q^*(a, b)$ of $\mathcal{A}_p(a, b)$ as follows: for all $\phi \in \mathcal{A}_p(a, b)$, we define the concave–convex (i.e. concave w.r.t. the first variable, convex w.r.t. the second variable, see Appendix 4.2) function $\tilde{\phi} : \text{dom}(\phi) \rightarrow \mathbb{R} \cup \{+\infty\}$ by setting:

$$\frac{1}{q} \tilde{\phi}(\rho, \mathbf{z}) := \sup_{\mathbf{w} \in \mathbb{R}^d} \left\{ \langle \mathbf{z}, \mathbf{w} \rangle - \frac{1}{p} \phi(\rho, \mathbf{w}) \right\}. \tag{11}$$

We will call the lower extension (see Appendix 4.2) of $\tilde{\phi}$ the *marginal conjugate* of ϕ and we will still denote it by $\tilde{\phi}$. We observe that $\tilde{\phi}$ is q -homogeneous with respect to the second variable and $\tilde{\phi}(\rho, \mathbf{z}) \geq 0$. We define:

$$\mathcal{A}_q^*(a, b) := \{ \tilde{\phi} : \tilde{\phi} \text{ is the marginal conjugate of } \phi, \phi \in \mathcal{A}_p(a, b) \}$$

and it is easy to check that $\text{int}(\text{dom}(\tilde{\phi})) = (a, b) \times \mathbb{R}^d$.

The following proposition can be proved exactly as Theorem 3.1 of [13], taking into account also Theorem 7 in the Appendix 4.2.

Proposition 1. (ϕ -norm) *Let $\phi \in \mathcal{A}_p(a, b)$. The following properties hold:*

1. *For every $\rho \in [a, b]$ such that $\{\rho\} \times \mathbb{R}^d \subset \text{dom}(\phi)$, the functions $\mathbf{w} \mapsto \phi(\rho, \mathbf{w})^{1/p}$ and $\mathbf{z} \mapsto \tilde{\phi}(\rho, \mathbf{z})^{1/q}$ are norms on \mathbb{R}^d each one dual of the other. We have:*

$$\begin{aligned} \|\mathbf{z}\|_{(\phi, \rho)^*} &:= \tilde{\phi}(\rho, \mathbf{z})^{1/q} = \sup_{\mathbf{w} \neq 0} \frac{\langle \mathbf{w}, \mathbf{z} \rangle}{\phi(\rho, \mathbf{w})^{1/p}}, \\ \|\mathbf{w}\|_{(\phi, \rho)} &:= \phi(\rho, \mathbf{w})^{1/p} = \sup_{\mathbf{z} \neq 0} \frac{\langle \mathbf{w}, \mathbf{z} \rangle}{\tilde{\phi}(\rho, \mathbf{z})^{1/q}}. \end{aligned} \tag{12}$$

2. *The restriction to $\text{dom}(\tilde{\phi})$ of the marginal conjugate $\tilde{\phi}$ of ϕ takes its values in $[0, +\infty)$ and it is a concave–convex function.*
3. *Given $\rho_0, \rho_1 \in \mathbb{R}$ with $[\rho_0, \rho_1] \times \mathbb{R}^d \subseteq \text{dom}(\phi)$, there exists a constant $C = C(\rho_0, \rho_1)$ such that for every $\rho \in [\rho_0, \rho_1]$ it holds:*

$$\begin{aligned} C^{-1} |\mathbf{w}|^p &\leq \phi(\rho, \mathbf{w}) \leq C |\mathbf{w}|^p, \\ C^{-1} |\mathbf{z}|^q &\leq \tilde{\phi}(\rho, \mathbf{z}) \leq C |\mathbf{z}|^q, \quad \forall \mathbf{w}, \mathbf{z} \in \mathbb{R}^d, \end{aligned}$$

where $|\mathbf{w}|$ denotes the euclidean norm of \mathbf{w} .

Moreover, a function ϕ belongs to $\mathcal{A}_p(a, b)$ if and only if it admits the dual representation formula

$$\frac{1}{p}\phi(\rho, \mathbf{w}) := \sup_{\mathbf{z} \in \mathbb{R}^d} \left\{ \langle \mathbf{z}, \mathbf{w} \rangle - \frac{1}{q}\tilde{\phi}(\rho, \mathbf{z}) \right\}, \tag{13}$$

where $\tilde{\phi} : \mathbb{R} \times \mathbb{R}^d \rightarrow [0, \infty)$ is (the lower extension of) a nonnegative concave-convex function which is q -homogeneous with respect to \mathbf{z} and $\text{int}(\text{dom}(\tilde{\phi})) = (a, b) \times \mathbb{R}^d$.

One of the most interesting example of admissible density function in $\mathcal{A}_p(a, b)$ is the following:

Definition 2. Let $p > 1$ and q its conjugate exponent. Let $h : \mathbb{R} \rightarrow [0, +\infty) \cup \{-\infty\}$ be an u.s.c. concave function with $\text{int}(\text{dom}(h)) = (a, b)$, $a, b \in \mathbb{R}$, $a < b$, $h(\rho) > 0$ for every $\rho \in (a, b)$. Define $\tilde{\phi}_h(\rho, \mathbf{z}) = h(\rho)|\mathbf{z}|^q$ on $\mathbb{R} \times \mathbb{R}^d$. We have that this is a concave-convex map which is q -homogeneous with respect to \mathbf{z} . Hence, it is the marginal conjugate of the l.s.c. convex map $\phi_h \in \mathcal{A}_p(a, b)$ defined by

$$\phi_h(\rho, \mathbf{w}) = \begin{cases} \frac{|\mathbf{w}|^p}{h(\rho)^{p-1}} & \text{if } \rho \in \text{dom}(h), h(\rho) \neq 0 \\ 0 & \text{if } h(\rho) = 0, \mathbf{w} = 0 \\ +\infty & \text{if } h(\rho) = 0, \mathbf{w} \neq 0 \text{ or } h(\rho) = -\infty. \end{cases} \tag{14}$$

Such function h is called *mobility function*.

The following proposition shows that every admissible function ϕ is bounded from above by an admissible function of the type (14).

Proposition 2. If $\phi \in \mathcal{A}_p(a, b)$, then there exists a concave function h such that $\text{int}(\text{dom}(h)) = (a, b)$, $h(r) > 0$ for every $r \in (a, b)$ and

$$\phi(r, \mathbf{w}) \leq \phi_h(r, \mathbf{w}). \tag{15}$$

Proof. Let us define

$$h(r) := \inf_{|\mathbf{z}|=1} \tilde{\phi}(r, \mathbf{z}),$$

where $\tilde{\phi}$ is defined in (11). By the q -homogeneity of $\tilde{\phi}$ we have

$$\tilde{\phi}(r, \mathbf{z}) \geq h(r)|\mathbf{z}|^q.$$

Then, by the representation (13) for ϕ and ϕ_h , we obtain

$$\frac{1}{p}\phi(r, \mathbf{w}) \leq \sup_{\mathbf{z} \in \mathbb{R}^d} \left\{ \langle \mathbf{w}, \mathbf{z} \rangle - \frac{1}{q}h(r)|\mathbf{z}|^q \right\} = \frac{1}{p}\phi_h(r, \mathbf{w}).$$

2.2. Action functional

Given an admissible action density function $\phi \in \mathcal{A}_p(a, b)$ and a reference measure γ on \mathbb{R}^d , we can define the corresponding action functional.

Definition 3. (*ϕ -Action functional*) Let $\gamma \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$ be a reference measure and $\phi \in \mathcal{A}_p(a, b)$.

For every $\mu \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ and $\mathbf{v} \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^d)$ we define

$$\Phi(\mu, \mathbf{v}|\gamma) := \begin{cases} \int_{\mathbb{R}^d} \phi(\rho, \mathbf{w}) \, d\gamma & \text{if } \mu = \rho\gamma \ll \gamma, \quad \mathbf{v} = \mathbf{w}\gamma \ll \gamma \\ +\infty & \text{otherwise} \end{cases} \quad (16)$$

It is a direct consequence of Definition 3 that

$$\Phi(\mu, \mathbf{v}) < +\infty \implies a \leq \rho(x) \leq b \text{ for } \gamma\text{-a.e. } x \in \mathbb{R}^d, \quad (17)$$

where $\mu = \rho\gamma$.

The property (17) will be crucial in order to prove the compactness of the solutions of the continuity equation with equibounded energy stated in Proposition 4.

We collect in the following theorem some properties of the action functional. The proof can be found in [13] (see also [2] for functionals defined on measures).

Theorem 1. (Properties of Φ) Let $\phi \in \mathcal{A}_p(a, b)$ and Φ as in Definition 3.

1. Lower semicontinuity. If three sequences $(\gamma_n)_{n \in \mathbb{N}} \subset \mathcal{M}_{\text{loc}}^+(\mathbb{R}^k)$, $(\mu_n)_{n \in \mathbb{N}} \subset \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$, $(\mathbf{v}_n)_{n \in \mathbb{N}} \subset \mathcal{M}_{\text{loc}}(\mathbb{R}^d, \mathbb{R}^d)$ weakly* converge to γ, μ, \mathbf{v} respectively, then $\Phi(\mu, \mathbf{v}|\gamma) \leq \liminf_{n \rightarrow +\infty} \Phi(\mu_n, \mathbf{v}_n|\gamma_n)$.
2. Monotonicity with respect to γ . Assume that $(0, 0) \in \text{dom}(\phi)$ (in this case by homogeneity we have $\phi(0, 0) = 0$) and let $\gamma_1, \gamma_2 \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^k)$ be such that $\gamma_1 \leq \gamma_2$. Then $\Phi(\mu, \mathbf{v}|\gamma_2) \leq \Phi(\mu, \mathbf{v}|\gamma_1)$ for every (μ, \mathbf{v}) such that $\text{supp}(\mu) \cup \text{supp}(\mathbf{v}) \subseteq \text{supp}(\gamma_i)$, $i = 1, 2$.
3. Monotonicity with respect to convolution. Let $k \in C_c^\infty(\mathbb{R}^d)$ be a convolution kernel satisfying $k(x) \geq 0$ for all $x \in \mathbb{R}^d$ and $\int_{\mathbb{R}^d} k(x) \, dx = 1$. Then $\Phi(\mu * k, \mathbf{v} * k|\gamma * k) \leq \Phi(\mu, \mathbf{v}|\gamma)$.

The following example shows that the statement on monotonicity with respect to the reference measure may fail if $(0, 0) \notin \text{dom}(\phi)$.

Example 1 (Non-monotonicity). Let $d = 1$. We define $\phi : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$ to be $\phi(r, \mathbf{v}) = |\mathbf{v}|^2$ if $r \in [3/2, 2]$ and $+\infty$ elsewhere. Define $\gamma_2 = 3/2\gamma_1 = \chi_{[1,2]}(x)\mathcal{L}^1$ and set $\mu = \mathbf{v} = \gamma_2 = 3/2\gamma_1$. Then

$$\begin{aligned} \Phi(\mu, \mathbf{v}|\gamma_2) &= \int_{\mathbb{R}} \phi(1, 1) \, d\gamma_2 = +\infty. \\ \Phi(\mu, \mathbf{v}|\gamma_1) &= \int_{\mathbb{R}} \phi(3/2, 3/2) \, d\gamma_1 = \frac{3}{2}. \end{aligned}$$

Hence $\gamma_1 < \gamma_2$ but $\Phi(\mu, \mathbf{v}|\gamma_1) < \Phi(\mu, \mathbf{v}|\gamma_2)$.

Remark 1. For a general convex function $\phi : \mathbb{R} \times \mathbb{R}^d \rightarrow [0, +\infty]$, according with the general theory of convex functionals defined on measures, in order to get the lower semi continuity with respect to the weak* convergence of measures, the integral functional of ϕ has to be defined as follows: for every $\mu \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ and $\mathbf{v} \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^d)$ such that $\text{supp}(\mu)$ and $\text{supp}(\mathbf{v})$ are contained in $\text{supp}(\gamma)$ we write their Lebesgue decomposition $\mu = \rho\gamma + \mu^\perp$, $\mathbf{v} = \mathbf{w}\gamma + \mathbf{v}^\perp$. Introducing a nonnegative Radon measure $\sigma \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$ such that $\mu^\perp \ll \sigma$ and $\mathbf{v}^\perp \ll \sigma$ (e.g. take $\sigma = |\mu^\perp| + |\mathbf{v}^\perp|$) and using the notation $\mu^\perp = \rho^\perp\sigma$ and $\mathbf{v}^\perp = \mathbf{w}^\perp\sigma$, we define the action functional Φ associated to ϕ by

$$\Phi(\mu, \mathbf{v}|\gamma) := \int_{\mathbb{R}^d} \phi(\rho, \mathbf{w}) \, d\gamma + \int_{\mathbb{R}^d} \phi^\infty(\rho^\perp, \mathbf{w}^\perp) \, d\sigma. \tag{18}$$

See the Appendix 4.2 for the definition of ϕ^∞ . Since ϕ^∞ is 1-homogeneous, (18) does not depend on σ .

If we assume that $\phi \in \mathcal{A}_p(a, b)$, then we can avoid this complicated definition, because we have $\phi^\infty(0, 0) = 0$ and $\phi^\infty(\rho, \mathbf{w}) = +\infty$ for $(\rho, \mathbf{w}) \neq (0, 0)$. Indeed, for every $\bar{\rho} \in \mathbb{R}$ such that $(\bar{\rho}, 0) \in \text{dom}(\phi)$, it holds

$$\begin{aligned} \phi^\infty(\rho, \mathbf{w}) &:= \lim_{\lambda \rightarrow +\infty} \frac{\phi(\bar{\rho} + \lambda\rho, \lambda\mathbf{w}) - \phi(\bar{\rho}, 0)}{\lambda} = \lim_{\lambda \rightarrow +\infty} \frac{\phi(\bar{\rho} + \lambda\rho, \lambda\mathbf{w})}{\lambda} \\ &= \lim_{\lambda \rightarrow +\infty} \lambda^{p-1} \phi(\bar{\rho} + \lambda\rho, \mathbf{w}). \end{aligned}$$

From the previous formula, for λ sufficiently large we have $(\bar{\rho} + \lambda\rho, \mathbf{w}) \notin \text{dom}(\phi)$.

Then $\int_{\mathbb{R}^d} \phi^\infty(\rho^\perp, \mathbf{w}^\perp) \, d\sigma < +\infty$ if and only if $\rho^\perp(x) = 0$ and $\mathbf{w}^\perp(x) = 0$ for σ -a.e. $x \in \mathbb{R}^d$. This implies that $\rho \ll \gamma$ and $\mathbf{v} \ll \gamma$ and the definition as in (18) coincides with the definition of Φ in (16).

2.3. Continuity equation

In this section we collect the basic facts on the measure solutions of the continuity equation. It is an adaptation of [3, 13], with the novelty that here we consider signed measures instead of non-negative measures.

Definition 4. Given $T > 0$, we consider the *continuity equation*:

$$\partial_t \mu_t + \text{div}(\mathbf{v}_t) = 0, \quad \text{in } \mathbb{R}^d \times (0, T), \tag{19}$$

where μ_t, \mathbf{v}_t are Borel families of measures in $\mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ and $\mathcal{M}_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^d)$ respectively, defined for $t \in (0, T)$ satisfying for all $R > 0$

$$\int_0^T |\mu_t|(B(0, R)) \, dt < +\infty, \quad V_R := \int_0^T |\mathbf{v}_t|(B(0, R)) \, dt < +\infty, \tag{20}$$

and the Eq. 19 holds in the sense of distributions, i.e.

$$\int_0^T \int_{\mathbb{R}^d} \partial_t \zeta(x, t) \, d\mu_t(x) \, dt + \int_0^T \int_{\mathbb{R}^d} \nabla_x(\zeta(x, t)) \, d\mathbf{v}_t(x) \, dt = 0, \quad (21)$$

for every $\zeta \in C_c^1(\mathbb{R}^d \times (0, T))$.

We recall that, thanks to the disintegration theorem, we can identify $(\mathbf{v}_t)_{t \in [0, T]}$ with the measure $\mathbf{v} = \int_0^T \mathbf{v}_t \, dt \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d \times (0, T); \mathbb{R}^d)$ defined by:

$$\langle \mathbf{v}, \zeta \rangle = \int_0^T \left(\int_{\mathbb{R}^d} \zeta(x, t) \, d\mathbf{v}_t \right) \, dt, \quad \forall \zeta \in C_c^0(\mathbb{R}^d \times (0, T); \mathbb{R}^d).$$

Similarly, we can identify $(\mu_t)_{t \in [0, T]}$ with a measure

$$\mu = \int_0^T \mu_t \, dt \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d \times (0, T)).$$

Lemma 1. *Let $T > 0$ and $(\mu_t, \mathbf{v}_t)_{t \in (0, T)}$ be a Borel family of measures satisfying (20) and (21). Then there exists a unique weakly* continuous curve $[0, T] \ni t \mapsto \tilde{\mu}_t \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ such that $\mu_t = \tilde{\mu}_t$ for \mathcal{L}^1 -a.e. $t \in (0, T)$; if $\zeta \in C_c^1(\mathbb{R}^d \times (0, T))$ and $t_1, t_2 \in [0, T]$ with $t_1 \leq t_2$ we have:*

$$\begin{aligned} \int_{\mathbb{R}^d} \zeta(t_2, x) \, d\mu_{t_2} - \int_{\mathbb{R}^d} \zeta(t_1, x) \, d\mu_{t_1} \\ = \int_{t_1}^{t_2} \left(\int_{\mathbb{R}^d} \partial_t \zeta(t, x) \, d\mu_t(x) + \int_{\mathbb{R}^d} \nabla_x(\zeta(t, x)) \, d\mathbf{v}_t(x) \right) \, dt. \end{aligned}$$

Moreover if $\tilde{\mu}_s(\mathbb{R}^d) \in \mathbb{R}$ for some $s \in [0, T]$ and $\lim_{R \rightarrow +\infty} R^{-1} V_R = 0$, then the total mass $\tilde{\mu}_t(\mathbb{R}^d) \in \mathbb{R}$ and is constant.

Definition 5. (Solution of continuity equation) Let $T > 0$, we denote by $\mathcal{CE}(0, T)$ the set of time-dependent measures $(\mu_t, \mathbf{v}_t)_{t \in [0, T]}$ such that

1. $t \mapsto \mu_t$ is weakly* continuous in $\mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ satisfying (20);
2. $(\mathbf{v}_t)_{t \in [0, T]}$ is a Borel family satisfying (20);
3. (μ, \mathbf{v}) satisfies (21).

Given $\mu^1, \mu^2 \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$, we denote the set of solutions connecting μ^1 to μ^2 (possibly empty) by

$$\mathcal{CE}(0, T, \mu^1 \rightarrow \mu^2) = \{(\mu, \mathbf{v}) \in \mathcal{CE}(0, T) : \mu_0 = \mu^1, \mu_T = \mu^2\}.$$

Given $\gamma \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$ reference measure and $\phi \in \mathcal{A}_p$, we denote by

$$\mathcal{CE}_{\phi,\gamma}(0, T; \mu^1 \rightarrow \mu^2) = \left\{ (\mu, \mathbf{v}) \in \mathcal{CE}(0, T; \mu^1 \rightarrow \mu^2) : \int_0^T \Phi(\mu_t, \mathbf{v}_t | \gamma) dt < +\infty \right\},$$

which is the set of solutions of the continuity equation connecting μ^1 to μ^2 with finite energy. We also use the notation

$$\mathcal{CE}_{\phi,\gamma}(0, T) := \left\{ (\mu, \mathbf{v}) \in \mathcal{CE}(0, T) : \int_0^T \Phi(\mu_t, \mathbf{v}_t | \gamma) dt < +\infty \right\}.$$

Lemma 2. *The following properties hold:*

1. (Time rescaling) Let $\tau : [0, T'] \rightarrow [0, T]$ be a strictly increasing absolutely continuous map with absolutely continuous inverse $s = \tau^{-1}$. Then (μ, \mathbf{v}) is a distributional solution of (21) iff $(\hat{\mu}, \hat{\mathbf{v}})$, where $\hat{\mu} = \mu \circ \tau$ and $\hat{\mathbf{v}} = \tau'(\mathbf{v} \circ \tau)$ is a distributional solution of (21) on $(0, T')$.
2. (Gluing solution) Let $(\mu^1, \mathbf{v}^1) \in \mathcal{CE}(0, T_1)$, $(\mu^2, \mathbf{v}^2) \in \mathcal{CE}(0, T_2)$ with $\mu_{T_1}^1 = \mu_0^2$. Then the new family $(\mu_t, \mathbf{v}_t)_{t \in (0, T_1 + T_2)}$, defined by $(\mu_t, \mathbf{v}_t) = (\mu_t^1, \mathbf{v}_t^1)$ for $0 \leq t \leq T_1$ and $(\mu_t, \mathbf{v}_t) = (\mu_{t-T_1}^2, \mathbf{v}_{t-T_1}^2)$ for $T_1 \leq t \leq T_2$, belongs to $\mathcal{CE}(0, T_1 + T_2)$.

2.3.1. Conservation of the mass for solutions with finite energy In this paragraph we prove that, under a condition on the generalized moments of the reference measure γ and for $\phi \in \mathcal{A}_p(a, b)$, the total (signed) mass conserves for solutions of the continuity equation with finite energy.

Definition 6. (Upper uniform concave bound) Let $\phi \in \mathcal{A}_p(a, b)$. Fixing $\bar{\rho} := (a + b)/2$ we use the notation

$$\|\mathbf{w}\| := \|\mathbf{w}\|_{(\phi, \bar{\rho})}, \quad \|\mathbf{z}\|_* := \|\mathbf{z}\|_{(\phi, \bar{\rho})_*}, \tag{22}$$

where the norms above (equivalents to the euclidean one) are defined in (12). We consider the set: $\mathcal{H} := \{g : \mathbb{R} \rightarrow \mathbb{R} \cup \{-\infty\} : g \text{ is u.s.c., concave, } g(\rho) \geq \tilde{\phi}(\rho, \mathbf{z}/\|\mathbf{z}\|_*) \forall \mathbf{z} \neq 0\}$. This set is nonempty, and we can define:

$$h(\rho) = \inf\{g(\rho) : g \in \mathcal{H}\},$$

which turns out to be the smallest u.s.c. concave function greater than or equal to $\sup\{\tilde{\phi}(\rho, \mathbf{z}) : \|\mathbf{z}\|_* = 1\}$. Since $\text{int}(\text{dom}(h)) = (a, b)$ we obtain that

$$h_{max} := \sup_{\rho \in \mathbb{R}} h(\rho) < +\infty. \tag{23}$$

By homogeneity property it is immediate to prove that

$$\tilde{\phi}(\rho, \mathbf{z}) \leq h(\rho) \|\mathbf{z}\|_*^q \quad \text{and} \quad \|\mathbf{w}\| \leq h(\rho)^{1/q} \phi(\rho, \mathbf{w})^{1/p}. \tag{24}$$

When ϕ is given as in Definition 2, we have $h(\rho) = C \cdot h(\rho)$, where $C := \max\{|z|/\|z\|_* : z \neq 0\}$, and $|\cdot|$ is the euclidean norm.

Definition 7. Let $\gamma \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$, $r \in \mathbb{R}$. We define the *generalized r -th momentum* $\tilde{m}_r(\gamma)$ of γ by setting:

$$\tilde{m}_r(\gamma) := \gamma(B(0, 1)) + \int_{\mathbb{R}^d \setminus B(0, 1)} |x|^r \, d\gamma(x).$$

We observe that if $\tilde{m}_r(\gamma) < +\infty$ then $\tilde{m}_s(\gamma) < +\infty$ for every $s \leq r$.

Proposition 3. (Mass conservation) *Let $p > 1$, q its conjugate exponent and $\phi \in \mathcal{A}_p(a, b)$. Let $r \in \mathbb{R}$ such that $r \geq -q$ and $\gamma \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$ be a reference measure satisfying $\tilde{m}_r(\gamma) < +\infty$.*

If $(\mu_t, \mathbf{v}_t)_{t \in [0, T]} \in \mathcal{CE}_{\phi, \gamma}(0, T)$ and $\mu_0(\mathbb{R}^d) \in \mathbb{R}$, then $\mu_t(\mathbb{R}^d) = \mu_0(\mathbb{R}^d)$ for every $t \in [0, T]$.

Proof. We consider a cutoff function $\zeta \in C_c^\infty(\mathbb{R}^d)$ such that $\zeta(x) = 1$ if $|x| \leq 1$, $\zeta(x) = 0$ if $|x| \geq 2$ and $|\nabla \zeta(x)| \leq 1$ for all $x \in \mathbb{R}^d$. We consider the family $\zeta_R(x) = \zeta(x/R)$, for $R > 0$, that obviously satisfies $|\nabla \zeta_R(x)| \leq 1/R$ for all $x \in \mathbb{R}^d$.

Using the notations of Definition 6, for every $t_1, t_2 \in [0, T]$, $t_1 < t_2$, by Proposition 1 and (24) we have

$$\begin{aligned} \left| \int_{\mathbb{R}^d} \zeta_R \, d\mu_{t_1} - \int_{\mathbb{R}^d} \zeta_R \, d\mu_{t_2} \right| &\leq \int_{t_1}^{t_2} \int_{\mathbb{R}^d} |\nabla \zeta_R \cdot \mathbf{w}_t| \, d\gamma \, dt \\ &\leq \int_{t_1}^{t_2} \int_{\mathbb{R}^d} \tilde{\phi}(\rho_t, \nabla \zeta_R)^{1/q} \phi(\rho_t, \mathbf{w}_t)^{1/p} \, d\gamma \, dt \\ &\leq \left(\int_{t_1}^{t_2} \int_{B_{2R} \setminus B_R} h(\rho_t) \|\nabla \zeta_R\|_*^q \, d\gamma \, dt \right)^{1/q} \\ &\quad \left(\int_{t_1}^{t_2} \int_{\mathbb{R}^d} \phi(\rho_t, \mathbf{w}_t) \, d\gamma \, dt \right)^{1/p}. \end{aligned}$$

Since $\int_0^T \Phi(\mu_t, \mathbf{v}_t | \gamma) \, dt < +\infty$, by (23) and the equivalence of $\|\cdot\|_*$ with the euclidean norm, the last inequality shows that there exists $C > 0$ such that

$$\left| \int_{\mathbb{R}^d} \zeta_R \, d\mu_{t_1} - \int_{\mathbb{R}^d} \zeta_R \, d\mu_{t_2} \right| \leq C \left(\frac{1}{R^q} \gamma(B_{2R} \setminus B_R) \right)^{1/q}. \tag{25}$$

Since $\tilde{m}_r(\gamma) < +\infty$ shows that $\lim_{R \rightarrow +\infty} R^r \gamma(B_{2R} \setminus B_R) = 0$ we have that $\lim_{R \rightarrow +\infty} \frac{1}{R^q} \gamma(B_{2R} \setminus B_R) = 0$ if $r \geq -q$. Then the conservation of the mean follows from (25).

Example 2. When $\phi \in \mathcal{A}_p(a, b)$ with $a < 0$ and $b > 0$, if $\gamma = \mathcal{L}^d$ and $d > q$ in general solutions of the continuity equations with finite energy could not conserve the mass.

Let $\varepsilon > 0$ such that $a + \varepsilon < 0$ and $b - \varepsilon > 0$ and consider an initial measure with compact support and mass different from 0, $\mu_0 = \rho_0 \mathcal{L}^d$, such that $a + \varepsilon \leq \rho_0 \leq b - \varepsilon$. We define the curve, for $t \geq 0$,

$$\mu_t := \rho_t \mathcal{L}^d, \quad \rho_t(x) := e^{-dt} \rho_0(e^{-t}x), \quad \mathbf{v}_t := \mathbf{w}_t \mathcal{L}^d = x \rho_t(x) \mathcal{L}^d. \quad (26)$$

It is easy to check that $(\mu, \mathbf{v}) \in \mathcal{CE}(0, +\infty)$, $\mu_t(\mathbb{R}^d) = \mu_0(\mathbb{R}^d)$ and $a + \varepsilon \leq \rho_t \leq b - \varepsilon$. By 3 of Proposition 1 we have that $\phi(\rho_t, \mathbf{w}_t) \leq C|\mathbf{w}_t|^p$. By a simple computation we obtain that

$$\begin{aligned} \int_{\mathbb{R}^d} |\mathbf{w}_t(x)|^p dx &= \int_{\mathbb{R}^d} |x|^p e^{-tdp} |\rho_0(e^{-t}x)|^p dx \\ &= \int_{\mathbb{R}^d} |y|^p e^{t((1-d)p+d)} |\rho_0(y)|^p dy \end{aligned}$$

and then

$$\int_0^{+\infty} \phi(\rho_t, \mathbf{w}_t) dx dt < +\infty$$

when $d > q$.

The curve (μ_t, \mathbf{v}_t) can be reparametrized between $[0, 1]$ setting $s = \frac{2}{\pi} \arctan t$, $t \in (0, +\infty)$ and $\eta_s = \rho_{\tan(\frac{\pi}{2}s)} = \rho_t$. It is not difficult to check that the energy is still finite and η_s connect μ_0 with the null measure.

2.3.2. Compactness for solutions with finite energy In this section we prove a compactness result for signed solutions of the continuity equation. This result is a useful tool in order to obtain existence of geodesics of the distance defined in the next Section and its lower semi-continuity with respect to weak* convergence.

Proposition 4. (Compactness) *Let $\phi \in \mathcal{A}_p(a, b)$ and $\gamma^n, \gamma \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$ be a sequence such that $\gamma^n \rightharpoonup^* \gamma$.*

If (μ^n, \mathbf{v}^n) is a sequence in $\mathcal{CE}_{\phi, \gamma^n}(0, T)$ satisfying

$$\sup_{n \in \mathbb{N}} \int_0^T \Phi(\mu_t^n, \mathbf{v}_t^n | \gamma^n) dt < +\infty, \quad (27)$$

then there exists a subsequence (still indexed by n) and a couple $(\mu, \mathbf{v}) \in \mathcal{CE}_{\phi, \gamma}(0, T)$ such that $\mu_t^n \rightharpoonup^ \mu_t$ in $\mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ for all $t \in [0, T]$, $\mathbf{v}^n \rightharpoonup^* \mathbf{v}$ in $\mathcal{M}_{\text{loc}}(\mathbb{R}^d \times (0, T); \mathbb{R}^d)$, and*

$$\int_0^T \Phi(\mu_t, \mathbf{v}_t | \gamma) dt \leq \liminf_{n \rightarrow +\infty} \int_0^T \Phi(\mu_t^n, \mathbf{v}_t^n | \gamma^n) dt. \quad (28)$$

If along the subsequence $\tilde{m}_{-q}(\gamma^n) < +\infty$ for all n and $\tilde{m}_{-q}(\gamma) < +\infty$, and $\mu_0^n(\mathbb{R}^d) \rightarrow \mu_0(\mathbb{R}^d) \in \mathbb{R}$, then $\mu_t^n(\mathbb{R}^d) \rightarrow \mu_t(\mathbb{R}^d)$ for every $t \in [0, T]$.

Proof. By (27), the definition of the action functional and (17) we have $\mu^n = \rho^n \gamma^n$, $\mathbf{v}^n = \mathbf{w}^n \gamma^n$ and $|\rho^n| \leq c := \max\{|a|, |b|\}$. Then there exists a subsequence (still indexed by n) and ρ such that $\rho^n \rightharpoonup \rho$ weakly in $L^1_{\text{loc}}(\mathbb{R}^d \times [0, T])$. On the other hand, by (24), for every bounded Borel set $B \subset \mathbb{R}^d$ and for every $t_1, t_2 \in [0, T], t_1 < t_2$ we have

$$\begin{aligned} \int_{t_1}^{t_2} \int_B \|\mathbf{w}^n\| \, d\gamma^n \, dt &\leq \int_{t_1}^{t_2} \int_B h(\rho^n)^{1/q} \phi(\rho^n, \mathbf{w}^n)^{1/p} \, d\gamma^n \, dt \\ &\leq \left(\int_{t_1}^{t_2} \int_B h(\rho^n) \, d\gamma^n \, dt \right)^{1/q} \left(\int_{t_1}^{t_2} \int_B \phi(\rho^n, \mathbf{w}^n) \, d\gamma^n \, dt \right)^{1/p}. \end{aligned}$$

By (23), (27) and the equivalence of $\|\cdot\|$ with the euclidean norm, the last inequality shows that there exist $C > 0$ such that

$$\int_{t_1}^{t_2} \int_B \|\mathbf{w}^n\| \, d\gamma^n \, dt \leq C((t_2 - t_1)\gamma^n(B))^{1/q},$$

By this estimate there exist $\mathbf{v} \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d \times [0, T], \mathbb{R}^d)$ and a subsequence such that $\mathbf{v}^n \rightharpoonup^* \mathbf{v}$. By the lower semicontinuity property of Theorem 1 we obtain (28). Reasoning as in the proof of Lemma 4.5 of [13] we obtain that (μ, \mathbf{v}) satisfies the continuity equation.

Finally, by Proposition 3 $\mu_t^n(\mathbb{R}^d)$ and $\mu_t(\mathbb{R}^d)$ do not depend on $t \in [0, T]$.

3. The modified Wasserstein distance

In this section we give the rigorous definition of the modified Wasserstein distance illustrated in the introduction. As already pointed out, we deal with the case of the distance induced by an action density function $\phi \in \mathcal{A}_p(a, b)$ for $a, b \in \mathbb{R}$, whereas we refer to [13] for the case $\phi \in \mathcal{A}_p(0, +\infty)$.

The proofs are almost all omitted because follows exactly as in [13, Sect. 5] from the results of the previous sections.

Definition 8. Given a reference measure $\gamma \in \mathcal{M}^+_{\text{loc}}(\mathbb{R}^d)$, an admissible action density function $\phi \in \mathcal{A}_p(a, b)$ and the corresponding action functional Φ of Definition 3, for $\mu^0, \mu^1 \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ we define

$$\mathcal{W}_{\phi, \gamma}(\mu^0, \mu^1) := \inf \left\{ \left(\int_0^1 \Phi(\mu_s, \mathbf{v}_s | \gamma) \, ds \right)^{1/p} : (\mu, \mathbf{v}) \in \mathcal{CE}_{\phi, \gamma}(0, 1; \mu^0 \rightarrow \mu^1) \right\}, \quad (29)$$

and $\mathcal{W}_{\phi,\gamma}(\mu^0, \mu^1) = +\infty$ if the set of connecting curves $\mathcal{CE}_{\phi,\gamma}(0, 1; \mu^0 \rightarrow \mu^1)$ is empty.

By using the compactness Proposition 4 and reasoning as in [13, Theorem 5.4], it is simple to obtain the existence of a minimizing curve in (29), when the set of connecting curves is not empty.

Proposition 5. (Existence and characterization of minimizers) *Given $\gamma \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$ and $\phi \in \mathcal{A}_p(a, b)$, for every $\mu_0, \mu_1 \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ such that $\mathcal{W}_{\phi,\gamma}(\mu_0, \mu_1) < +\infty$ there exists a minimizing couple (μ, \mathbf{v}) in (29) and the curve $(\mu_s)_{s \in [0,1]}$ satisfies*

$$\mathcal{W}_{\phi,\gamma}(\mu_t, \mu_s) = |t - s| \mathcal{W}_{\phi,\gamma}(\mu_0, \mu_1) \quad \forall s, t \in [0, 1]. \tag{30}$$

We have the characterization

$$\mathcal{W}_{\phi,\gamma}(\mu^0, \mu^1) = \inf \left\{ \int_0^1 (\Phi(\mu_s, \mathbf{v}_s | \gamma))^{1/p} ds : \right. \\ \left. (\mu, \mathbf{v}) \in \mathcal{CE}(0, 1; \mu^0 \rightarrow \mu^1) \right\}. \tag{31}$$

The property (30) means that the curve $t \mapsto \mu_t$ is a constant speed geodesic for $\mathcal{W}_{\phi,\gamma}$.

Using the properties of the action functional Φ and of the solutions of the continuity equation stated in Sect. 2 we are able to prove, following the arguments of [13] the fundamental Theorem stating the basic metric and topological properties of $\mathcal{W}_{\phi,\gamma}$.

Theorem 2. (Basic properties of $\mathcal{W}_{\phi,\gamma}$) *Let $\gamma \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$, $\phi \in \mathcal{A}_p(a, b)$. Then the application on measures $\mathcal{W}_{\phi,\gamma} : \mathcal{M}_{\text{loc}}(\mathbb{R}^d) \times \mathcal{M}_{\text{loc}}(\mathbb{R}^d) \rightarrow [0, +\infty]$ defined in (29) satisfies the following properties:*

- (a) (Metric character) $\mathcal{W}_{\phi,\gamma}$ is a pseudo-distance on $\mathcal{M}_{\text{loc}}(\mathbb{R}^d)$; i.e. $\mathcal{W}_{\phi,\gamma}$ satisfies the axiom of the distance but can assume the value $+\infty$.
- (b) (Topological properties) The topology induced by $\mathcal{W}_{\phi,\gamma}$ on $\mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ is stronger than or equivalent to the weak* one. Bounded sets with respect to $\mathcal{W}_{\phi,\gamma}$ are weakly* relatively compact.
- (c) (Lower semicontinuity) If $\gamma^n \rightharpoonup^* \gamma$ in $\mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$, $\mu_0^n \rightharpoonup^* \mu_0$, $\mu_1^n \rightharpoonup^* \mu_1$ in $\mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ and $\phi^n \in \mathcal{A}_p(a, b)$, is a sequence such that $\phi^n \leq \phi^{n+1}$ and ϕ^n pointwise converges to ϕ , then

$$\liminf_{n \rightarrow +\infty} \mathcal{W}_{\phi^n, \gamma^n}(\mu_0^n, \mu_1^n) \geq \mathcal{W}_{\phi,\gamma}(\mu_0, \mu_1). \tag{32}$$

- (d) (Completeness) The space $\mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ endowed with the pseudo-distance $\mathcal{W}_{\phi,\gamma}$ is complete.

- (e) (Connected components and mass conservation) Given a measure $\sigma \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$, the space $\mathcal{M}[\sigma] := \{\mu \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d) : \mathcal{W}_{\phi, \gamma}(\mu, \sigma) < +\infty\}$ is a complete geodesic metric space (every two measures in $\mathcal{M}[\sigma]$ can be connected by a constant speed geodesic). If $\tilde{m}_{-q}(\gamma) < +\infty$ then $\mu(\mathbb{R}^d) = \sigma(\mathbb{R}^d)$ for every $\mu \in \mathcal{M}[\sigma]$.
- (f) (Convexity) The map $(\mu^0, \mu^1) \mapsto \mathcal{W}_{\phi, \gamma}^p(\mu^0, \mu^1)$ is convex with respect to the linear structure of $\mathcal{M}_{\text{loc}}(\mathbb{R}^d)$, i.e. for every $\mu_i^j \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$, $i, j = 0, 1$, and $\tau \in [0, 1]$, if $\mu_i^\tau = (1 - \tau)\mu_i^0 + \tau\mu_i^1$, then

$$\mathcal{W}_{\phi, \gamma}^p(\mu_0^\tau, \mu_1^\tau) \leq (1 - \tau)\mathcal{W}_{\phi, \gamma}^p(\mu_0^0, \mu_1^0) + \tau\mathcal{W}_{\phi, \gamma}^p(\mu_0^1, \mu_1^1). \quad (33)$$

If ϕ is strictly convex, then for $\mu_0, \mu_1 \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ with $\mathcal{W}_{\phi, \gamma}(\mu_0, \mu_1) < +\infty$ the minimizer $(\mu, \mathbf{v}) \in \mathcal{CE}_{\phi, \gamma}(0, 1; \mu_0 \rightarrow \mu_1)$ of (29) is unique.

- (g) (Monotonicity) If $\phi_1 \leq \phi_2$ then

$$\mathcal{W}_{\phi_1, \gamma}(\mu_0, \mu_1) \leq \mathcal{W}_{\phi_2, \gamma}(\mu_0, \mu_1),$$

for every $\mu_0, \mu_1 \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$.

Moreover, if $(0, 0) \in \text{dom}(\phi_i)$, $i = 1, 2$ and $\gamma_1 \leq \gamma_2$ then

$$\mathcal{W}_{\phi_1, \gamma_2}(\mu_0, \mu_1) \leq \mathcal{W}_{\phi_2, \gamma_1}(\mu_0, \mu_1),$$

for every $\mu_0, \mu_1 \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$.

- (h) (Approximation by convolution) Let $k \in C_c^\infty(\mathbb{R}^d)$ be a nonnegative convolution kernel, with $\int_{\mathbb{R}^d} k(x) \, dx = 1$ and $\text{supp}(k) = \bar{B}_1(0)$, and let $k_\varepsilon(x) := \varepsilon^{-d}k(x/\varepsilon)$. For every $\mu_0, \mu_1 \in \mathcal{M}(\mathbb{R}^d)$

$$\begin{aligned} \mathcal{W}_{\phi, \gamma * k_\varepsilon}(\mu_0 * k_\varepsilon, \mu_1 * k_\varepsilon) &\leq \mathcal{W}_{\phi, \gamma}(\mu_0, \mu_1) \quad \forall \varepsilon > 0; \\ \lim_{\varepsilon \rightarrow 0} \mathcal{W}_{\phi, \gamma * k_\varepsilon}(\mu_0 * k_\varepsilon, \mu_1 * k_\varepsilon) &= \mathcal{W}_{\phi, \gamma}(\mu_0, \mu_1). \end{aligned}$$

- (i) (Necessary condition for the finiteness of the distance) If $\mathcal{W}_{\phi, \gamma}(\mu_0, \mu_1) < +\infty$ then $\mu_j = u_j \gamma \ll \gamma$, $j = 0, 1$, and $a \leq u_j(x) \leq b$ for γ -a.e. $x \in \mathbb{R}^d$.

Proof. (a) The verification of the axioms of the distance is straightforward except for the triangular inequality where we use the gluing of solutions of Lemma 2 and the characterization (31).

- (b) In order to prove the topological property, reasoning as in the proof of Proposition 3 we obtain that

$$\left| \int_{\mathbb{R}^d} \zeta \, d\mu_1 - \int_{\mathbb{R}^d} \zeta \, d\mu_0 \right| \leq \sup |\nabla \zeta| (\text{h}_{\max} \gamma(\text{supp}(\zeta)))^{1/q} \mathcal{W}_{\phi, \gamma}(\mu_0, \mu_1) \quad (34)$$

for every $\zeta \in C_c^1(\mathbb{R}^d)$. Since $C_c^1(\mathbb{R}^d)$ is dense in $C_c(\mathbb{R}^d)$ we obtain the assertion on the topology induced by the distance and on the relative compactness.

- (c) This property is proved exactly as Theorem 5.6 of [13] by using the compactness Proposition 4.
- (d) The result can be proved as in Theorem 5.7 of [13] ad using the above semi-continuity property.

- (e) Same as above. The final assertion about the equality of the signed mass follows from Proposition 3.
- (f) Can be proved exactly as Theorem 5.11 of [13].
- (g) Follows from the definition of Φ and 2. of Theorem 1.
- (h) Follows from items 1. and 3. of Theorem 1.
- (i) Follows from (17), the weak* continuity of the connecting curves in $\mathcal{CE}_{\phi,\gamma}(0, 1; \mu_0 \rightarrow \mu_1)$ and the lower semi continuity of Φ with respect to the weak* convergence.

The following proposition deals with a control of the moments and a comparison between the convergence with respect to $\mathcal{W}_{\phi,\gamma}$ and the standard Wasserstein distance defined in (1).

Proposition 6. (Moments estimate) *Let $\gamma \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$ be satisfying $\tilde{m}_r(\gamma) < +\infty$ for some $r \in \mathbb{R}$ and $\phi \in \mathcal{A}_p(a, b)$.*

If $\mu_0, \mu_1 \in \mathcal{M}_{\text{loc}}(\mathbb{R}^d)$ satisfy $\mathcal{W}_{\phi,\gamma}(\mu_0, \mu_1) < +\infty$, then, setting $C := \max\{|a|, |b|\}$, we have

$$\tilde{m}_\delta(|\mu_i|) \leq C \tilde{m}_r(\gamma), \quad \text{for } i = 0, 1, \quad \forall \delta \leq r. \tag{35}$$

If $r \geq 1$ and $a \geq 0$ then the convergence with respect to $\mathcal{W}_{\phi,\gamma}$ in $\mathcal{M}[\sigma]$, for some non-negative measure σ satisfying $\sigma(\mathbb{R}^d) < +\infty$, implies the convergence with respect to the r -Wasserstein distance W_r .

Proof. Denoting by $1 \vee |x| = \max\{1, |x|\}$, given $C = \max\{|a|, |b|\}$, $\delta \leq r$ and a Borel set $A \subset \mathbb{R}^d$, by (17), we obtain

$$\begin{aligned} \int_A (1 \vee |x|)^\delta d|\mu_i|(x) &= \int_A (1 \vee |x|)^\delta |\rho_i(x)| d\gamma(x) \\ &\leq C \int_A (1 \vee |x|)^\delta d\gamma(x) \leq C \int_A (1 \vee |x|)^r d\gamma(x). \end{aligned} \tag{36}$$

Choosing $A = \mathbb{R}^d$ in (36) we obtain (35).

If μ_n is a sequence in $\mathcal{M}[\sigma]$ converging to μ with respect to $\mathcal{W}_{\phi,\gamma}$, then, by Proposition 2, μ_n weakly* converges to μ and $\mu_n(\mathbb{R}^d) = \mu(\mathbb{R}^d) = \mu(\sigma)$ because of the assumption on the moment of γ and $r \geq 1$. By (36) with $\delta = 0$ we have that the sequence μ_n is tight and then μ_n narrowly converges to μ . Since (36) implies that the r -moments of μ_n are uniformly equiintegrable we obtain that (see Lemma 5.1.7 of [3]) $\tilde{m}_r(\mu_n)$ converges to $\tilde{m}_r(\mu)$ and we conclude.

In particular the previous Proposition applies to the case $\gamma(\mathbb{R}^d) < +\infty$. In the next proposition we state a simple comparison with the standard Wasserstein distance (1).

Proposition 7. (Comparison with Wasserstein distance) *Let $p > 1$, $\phi \in \mathcal{A}_p(0, M)$, $\Omega \subset \mathbb{R}^d$ an open convex set and $\gamma_\Omega = \chi_\Omega \mathcal{L}^d$. If $\mu_i, i = 0, 1$, are two absolutely continuous measures with respect to γ_Ω , $\mu_i = \rho_i \gamma_\Omega$, such that $0 \leq \rho_i(x) \leq M' <$*

$M, \tilde{m}_p(\mu_i) < +\infty$ for $i = 0, 1$ and $\mu_0(\mathbb{R}^d) = \mu_1(\mathbb{R}^d)$, then there exists a constant C , depending only on M', ϕ and p , such that

$$\mathcal{W}_{\phi, \gamma_\Omega}(\mu_0, \mu_1) \leq C W_p(\mu_0, \mu_1) < +\infty, \tag{37}$$

where W_p denotes the standard p -Wasserstein distance.

Proof. Let h be given by Proposition 2. Since h is concave and positive on $(0, M)$, we have that

$$h(\rho) \geq \frac{h(M')}{M'} \rho, \quad \forall \rho \in (0, M'),$$

and, consequently,

$$\phi(\rho, \mathbf{w}) \leq \frac{|\mathbf{w}|^p}{h(\rho)^{p-1}} \leq \left(\frac{M'}{h(M')} \right)^{p-1} \frac{|\mathbf{w}|^p}{\rho^{p-1}} \quad \forall \rho \in (0, M'). \tag{38}$$

Since the p -moments of μ_0 and μ_1 are finite, taking the geodesic interpolant μ_t between μ_0 and μ_1 for the standard p -Wasserstein distance $W_p(\mu_0, \mu_1)$, and denoting by ρ_t the density of μ_t , we have that $\rho_t \leq M'$ (see the proof of [13, Theorem 5.24]). Since Ω is convex, the support of μ_t belongs to $\overline{\Omega}$ and, denoting by $\psi(\rho, \mathbf{w}) := \frac{|\mathbf{w}|^p}{\rho^{p-1}}$, we have that $\mathcal{W}_{\psi, \gamma_\Omega} = W_p$ for all the measures with support in $\overline{\Omega}$. Then, by (38), and recalling (g) in Theorem 2 we obtain (37).

4. Measures at finite distance and convergence

In this section we give sufficient conditions for the finiteness of the distance between two measures. We study also the relation between the convergence with respect to the distance and the weak-* one. The first result concerns measures defined on the whole space \mathbb{R}^d with the reference measure $\gamma = \mathcal{L}^d$, whereas the second one deals with measures defined on a bounded convex domain Ω with the reference measure $\gamma_\Omega = \mathcal{L}^d|_\Omega$.

4.1. The case of reference measure \mathcal{L}^d

Theorem 3. (Connectivity in \mathbb{R}^d) *Let $p > 1$ and $\phi \in \mathcal{A}_p(0, M)$. If $\mu_i, i = 0, 1$, are two absolutely continuous measures $\mu_i = \rho_i \mathcal{L}^d$, such that $0 \leq \rho_i(x) \leq M, \tilde{m}_p(\mu_i) < +\infty$ for $i = 0, 1$ and $\mu_0(\mathbb{R}^d) = \mu_1(\mathbb{R}^d)$, then there exists a constant $C > 0$ depending only on ϕ, d and p such that*

$$\mathcal{W}_{\phi, \mathcal{L}^d}(\mu_0, \mu_1) \leq C(\tilde{m}_p(\mu_0) + \tilde{m}_p(\mu_1)) < +\infty. \tag{39}$$

We observe that the inequality (39) holds in the case of the standard Wasserstein distance (it is a very easy consequence of the definition (1)).

Proof. Let h be given by Proposition 2. Since h is concave and non-negative, there exists $\tilde{h} : [0, M] \rightarrow [0, +\infty)$ of the form $\tilde{h}(\rho) = A\rho(M/B - B\rho)$ for $A, B > 0$ such that $\tilde{h}(\rho) \leq h(\rho)$ in $[0, M]$. Hence $\mathcal{W}_{\phi, \mathcal{L}^d}(\mu_0, \mu_1) \leq \mathcal{W}_{\phi_h, \mathcal{L}^d}(\mu_0, \mu_1) \leq \mathcal{W}_{\phi_{\tilde{h}}, \mathcal{L}^d}(\mu_0, \mu_1)$. Thanks to this observation, it is sufficient to prove the result under the assumption that $M = 1, h(\rho) = \rho(1 - \rho)$ and $0 \leq \rho_i(x) \leq 1$, for $i = 0, 1$.

Recalling the definition of push-forward of measures (see Appendix 4.2) and defining

$$\tilde{\mu}_i = \tilde{\rho}_i \mathcal{L}^d = 2\text{Id}_{\#}\mu_i, \tag{40}$$

where Id denotes the identity map in \mathbb{R}^d , we prove that there exists a constant $C_{p,d}$, depending only on p and d , such that

$$W_{\phi_h, \mathcal{L}^d}(\mu_i, \tilde{\mu}_i) < C_{p,d} \tilde{m}_p(\mu_i) \quad \text{for } i = 0, 1. \tag{41}$$

Indeed, for $t \in [0, 1]$, taking $T_t(x) := (1 + t^p)x$ and $\mu_t := (T_t)_{\#}\mu_i = \rho_t \mathcal{L}^d$, by (60) we have that $\rho_t(y) = \frac{1}{(1+t^p)^d} \rho_i\left(\frac{y}{1+t^p}\right)$. Defining $\mathbf{v}_t(x) := \dot{T}_t \circ T_t^{-1}(x) = \frac{(pt^{p-1})}{1+t^p} x$, and $\mathbf{w}_t = \mathbf{v}_t \rho_t, \mathbf{v}_t = \mathbf{w}_t \mathcal{L}^d$ it is easy to check that $(\mu_t, \mathbf{v}_t)_{t \in (0,1)} \in \mathcal{CE}(0, 1; \mu_i \rightarrow \tilde{\mu}_i)$. By elementary computations, using the definition of μ_t and \mathbf{v}_t , we have

$$\begin{aligned} & \int_0^1 \int_{\mathbb{R}^d} \frac{|\mathbf{w}_t(x)|^p}{(\rho_t(x)(1 - \rho_t(x)))^{p-1}} dx dt \\ &= \int_0^1 \int_{\mathbb{R}^d} \frac{|\mathbf{v}_t(x)|^p \rho_t(x)}{(1 - \rho_t(x))^{p-1}} dx dt \\ &= \int_0^1 \int_{\mathbb{R}^d} \frac{|\mathbf{v}_t(x)|^p}{(1 - \rho_t(x))^{p-1}} d\mu_t(x) dt \\ &= \int_0^1 \int_{\mathbb{R}^d} \frac{|\mathbf{v}_t(T_t(x))|^p}{(1 - \rho_t(T_t(x)))^{p-1}} d\mu_i(x) dt \\ &= \int_0^1 \int_{\mathbb{R}^d} \frac{(pt^{p-1})^p |x|^p}{(1 - (1 + t^p)^{-d} \rho_i(x))^{p-1}} d\mu_i(x) dt. \end{aligned}$$

Since $\rho_i(x) \leq 1$ and $(1 + t^p)^d \geq 1 + dt^p$ we have

$$\frac{1}{1 - (1 + t^p)^{-d} \rho_i(x)} \leq \frac{1}{1 - (1 + t^p)^{-d}} = \frac{(1 + t^p)^d}{(1 + t^p)^d - 1} \leq \frac{(1 + t^p)^d}{dt^p}.$$

Then

$$\begin{aligned} & \int_0^1 \int_{\mathbb{R}^d} \frac{(pt^{p-1})^p |x|^p}{(1 - (1 + t^p)^{-d} \rho_i(x))^{p-1}} \, d\mu_i(x) \, dt \\ & \leq \int_0^1 \int_{\mathbb{R}^d} \frac{(pt^{p-1})^p (1 + t^p)^{d(p-1)}}{(dt^p)^{p-1}} |x|^p \, d\mu_i(x) \, dt \\ & \leq \tilde{m}_p(\mu_i) \int_0^1 p^p d^{1-p} (1 + t^p)^{d(p-1)} \, dt, \end{aligned}$$

and (41) follows with $C_{p,d} = \int_0^1 p^p d^{1-p} (1 + t^p)^{d(p-1)} \, dt$.

Finally, by the triangular inequality, we have

$$\mathcal{W}_{\phi_h, \mathcal{L}^d}(\mu_0, \mu_1) \leq \mathcal{W}_{\phi_h, \mathcal{L}^d}(\mu_0, \tilde{\mu}_0) + \mathcal{W}_{\phi_h, \mathcal{L}^d}(\tilde{\mu}_0, \tilde{\mu}_1) + \mathcal{W}_{\phi_h, \mathcal{L}^d}(\tilde{\mu}_1, \mu_1). \tag{42}$$

Since by (60) we have $\tilde{\rho}_i(x) = 2^{-d} \rho_i(x/2) \leq 2^{-d}$ and $\tilde{m}_p(\tilde{\mu}_i) = 2^p \tilde{m}_p(\mu_i) < +\infty$, by Proposition 7 applied to $\tilde{\mu}_0, \tilde{\mu}_1$, and observing that $W_p(\tilde{\mu}_0, \tilde{\mu}_1) \leq \tilde{m}_p(\tilde{\mu}_0) + \tilde{m}_p(\tilde{\mu}_1)$ (it is a simple consequence of the definition (1)), by (41) and (42) we obtain (39).

Given $M > 0$ and $c > 0$ we define the set of measures

$$\begin{aligned} \mathcal{M}_{M,c}^+(\mathbb{R}^d) & := \left\{ \mu \in \mathcal{M}^+(\mathbb{R}^d) : \right. \\ & \left. \mu = \rho \mathcal{L}^d, 0 \leq \rho \leq M, \mu(\mathbb{R}^d) = c, \tilde{m}_p(\mu) < +\infty \right\}. \end{aligned}$$

Theorem 4. *Let $p > 1$ and $\phi \in \mathcal{A}_p(0, M)$. If $(\mu_n)_{n \in \mathbb{N}}$ is a sequence in $\mathcal{M}_{M,c}^+(\mathbb{R}^d)$ weakly- $*$ convergent to $\mu \in \mathcal{M}_{M,c}^+(\mathbb{R}^d)$, such that*

$$\tilde{m}_p(\mu_n) \rightarrow \tilde{m}_p(\mu), \tag{43}$$

then

$$\lim_{n \rightarrow +\infty} \mathcal{W}_{\phi, \mathcal{L}^d}(\mu_n, \mu) = 0.$$

Proof. Let $\bar{\mu} = \bar{\rho} \mathcal{L}^d \in \mathcal{M}_{M,c}^+$ be a fixed auxiliary measure such that $M' := \sup \bar{\rho} < M$.

For every $\lambda \in (0, 1)$, we define the convex combinations $\mu_n^\lambda := (1 - \lambda)\mu_n + \lambda \bar{\mu}$ and $\mu^\lambda := (1 - \lambda)\mu + \lambda \bar{\mu}$. Denoting by ρ_n^λ the density of μ_n^λ with respect to \mathcal{L}^d we have $\rho_n^\lambda \leq 1 - \lambda(M - M')$. By Proposition 7 and the convexity of the p -power of the standard p -Wasserstein distance (Theorem 2 (f)) in the case $\phi(\rho, \mathbf{w}) = |\mathbf{w}|^p / \rho^{p-1}$ or [26]) we have

$$\mathcal{W}_{\phi, \mathcal{L}^d}^p(\mu_n^\lambda, \mu^\lambda) \leq C W_p^p(\mu_n^\lambda, \mu^\lambda) \leq C(1 - \lambda) W_p^p(\mu_n, \mu). \tag{44}$$

By the convergence of the p -moments (43) and the weak- $*$ convergence we have (see [3] or [26])

$$\lim_{n \rightarrow +\infty} W_p(\mu_n, \mu) = 0. \tag{45}$$

Moreover for the convexity of $\mathcal{W}_{\phi, \mathcal{L}^d}^p$ (Theorem 2 (f)) we have

$$\mathcal{W}_{\phi, \mathcal{L}^d}^p(\mu_n, \mu_n^\lambda) \leq \lambda \mathcal{W}_{\phi, \mathcal{L}^d}^p(\mu_n, \bar{\mu}), \quad \mathcal{W}_{\phi, \mathcal{L}^d}^p(\mu, \mu^\lambda) \leq \lambda \mathcal{W}_{\phi, \mathcal{L}^d}^p(\mu, \bar{\mu}). \tag{46}$$

Since

$$\mathcal{W}_{\phi, \mathcal{L}^d}(\mu_n, \mu) \leq \mathcal{W}_{\phi, \mathcal{L}^d}(\mu_n, \mu_n^\lambda) + \mathcal{W}_{\phi, \mathcal{L}^d}(\mu_n^\lambda, \mu^\lambda) + \mathcal{W}_{\phi, \mathcal{L}^d}(\mu^\lambda, \mu), \tag{47}$$

by (44), (45) and (46) we have

$$\limsup_{n \rightarrow +\infty} \mathcal{W}_{\phi, \mathcal{L}^d}(\mu_n, \mu) \leq \lambda^{1/p} \left(\sup_n \mathcal{W}_{\phi, \mathcal{L}^d}(\mu_n, \bar{\mu}) + \mathcal{W}_{\phi, \mathcal{L}^d}(\mu, \bar{\mu}) \right). \tag{48}$$

By (43) and Theorem 3 we obtain

$$\sup_n \mathcal{W}_{\phi, \mathcal{L}^d}(\mu_n, \bar{\mu}) < +\infty. \tag{49}$$

Since $\lambda > 0$ is arbitrary, (48) and (49) imply

$$\limsup_{n \rightarrow +\infty} \mathcal{W}_{\phi, \mathcal{L}^d}(\mu_n, \mu) = 0$$

and we conclude.

We recall that the convergence with respect to the standard Wasserstein distance W_p is equivalent to the weak- $*$ convergence and the convergence of the p -moments \tilde{m}_p (see [26] or [3]). Consequently, Theorem 4 states that the convergence with respect to W_p in $\mathcal{M}_{M,c}^+(\mathbb{R}^d)$ implies the convergence with respect to $\mathcal{W}_{\phi, \mathcal{L}^d}$ for every $\phi \in \mathcal{A}_p(0, M)$. We observe that this property is not true in the case of $\phi \in \mathcal{A}_p(0, +\infty)$, where only a result like Proposition 7 hold (see Theorem 5.24 of [13]).

4.2. The case of the reference measure $\chi_\Omega \mathcal{L}^d$ with Ω bounded convex

When the reference measure is $\gamma_\Omega := \chi_\Omega \mathcal{L}^d$, where Ω is a bounded convex smooth domain, we have the following result of finiteness of the distance and of boundedness of the space of admissible measures.

Theorem 5. *Let $\phi \in \mathcal{A}_2(a, b)$ and $\gamma_\Omega := \chi_\Omega \mathcal{L}^d$ with $\Omega \subset \mathbb{R}^d$ a bounded convex smooth domain. For every $c \in (a \mathcal{L}^d(\Omega), b \mathcal{L}^d(\Omega))$ we define the set of measures*

$$\mathcal{M}_{(a,b),c}(\Omega) := \{\mu \in \mathcal{M}(\bar{\Omega}) : \mu = \rho \gamma_\Omega, a \leq \rho \leq b, \mu(\bar{\Omega}) = c\}.$$

The space $\mathcal{M}_{(a,b),c}(\Omega)$ endowed with the distance $\mathcal{W}_{\phi, \gamma_\Omega}$ is bounded. In particular $\mathcal{W}_{\phi, \gamma_\Omega}(\mu_0, \mu_1) < +\infty$ for every $\mu_0, \mu_1 \in \mathcal{M}_{(a,b),c}(\Omega)$.

Proof. Defining $\mu_\infty := \frac{c}{\mathcal{L}^d(\Omega)}\gamma_\Omega$, we prove that

$$\sup_{\mu_0 \in \mathcal{M}_{(a,b),c}(\Omega)} \mathcal{W}_{\phi,\gamma_\Omega}(\mu_0, \mu_\infty) < +\infty. \tag{50}$$

Let h be given by Proposition 2.

For $\mu_0 = \rho_0\gamma_\Omega \in \mathcal{M}_{(a,b),c}(\Omega)$, let $\rho : (0, +\infty) \times \Omega \rightarrow \mathbb{R}$ be the solution of Cauchy-Neumann problem for the heat equation

$$\begin{cases} \partial_t \rho - \Delta \rho = 0 & \text{in } (0, +\infty) \times \Omega \\ \rho(0, \cdot) = \rho_0 & \text{in } \Omega \\ \nabla \rho \cdot \mathbf{n} = 0 & \text{on } (0, \infty) \times \partial\Omega. \end{cases} \tag{51}$$

We use the notation $\rho_t := \rho(t, \cdot)$ and $S_t(\mu_0) := \rho_t\gamma_\Omega$.

Defining the convex function $U : (a, b) \rightarrow \mathbb{R}$ by

$$U''(r) = \frac{1}{h(r)}, \quad U'((a+b)/2) = 0, \quad U((a+b)/2) = 0 \tag{52}$$

and the entropy functional

$$\mathcal{U}(\rho) = \int_{\Omega} U(\rho(x)) \, dx,$$

we have the following entropy dissipation inequality

$$\mathcal{U}(\rho_T) - \mathcal{U}(\rho_0) \leq - \int_0^T \int_{\Omega} \frac{|\nabla \rho_s|^2}{h(\rho_s)} \, dx \, ds. \tag{53}$$

The inequality (53) can be obtained, in the case of smooth initial datum, with a simple computation and, in the general case, by a convolution approximation argument.

By Lemma 3 in the Appendix, observing that in our case $\rho_\infty = \frac{c}{\mathcal{L}^d(\Omega)}$, we can prove that there exists $T > 0$, independent on μ_0 , such that

$$\rho_t \leq \rho_\infty + \frac{b - \rho_\infty}{2}, \quad \forall t \geq T. \tag{54}$$

By the triangular inequality we have that

$$\mathcal{W}_{\phi,\gamma_\Omega}(\mu_0, \mu_\infty) \leq \mathcal{W}_{\phi,\gamma_\Omega}(\mu_0, S_T(\mu_0)) + \mathcal{W}_{\phi,\gamma_\Omega}(S_T(\mu_0), \mu_\infty). \tag{55}$$

Since h is concave and Ω is bounded, it is not difficult to see that \mathcal{U} is bounded in $\mathcal{M}_{(a,b),c}(\Omega)$, and recalling (15) we have $\mathcal{W}_{\phi,\gamma_\Omega}(\mu_0, S_T(\mu_0)) \leq \int_0^T \int_{\Omega} \frac{|\nabla \rho_s|^2}{h(\rho_s)} \, dx \, ds$, consequently (53) implies that

$$\sup_{\mu_0 \in \mathcal{M}_{(a,b),c}(\Omega)} \mathcal{W}_{\phi,\gamma_\Omega}(\mu_0, S_T(\mu_0)) < +\infty. \tag{56}$$

Since

$$\mathcal{W}_{\phi, \gamma\Omega}(\mu, \nu) = \mathcal{W}_{\tilde{\phi}, \gamma\Omega}(\mu - a\gamma\Omega, \nu - a\gamma\Omega), \text{ where } \tilde{\phi}(r, \mathbf{w}) := \phi(r + a, \mathbf{w}), \tag{57}$$

considering the new densities $\tilde{\rho} := \rho - a$, and using (54), by Proposition 7 we obtain

$$\begin{aligned} & \sup_{\mu_0 \in \mathcal{M}_{(a,b),c}(\Omega)} \mathcal{W}_{\phi, \gamma\Omega}(S_T(\mu_0), \mu_\infty) \\ & \leq C \sup_{\mu_0 \in \mathcal{M}_{(a,b),c}(\Omega)} W_2(S_T(\mu_0) - a\gamma\Omega, \mu_\infty - a\gamma\Omega) < +\infty, \end{aligned} \tag{58}$$

because of the boundedness of the Wasserstein distance on the set of measures defined on the bounded convex set Ω . Finally, we conclude by (55), (56) and (58).

Also in this case, following the proof of Theorem 4, and using the equality (57), Proposition 7 and Theorem 5, we can prove the following theorem.

Theorem 6. *Let $\phi \in \mathcal{A}_2(a, b)$ and $\gamma\Omega := \chi_\Omega \mathcal{L}^d$ with $\Omega \subset \mathbb{R}^d$ a bounded convex smooth domain. If $(\mu_n)_{n \in \mathbb{N}}$ is a sequence in $\mathcal{M}_{(a,b),c}(\Omega)$ weakly-* convergent to $\mu \in \mathcal{M}_{(a,b),c}(\Omega)$, then*

$$\lim_{n \rightarrow +\infty} \mathcal{W}_{\phi, \gamma\Omega}(\mu_n, \mu) = 0.$$

We recall that the space of non-negative measures with fixed mass $c > 0$, supported on a bounded convex open set, is bounded with respect to the standard Wasserstein distance (easy consequence of the definition), and the convergence with respect to the standard Wasserstein distance is equivalent to the weak* one on this set. Theorems 5 and 6 state that the analogous properties hold for the space $\mathcal{M}_{(a,b),c}(\Omega)$ endowed with the distance $\mathcal{W}_{\phi, \gamma\Omega}$.

Appendix

Notation

In this appendix we collect some notation used in the paper.

Let X be a topological space, $A \subset X$, $f : X \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be a function. We denote by:

$\text{int}(A), \bar{A}, \partial A$	the <i>interior</i> , the <i>closure</i> and the <i>boundary</i> of A , respectively;
$\chi_A : X \rightarrow \{0, 1\}$	the <i>characteristic function</i> of A , namely $\chi_A(x) = 1$ if $x \in A$, $\chi_A(x) = 0$ if $x \notin A$;
$\text{dom}(f) := \{x \in X : f(x) \in \mathbb{R}\}$	the (<i>effective</i>) <i>domain</i> of f ;
$\text{epi}(f) := \{(x, \alpha) \in X \times \mathbb{R} : \alpha \geq f(x)\}$	the <i>epigraph</i> of f ;
$\text{hypo}(f) := \{(x, \beta) \in X \times \mathbb{R} : \beta \leq f(x)\}$	the <i>hypograph</i> of f .
\mathcal{L}^d	the <i>Lebesgue measure</i> on \mathbb{R}^d ;
$\mathcal{M}_{\text{loc}}(\mathbb{R}^d)$	the set of <i>signed Radon measures</i> on \mathbb{R}^d ;
$\mathcal{M}_{\text{loc}}^+(\mathbb{R}^d)$	the set of <i>non-negative Radon measures</i> on \mathbb{R}^d ;
$\mathcal{M}_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^h)$	the set of \mathbb{R}^h -valued Radon measures on \mathbb{R}^d .

We say that f is *lower semicontinuous* or *l.s.c.* (resp. *upper semicontinuous* or *u.s.c.*) iff $\text{epi}(f)$ (resp. $\text{hypo}(f)$) is closed in $X \times \mathbb{R}$. If (X, d) is a metric space, this is equivalent to say that f is l.s.c. (resp. u.s.c.) iff $f(x) \leq \liminf_{y \rightarrow x} f(y)$ (resp. $f(x) \geq \limsup_{y \rightarrow x} f(y)$).

Push-forward of measures Given a Borel measure μ on a topological space X , and a Borel map $T : X \rightarrow Y$, with values in a topological space Y , we define the image measure of μ through the map T , denoted by $\nu = T_{\#}\mu$, by $\nu(B) := \mu(T^{-1}(B))$, for any Borel measurable set $B \subset Y$, or equivalently

$$\int_Y \zeta(y) d\nu(y) = \int_X \zeta(T(x)) d\mu(x), \quad \forall \zeta \in C_b^0(Y). \quad (59)$$

If X and Y are domains of \mathbb{R}^d , the map T is sufficiently smooth and the measures μ and ν are absolutely continuous with respect to Lebesgue measure with densities $\tilde{\rho}$ and ρ respectively, then $\nu = T_{\#}\mu$ is equivalent, by the change of variables theorem, to

$$\rho(T(x)) \det(DT(x)) = \tilde{\rho}(x). \quad (60)$$

The formula (60) for the densities holds in a more general setting (see [3, Lemma 5.5.3]).

Convex analysis

In this appendix we recall some concepts from convex analysis, our main reference is [24].

Definition 9. (*Recession functional*) Let $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper convex function. The *recession functional* f^∞ of f is the positively homogeneous proper convex function defined by (cf. [24, Theorem 8.5, p. 66]):

$$f^\infty(y) := \sup\{f(x + y) - f(x) : x \in \text{dom } f\}.$$

If f is l.s.c, then f^∞ is l.s.c., and for any $x \in \text{dom}(f)$ it holds:

$$f^\infty(y) := \lim_{\lambda \rightarrow +\infty} \frac{f(x + \lambda y) - f(x)}{\lambda}.$$

We have that:

1. if $0 \in \text{dom}(f)$, it holds $f^\infty(y) := \lim_{\lambda \rightarrow +\infty} \frac{f(\lambda y)}{\lambda}$ for all $y \in \mathbb{R}^N$.
2. if $0 \notin \text{dom}(f)$, it holds $f^\infty(y) := \lim_{\lambda \rightarrow +\infty} \frac{f(\lambda y)}{\lambda}$ for all $y \in \text{dom}(f)$.

Definition 10. (*Concave–convex functions*) Let $C \subset \mathbb{R}^k, D \subset \mathbb{R}^d$ be convex sets, and $\tilde{f} : C \times D \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be a function. We will call \tilde{f} a *concave–convex* function if:

1. for each $z \in D$ the map $r \mapsto \tilde{f}(r, z)$ is concave,
2. for each $r \in C$ the map $z \mapsto \tilde{f}(r, z)$ is convex.

Given a concave–convex function $\tilde{f} : C \times D \rightarrow \mathbb{R}$, we define its *lower extension* $\tilde{f}_1 : \mathbb{R}^k \times \mathbb{R}^d \rightarrow \mathbb{R} \cup \{\pm\infty\}$ by setting:

$$\tilde{f}_1(r, z) = \begin{cases} \tilde{f}(r, z) & \text{if } r \in C, z \in D \\ +\infty & \text{if } r \in C, z \notin D \\ -\infty & \text{if } r \notin C \end{cases}$$

\tilde{f}_1 is still a concave–convex function.

Theorem 7. (*Partial Legendre*) Let $f : \mathbb{R}^k \times \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be convex and l.s.c. Then the function defined by:

$$\tilde{f}(r, z) := \sup_{w \in \mathbb{R}^d} [\langle z, w \rangle - f(r, w)]$$

is a concave–convex function from $\mathbb{R}^k \times \mathbb{R}^d$ to $\mathbb{R} \cup \{\pm\infty\}$. For every fixed r , the function $z \mapsto \tilde{f}(r, z)$ is l.s.c. Conversely, given any concave–convex function $\tilde{f} : \mathbb{R}^k \times \mathbb{R}^d \rightarrow \mathbb{R} \cup \{\pm\infty\}$, the function defined by:

$$f(r, w) := \sup_{z \in \mathbb{R}^d} [\langle z, w \rangle - \tilde{f}(r, z)]$$

is a convex map and for every fixed r , the function $w \mapsto f(r, w)$ is l.s.c. Moreover, if $\text{dom}(\tilde{f}) = C \times D$ and \tilde{f} agrees with its lower extension, then f is l.s.c.

Proof. See [24, Theorem 33.1].

Decay for heat equation

In this appendix we recall a standard result on the asymptotic behavior of the heat equation. Since it seems not simple to find it in this form, we also give a proof.

Lemma 3. *Let Ω be a convex smooth domain of \mathbb{R}^d . If $\rho_0 : \Omega \rightarrow [a, b]$, and $\rho : (0, +\infty) \times \Omega \rightarrow \mathbb{R}$ denotes the solution of the problem*

$$\begin{cases} \partial_t \rho - \Delta \rho = 0 & \text{in } (0, +\infty) \times \Omega \\ \rho(0, \cdot) = \rho_0 & \text{in } \Omega \\ \nabla \rho \cdot \mathbf{n} = 0 & \text{on } (0, \infty) \times \partial\Omega, \end{cases} \tag{61}$$

then there exist two constants $C >$ and $\lambda > 0$, depending only on a, b and Ω such that

$$\|\rho_s - \rho_\infty\|_{L^\infty(\Omega)} \leq C e^{-\lambda s}, \quad \forall s \geq 0, \tag{62}$$

where $\rho_s := \rho(s, \cdot)$ and $\rho_\infty := \frac{1}{\mathcal{L}^d(\Omega)} \int_\Omega \rho_0(x) \, dx$.

Proof. Since $\partial_t(\rho_t - \rho_\infty) - \Delta(\rho_t - \rho_\infty) = 0$ with homogeneous Neumann boundary conditions, multiplying this equation by $\rho_t - \rho_\infty$ and integrating by parts we obtain the identity

$$\frac{d}{dt} \|\rho_t - \rho_\infty\|_{L^2(\Omega)}^2 + 2\|\nabla \rho_t\|_{L^2(\Omega)}^2 = 0. \tag{63}$$

By Poincaré’s inequality, there exists a constant C_P depending only on Ω such that

$$\|\nabla \rho_t\|_{L^2(\Omega)}^2 \geq C_P \|\rho_t - \rho_\infty\|_{L^2(\Omega)}^2, \tag{64}$$

and from (63) we immediately obtain the $L^2(\Omega)$ exponential decay

$$\|\rho_t - \rho_\infty\|_{L^2(\Omega)} \leq e^{-C_P t} \|\rho_0 - \rho_\infty\|_{L^2(\Omega)}, \quad \forall t \geq 0. \tag{65}$$

The $L^2(\Omega) - W^{1,\infty}(\Omega)$ interpolation inequality (see for instance [6, Complements of Chapter IX] or [23]), states that there exist a constant C depending only on Ω such that

$$\|\rho_t - \rho_\infty\|_{L^\infty(\Omega)} \leq C \|\rho_t - \rho_\infty\|_{L^2(\Omega)}^{2/(d+2)} \|\rho_t - \rho_\infty\|_{W^{1,\infty}(\Omega)}^{d/(d+2)} \quad \forall t \geq 0. \tag{66}$$

In order to get a uniform bound of the L^∞ norm of the gradient, we define $v(t, x) := \rho_t^2(x) + t|\nabla \rho_t(x)|^2$, which solves the problem

$$\begin{cases} \partial_t v - \Delta v \leq 0 & \text{in } (0, +\infty) \times \Omega \\ v(0, \cdot) = \rho_0^2 & \text{in } \Omega \\ \nabla v \cdot \mathbf{n} \leq 0 & \text{on } (0, \infty) \times \partial\Omega. \end{cases} \tag{67}$$

Indeed, by a simple computation we have that v satisfies the partial differential inequality in (67). In order to obtain the boundary condition satisfied by v we have $\nabla v \cdot \mathbf{n} = \nabla \rho^2 \cdot \mathbf{n} + t \nabla |\nabla \rho|^2 \cdot \mathbf{n} = t \nabla |\nabla \rho|^2 \cdot \mathbf{n}$ because of the boundary condition in (61). Moreover, by the smoothness and the convexity of Ω , we have that $\nabla |\nabla \rho|^2 \cdot \mathbf{n} \leq 0$ (see for instance [19, Lemma 5.1]).

The maximum principle for problem (67) (see for instance [17]) states that $v(t, x) \leq \|\rho_0^2\|_{L^\infty(\Omega)}$. In particular we have

$$\sqrt{t}\|\nabla\rho_t\|_{L^\infty(\Omega)} \leq \|\rho_0\|_{L^\infty(\Omega)} \leq \max(|a|, |b|). \quad (68)$$

The inequality (62) follows from (66) and (68) (for $t \geq 1$ for instance) and (65), recalling that $\|\rho_t - \rho_\infty\|_{L^\infty(\Omega)} \leq 2 \max(|a|, |b|)$.

Acknowledgement. The authors would like to thank Giuseppe Savaré for useful suggestions on this work.

References

- [1] Ambrosio, L., Buttazzo, G.: Weak lower semicontinuous envelope of functionals defined on a space of measures. *Ann. Mat. Pura Appl.* **150**, 311–339 (1988)
- [2] Ambrosio, L., Fusco, N., Pallara, D.: *Functions of Bounded Variation and Free Discontinuity Problems*. Oxford Mathematical Monographs. Clarendon Press, Oxford (2000)
- [3] Ambrosio, L., Gigli, N., Savaré, G.: *Gradient Flows in Metric Spaces and in the Space of Probability Measures*. Birkhäuser Verlag, Basel (2005)
- [4] Beneamou, J.-D., Brenier, Y.: A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem. *Numer. Math.* **84**, 375–393 (2000)
- [5] Bertozzi, A.: The mathematics of moving contact lines in thin liquid films. *Not. Am. Math. Soc.* **45**, 689–697 (1998)
- [6] Brezis, H.: *Analyse fonctionnelle*. Masson, Paris (1983)
- [7] Burger, M., Di Francesco, M.: Large time behavior of nonlocal aggregation models with nonlinear diffusion. *Netw. Heterog. Media* **3**, 749–785 (2008)
- [8] Burger, M., di Francesco, M., Dolak, Y.: The Keller-Segel model for chemotaxis with prevention of overcrowding: linear vs. nonlinear diffusion. *SIAM J. Math. Anal.* **38**, 1288–1315 (2006)
- [9] Carrillo, J.A., Rosado, J., Salvarani, F.: 1D nonlinear Fokker-Planck equations for fermions and bosons. *Appl. Math. Lett.* **21**, 148–154 (2008)
- [10] Carrillo, J.A., Laurençot, P., Rosado, J.: Fermi-Dirac-Fokker-Planck equation: well-posedness & long-time asymptotics. *J. Differ. Equ.* **247**, 2209–2234 (2009)
- [11] Carrillo, J.A., Lisini, S., Savaré, G., Slepčev, D.: Nonlinear mobility continuity equations and generalized displacement convexity. *J. Funct. Anal.* **258**, 1273–1309 (2010)
- [12] Di Francesco, M., Rosado, J.: Fully parabolic Keller-Segel model for chemotaxis with prevention of overcrowding. *Nonlinearity* **21**, 2715–2730 (2008)
- [13] Dolbeault, J., Nazaret, B., Savaré, G.: A new class of transport distances between measures. *Calc. Var. Partial Differ. Equ.* **34**, 193–231 (2009)
- [14] Elliott, C.M., Garcke, H.: On the Cahn-Hilliard equation with degenerate mobility. *SIAM J. Math. Anal.* **27**, 404–423 (1996)
- [15] Frank, T.D.: Classical Langevin equations for the free electron gas and blackbody radiation. *J. Phys. A* **37**, 3561–3567 (2004)
- [16] Frank, T.D.: *Nonlinear Fokker-Planck Equations*. Springer Series in Synergetics. Springer, Berlin (2005)
- [17] Friedman, A.: *Partial Differential Equations of Parabolic Type*. Prentice-Hall, Englewood Cliffs (1964)
- [18] Giacomin, G., Lebowitz, J.: Phase segregation dynamics in particle systems with long range interactions. I. Macroscopic limits. *J. Stat. Phys.* **87**, 37–61 (1997)

-
- [19] Gianazza, U., Savaré, G., Toscani, G.: The Wasserstein gradient flow of the Fisher information and the quantum drift-diffusion equation. *Arch. Rational. Mech. Anal.* **194**, 133–220 (2009)
 - [20] Kaniadakis, G.: Generalized Boltzmann equation describing the dynamics of bosons and fermions. *Phys. Lett. A* **203**, 229–234 (1995)
 - [21] Kaniadakis, G., Quarati, P.: Kinetic equation for classical particles obeying an exclusion principle. *Phys. Rev. E* **48**, 4263–4270 (1993)
 - [22] Lisini, S., Matthes, D., Savaré, G.: (paper in preparation) (2010)
 - [23] Nirenberg, L.: On elliptic partial differential equations. *Ann. Scuola Norm. Sup. Pisa* **3**, 115–162 (1959)
 - [24] Rockafellar, R.T.: *Convex Analysis*. Princeton University, Princeton (1970)
 - [25] Slepčev, D.: Coarsening in nonlocal interfacial systems. *SIAM J. Math. Anal.* **40**, 1029–1048 (2008)
 - [26] Villani, C.: *Topics in Optimal Transportation*. Graduate Studies in Mathematics, vol. 58. American Mathematical Society, Providence (2003)
 - [27] Villani, C.: *Optimal Transport. Old and New*. Grundlehren der Mathematischen Wissenschaften, 338. Springer-Verlag, Berlin (2009)