Neuroradiology (2020) 62:1017
https://doi.org/10.1007/s00234-020-02468-y

CORRECTION

Correction to: Spatial coefficient of variation applied to arterial spin labeling MRI may contribute to predict surgical revascularization outcomes in pediatric moyamoya vasculopathy

Domenico Tortora[1] · Camilla Scavetta[2] · Giacomo Rebella[2] · Marta Bertamino[3] · Marcello Scala[4] · Thea Giacomini[5] · Giovanni Morana[1] · Marco Pavanello[4] · Andrea Rossi[1] · Mariasavina Severino[1]

Published online: 4 June 2020

Correction to: Neuroradiology (2020)
https://doi.org/10.1007/s00234-020-02446-4

This article was published online with incorrect alignment in Table 4. Column and rows are out of order. The correct Table 4 is presented here. The original article has been corrected.

Table 4 Correlations between perfusion parameters evaluated at each time points and long-term clinical outcome at 24 months after surgery

Perfusion Parameter	Pre-surgery		6 months		12 months		24 months	
	Rho	*P*	Rho	*P*	Rho	*P*	Rho	*P*
ASL-sCoV	−0.621	*<.001*	−0.551	*.001*	−0.487	*.003*	−0.389	*.007*
ASL-CBF	−0.117	.492	0.163	.371	0.217	.112	0.417	*.028*
DSC-CBF	0.258	.153	0.276	.202	0.228	.333	0.558	*.014*
DSC-TTP	−0.301	.041	−0.513	*.012*	−0.426	*.027*	−0.501	*.002*

Rho indicates Spearman's rho coefficient; *P* indicates significance level of the correlation analysis; the values reported in italics are statistically significant

The online version of the original article can be found at https://doi.org/10.1007/s00234-020-02446-4

✉ Andrea Rossi
andrearossi@gaslini.org

[1] Neuroradiology Unit, IRCCS Istituto Giannina Gaslini, via Gaslini 5, 16147 Genoa, Italy

[2] Radiology Section, Department of Health Sciences (DISSAL), University of Genoa, Genoa, Italy

[3] Rehabilitation Unit, IRCCS Istituto Giannina Gaslini, Genoa, Italy

[4] Neurosurgery Unit, IRCCS Istituto Giannina Gaslini, Genoa, Italy

[5] Neuropsychiatry Unit, IRCCS Istituto Giannina Gaslini, Genoa, Italy