

E. Olavi Kajander · Neva Ciftcioglu · Katja Aho
Enrique Garcia-Cuerpo

Characteristics of nanobacteria and their possible role in stone formation

Received: 22 March 2002 / Accepted: 10 February 2003 / Published online: 27 March 2003
© Springer-Verlag 2003

Abstract Kidney stone formation is a multifactorial disease in which the defence mechanisms and risk factors are imbalanced in favour of stone formation. We have proposed a novel infectious agent, mineral forming nanobacteria (NB), to be active nidi that attach to, invade and damage the urinary epithelium of collecting ducts and papilla forming the calcium phosphate center(s) found in most kidney stones. Stone formation may proceed in urine supersaturated with calcium phosphate, calcium oxalate and uric acid/urate under the influence of crystallization promoters and inhibitors. Our hypothesis underlines the role of active nidi: even supersaturated urine requires nidi for crystallization to appear.

Keywords Nanobacteria · Calcification · Kidney stones · Nephrolithiasis · Nephrocalcinosis

Introduction

The current probability of becoming affected by kidney stone disease in the western world ranges geographically from 5% to 13% [39]. The disease has numerous risk factors: genetic hereditary diseases, such as hyperoxaluria and Dent's disease [41], metabolic diseases, such as

hyperparathyreosis, life style and diet (eating, drinking) [50]. However, all of these factors cannot fully explain calcium stone formation. Ideas for risk factors and treatment or prevention of the disease have changed much in the recent years. However, nobody knows how the majority of cases without any metabolic/biochemical disorder should be treated to prevent the recurrence of stone attack. There was hope that reducing calcium from diet would help, but this only increased stone recurrence [31]. The hope placed on oxalate-eating bacteria introduced into the gut [43] has diminished because the immune system and antibiotic treatments may destroy the bacteria. Current knowledge on biochemical markers for risk factors and beneficial factors in urine is controversial. A consensus exists that urinary supersaturation is bad and high concentrations of citrate and possibly magnesium [42] are good. Otherwise, the etiological factors and thus mechanisms and treatment strategies remain poorly understood.

Characteristics of nanobacteria

Nanobacteria (NB) were discovered as a contaminating agent in cell culture over 10 years ago. Despite visible biomass present on a cell culture dish, standard microbiological methods failed to detect any known microbe [24]. Culture studies indicated that the novel agent was apparently indefinitely passagable in cell culture medium (beyond 10 years) and could adapt to growing in plain DMEM or RPMI-1640. Omitting serum supplementation resulted in larger cells [10, 22] being formed inside cavities formed by thick apatite layers (Fig. 1a, c, d). In old cultures, large colonies with slimy, but only slightly mineralized walls (arrows in Fig. 1b) were observed indicating social behaviour. Figure 1b also shows the release of tiny forms of NB from the colony. Further studies with electron microscopy revealed mineralized igloos consisting mainly of carbonate apatite (Fig. 1d). Such igloos could grow in size, bud-off new ones and fuse with others to form stones visible to the naked eye (Fig. 2a).


E. O. Kajander (✉) · K. Aho
Department of Biochemistry, University of Kuopio,
P.O.Box 1627, 70211 Kuopio, Finland
E-mail: olavi.kajander@uku.fi
Tel.: + 358-17-163060
Fax: + 358-172811510

E. O. Kajander
Nanobac Oy, Kuopio, Finland

N. Ciftcioglu
USRA-NASA Johnson Space Center, Houston, Texas

E. Garcia-Cuerpo
Hospital Ramon y Cajal, Universidad de Alcala, Madrid, Spain

Fig. 1 a Nanobacteria (NB) cultured without serum in DMEM, a phase contrast microscopy image, magnification 800 \times . The *arrow* shows a mineralized colony with two large organisms inside a mineralized igloo. **b** NB biofilm in DMEM. The micrograph shows a large community with relatively thin walls, see the two *arrows*. Small particles have been released outside the community. Magnification 800 \times . **c** Transmission electron microscopy (TEM) micrograph of a section of NB igloo. The micrograph shows an igloo similar to that in **a**. A layer of apatite crystals is evident on the surface. *Bar* = 0.2 μm . **d** Scanning electron microscopy (SEM) micrograph of igloos. *Bar* = 1 μm



Mineral growth took place as concentric layers of elongated nanoscale crystals of apatite with intervening pore structures (Fig. 2b). What was the source of this growth?


Cultures established with some fetal bovine serum (FBS) batches revealed growth of the mineralizing agent as dense particles having a diameter of 80 nm to 500 nm (Fig. 3a–d). Gamma irradiation abolished culture positivity. When the gamma-irradiated serum culture was inoculated with unirradiated positive serum or cultured agent, growth was restored. In this way, the source of the agent was tracked to ‘sterile’ FBS used as a supplement for culture medium [24]. Over 80% of tested commercial FBS batches from many different manufacturers were positive for NB [26]. NB were found in a significant number of humans as well, in serum and/or in urine. To date, NB have been isolated from bovine serum and from human serum, urine, kidney stones, dental stones and tissue samples [12, 13, 19, 24, 38, 49]. Interestingly, NB show improved growth properties in artificial urine [6]. The general characteristics of NB are given in Table 1. The compilation of data is based on research results obtained by the present authors on a standard “strain” (SeraLab901045). Intrastrain differences due to prolonged culture have been found in cytotoxicity [8] and in the kinetics of elimination into urine after intravenous injection into rats [1].

In addition to culture methods [9], several other diagnostic tools have been developed for the identification

of NB. One of the most powerful methods is transmission electron microscopy (TEM). TEM sample preparation for negative staining takes only a few minutes and allows for the detection of NB as dark particles, due to their apatite content [33]. As shown in Fig. 3, NB culture sample dried on a carbon-coated grid can be inspected either without staining, or after staining with 2% uranyl acetate. The latter reveals slimy material around the particles (Fig. 3c,d). The most powerful tool is a novel technique in which an unstained grid is incubated for 20 min with colloidal gold-labelled anti-NB antibody, washed, dried and inspected (Fig. 3b). The antibody reveals its target (NB surface epitope) inside of the slimy material on the NB surface. These techniques allow imaging at high resolution without any fixation steps. NB have apatite mineral as a structural support, which makes them visible and so robust that fixation is unnecessary for TEM. This is a unique feature of NB, which allows fast and specific diagnosis using electron microscopy.

As shown in Table 1, many properties of NB are rare and extreme. Relatively tiny mineral-associated microbes have been found by geologists [17, 48]. Older findings link such small forms to cancer [51]. Many properties of NB support the theory that they might be primitive life forms [25]. The extremophilic characteristics of NB would be beneficial in surviving hostile conditions [4, 23]. Several groups are researching NB and

Fig. 2 **a** SEM micrograph of cultured NB showing budding and fusion of spheroids, forming a large stone. *Bar* = 10 μm . **b** Detail of the previous area showing fractured top surface. Highly organized architecture with tiny elongated apatite crystals is evident. The *arrow* shows an apparent pore between adjacent crystals. *Bar* = 1 μm . **c** SEM micrograph of NB cultured from human kidney stone. The *arrow* points to a broken spheroid which has a highly organized lamellar wall. *Bar* = 10 μm . **d** SEM micrograph of a fractured human kidney stone revealing spheroid particles with a similar structure to **c**. *Bar* = 1 μm



have succeeded in the detection or culture of NB or NB-like forms [6, 15, 18, 19, 32, 38, 49]. However, the concept that NB are living organisms is controversial as long as their putative nucleic acid is not sequenced.


How nanobacteria are involved in stone formation?

The theory of NB-linked stone formation [28] is based on: (1) finding NB in kidney stones (Fig. 2d), (2) finding similar forms of NB in kidney stones culture (Fig. 2c), (3) in vitro calcific stone formation by NB (Fig. 1c, d, 2a, b), and (4) kidney stone formation after NB inoculation to rat kidneys. In a study by Ciftcioglu et al. [13], 70 out of 72 kidney stones contained NB. The presence of NB was independent of the stone type, although apatite stones gave the highest immunopositivity. Kidney stones were crushed, treated with 1 M HCl followed by neutralization and then analyzed using culture, immunological methods and electron microscopy. Surprisingly, kidney stones contained NB that started to replicate under culture conditions and formed calcium phosphate stones in vitro. In addition, NB were able to

produce stony colonies in modified Loeffler medium, cause intra- and extracellular calcium deposits and cell damage in many cultured cell lines [8, 11]. Importantly, dose-dependent kidney stone formation was observed within 1 month in rats after injection of NB using the translumbar, percutaneous renal puncture method [18]. Despite the small number of experimental animals ($n=4$), the result provides evidence that kidney stone formation can be caused by introducing NB into kidney.

The suggested involvement of NB in kidney stone formation [28] supports the observations made by Carr and Randall [7, 40]. Carr's concretions are small shiny deposits of calcium phosphate in kidney lymphatics and collecting ducts. Calcium phosphate formed above the collecting duct might induce heterogeneous nucleation of calcium oxalate at lower levels of the renal collection system [20] and be a risk factor for Randall's plaque formation [30]. Randall described calcium-containing plaques in the kidney papilla [40]. According to his hypothesis, the formation of kidney stone starts from these plaques due to a primary lesion in the tissue. Cell culture experiments have revealed that the adherence and internalization of calcium oxalate crystals into cultured

Fig. 3 **a** TEM micrograph of a sample from NB culture in DMEM supplemented with 10% FBS. Negative staining technique omitting the stain. *Bar* = 1 μm . **b** TEM micrograph of NB culture sample incubated for 20 min with colloidal gold-conjugated anti-NB 8D10 monoclonal antibody, otherwise as in **a**. Three approximately 20 nm-sized gold-particles are indicated by the *arrow*. There is highly significant binding of the gold-labelled antibody to NB. *Bar* = 0.12 μm . **c** TEM micrograph of NB subjected to negative staining with 2% uranyl acetate. *Bar* = 1 μm . **d** TEM image as in **c** at high magnification. Mucus-protein layer surrounding NB can be seen (see *arrow* in **c**). Such a sticky layer apparently helps NB to adhere on surfaces, to grow as a social colony or biofilm, and promotes apatite crystallization. *Bar* = 0.1 μm



cells is an active process potentially important for kidney stone formation [29]. Recent studies on NB have produced findings suggesting that NB might be calcium phosphate nuclei for kidney stone formation. NB are renotrophic, as reported from rabbit experiments using radiolabelled NB [3]. They are eliminated from the circulation through excretion into the urine [1, 3]. NB were found to adhere, invade and damage cells in collecting tubuli and the papillary area in the rat and rabbit models [3; Kajander, unpublished data]. NB colonization could lead to the accumulation of calcium deposits on the lesions and trigger stone formation as described by Randall.

Nanoscale biocrystallization by nanobacteria

Many kidney stones have a core composed of apatitic spheroids (Fig. 2d). NB cultured from human kidney stones formed apatitic spheroid particles in vitro with a similar architecture to that in kidney stone; observe the broken spheroids in Fig. 2c, d. Tiny nanoscale crystals show a highly organized structure resembling the nanosphere structures found in pearls and other calcium

carbonate formations in living bodies. This suggests a protein mediated mineral growth mechanism. Pearls are known to grow in this way. Figure 3c, d indicate that the biomineralization process of NB takes place inside a mucus-protein matrix around the NB that is detectable with uranyl acetate staining. This suggests that nanoscale apatite crystals grow as a result of mucus-protein mediated crystal formation on the surface of NB. This theory was first introduced by Vali et al. [49]. Crystalluria appears to form at a lower urinary ionic concentration in stone formers [16], suggesting higher crystallization potency in these individuals, i.e., active nuclei or weaker crystallization inhibitor activity. NB are transportable apatitic nuclei from blood into kidney tissue and urine [3], and their active role in crystallization may explain the observation above [13, 22, 28].

Biomineralization is an effective process: apatite formation in vitro stopped only when the calcium level decreased by 50% from 1.8 to 0.9 mM and the phosphate levels fell to near zero [13]. NB can use dolomite [12] and synthetic apatite (Kajander, unpublished observation) as a calcium source. NB-induced biomineralization is dependent on the presence of oxygen [22, 24]. Gamma irradiation at doses that prevented the

Table 1 General characteristics and behaviour of nanobacteria (NB)

Morphology	<p>Stained Gram-negative, sterile-filterable (0.22 μm pore-size), bacteria-like particles with varying amounts of a carbonate apatite coat</p> <p>Size of individual NB ranges from 80 nm to 500 nm</p> <p>By light and electron microscopy, apatite “igloos” have a central chamber occupied by one or more NB</p> <p>Under low nutrient conditions (e.g., serum-free), NB tend to form microscopic colonies in liquid media surrounded by a thick coat of calcium apatite; calcified colonies can approach >1 mm in size</p> <p>Exhibit budding and fragmentation, social behaviour, and communities reminiscent of biofilms, but with unique characteristics consistent with that of extremophiles; withstand 90°C for 1 h, 15 kGy gamma irradiation, 5% NaCl</p>
Growth and metabolism	<p>Serum forms have a generation time of about 3 days</p> <p>Serum-free forms double about every 6 days</p> <p>Can be passaged indefinitely in DMEM with or without serum</p> <p>Metabolism is 10,000 times slower than in <i>E. coli</i></p> <p>Incorporate uridine and methionine into DNA and protein, respectively</p> <p>Grow best under aerobic conditions: 5% CO₂:95% air</p> <p>Inhibitors of nucleic acid synthesis, 5-fluorouracil and cytosine arabinoside, inhibit NB growth</p> <p>Tetracycline, an apatite-binding protein synthesis inhibitor, inhibits NB growth at therapeutically achievable blood levels, as do trimethoprim, sulfamethoxazole, nitrofurantoin and ampicillin; at supra-pharmacologic levels, aminoglycosides also inhibit growth</p> <p>Calcium chelators, such as EGTA and citrate, inhibit growth in vitro</p> <p>Bisphosphonates are highly nanobactericidal</p>
Structure	<p>NB biomass contains novel proteins and “tough” polysaccharides</p> <p>Over 30 proteins have been found by SDS-PAGE</p> <p>One of these proteins is a bacterial porin protein</p> <p>Muramic acid, a major component of bacterial peptidoglycans, was identified</p> <p>The 16S rDNA of NB obtained with PCR places it in the alpha-2 subgroup of proteobacteria; further proof for nucleic acids are needed, since many data indicate that nucleic acids are modified and PCR methods may not work well</p>
Detection	<p>Monoclonal antibodies to the nanobacterial porin protein and peptidoglycan recognize intact NB as shown by immunogold labeling</p> <p>Hoechst DNA fluorochrome stains NB</p> <p>Demineralization of NB enhances their endotoxin positivity in the <i>Limulus</i> amoebocyte lysate assay</p> <p>Monoclonal antibodies to chlamydial lipopolysaccharide (endotoxin) react with NB</p>
Effect on cells	<p>Some, but not all isolates of NB exhibit cytotoxicity to mammalian cells in vitro</p> <p>NB can bind to mammalian cells in vitro and be internalized by endocytosis</p> <p>In human and animal tissues, transmission electron microscopy has revealed intracellular putative NB</p> <p>I.v. administered NB were excreted to urine in rodents. In rats they caused apoptosis and sloughing of renal epithelium in collecting ducts and papilla</p>

replication of NB, abolished the biomineralization [14, 24]. Biomineralization was abolished with several antibiotics and antimetabolites that showed a nanobactericidal effect at concentrations relevant for human therapy [14]. Further proof that biomineralization by NB are a biological phenomenon related to being a living entity came from recent experiments with light. Low intensity light treatment (without thermal effects) at certain wavelengths stimulated NB replication as detected by particle numbers, incorporation of uridine and electron microscopy. Concomitantly, light stimulated apatite formation as detected by ⁸⁵-strontium incorporation [45]. Synthetic apatite did not respond. Biostimulation by light treatment is a general phenomenon observed in living entities from bacteria to mammalian cells with the used light treatment [44, 46].

Macromolecule-calcium phosphate mineral complexes have also been recently observed in human and animal circulation by other researchers [34, 35, 36, 37]. Price hypothesized that these particles cause soft tissue calcification, such as atherosclerosis and kidney calcification. The source of Price’s particles was an enigma. These high molecular weight complexes of calcium

phosphate together with proteinaceous calcification inhibitors were circulating in rats (subjected to atherogenic treatments) after a single subcutaneous dose of etidronate [36]. The maximum concentration of complexes was observed at 6 h after the drug dose and complexes were cleared from circulation within 24 h after injection. The route of elimination was not studied. The presence of the protein-mineral complex increased total serum calcium and phosphate 1.8- and 1.6-fold, respectively, after a dose of 8 mg/100 g body weight etidronate, and even more with higher doses [36]. It was suggested that the complex originates due to the inhibition of bone mineralization by etidronate [36]. These findings confirm our detection of high molecular weight mineral-protein complexes containing calcium phosphate in serum (NB). We have shown that the treatment of such complexes inside calcific biofilms or stones with bisphosphonates, chelating agents and some antibiotics, resulted in the release of destroyed particles into the medium [14]. We propose an explanation for the appearance of complexes as described by Price et al.: bisphosphonate administration causes the destruction of NB into “popcorn-like” floating particulate debris in a

few hours [14], and these detached particles could appear in the blood until removed by the reticuloendothelial system. Bisphosphonates and chelating agents, either alone or together with antibiotics, might thus be useful agents in the treatment of pathological calcification, whether in the form of atherosclerosis or stone formation. In fact, a recent summary advocates bisphosphonate treatment for stopping or preventing atherosclerosis [52], and one earlier report has shown bisphosphonate therapy to decrease the recurrence of kidney stones [5]. Larger studies are warranted, because this approach might have deep implications in the treatment of recurrent kidney stones, nephrocalcinosis and atherosclerosis.

Cisar et al. [15] were able to culture NB-like apatitic particles from human saliva and dental plaques. They reported unsuccessful DNA extraction, failed PCR detection due to bacterial contamination and a negative result from protein isolation although some protein bands were obtained. Their conclusion was that the particles were self-replicating inorganic apatite. The use of positive and negative controls and methods to identify NB could have been used in their study to confirm or exclude the presence of NB, but were not performed. Interpretation should not be based only on failed nucleic acid results. Nucleic acid research on NB has many problems, e.g., nucleic acid extraction is difficult due to apatite and extracted DNA-like material has inhibited the amplification of exogenous bacterial DNA in PCR methods. More effort should be made for the characterization of NB.

Ongoing research on nanobacteria

Ongoing research aims at solving the mystery behind nanoscale biomineralization: What are NB? What are their survival and growth strategies? How do they mineralize and what is their role in kidney stone disease and other calcifications? Does their eradication prevent stone formation? Research is now being carried out by an increasing number of researchers, among others, in the Mayo Clinic and NASA. Effective eradication therapies may arise as a consequence of such international research efforts. New approaches in the treatment and prevention of kidney stones could significantly reduce health care costs and increase the quality of life. Emerging knowledge on the drug sensitivity of nanobacteria/kidney stone forming units [14] suggests that novel treatment strategies could be based on a combination therapy using "old" drugs. A major hindrance in adopting such new therapies can be the reluctance of drug companies to carry out the necessary but expensive clinical studies with generic drugs.

Epidemiological studies are important for determining the prevalence of NB in various populations and diseases. Serum prevalence of the antigen in adult Finnish volunteers is about 5% [26]. Our recent collaboration with Holmberg et al. from Uppsala revealed that

about 14% of healthy Swedish blood donors have antibodies to the agent [21]. Furthermore, gamma globulin preparations pooled from thousands of healthy volunteers revealed NB antibodies [2]. In some disease states, e.g. atherosclerosis and hemodialysis, NB markers, antibodies and antigen, can be found in the serum/urine in the majority of cases (Kajander, unpublished data).

Whether NB are bacteria, mineral autocatalytic aggregates or self-replicating biological particles, they should be regarded as an infectious agent which can be involved in the pathogenesis of pathological calcifications. Exposure to NB can cause an immune response and may result in chronic bacteremia. One accidental exposure to NB during laboratory work has been monitored. The exposure was followed by the development of antibodies against NB, antibody levels remaining high for several years after the accident (Kajander, unpublished data). This finding suggests that NB may cause chronic infection without immediate clinical symptoms. It has been estimated that the growth of a 3 mm thick layer of calcium oxalate takes approximately 2.7 years, based on crystal growth rate [47]. For this and other reasons, exposure to NB infection might have serious consequences several years after exposure. It is suspected that biopharmaceuticals might be contaminated via FBS [26] and some viral vaccines were found to contain NB [27]. This possibility should be kept in mind and efforts should be made to determine the role of NB in the etiology of kidney stones and pathological calcification, diseases with an apparently increasing prevalence.

Conclusions

NB remain controversial agents that mediate apatite nucleation and crystal growth. They are renotropic, cause apoptotic cell death, are present in human kidney stones and occasionally in urine. They may trigger renal pathology involving damage to tubular epithelium, biomineralization, and perhaps tubule obstruction and chronic infection resulting in defective tissue repair and stone formation.

Acknowledgments This study was supported by grants from the Sigrid Juselius Foundation, TEKES and the Research Council for Biosciences and Environment, Finnish Academy.

References

1. Aho K (2002) In vivo biodistribution and toxicity of nanobacteria in rodents. In: International Nanobacteria Minisymposium, March 8 2001, Kuopio. Book of abstracts, Kuopio University Press, Kuopio, (in press). Abstract at <http://www.nanobac.com/nbminisymp080301/page9.html>
2. Aho K, Kajander EO, Ciftcioglu N, Hjelle JT, Miller-Hjelle MA (2002) Screening of human gamma globulin products for nanobacteria markers. American Society for Microbiology, 102nd General Meeting; May 19–23 2002, Salt Lake City. Abstract Y-10, p. 507.

3. Akerman KK, Kuikka JT, Ciftcioglu N, Parkkinen J, Bergstrom KA, Kuronen I, Kajander EO (1997) Radiolabeling and in vivo distribution of nanobacteria in rabbit. *Proc SPIE Int Soc Opt Eng* 3111:436
4. Bjorklund M, Ciftcioglu N, Kajander EO (1998) Extraordinary survival of nanobacteria under extreme conditions. *Proc SPIE Int Soc Opt Eng* 3441:123
5. Bone HG3rd, Zerwekh JE, Britton F, Pak CY (1979) Treatment of calcium urolithiasis with diphosphonate: efficacy and hazards. *J Urol* 121:568
6. Burton SK (2001) Nanobacteria growth characteristics and biofilm formation. Proceedings American Society of Microbiology 101st General Meeting, May 20–24, 2001, Orlando. Abstract I-98, p 437
7. Carr RJ (1954) A new theory of the formation of renal calculi. *Brit J Urol* 26:105
8. Ciftcioglu N, Kajander EO (1998) Interaction of nanobacteria with cultured mammalian cells. *Pathophysiology* 4:259
9. Ciftcioglu N, Kajander EO (2000) Growth factors for nanobacteria. *Proc SPIE Int Soc Opt Eng* 3755:113
10. Ciftcioglu N, Pelttari A, Kajander EO (1997) Extraordinary growth phases of nanobacteria isolated from mammalian blood. *Proc SPIE Int Soc Opt Eng* 3111:429
11. Ciftcioglu N, Bjorklund M, Kajander EO (1998) Stone formation and calcification by nanobacteria in human body. *Proc SPIE Int Soc Opt Eng* 3441:105
12. Ciftcioglu N, Ciftcioglu V, Vali H, Turcott E, Kajander EO (1998) Sedimentary rocks in our mouth: dental pulp stones made by nanobacteria. *Proc SPIE Int Soc Opt Eng* 3441:130
13. Ciftcioglu N, Bjorklund M, Kuorikoski K, Bergstrom K, Kajander EO (1999) Nanobacteria: an infectious cause for kidney stone formation. *Kidney Int* 56:1893
14. Ciftcioglu N, Miller-Hjelle MA, Hjelle JT, Kajander EO (2002) Inhibition of nanobacteria by antimicrobial drugs as measured by a modified microdilution method. *Antimicrob Agents Chemother* 46:2077
15. Cisar JO, Xu D-Q, Thompson J, Swaim W, Hu L, Kopecko DJ (2000) An alternative interpretation of nanobacteria-induced biomineralization. *Proc Natl Acad Sci U S A* 97:11511
16. Fan J, Chandhoke PS (1999) Examination of crystalluria in freshly voided urines of recurrent calcium stone formers and normal individuals using a new filter technique. *J Urol* 161:1685
17. Folk RL (1993) SEM imaging of bacteria and nannobacteria in carbonate sediments and rocks. *J Sediment Petrol* 63:990
18. Garcia Cuerpo E, Kajander EO, Ciftcioglu N, Lovaco Castellano F, Correa C, Gonzalez J, Mampaso F, Liano F, Garcia de Gabiola E, Escudero Barrileiro A (2000): Nanobacteria. Un modelo de neo-litogenesis experimental. *Arch Esp Urol* 53:291
19. Hjelle JT, Miller-Hjelle MA, Poxton IR, Kajander EO, Ciftcioglu N, Jones ML, Caughey RC, Brown R, Millikin PD, Darras FS (2000) Endotoxin and nanobacteria in polycystic kidney disease. *Kidney Int* 57:2360
20. Hojgaard I, Fornander AM, Nilsson MA, Tiselius HG (1999) Crystallization during volume reduction of solutions with a composition corresponding to that in the collecting duct: the influence of hydroxyapatite seed crystals and urinary macromolecules. *Urol Res* 27:417
21. Holmberg M (2002) Serological evidence for nanobacteria prevalence suggests a possible zoonosis. In: International Nanobacteria Minisymposium, March 8 2001, Kuopio. Book of abstracts, Kuopio University Press, Kuopio, (in press). Abstract at <http://www.nanobac.com/nbminisymph080301/page10.html>
22. Kajander EO, Ciftcioglu N (1998) Nanobacteria: an alternative mechanism for pathogenic intra- and extracellular calcification and stone formation. *Proc Natl Acad Sci U S A* 95:8274
23. Kajander EO, Ciftcioglu N (2000) Nanobacteria as extremophiles. *Proc SPIE Int Soc Opt Eng* 3755:106
24. Kajander EO, Kuronen I, Akerman K, Pelttari A, Ciftcioglu N (1997) Nanobacteria from blood, the smallest culturable autonomously replicating agent on Earth. *Proc SPIE Int Soc Opt Eng* 3111:420
25. Kajander EO, Bjorklund M, Ciftcioglu N (1998) Mineralization by nanobacteria. *Proc SPIE Int Soc Opt Eng* 3441:86
26. Kajander EO, Heinonen T, Kuronen I, Luoto K, Ciftcioglu N (1999) Nanobacteria, mycoplasma and bacterial L-forms; problems for sterile filtration. In: Wirtanen G, Salo S, Mikkola A (eds) 30th R3-Nordic Contamination Control Symposium, Helsinki, Finland, May 30–June 2, 1999. VTT Technical Research Centre of Finland, Espoo, p 279
27. Kajander EO, Ciftcioglu N, Aho K (2001) Detection of nanobacteria in viral vaccines. Proceedings of the American Society of Microbiology, 101st General Meeting, 20–24 May 2001, Orlando. Abstract Y-3, p 736.
28. Kajander EO, Ciftcioglu N, Miller-Hjelle MA, Hjelle JT (2001) Nanobacteria: controversial pathogens in nephrolithiasis and polycystic kidney disease. *Curr Opin Nephrol Hypertens* 10:445
29. Lieske JC, Toback FG (2000) Renal cell-urinary crystal interactions. *Curr Opin Nephrol Hypertens* 9:349
30. Low RK, Stoller ML, Schreiber CK (2000) Metabolic and urinary risk factors associated with Randall's papillary plaques. *J Endourol* 14:507
31. Martini LA, Wood RJ (2000) Should dietary calcium and protein be restricted in patients with nephrolithiasis? *Nutr Rev* 58:111
32. McKay DS, Gibson EKJr, Thomas-Keptra KL, Vali H, Romanek CS, Clemett SJ, Chillier XD, Maechling CR, Zare RN (1996) Search for past life on Mars: Possible relic biogenic activity in Martian meteorite ALH840001. *Science* 273:924
33. Miller-Hjelle MA, Hjelle JT, Ciftcioglu N, Kajander EO (2003). Nanobacteria: methods for growth and identification of this recently discovered calciferous microbe. In: Olson W (ed) Rapid analytical microbiology: the chemistry and physics of microbial identification. Sue Horwood, Storrington (in press)
34. Price PA, Buckley JR, Williamson MK (2001) The amino bisphosphonate ibandronate prevents vitamin D toxicity and inhibits vitamin D-induced calcification of arteries, cartilage, lungs and kidneys in rats. *J Nutr* 131:2910
35. Price PA, Faus SA, Williamson MK (2001) Bisphosphonates alendronate and ibandronate inhibit artery calcification at doses comparable to those that inhibit bone resorption. *Arterioscler Thromb Vasc Biol* 21:817
36. Price PA, Thomas GR, Pardini AW, Figueira WF, Caputo JM, Williamson MK (2002) Discovery of a high molecular weight complex of calcium, phosphate, fetuin, and matrix γ -carboxyglutamic acid protein in the serum of etidronate-treated rats. *J Biol Chem* 277:3926
37. Price PA, Omid N, Than TN, Williamson MK (2002) The amino bisphosphonate ibandronate prevents calciphylaxis in the rat at doses that inhibit bone resorption. *Calcif Tissue Int* 71:356
38. Puskás L (2002) Detection of nanobacteria in human atherosclerotic plaques. In: International Nanobacteria Minisymposium, March 8 2001, Kuopio. Book of abstracts, Kuopio University Press, Kuopio, (in press). Abstract at <http://www.nanobac.com/nbminisymph080301/page13.html>
39. Ramello A, Vitale C, Marangella M (2000) Epidemiology of nephrolithiasis. *J Nephrol* 13:S45
40. Randall A (1937) The origin and growth of renal calculi. *Ann Surg* 105:1009
41. Scheinmann SJ (1999) Nephrolithiasis. *Semin Nephrol* 19:381
42. Schwartz BF, Bruce J, Leslie S, Stoller ML (2001) Rethinking the role of urinary magnesium in calcium urolithiasis. *J Endourol* 15:233
43. Sidhu H, Schmidt ME, Cornelius JG, Thamilselvan S, Khan SR, Hesse A, Peck AB (1999) Direct correlation between hyperoxaluria/oxalate stone disease and the absence of the gastrointestinal tract-dwelling bacterium *Oxalobacter formigenes*: possible prevention by gut recolonization or enzyme replacement therapy. *J Am Soc Nephrol* 10 (Suppl 14):S334

44. Sommer AP, Pinheiro AL, Mester AR, Franke RP, Whelan HT (2001) Biostimulatory windows in low-intensity laser activation: lasers, scanners and NASA's light-emitting diode array system. *J Clin Laser Med Surg* 19:29
45. Sommer AP, Hassinen HI, Kajander EO (2002) Light induced replication of nanobacteria-a preliminary report. *J Clin Laser Med Surg* 20:241
46. Sommer AP, Oron U, Kajander EO, Mester AR (2002) Stressed cells survive better with light. *J Proteome Res* 1:475
47. Söhnle O, Grases F (1993) Fine structure of calcium oxalate monohydrate renal calculi. *Nephron* 63:176
48. Uwins PJR, Webb RI, Taylor AP (1998) Novel nano-organisms from Australian sandstones. *Am Mineral* 83:1541
49. Vali H, McKee MD, Ciftcioglu N, Sears SK, Plows FL, Chevet E, Ghiasi P, Plavsic M, Kajander EO, Zare RN (2001) Nanoforms: a new type of protein-associated mineralization. *Geochim Cosmochim Acta* 65:63
50. Wahl C, Hess B (2000) Kidney calculi – is nutrition a trigger or treatment? *Ther Umsch* 57:138
51. Wainwright M (1999) Nanobacteria and associated 'elementary bodies' in human disease and cancer. *Microbiology* 145:2623
52. Ylitalo R (2002) Bisphosphonates and atherosclerosis. *Gen Pharmacol* 35:287