

Carleman Estimates with No Lower-Order Terms for General Riemann Wave Equations. Global Uniqueness and Observability in One Shot*

Roberto Triggiani¹ and P. F. Yao²

¹Department of Mathematics, University of Virginia,
Charlottesville, VA 22904, USA

²Institute of Systems Science, Academy of Mathematics and Systems Science,
Chinese Academy of Sciences, Beijing 100080, People's Republic of China

Abstract. This paper considers a fully general (Riemann) wave equation on a finite-dimensional Riemannian manifold, with energy level ($H^1 \times L_2$)-terms, under essentially minimal smoothness assumptions on the variable (in time and space) coefficients. The paper provides Carleman-type inequalities: first pointwise, for C^2 -solutions, then in integral form for $H^{1,1}(Q)$ -solutions. The aim of the present approach is to provide Carleman inequalities which do *not* contain lower-order terms, a distinguishing feature over most of the literature. Accordingly, global uniqueness results for overdetermined problems as well as Continuous Observability/Uniform Stabilization inequalities follow in one shot, as a part of the same stream of arguments. Constants in the estimates are, therefore, generally explicit. The paper emphasizes the more challenging pure Neumann B.C. case. The paper is a generalization from the Euclidean to the Riemannian setting of [LTZ] in the more difficult case of purely Neumann B.C., and of [KK1] in the case of Dirichlet B.C. The approach is Riemann geometric, but different from—indeed, more flexible than—the one in [LTY1].

Key Words. Riemann wave, Carleman estimates, Observability.

AMS Classification. 35, 49, 53, 93.

* The research by R. Triggiani was partially supported by the National Science Foundation under Grant DMS-9804056 and the Army Research Office under Grant DAA4-96-1-0059. The research by P. F. Yao was performed while he was visiting the Mathematics Department, University of Virginia, Charlottesville, VA 22904, USA, and he was partially supported by NSF Grant DMS-9804056, and by the National Science Foundation of China.

1. Introduction

Throughout this paper M is a finite-dimensional Riemannian manifold with metric $g(\cdot, \cdot) = \langle \cdot, \cdot \rangle$ and squared norm $|X|^2 = g(X, X)$, and Ω is an *open bounded*, connected, compact set of M with smooth boundary $\Gamma = \Gamma_0 \cup \Gamma_1$. We let n denote the outward unit normal field along the boundary Γ . Further, we denote by Δ the Laplace (Laplace–Beltrami) operator on the manifold M and by D the Levi–Civita connection on M [D], [Le].

In this paper we study the following wave equation with energy level terms on Ω :

$$w_{tt} = \Delta w + F(w) + f \quad \text{in } (0, T] \times \Omega \quad (1.1)$$

under the following standing assumptions: the energy-level differential term

$$F(w) \equiv \langle P(t, x), Dw \rangle + p_1(t, x)w_t + p_0(t, x)w, \quad (1.2a)$$

where p_0, p_1 are functions on $[0, T] \times \Omega$, and $P(t)$ is a vector field on M for $t > 0$, satisfies the following estimate: there exists a constant $C_T > 0$ such that

$$|F(w)|^2 \leq C_T \{|Dw|^2 + w_t^2 + w^2\}, \quad x, t \in Q, \quad (1.2b)$$

where $Dw = \nabla_g w$ for the scalar function w . So Dw is a vector field, $Dw \in \mathcal{X}(M) =$ the set of all vector fields on M . Two vertical bars $|\cdot|$ denote the norm in the tensor space T_x or its completion $L^2(\Omega, T)$ [He]. Furthermore, we assume throughout that the forcing term f in (1.1) satisfies

$$f \in L_2(0, T; \Omega) \equiv L_2(Q); \quad \int_Q f^2 dQ < \infty, \quad (1.3)$$

where $dQ = d\Omega dt$, and $d\Omega$ is the volume element of the manifold M in its Riemann metric g .

Remark 1.1. Property (1.2b) is fulfilled if $P \in L_\infty(0, T; \Lambda)$ [He], $p_0, p_1 \in L_\infty(Q)$. In effect, one could relax the regularity assumption on the lower-order coefficient p_0 to read $p_0 \in L_p(Q)$ for $p = \dim \Omega + 1$, see Remark 1.1.1 of [LTZ], by using a Sobolev embedding theorem.

Main Assumptions. In addition to the standing assumptions (1.2b), (1.3) on the first-order operator F and the forcing term f , the following assumption is postulated throughout Section 9 of this paper.

(A.1) There exists a function $d: \bar{\Omega} \Rightarrow \mathbb{R}$ of class C^3 that is strictly convex in the metric g . This means that the Hessian D^2d (a two-order tensor) satisfies $D^2d(X, X) > 0$, $\forall x \in \bar{\Omega}, \forall X \in M_x$. By translation and rescaling, we can always achieve (see Remark 1.2 below) that $d(x)$ satisfies the following conditions:

$$\begin{aligned} D^2d(X, X) &\equiv \langle D_X(Dd), X \rangle_g \geq 2|X|_g^2, & \forall x \in \Omega, \quad \forall X \in M_x, \\ \min_{\bar{\Omega}} d(x) &\equiv m > 0, \end{aligned} \quad (1.4)$$

where, as we said, D^2d is the Hessian of d (a second-order tensor) and where M_x is the tangent space at $x \in \Omega$. In (1.4), we think of m as arbitrarily small, in order not to deteriorate the threshold time T_0 in (1.6b) below.

A working assumption throughout Section 9, to be later relaxed in Section 10, is that $d(x)$ has no critical point on $\bar{\Omega}$:

(A.2)

$$\inf_{x \in \bar{\Omega}} |Dd| = p > 0, \quad \text{so that we can take } \kappa = \inf_{\bar{\Omega}} \frac{|Dd|^2}{d} > 4 \quad (1.5)$$

by rescaling, see Remark 1.2 below. The above setting is kept throughout Chapter 9.

Pseudo-Convex Function. Having chosen, on the strength of assumption (A.1), a strictly convex potential function $d(x)$ satisfying condition (1.4), we next introduce the function $\varphi: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ of class C^3 by setting

$$\varphi(x, t) = d(x) - c \left(t - \frac{T}{2} \right)^2, \quad 0 \leq t \leq T, \quad x \in \Omega, \quad (1.6a)$$

where $T > 0$ and $0 < c < 1$ are selected as follows. We define first T_0 by setting, as usual,

$$T_0^2 = 4 \max_{x \in \bar{\Omega}} d(x). \quad (1.6b)$$

Let $T > T_0$ be given. By (1.6b), there exists $\delta > 0$ such that

$$T^2 > 4 \max_{x \in \bar{\Omega}} d(x) + 4\delta. \quad (1.6c)$$

For this $\delta > 0$, there exists a constant c , $0 < c < 1$, such that

$$cT^2 > 4 \max_{x \in \bar{\Omega}} d(x) + 4\delta. \quad (1.6d)$$

Henceforth, let $\varphi(x, t)$ be defined by (1.6a) with T and c chosen as described above, unless otherwise explicitly noted. Such a function $\varphi(x, t)$ has the following properties:

(a) For the constant $\delta > 0$, fixed in (1.6c), we have via (1.6d) that

$$\begin{aligned} \varphi(x, 0) = \varphi(x, T) = d(x) - c \frac{T^2}{4} &\leq \max_{\bar{\Omega}} d(x) - c \frac{T^2}{4} \leq -\delta, \\ \text{uniformly in } x \in \bar{\Omega}. \end{aligned} \quad (1.7)$$

(b) There are t_0 and t_1 , with $0 < t_0 < T/2 < t_1 < T$, such that

$$\min_{x \in \bar{\Omega}, t \in [t_0, t_1]} \varphi(x, t) \geq \sigma, \quad 0 < \sigma < m, \quad (1.8)$$

since $\varphi(x, T/2) = d(x) \geq m > 0$, under the present choice, with m given by (1.4).

Remark 1.2. If we rescale the original function $d(x)$ ($d(x) \geq m > 0$), i.e., if we replace $d(x)$ by $d_{\text{new}}(x) = ad(x)$ for a constant $a > 1$, then

$$\left\{ \begin{array}{l} Dd_{\text{new}} = aDd; \quad |Dd_{\text{new}}|^2 = a^2|Dd|^2; \quad D^2d_{\text{new}} = aD^2d; \\ T_{0,\text{new}}^2 = aT_0^2; \quad T_{\text{new}}^2 = aT^2; \quad D^2d_{\text{new}}(Dd_{\text{new}}, Dd_{\text{new}}) = a^3D^2d(Dd, Dd); \\ \text{the coercivity lower bound 2 in (1.4) is replaced by } 2a > 2; \\ \kappa_{\text{new}} = \inf_{\Omega} \frac{|Dd_{\text{new}}|^2}{d_{\text{new}}} = a\kappa = a \inf_{\Omega} \frac{|Dd|^2}{d}. \end{array} \right.$$

Hence, condition (1.5, right) can always be achieved by rescaling, if $d(x)$ has no critical points in $\bar{\Omega}$. The importance of condition $\kappa > 4$ in (1.5) will be seen in (1.21) below, in asserting that

$$f(c) \equiv (5 + 3c)\kappa - 4(1 + 7c) \geq 0 \quad \text{for all } c < 1 \text{ and } c \text{ near } 1. \quad (1.9)$$

Indeed, $f(c = 1) > 0$ by (1.5, right), and hence (1.9) holds true. Thus, c in (1.6d) can be taken arbitrarily close to 1, as needed.

Consequences of Assumptions (A.1), (A.2) and of the Scaling Condition in (1.5). Let $d(x)$ be the strictly convex (potential) function provided by assumptions (A.1), (A.2) and subject to the translation/scaling conditions in (1.4), (1.5). It then follows that by choosing the function

$$\alpha(x) \equiv \Delta d(x) - c - 1, \quad (1.10)$$

the following two properties, (p₁), (p₂), hold true (see proof of (1.12) below, while (1.11) is immediate):

$$\begin{array}{l} \text{(p}_1\text{)} \quad \Delta d(x) - 2c - \alpha(x) \equiv 1 - c > 0, \quad \forall x \in \bar{\Omega}; \\ \text{(p}_2\text{)} \end{array} \quad (1.11)$$

$$\begin{aligned} & [2c + \Delta d - \alpha]|Dd|^2 + 2D^2d(Dd, Dd) - 4c^2(\Delta d + 6c - \alpha) \left(t - \frac{T}{2} \right)^2 \\ & \geq 4(1 + 7c)\varphi^*(x, t), \quad \forall t, x \in Q = [0, T] \times \Omega, \end{aligned} \quad (1.12)$$

where $\varphi^*(x, t)$ is the function defined by

$$\varphi^*(x, t) = d(x) - c^2 \left(t - \frac{T}{2} \right)^2, \quad x \in \Omega, \quad 0 \leq t \leq T. \quad (1.13)$$

Since $0 < c < 1$, we note via (1.6a), (1.13) that

$$\varphi^*(x, t) \geq \varphi(x, t), \quad x \in \Omega, \quad 0 \leq t \leq T. \quad (1.14)$$

Next, we define two sets, subsets of $\Omega \times [0, T]$:

$$Q(\sigma) \equiv \{(x, t): x \in \Omega, 0 \leq t \leq T, \varphi(x, t) \geq \sigma > 0\}, \quad (1.15)$$

$$Q^*(\sigma^*) \equiv \{(x, t): x \in \Omega, 0 \leq t \leq T, \varphi^*(x, t) \geq \sigma^* > 0\}, \quad 0 < \sigma^* < \sigma, \quad (1.16)$$

for a constant σ^* chosen to satisfy $0 < \sigma^* < \sigma$, with $\sigma < m$ defined in (1.8). Indeed, recalling (1.8), we readily have

$$[t_0, t_1] \times \Omega \subset Q(\sigma) \subset Q^*(\sigma^*) \subset [0, T] \times \Omega. \quad (1.17)$$

Proof of (1.12). By virtue of the choice (1.10) for α and assumption (1.5), we compute via (1.10), (1.5), (1.9), for all $(t, x) \in Q$:

$$\begin{aligned} [2c + \Delta d - \alpha]|Dd|^2 + 2D^2d(Dd, Dd) - 4c^2(\Delta d + 6c - \alpha) \left(t - \frac{T}{2}\right)^2 \\ = (1 + 3c)|Dd|^2 + 2D^2d(Dd, Dd) - 4c^2(1 + 7c) \left(t - \frac{T}{2}\right)^2 \end{aligned} \quad (1.18)$$

$$\text{(by (1.4))} \geq (5 + 3c)|Dd|^2 - 4c^2(1 + 7c) \left(t - \frac{T}{2}\right)^2 \quad (1.19)$$

$$\text{(by (1.5))} \geq (5 + 3c)\kappa d(x) - 4c^2(1 + 7c) \left(t - \frac{T}{2}\right)^2 \quad (1.20)$$

$$\text{(by (1.9))} \geq 4(1 + 7c) \left[d(x) - c^2 \left(t - \frac{T}{2}\right)^2 \right] \equiv 4(1 + 7c)\varphi^*(x, t), \quad (1.21)$$

recalling (1.13), and (1.12) is proved, as desired.

Critical Property to Absorb Lower-Order Terms. We shall see in Section 4, (4.20) below, that the following property, derived from (1.12), is critical in the present approach in absorbing lower-order terms in w : under assumptions (A.1), (A.2) and for the choice of α in (1.10) we have:

(p'2)

$$\begin{aligned} [2c + \Delta d - \alpha]|Dd|^2 + 2D^2d(Dd, Dd) - 4c^2(\Delta d + 6c - \alpha) \left(t - \frac{T}{2}\right)^2 \\ \text{(by (1.12))} \geq 4(1 + 7c)\varphi^*(x, t) \geq 4(1 + 7c)\sigma^* = \beta_1 > 0, \\ \forall (x, t) \in Q^*(\sigma^*), \end{aligned} \quad (1.22)$$

recalling the definition of the set $Q^*(\sigma^*)$ in (1.16).

The term on the left-hand side of (1.22) is precisely the term that will arise as a coefficient in front of the lower-order term, see (4.5), (4.7), (4.8) below.

2. Main Results of the Paper. Literature

Throughout this paper the “energy” is given by

$$E(t) = \int_{\Omega} [w_t^2 + |Dw|^2 + w^2] d\Omega, \quad (2.1)$$

see (5.4). This is the quantity which occurs in the estimates.

Main Results of the Paper. We hereby summarize the main results of the present paper on the Riemann wave equation (1.1), with the understanding that *the standing hypotheses (1.2) on F and (1.3) on f are in force throughout.*

(1) *Results under assumptions (A.1) = (1.4) and (A.2) = (1.5).* These include, in order of presentation:

(1a) Carleman-type estimates which contain no lower-order terms (l.o.t.) and are valid for $H^{2,2}(Q)$ -solutions of (1.1) (with no B.C.). The first version thereof is Theorem 5.1 (Section 5), while the second, final, version is Theorem 6.1 (Section 6). Their proof rests on the fundamental lemma of Section 3, which provides a *pointwise* Carleman-type estimate (at each (t, x)) for C^2 -solutions of (1.1), with *controlled sign* on the coefficient of the l.o.t., at least in a suitable set in space and time of $\Omega \times [0, T]$. This set is $Q^*(\sigma^*)$ under assumptions (A.1) and (A.2).

(1b) Extension of Theorems 5.1 and 6.1 to the $H^{1,1}(Q)$ -solution of (1.1), by a regularizing procedure. This is done in Section 7.

(1c) A Continuous Observability Inequality—a fortiori a global uniqueness theorem—for $H^{1,1}(Q)$ -solutions, as given in Theorem 8.1 (Section 8) in the case of pure homogeneous Dirichlet B.C.; and as given in Theorems 9.1 and 9.3 (Section 9) in the case of pure homogeneous Neumann B.C.

(2) *Extension of the results of part (1) to a setting that removes assumption (A.2) = (1.5).* This is done, as in Section 10 of [LTZ], by considering Ω as the union of two suitable overlapping subdomains Ω_1 and Ω_2 : $\Omega = \Omega_1 \cup \Omega_2$, and assuming the existence of two smooth strictly convex functions d_1 and d_2 on Ω (as in (A.1) = (1.4)), with the additional feature that d_i has no critical point on Ω_i , thereby eliminating assumption (A.2), at least on each Ω_i . Accordingly, the main result of the present paper is Theorem 10.1.1. Only some key aspects of the proof of this theorem will be given explicitly in Section 10—those dealing with the elimination of the l.o.t. in the final Carleman estimates—as the proof follows rather faithfully the one given in Section 10 of [LTZ].

Literature. To begin with, regarding the generality of model (1.1), we notice that such a Riemann wave equation includes, in particular, a general second-order hyperbolic equation defined on a Euclidean bounded domain, with principal part coefficients $a_{ij}(x)$ variable in space, and coefficients of energy level terms variable in both time and space [LTY1], [LTY2]. The case with no energy level terms was dealt with in [Y], where the geometric method was introduced. Here in these aforementioned references, the transformation $\{g_{ij}(x)\} = \{a_{ij}(x)\}^{-1}$ (two $n \times n$ positive symmetric matrices), gives the coefficients of the Riemann metric g . Indeed, one may more generally start with a general second-order elliptic operator on the manifold M , and change it into the Laplace–Beltrami operator as in (1.1), through a similar transformation based on the coefficients. Having assessed the generality of our model, we may next state our goal: the main goal of the present paper is then to obtain Carleman-type inequalities for the aforementioned general model (1.1), with the additional requirement that they do not contain l.o.t. This is a most desirable feat, which is in contrast with much of the large literature on this subject, from the canonical equations of the mid-80s [LT3], [LT8], [Lio1], [Lio2] to the very general geometric optics approach in [BLR], [Lit1], and [Lit2] to the earlier approaches to Carleman estimates on Euclidean setting [T1], [LT5] to the Riemann geo-

metric generalization in [LTY1] and [LTY2]. The advantages of eliminating l.o.t. in the Carleman estimates are then several and substantial. To begin with, as a consequence, this approach permits one to obtain global uniqueness results of overdetermined problems, as well as the much stronger continuous observability/uniform stabilization inequalities in one shot, as part of the same stream of arguments. This is in contrast with much of the literature on the subject, where elimination of l.o.t. from the estimate would require appealing to, or invoking or assuming a global uniqueness result, not a readily-available task in the presence of nonsmooth coefficients, particularly with $L_\infty(Q)$ -coefficients in time and space at the energy level. By contrast the present approach establishes—rather than assumes—global uniqueness results, precisely in the form needed by the COI/US inequalities, which compare very favorably with the literature [Ho1], [Ho2], [Ho3], [I], [Lit3], [T2], [T5]. An additional bonus of the present approach is that the final COI/US inequalities are obtained with explicit constants, certainly so in the case of Dirichlet B.C., and also in the case of Neumann B.C., if one tolerates geometrical conditions on the controlled/observed portion Γ_1 of the boundary. Knowledge of explicit constants in these estimates plays a positive role in dealing with semilinear problems [LT9], [Z1], [Z2], [Z3] as well as in estimating the minimal norm of the steering controls. Only a few recent references [T3], [IY], [LTZ] have addressed successfully the issue of obtaining Carleman/COI/US estimates with no l.o.t. for control theoretic purposes. Both [IY] and its more general version [LTZ], done simultaneously and independently, refer, however, to the case of the *Euclidean* Laplacian, with particular emphasis on the more challenging purely Neumann case. Also [KK], which refers to $H^{2,2}(Q)$ -solutions and includes the Dirichlet, or Dirichlet/Neumann, but excludes the purely Neumann, case, refers to the Euclidean Laplacian. All are inspired by a pointwise Carleman-type estimate [LRS, Lemma 1, p. 124]: indeed, [KK] and [IY] invoke such a pointwise Carleman estimate, while [LTZ] provides a new version of it, more suitable for the purely Neumann B.C. case, in order to relax geometrical conditions. The present paper is a faithful extension of [LTZ] from the Euclidean to the Riemannian setting. In particular, the preliminary fundamental Lemma 3.1 of Section 3 is the present Riemannian counterpart of Lemma 3.1 in [LTZ] in the Euclidean setting, in turn related—as recalled above—to Lemma 1 on p. 124 of [LRS]. One may give several classes of examples where all the necessary assumptions—particularly in the purely Neumann B.C. case—are satisfied. One such attractive class is given in Appendix B, which is the Riemannian counterpart of the result given in the Euclidean setting in Appendix A, Theorem A.4.1, of [LTZ]. Further information on the literature is given in Section 2.3 of [LTZ], to which we refer for lack of space. For additional references on the problems here considered, we cite [BGW], [FI], [GL], [LT2], [LTY3], and [LiTa]. Regularity results of the Neumann mixed problem are given in [LT7], [LT10], [LT11], and [T4].

3. A Fundamental Lemma

The following lemma is the key starting point of our analysis. It gives a pointwise estimate which is the counterpart, in the present Riemann metric, of Lemma 3.1 of [LTZ]. Its proof is a parallel development, however, in the Riemann metric, of that given in [LTZ] in the Euclidean metric. To streamline the treatment and write the resulting computations in

a more concise form, we introduce some notation as well as some operators. We recall from Section 1 that D denotes the Levi–Civita connection of (M, g) and that $\mathcal{X}(M)$ is the set of all vector fields on M . Here below, we let $f(t, x), h(t, x) \in C^1(\mathbb{R} \times M)$ and $X \in \mathcal{X}(M)$. Then [GPV]

$$\operatorname{div} X = \sum_{i=1}^N [D_{\partial/\partial x_i} X]_i, \quad \text{with the property} \quad \operatorname{div}(fX) = f \operatorname{div} X + X(f),$$

$$\text{satisfying Green's formula} \quad \int_{\Omega} \operatorname{div} X \, d\Omega = \int_{\Gamma} \langle X, n \rangle \, d\Gamma \quad (3.0)$$

is the divergence operator on the manifold (M, g) . At this point, we set by definition

$$\hat{D}f \equiv (f_t, -Df); \quad \widehat{\operatorname{div}}(h, X) \equiv h_t + \operatorname{div} X; \quad (3.1)$$

$$(h, X)(f) \equiv hf_t + X(f); \quad \mathcal{A}w \equiv \widehat{\operatorname{div}} \hat{D}w, \quad (3.2)$$

where the notation “(function, vector field)” means simply a pair, in that order, of the function and the vector field noted.

Lemma 3.0. *As a consequence of the above definitions, we have the following identities:*

$$(a) \quad \mathcal{A}w = \widehat{\operatorname{div}}(w_t, -Dw) = w_{tt} - \operatorname{div}(Dw) = w_{tt} - \Delta w; \quad (3.3)$$

$$(b) \quad \widehat{\operatorname{div}}[f(h, X)] = \widehat{\operatorname{div}}(fh, fX) = (fh)_t + \operatorname{div}(fX) \quad (3.4)$$

$$\text{(by (3.0))} = f(h_t + \operatorname{div} X) + hf_t + X(f) \quad (3.5)$$

$$= f \widehat{\operatorname{div}}(h, X) + (h, X)(f); \quad (3.6)$$

$$(c) \quad \begin{cases} (\hat{D}f)(h) = (f_t, -Df)(h) = f_t h_t - Df(h), & (3.7a) \\ (\hat{D}h)(h) = h_t^2 - |Dh|^2; & (3.7b) \end{cases}$$

$$(d) \quad \widehat{\operatorname{div}} f(X) = f \widehat{\operatorname{div}} X + X(f). \quad (3.8)$$

Proof. The steps in (3.3) use: definition (3.2) for \mathcal{A} ; definition (3.1) for $\widehat{\operatorname{div}}$; and $\operatorname{div} Dw = \operatorname{div} \nabla w = \Delta w$ for the Laplace–Beltrami operator. Next, the steps from (3.4)–(3.6) use: (3.1) twice for $\widehat{\operatorname{div}}$; (3.0); and (3.2, left). Finally, (3.1) for \hat{D} and (3.2, left) prove (3.7). \square

Below, to facilitate the comparison with [LTZ], we generally adhere to its notation.

Lemma 3.1. *Let*

$$w(t, x) \in C^2(\mathbb{R} \times M); \quad \ell(t, x) \in C^2(\mathbb{R} \times M);$$

$$\psi(t, x) \in C^2 \quad \text{in } t \text{ and } C^1 \text{ in } x \quad (3.9)$$

be three given functions. Set

$$\theta(t, x) = e^{\ell(t, x)}; \quad v(t, x) = \theta(t, x)w(t, x) = e^{\ell(t, x)}w(t, x),$$

$$(t, x) \in Q = (0, T] \times \Omega; \quad (3.10a)$$

$$\theta_t = \ell_t \theta; \quad D\theta = \theta D\ell; \quad v_t = \ell_t v + \theta w_t; \quad Dv = \theta Dw + v D\ell. \quad (3.10b)$$

Then, with reference to $\mathcal{A}w = w_{tt} - \Delta w$ in (3.3), the following pointwise inequality holds true:

$$\begin{aligned} & \theta^2(\mathcal{A}w)^2 + (\psi_t v^2)_t + 2\widehat{\operatorname{div}}\{[(a + \psi)v^2 - v_t^2 + |Dv|^2]\hat{D}\ell + [2\hat{D}\ell(v) - \psi v]\hat{D}v\} \\ & \geq -8v_t D\ell_t(v) + 2[\Delta\ell + \ell_{tt} - \psi]v_t^2 + 2(\mathcal{A}\ell + \psi)|Dv|^2 \\ & \quad + 2v\langle D\psi, Dv \rangle + 4D^2\ell(Dv, Dv) \\ & \quad + \{2\widehat{\operatorname{div}}[(a + \psi)\hat{D}\ell] + \psi_{tt} + 2a\psi\}v^2, \end{aligned} \quad (3.11)$$

where we have set (via (3.7b) with $h = \ell$)

$$a \equiv \hat{D}\ell(\ell) - \mathcal{A}\ell - \psi = \ell_t^2 - |D\ell|^2 - \ell_{tt} + \Delta\ell - \psi, \quad (3.12)$$

and $D^2\ell(\cdot, \cdot)$ is the Hessian of ℓ (a second-order tensor).

Proof. Step 1. First, by (3.1) on \hat{D} and (3.10a, b), we have

$$\hat{D}\theta = (\theta_t, -D\theta) = \theta(\ell_t, -D\ell) = \theta\hat{D}\ell; \quad (3.13)$$

$$\begin{aligned} \hat{D}v &= (v_t, -Dv) = (\ell_t v + \theta w_t, -\theta Dw - vD\ell) \\ &= \theta(w_t, -Dw) + v(\ell_t, -D\ell) = \theta\hat{D}w + v\hat{D}\ell. \end{aligned} \quad (3.14)$$

The goal of this step is to show the following result (counterpart of (3.11) of [LTZ]):

$$\theta^2(\mathcal{A}w)^2 = \{\mathcal{A}v + [\hat{D}\ell(\ell) - \mathcal{A}\ell]v - 2\hat{D}v(\ell)\}^2 \quad (3.15)$$

$$= (I_1 + I_2 + I_3)^2 \quad (3.16)$$

$$\geq 2[I_1 I_2 + I_1 I_3 + I_2 I_3], \quad (3.17)$$

where we have set (as in (3.10) of [LTZ]), via (3.12),

$$\begin{cases} I_1 \equiv \mathcal{A}v + [\hat{D}\ell(\ell) - \mathcal{A}\ell]v = \mathcal{A}v + av = v_{tt} - \Delta v + av; \\ I_2 \equiv -2\hat{D}v(\ell) = -2\hat{D}\ell(v), \quad I_3 \equiv \psi v. \end{cases} \quad (3.18)$$

$$(3.19)$$

Proof of (3.16). To this end, we first prove that

$$\mathcal{A}v = \theta\mathcal{A}w + v\mathcal{A}\ell + 2\hat{D}v(\ell) - v\hat{D}\ell(\ell). \quad (3.20)$$

Indeed, to establish (3.20), we first compute, by (3.2, right), (3.14), (3.8),

$$\mathcal{A}v = \widehat{\operatorname{div}} \hat{D}v = \widehat{\operatorname{div}}(\theta\hat{D}w) + \widehat{\operatorname{div}}(v\hat{D}\ell) \quad (3.21)$$

$$= \theta\widehat{\operatorname{div}} \hat{D}w + v\widehat{\operatorname{div}} \hat{D}\ell + \hat{D}w(\theta) + \hat{D}\ell(v) \quad (3.22)$$

$$= \theta\mathcal{A}w + v\mathcal{A}\ell + \hat{D}w(\theta) + \hat{D}\ell(v). \quad (3.23)$$

Regarding the last two terms in (3.23), we obtain, via (3.7a) applied repeatedly, (3.10a, b), and (3.7b),

$$\hat{D}w(\theta) = w_t\theta_t - Dw(\theta) = \theta(w_t\ell_t - Dw(\ell)) = \theta\hat{D}w(\ell), \quad (3.24)$$

$$\begin{aligned} \hat{D}\ell(v) &= \ell_t v_t - \langle D\ell, Dv \rangle = \ell_t(\ell_t v + \theta w_t) - \langle D\ell, \theta Dw + vD\ell \rangle \\ &= \theta[\ell_t w_t - Dw(\ell)] + v[\ell_t^2 - D\ell(\ell)] \end{aligned} \quad (3.25)$$

$$= \theta\hat{D}w(\ell) + v\hat{D}\ell(\ell). \quad (3.26)$$

Substituting (3.24) and (3.26) into the last two terms of (3.23) yields

$$\mathcal{A}v = \theta \mathcal{A}w + v \mathcal{A}\ell + 2[\theta \hat{D}w(\ell) + v \hat{D}\ell] - v \hat{D}\ell(\ell) \quad (3.27)$$

(by (3.7a))

$$= \theta \mathcal{A}w + v \mathcal{A}\ell + 2\hat{D}v(\ell) - v \hat{D}\ell(\ell), \quad (3.28)$$

and (3.28) proves (3.20), as claimed.

Finally, we use (3.23) in the form $\theta \mathcal{A}w = \mathcal{A}v + v[\hat{D}\ell(\ell) - \mathcal{A}\ell] - 2\hat{D}v(\ell)$, where $\hat{D}v(\ell) = \hat{D}\ell(v)$, to obtain (3.15), (3.16).

Step 2. With reference to (3.18), (3.19), the goal of this step is to show the following result (counterpart of (3.12) of [LTZ]):

$$\begin{aligned} I_1 I_2 &= \widehat{\operatorname{div}}[(av^2 - v_t^2 + |Dv|^2)\hat{D}\ell + 2\hat{D}\ell(v)\hat{D}v] + (\widehat{\operatorname{div}}(a\hat{D}\ell))v^2 \\ &\quad - 4v_t D\ell_t(v) + (\ell_{tt} + \Delta\ell)v_t^2 + (\mathcal{A}\ell)|Dv|^2 + 2D^2\ell(Dv, Dv). \end{aligned} \quad (3.29)$$

Proof of (3.29). By recalling I_1, I_2 from (3.18), (3.19), we compute, via (3.2) on \mathcal{A} ,

$$I_1 I_2 = -2(\mathcal{A}v + av)\hat{D}v(\ell) = -2(\widehat{\operatorname{div}}(\hat{D}v))\hat{D}v(\ell) - 2av\hat{D}v(\ell). \quad (3.30)$$

By (3.8) with $X = \hat{D}v$ and $f = \hat{D}v(\ell)$, we have

$$(\widehat{\operatorname{div}}(\hat{D}v))\hat{D}v(\ell) = \widehat{\operatorname{div}}(\hat{D}v(\ell)\hat{D}v) - \hat{D}v(\hat{D}v(\ell)). \quad (3.31)$$

Then (3.31) used in (3.30) along with $2v\hat{D}v(\ell) = \hat{D}\ell(v^2)$, yields, with $\hat{D}v(\ell) = \hat{D}\ell(v)$,

$$I_1 I_2 = -2\widehat{\operatorname{div}}(\hat{D}v(\ell)\hat{D}v) + 2\hat{D}v(\hat{D}v(\ell)) - a\hat{D}\ell(v^2) \quad (3.32)$$

$$\begin{aligned} &= -\widehat{\operatorname{div}}(2\hat{D}\ell(v)\hat{D}v + av^2\hat{D}\ell) + \widehat{\operatorname{div}}(av^2\hat{D}\ell) \\ &\quad + 2\hat{D}v(\hat{D}v(\ell)) - a\hat{D}\ell(v^2) \end{aligned} \quad (3.33)$$

(by (3.8))

$$\begin{aligned} &= -\widehat{\operatorname{div}}(2\hat{D}\ell(v)\hat{D}v + av^2\hat{D}\ell) + v^2\widehat{\operatorname{div}}(a\hat{D}\ell) \\ &\quad + \cancel{a\hat{D}\ell(v^2)} + 2\hat{D}v(\hat{D}v(\ell)) - \cancel{a\hat{D}\ell(v^2)}, \end{aligned} \quad (3.34)$$

after applying, in the last step, (3.8) with $f = v^2$ and $X = a\hat{D}\ell$.

We next evaluate the term $2\hat{D}v(\hat{D}\ell(v))$ in (3.34). To this end, by (3.7a) with $f = v$, $h = \hat{D}\ell(v) = \hat{D}v(\ell)$, one first has

$$2\hat{D}v(\hat{D}\ell(v)) = 2v_t(\hat{D}\ell(v))_t - 2Dv(\hat{D}\ell(v)) \quad (3.35)$$

$$= 2v_t[\hat{D}\ell_t(v) + \hat{D}\ell(v_t)] - 2Dv(\ell_t v_t - D\ell(v)) \quad (3.36)$$

$$\begin{aligned} &= 2v_t\hat{D}\ell_t(v) + 2v_t\hat{D}\ell(v_t) - 2\ell_t Dv(v_t) \\ &\quad - 2v_t Dv(\ell_t) + 2Dv(D\ell(v)) \end{aligned} \quad (3.37)$$

(by (3.7a))

$$= 2v_t[\ell_{tt}v_t - D\ell_t(v)] + 2v_t[\ell_t v_{tt} - D\ell(v_t)] \quad (3.38)$$

$$- 2v_t D\ell_t(v) - \ell_t(|Dv|^2)_t + 2Dv(D\ell(v)), \quad (3.39)$$

where $D\ell_t(v) = Dv(\ell_t)$. By using a key property of the Levi-Civita connection [Le], we compute the last term of (3.39) as follows:

$$Dv(D\ell(v)) = Dv\langle D\ell, Dv \rangle = X\langle Y, Z \rangle = \langle D_X Y, Z \rangle + \langle Y, D_X Z \rangle \quad (3.40)$$

$$= \langle D_{Dv}(D\ell), Dv \rangle + \langle D\ell, D_{Dv}(Dv) \rangle \quad (3.41)$$

(by (1.4))

$$= D^2\ell(Dv, Dv) + \langle D\ell, D_{Dv}(Dv) \rangle, \quad (3.42)$$

where, in the last step, we have used the definition in (1.4) of the Hessian $D^2\ell$. It is shown in Appendix A that

$$\langle D\ell, D_{Dv}(Dv) \rangle = \frac{1}{2}D\ell(|Dv|^2). \quad (3.43)$$

Using (3.43) in (3.42), we obtain

$$2Dv(D\ell(v)) = 2D^2\ell(Dv, Dv) + D\ell(|Dv|^2). \quad (3.44)$$

Substituting (3.44) for the last term in (3.39) yields

$$\begin{aligned} 2\hat{D}v(\hat{D}\ell(v)) &= 2v_t^2\ell_{tt} - 4v_t D\ell_t(v) + \hat{D}\ell(v_t^2) - \ell_t(|Dv|^2)_t \\ &\quad + D\ell(|Dv|^2) + 2D^2\ell(Dv, Dv) \end{aligned} \quad (3.45)$$

(by (3.7a))

$$= 2v_t^2\ell_{tt} - 4v_t D\ell_t(v) + \hat{D}\ell(v_t^2 - |Dv|^2) + 2D^2\ell(Dv, Dv) \quad (3.46)$$

by invoking, in the last step, (3.7a) with $f = \ell$ and $h = |Dv|^2$ on the fourth and fifth terms in (3.45). Next, recalling (3.8) with $f = v_t^2 - |Dv|^2$ and $X = \hat{D}\ell$, we obtain

$$\begin{aligned} \widehat{\text{div}}[(v_t^2 - |Dv|^2)\hat{D}\ell] &= (v_t^2 - |Dv|^2)\widehat{\text{div}}\hat{D}\ell + \hat{D}\ell(v_t^2 - |Dv|^2) \\ & \text{(by (3.2))} = (v_t^2 - |Dv|^2)\mathcal{A}\ell + \hat{D}\ell(v_t^2 - |Dv|^2) \end{aligned} \quad (3.47)$$

$$\text{(by (3.3))} = v_t^2(\ell_{tt} - \Delta\ell) - |Dv|^2\mathcal{A}\ell + \hat{D}\ell(v_t^2 - |Dv|^2). \quad (3.48)$$

Substituting the last term in (3.48) for the third term in (3.46), we obtain

$$\begin{aligned} 2\hat{D}v(\hat{D}\ell(v)) &= v_t^2(\ell_{tt} + \Delta\ell) + |Dv|^2\mathcal{A}\ell + \widehat{\text{div}}[(v_t^2 - |Dv|^2)\hat{D}\ell] \\ &\quad - 4v_t D\ell_t(v) + 2D^2\ell(Dv, Dv), \end{aligned} \quad (3.49)$$

which was our present objective. Substituting (3.49) into (3.34) then yields

$$\begin{aligned} I_1 I_2 &= -\widehat{\operatorname{div}}[(av^2 - v_t^2 + |Dv|^2)\hat{D}\ell + 2\hat{D}\ell(v)\hat{D}v] + v^2 \widehat{\operatorname{div}}(a\hat{D}\ell) \\ &\quad - 4v_t D\ell_t(v) + (\ell_{tt} + \Delta\ell)v_t^2 + (\mathcal{A}\ell)|Dv|^2 + 2D^2\ell(Dv, Dv), \end{aligned} \quad (3.50)$$

which is precisely (3.29), as desired.

Step 3. The goal of this step is to show that

$$\begin{aligned} I_1 I_3 &= \widehat{\operatorname{div}}(\psi v \hat{D}v) - \frac{1}{2}(v^2 \psi_t)_t + (\frac{1}{2}\psi_{tt} + a\psi)v^2 \\ &\quad + v\langle Dv, D\psi \rangle + \psi(|Dv|^2 - v_t^2). \end{aligned} \quad (3.51)$$

Proof of (3.51). By (3.18), (3.19), we compute, via (3.2) on \mathcal{A} ,

$$I_1 I_3 = (\mathcal{A}v + av)\psi v = (\widehat{\operatorname{div}} \hat{D}v)\psi v + a\psi v^2 \quad (3.52)$$

(by (3.8))

$$= \widehat{\operatorname{div}}(\psi v \hat{D}v) - \hat{D}v(\psi v) + a\psi v^2 \quad (3.53)$$

(by (3.7a))

$$= \widehat{\operatorname{div}}(\psi v \hat{D}v) - [v_t(\psi v)_t - Dv(\psi v)] + a\psi v^2 \quad (3.54)$$

$$= \widehat{\operatorname{div}}(\psi v \hat{D}v) - [v_t(\psi_t v + \psi v_t) - (\psi Dv(v) + v Dv(\psi))] + a\psi v^2 \quad (3.55)$$

$$= \widehat{\operatorname{div}}(\psi v \hat{D}v) - v(v_t \psi_t - Dv(\psi)) - \psi(v_t^2 - Dv(v)) + a\psi v^2. \quad (3.56)$$

Using

$$[-vv_t \psi_t + a\psi v^2] = -\frac{1}{2}(v^2 \psi_t)_t + (\frac{1}{2}\psi_{tt} + a\psi)v^2 \quad \text{and} \quad Dv(v) = |Dv|^2, \quad (3.57)$$

in (3.56), we finally obtain

$$I_1 I_3 = \widehat{\operatorname{div}}(\psi v \hat{D}v) - \frac{1}{2}(v^2 \psi_t)_t + (\frac{1}{2}\psi_{tt} + a\psi)v^2 + v Dv(\psi) - \psi(v_t^2 - |Dv|^2), \quad (3.58)$$

which is precisely (3.51), as desired.

Step 4. By (3.19), we readily have, via (3.8) with $f = v^2$, $X = \psi \hat{D}\ell$,

$$I_2 I_3 = -2\psi v \hat{D}\ell(v) = -\widehat{\operatorname{div}}(\psi v^2 \hat{D}\ell) + v^2 \widehat{\operatorname{div}}(\psi \hat{D}\ell), \quad (3.59)$$

since $\hat{D}\ell(v^2) = 2v\hat{D}\ell(v)$, as noted below (3.31).

Step 5. Inserting (3.29), (3.51), (3.58) in (3.17) yields the desired inequality (3.11). The proof of Lemma 3.1 is complete. \square

4. A Basic Pointwise Inequality

We now make suitable choices of the functions $\ell(t, x)$ and $\psi(x)$ involved in Lemma 3.1, thus obtaining the perfect counterpart of Theorem 4.1 of [LTZ].

Theorem 4.1. *Let*

$$w(t, x) \in C^2(\mathbb{R}_t \times \Omega), \quad d(x) \in C^3(\Omega), \quad \alpha(x) \in C^1(\Omega) \quad (4.1)$$

be three given functions (at this stage, w need not be a solution of (1.1a), d need not be the function provided by the assumptions (A.1) and (A.2), and α need not be given by (1.10)). If $\tau > 0$ is a parameter, we introduce the functions

$$\ell(t, x) = \tau \left[d(x) - c \left(t - \frac{T}{2} \right)^2 \right] = \tau \varphi(t, x); \quad (4.2)$$

$$\psi(x) \equiv \tau \alpha(x); \quad \theta(t, x) = e^{\ell(t, x)} = e^{\tau \varphi(t, x)}, \quad (4.3)$$

where $\varphi(t, x)$ is defined consistently with (1.6a), with constant $0 < c < 1$ selected as in (1.6d). Then, with the above choices, Lemma 3.1 specializes as follows:

$$\begin{aligned} \psi_t &= 0; & \ell_t &= -2\tau c \left(t - \frac{T}{2} \right); & \ell_{tt} &= -2c\tau; & D\ell &= \tau Dd; \\ D\psi &= \tau D\alpha; & D\ell_t &= 0; & \Delta\ell &= \tau \Delta d; & \Delta\ell &= \tau \Delta d, \end{aligned} \quad (4.4)$$

so that the pointwise estimate (3.11) becomes, with $\mathcal{A}w = w_{tt} - \Delta w$ in (3.3), and $v = \theta w$,

$$\begin{aligned} &\theta^2 (\mathcal{A}w)^2 + 2 \widehat{\text{div}} \{ [(a + \psi)v^2 - v_t^2 + |Dv|^2] \hat{D}\ell + [2\hat{D}\ell(v) - \psi v] \hat{D}v \} \\ &\geq 2\tau (\Delta d - 2c - \alpha) v_t^2 + 2\tau \left(\alpha - 2c - \Delta d - \frac{\varepsilon}{2\tau} \right) |Dv|^2 \\ &\quad + 4\tau D^2 d(Dv, Dv) + \theta^2 \tilde{B}w^2, \end{aligned} \quad (4.5)$$

where $\varepsilon > 0$ and

$$\begin{aligned} a &= \tau^2 \left[4c^2 \left(t - \frac{T}{2} \right)^2 - |Dd|^2 \right] + 2c\tau + \tau \Delta d - \psi \\ &= \tau^2 \left[4c^2 \left(t - \frac{T}{2} \right)^2 - |Dd|^2 \right] + \mathcal{O}(\tau); \end{aligned} \quad (4.6)$$

$$\begin{cases} \tilde{B} = 2\beta(x, t)\tau^3 + \mathcal{O}(\tau^2); \\ \beta(x, t) = (2c + \Delta d - \alpha)|Dd|^2 + 2D^2 d(Dd, Dd) \\ \quad - 4c^2(\Delta d + 6c - \alpha) \left(t - \frac{T}{2} \right)^2. \end{cases} \quad (4.7)$$

$$\quad (4.8)$$

Proof. We only need to compute the coefficients of all terms on the right-hand side of inequality (3.11), as well as the term a in (3.12), using the choice of functions made in (4.2), (4.3). This task is a direct computation which yields (4.4) and (4.6) at once. Then (4.3), (4.4), (3.3) give

$$\ell_{tt} + \Delta\ell - \psi = \tau(\Delta d - 2c - \alpha); \quad \mathcal{A}\ell + \psi = \ell_{tt} - \Delta\ell + \psi = \tau(\alpha - 2c - \Delta d). \quad (4.9)$$

We then verify (4.7), (4.8) for the coefficient \tilde{B} of $v^2 = \theta^2 w^2$ in (4.5), which is defined in (3.13). Recalling (3.8), (3.7a), (3.2), (3.3), we find, since $\psi_t \equiv 0$ by (4.4),

$$\begin{aligned} \tilde{B} &\equiv 2\widehat{\operatorname{div}}[(\alpha + \psi)\hat{D}\ell] + \psi_{tt} + 2a\psi = 2(a + \psi)\widehat{\operatorname{div}}\hat{D}\ell \\ &\quad + 2\hat{D}\ell(\alpha + \psi) + 2a\psi \end{aligned} \quad (4.10)$$

$$= \{2(a + \psi)(\ell_{tt} - \Delta\ell)\} + \{2(a_t + \psi_t)\ell_t - 2D\ell(a + \psi)\} + 2a\psi. \quad (4.11)$$

By (4.6) on a , (4.4) on ℓ_{tt} , and $\Delta\ell$, we find

$$\begin{aligned} (a + \psi)(\ell_{tt} - \Delta\ell) &= \left\{ \tau^2 \left[4c^2 \left(t - \frac{T}{2} \right)^2 - |Dd|^2 \right] + \mathcal{O}(\tau) + \psi \right\} \\ &\quad \times (-2c\tau - \tau\Delta d); \end{aligned} \quad (4.12)$$

$$(a_t + \psi_t)\ell_t = \left\{ \tau^2 \left[8c^2 \left(t - \frac{T}{2} \right) \right] + \mathcal{O}(\tau) \right\} \left\{ -2\tau c \left(t - \frac{T}{2} \right) \right\}; \quad (4.13)$$

$$\begin{aligned} D\ell(a) &= \tau Dd(a) = \tau Dd \left\{ \tau^2 4c^2 \left(t - \frac{T}{2} \right)^2 - \tau^2 |Dd|^2 + \mathcal{O}(\tau) \right\} \\ &= \tau^3 Dd(|Dd|^2) + \mathcal{O}(\tau), \end{aligned} \quad (4.14)$$

where by a property of the Levi-Civita connection [Le], and the definition of D^2d in (1.4), one has

$$Dd(|Dd|^2) = Dd\langle Dd, Dd \rangle = 2\langle D_{Dd}(Dd), Dd \rangle = 2D^2d\langle Dd, Dd \rangle. \quad (4.15)$$

Inserting (4.12)–(4.15) along with $D\ell(\psi) = \tau Dd(\psi)$ into (4.11), we obtain

$$\begin{aligned} \tilde{B} &= 2\tau^3(2c + \Delta d - \alpha)|Dd|^2 + 4\tau^3 D^2d\langle Dd, Dd \rangle \\ &\quad + 8\tau^3 c^2(\alpha - 6c - \Delta d) \left(t - \frac{T}{2} \right)^2 + \mathcal{O}(\tau^2), \end{aligned} \quad (4.16)$$

and (4.7), (4.8) are proved, as desired.

We finally verify (4.5). To this end, we insert formulas (4.4), (4.9, left), and (4.15) into the right-hand side of inequality (3.11), use $D\psi = \tau D\alpha$ from (4.4) as well as

$$|v\langle D\psi, Dv \rangle| \leq \varepsilon |Dv|^2 + \frac{\tau}{4\varepsilon} |D\alpha|^2 v^2, \quad (4.17)$$

and obtain (4.5), as desired. \square

Inequality (4.5), as well as formula (4.7), (4.8) for \tilde{B} are exactly the same as those obtained in Theorem 4.1, (4.6), and (4.8) of [LTZ] in the Euclidean case. Thus, we next proceed following [LTZ]. Namely, the pointwise estimate of interest in Corollary 4.2 below is obtained for function $d(x) \in C^3(\bar{\Omega})$, $\alpha \in C^1(\bar{\Omega})$, such that the following three

inequalities hold true:

$$(H.1) \quad \Delta d - 2c - \alpha \geq \rho > 0, \quad \forall x \in \bar{\Omega}; \quad (4.18)$$

$$(H.2) \quad D^2 d(X, X) \geq 2|X|_g^2, \quad \forall x \in \Omega, \quad \forall X \in M_x; \quad (4.19)$$

$$(H.3) \quad [2c + \Delta d - \alpha]|Dd|^2 + 2D^2 d(Dd, Dd) - 4c^2(\Delta d + 6c - \alpha) \left(t - \frac{T}{2}\right)^2 \geq \tilde{\beta} > 0, \quad \forall (t, x) \in Q^*(\sigma^*), \quad (4.20)$$

where $Q^*(\sigma^*)$ is the subset of $[0, T] \times \Omega$ defined in (1.16).

As noted in Section 1, all three of these inequalities hold true, in particular, in the case of our interest under assumptions (A.1) and (A.2) (subject to the translation/rescaling conditions in (1.4), (1.5)) and the choice $\alpha(x) = \Delta d(x) - c - 1$ in (1.10), to obtain properties (p₁) = (1.11), which is (H.1) = (4.18) with $\rho = 1 - c > 0$ —as well as (p'₂) = (1.22)—which is (H.3) = (4.20) with $\tilde{\beta} = 4(1 + 7c)\sigma^* > 0$. This is the content of the next corollary.

Corollary 4.2. *With $0 < c < 1$ chosen in (1.6d), let $d(x) \in C^3(\bar{\Omega})$ and $\alpha(x) \in C^1(\bar{\Omega})$ be two functions such that inequalities (H.1) = (4.18), (H.2) = (4.19), (H.3) = (4.20) hold true. As seen in Section 1, (1.11)–(1.22), this is the case, in particular if $d(x)$ is the strictly convex function provided by assumptions (A.1) and (A.2) and suitably rescaled to achieve the conditions in (1.4), (1.5), and then $\alpha(x)$ is chosen to be (see (1.10))*

$$\alpha(x) \equiv \Delta d(x) - c - 1, \quad (4.21)$$

in which case $\rho = 1 - c > 0$ and $\tilde{\beta} = 4(1 + 7c)\sigma^* > 0$, see (1.22). Let $w \in C^2(\mathbb{R}_t \times \Omega)$. Then, with such choices in (4.2), (4.3) for $\ell(t, x)$ and $\psi(x)$, respectively, Theorem 4.1 specializes to the following results for all $T > T_0$: for all $\varepsilon > 0$ suitably small and all $\tau \geq 1$, the following inequality holds true:

$$\begin{aligned} & \theta^2 (Aw)^2 + 2 \widehat{\text{div}}\{[(a + \psi)v^2 - v_t^2 + |Dv|^2]\hat{D}\ell + [2\hat{D}\ell(v) - \psi v]\hat{D}v\} \\ & \geq \tau \varepsilon \rho \theta^2 [w_t^2 + |Dw|^2] + B\theta^2 w^2, \quad \forall t \in [0, T], \quad \forall x \in \bar{\Omega}, \end{aligned} \quad (4.22)$$

where, recalling \tilde{B} from (4.7), (4.8), we have, for $0 < \varepsilon < \min\{2\rho, 1\}$,

$$\begin{cases} B = \tilde{B} - 2\varepsilon \rho \tau^3 (\varphi_t^2 + |D\varphi|^2) \geq \tilde{B} - 2\varepsilon \rho \tau^3 r, & (4.23a) \end{cases}$$

$$\begin{cases} r = \max_{\bar{Q}} (\varphi_t^2 + |D\varphi|^2), \quad |D\varphi| = |Dd|. & (4.23b) \end{cases}$$

Thus, for $\varepsilon > 0$ suitably small, the constant $\beta = \beta_\varepsilon \equiv \tilde{\beta} - \varepsilon \rho r$ is positive via (4.20), and recalling (4.7), (4.8), (4.20), we obtain, from (4.23),

$$\begin{cases} Bw^2 \geq [2\tau^3 \beta + \mathcal{O}(\tau^2)]w^2, \quad \forall (t, x) \in Q^*(\sigma^*); & (4.24) \end{cases}$$

$$\begin{cases} \beta = \beta_\varepsilon = [\tilde{\beta} - \varepsilon \rho r] > 0, \quad B = \mathcal{O}(\tau^3), \quad \forall (x, t) \in [0, T] \times \Omega, & (4.25) \end{cases}$$

where $Q^*(\sigma^*)$ is the subset of Q defined by (1.16).

Proof. First, preliminarily, we note that by (4.21), taking $0 < \varepsilon < \min\{2\rho, 1\}$, we obtain for $2\tau \geq 1$,

$$\begin{aligned} \alpha - 2c - \Delta d - \frac{\varepsilon}{2\tau} &= -1 - 3c - \frac{\varepsilon}{2\tau} \geq -3 - c, \\ \text{for } 0 < \varepsilon < [2\rho, 1], \quad 2\tau &\geq 1. \end{aligned} \quad (4.26)$$

Moreover, the formulas

$$v_t = \tau\varphi_t v + \theta w_t; \quad Dv = \tau v Dd + \theta Dw, \quad v = \theta w, \quad (4.27)$$

obtained by $v = \theta w$ and (3.10) imply the inequalities (see (4.28) of [LTZ])

$$\begin{aligned} 2v_t^2 &\geq \theta^2 w_t^2 - 2\tau^2 \varphi_t^2 v^2; \quad 2|Dv|^2 \geq \theta^2 |Dw|^2 - 2\tau^2 |Dd|^2 v^2, \\ v^2 &= \theta^2 w^2. \end{aligned} \quad (4.28)$$

After these preliminaries, we return to the right-hand side of (4.5), use here (4.21), (4.26), (1.4), and (4.28) and obtain $0 < \varepsilon < 1$:

$$\begin{aligned} \text{RHS of (4.5)} &\equiv 2\tau(\Delta d - 2c - \alpha)v_t^2 + 2\tau\left(\alpha - 2c - \Delta d - \frac{\varepsilon}{2\tau}\right)|Dv|^2 \\ &\quad + 4\tau D^2 d(Dv, Dv) + \theta^2 \tilde{B}w^2 \end{aligned} \quad (4.29)$$

(by (4.21), (4.26), (1.4))

$$\geq 2\tau\rho v_t^2 + 2\tau\rho|Dv|^2 + \theta^2 \tilde{B}w^2 \quad (4.30)$$

(by (4.28))

$$\begin{aligned} &\geq \tau\rho\{[\theta^2 w_t^2 - 2\tau^2 \varphi_t^2 \theta^2 w^2] + [\theta^2 |Dw|^2 - 2\tau^2 |Dd|^2 \theta^2 w^2]\} \\ &\quad + \tilde{B}\theta^2 w^2 \end{aligned} \quad (4.31)$$

$$\geq \tau\varepsilon\rho\theta^2\{w_t^2 + |Dw|^2\} + [\tilde{B} - 2\varepsilon\rho\tau^3(\varphi_t^2 + |Dd|^2)]\theta^2 w^2, \quad (4.32)$$

and (4.32) establishes (4.22) and (4.23a, left). Then (4.23a, right) follows via (4.23d). Finally, (4.7) and (4.23a) show (4.24), (4.25), except for the positive sign of $\tilde{\beta}$. We finally show that $\tilde{\beta} > 0$ in the set $Q^*(\sigma^*)$ defined in (1.16) and this will yield $\beta > 0$ in $Q^*(\sigma^*)$ for $\varepsilon > 0$ sufficiently small. By definition (4.8), we compute, via (4.21), (1.4a), (1.4b), (1.6a) (same computations as in (1.18)–(1.22), repeated for convenience),

$$\begin{aligned} \tilde{\beta}(x, t) &= [2c + \Delta d - \alpha]|Dd|^2 + 2D^2 d(Dd, Dd) \\ &\quad - 4c^2[\Delta d + 6c - \alpha]\left(t - \frac{T}{2}\right)^2 \end{aligned} \quad (4.33)$$

(by (4.21))

$$\begin{aligned} &= [2c + (1 + c)]|Dd|^2 + 2D^2 d(Dd, Dd) \\ &\quad - 4c^2[(1 + c) + 6c]\left(t - \frac{T}{2}\right)^2 \end{aligned} \quad (4.34)$$

(by (1.4))

$$\geq (5 + 3c)|Dd|^2 - (1 + 7c)4c^2 \left(t - \frac{T}{2}\right)^2 \quad (4.35)$$

(by (1.5))

$$\geq (5 + 3c)\kappa d(x) - (1 + 7c)4c^2 \left(t - \frac{T}{2}\right)^2 \quad (4.36)$$

$$\geq 4c(1 + 7c) \left[d(x) - c^2 \left(t - \frac{T}{2}\right)^2 \right] \quad (4.37)$$

(by (1.6a))

$$\geq 4c(1 + 7c)\varphi^*(x, t) \geq 4c(1 + 7c)\sigma^* > 0, \quad \forall (t, x) \in Q^*(\sigma^*), \quad (4.38)$$

where the set $Q^*(\sigma^*)$ is the subset of $Q = [0, T] \times \Omega$ defined by (1.16). Thus, Corollary 4.2 is proved. \square

We complete this section by providing one Green identity and one notational simplification, to be invoked in the subsequent Section 5.

Claim 1. For any function $f \in C^1(\mathbb{R}_t \times M)$ and any vector field $X \in \mathcal{X}(M)$, we have the following Green formula via (3.1) and Green formula for $\operatorname{div} X$ in (3.0):

$$\int_Q \widehat{\operatorname{div}}(f, X) dQ = \int_\Omega \int_0^T f_t dt d\Omega + \int_0^T \int_\Omega \operatorname{div} X d\Omega dt \quad (4.39)$$

$$= \left[\int_\Omega f d\Omega \right]_0^T + \int_\Sigma \langle X, n \rangle d\Sigma. \quad (4.40)$$

Claim 2. With reference to the second term on the left-hand side of inequality (4.22), set, via (3.1) for ℓ defined in (4.2), so that $D\ell = \tau Dd$,

$$f_1 \equiv (a + \psi)v^2 - v_t^2 + |Dv|^2, \quad (4.41)$$

$$f_2 \equiv 2\hat{D}\ell(v) - \psi v = 2\ell_t v_t - 2\tau Dd(v) - \psi v \quad (4.42a)$$

$$= 2\ell_t v_t - 2\tau \langle Dv, Dd \rangle - \psi v. \quad (4.42b)$$

Then we have, by (4.31), (4.32),

$$\begin{aligned} & 2\widehat{\operatorname{div}}\{[(a + \psi)v^2 - v_t^2 + |Dv|^2]\hat{D}\ell + [2\hat{D}\ell(v) - \psi v]\hat{D}v\} \\ &= 2\widehat{\operatorname{div}}\{f_1 \hat{D}\ell + f_2 \hat{D}v\} = 2\widehat{\operatorname{div}}\{f_1(\ell_t, -D\ell) + f_2(v_t, -Dv)\} \\ &= 2\widehat{\operatorname{div}}(f_1 \ell_t + f_2 v_t, -f_1 D\ell - f_2 Dv). \end{aligned} \quad (4.43)$$

Hence, by (4.33) and (4.30),

$$\begin{aligned}
& 2 \int_Q \widehat{\operatorname{div}}\{[(a + \psi)v^2 - v_t^2 + |Dv|^2]\hat{D}\ell + [2\hat{D}\ell(v) - \psi v]\hat{D}v\} dQ \\
&= 2 \int_Q \widehat{\operatorname{div}}(f_1\ell_t + f_2v_t, -f_1D\ell - f_2Dv) dQ \\
&\text{(by (4.30))} = 2 \left[\int_{\Omega} (f_1\ell_t + f_2v_t) d\Omega \right]_0^T - 2 \int_{\Sigma} \langle f_1D\ell + f_2Dv, n \rangle d\Sigma. \quad (4.44)
\end{aligned}$$

5. Carleman Estimates for Smooth Solutions of (1.1a). First Version

The next key result yields a Carleman-type estimate.

Theorem 5.1. *With $0 < c < 1$ chosen in (1.6a), let $d(x) \in C^3(\bar{\Omega})$, $\alpha(x) \in C^1(\bar{\Omega})$ be two functions such that inequalities (H.1) = (4.18), (H.2) = (4.19), (H.3) = (4.20) hold true. As seen in Section 1, (1.11)–(1.22), this is the case, in particular, if $d(x)$ is the strictly convex function provided by assumptions (A.1) and (A.2), and then $\alpha(x) = \Delta d(x) - c - 1$, as in (4.21) = (1.11). Let $\varphi(x, t)$ be the pseudo-convex function defined by (1.6). Let $w \in C^2(\mathbb{R}_t \times \Omega)$ be a solution of (1.1) (and no B.C.) under the standing assumptions (1.2) for $F(w)$ and (1.3) for f . Then the following one-parameter family of estimates holds true, with $\rho > 0$ (see (4.18)), $\rho = 1 - c$ under (A.1), (A.2)), and $\beta > 0$ (see (4.25)):*

(i) *For all $\tau > 0$ sufficiently large and any $\varepsilon > 0$ small,*

$$\begin{aligned}
& BT|_{\Sigma} + 2 \int_0^T \int_{\Omega} e^{2\tau\varphi} f^2 dQ + C_{1,T} e^{2\tau\sigma} \int_0^T \int_{\Omega} w^2 dQ \\
&\geq [\tau\varepsilon\rho - 2C_T] \int_0^T \int_{\Omega} e^{2\tau\varphi} [w_t^2 + |Dw|^2] dQ \\
&\quad + (2\tau^3\beta + \mathcal{O}(\tau^2) - 2C_T) \int_{Q(\sigma)} e^{2\tau\varphi} w^2 dx dt \\
&\quad - C_T \tau^3 e^{-2\tau\delta} [E(0) + E(T)], \quad (5.1)
\end{aligned}$$

where $Q(\sigma)$ is the subset of $[0, T] \times \Omega$ defined by (1.15), and where we recall from (4.25, left) that β depends on ε .

(ii) *For all $\tau > 0$ sufficiently large and any $\varepsilon > 0$ small,*

$$\begin{aligned}
& BT|_{\Sigma} + 2 \int_0^T \int_{\Omega} e^{\tau\varphi} f^2 dQ \\
&\geq [\tau\varepsilon\rho - 2C_T] e^{2\tau\sigma} \int_{t_0}^{t_1} [w_t^2 + |Dw|^2] d\Omega dt \\
&\quad - C_{1,T} e^{2\tau\sigma} \int_0^T E(t) dt - C_T \tau^3 e^{-2\tau\delta} [E(0) + E(T)]. \quad (5.2)
\end{aligned}$$

Here, $\delta > 0$, $\sigma > 0$, and $\sigma > -\delta$ are the constants in (1.6c), (1.8), while C_T is a positive constant depending on T and d . Moreover, the boundary terms $BT|_{\Sigma}$, $\Sigma = [0, T] \times \Gamma$, are defined by

$$BT|_{\Sigma} = -2 \int_{\Sigma} [f_1 \langle D\ell, n \rangle + f_2 \langle Dv, n \rangle] d\Sigma, \quad (5.3)$$

with f_1 and f_2 defined by (4.31), (4.32), for which an explicit formula in terms of w will be provided in Proposition 5.2 below. Moreover, as in (2.1), we have set

$$E(t) = \int_{\Omega} [w_t^2 + |Dw|^2 + w^2] d\Omega. \quad (5.4)$$

(iii) The above inequality may be extended to all solutions

$$w \in H^{2,2}(Q) \equiv L_2(0, T; H^2(\Omega)) \cap H^2(0, T; L_2(\Omega)).$$

Proof. Step 1. (i) By (3.3), we rewrite (1.1) as

$$w_{tt} - \Delta w = \mathcal{A}w = F(w) + f. \quad (5.5)$$

Next, with $w \in C^2(\mathbb{R}_t \times \Omega)$, we return to inequality (4.22) of Corollary 4.2 and integrate it over $Q = (0, T] \times \Omega$. Recalling Claim 2 above, (4.31), (4.32), (4.34), we obtain

$$\begin{aligned} & \int_Q \theta^2 (\mathcal{A}w)^2 dQ + 2 \int_Q \widehat{\text{div}}\{[(a + \psi)v^2 - v_t^2 + |Dv|^2] \hat{D}\ell + [2\hat{D}\ell(v) - \psi v] \hat{D}v\} dQ \\ &= \int_Q \theta^2 (\mathcal{A}w)^2 dQ + 2 \left[\int_{\Omega} (f_1 \ell_t + f_2 v_t) d\Omega \right]_0^T \\ & \quad - 2 \int_{\Sigma} f_1 \langle D\ell, n \rangle d\Sigma - 2 \int_{\Sigma} f_2 \langle Dv, n \rangle d\Sigma \end{aligned} \quad (5.6)$$

$$\geq \tau \varepsilon \rho \int_Q \theta^2 [w_t^2 + |Dw|^2] dQ + \int_Q \theta^2 Bw^2 dQ. \quad (5.7)$$

We now analyze and estimate the terms in (5.7). First, since $Q \equiv Q^*(\sigma^*) \cup [Q^*(\sigma^*)]^c$, where $[\]^c$ denotes complement in Q , we obtain, via (4.24),

$$\begin{aligned} \int_Q \theta^2 Bw^2 dQ &= \int_{Q^*(\sigma^*)} \theta^2 Bw^2 dx dt + \int_{[Q^*(\sigma^*)]^c} \theta^2 Bw^2 dx dt \\ &\geq [2\tau^3 \beta + \mathcal{O}(\tau^2)] \int_{Q^*(\sigma^*)} \theta^2 w^2 dx dt + \int_{[Q^*(\sigma^*)]^c} \theta^2 Bw^2 dx dt. \end{aligned} \quad (5.8)$$

Next, via (5.5) and (1.2b), we obtain

$$\begin{aligned} \int_Q \theta^2 (\mathcal{A}w)^2 dQ &= \int_Q \theta^2 [F(w) + f]^2 dQ \\ &\leq 2C_T \int_Q \theta^2 [w_t^2 + |Dw|^2 + w^2] dQ + 2 \int_Q \theta^2 f^2 dQ. \end{aligned} \quad (5.9)$$

Inserting (5.8), (5.9) in (5.7), we obtain, via (5.3),

$$\begin{aligned}
& BT|_{\Sigma} + 2 \int_Q \theta^2 f^2 dQ - \int_{[Q^*(\sigma^*)]^c} \theta^2 B w^2 dx dt \\
& \geq [\tau \varepsilon \rho - 2C_T] \int_Q \theta^2 [w_t^2 + |Dw|^2] dQ + [2\tau^3 \beta + \mathcal{O}(\tau^2)] \int_{Q^*(\sigma^*)} \theta^2 w^2 dx dt \\
& \quad - 2C_T \int_Q \theta^2 w^2 dQ - 2 \left[\int_{\Omega} (f_1 \ell_t + f_2 v_t) d\Omega \right]_0^T. \tag{5.10}
\end{aligned}$$

Step 2. By (4.25), we have $B = \mathcal{O}(\tau^3)$ on $(0, T] \times \Omega$. Moreover, we have that $\varphi \leq \sigma^*$ on $[Q^*(\sigma^*)]^c$ by (1.14) and the very definition (1.16). Finally, we have chosen $0 < \sigma^* < \sigma$ below (1.16) and $[Q^*(\sigma^*)]^c$ is a subset of $[0, T] \times \Omega$. Hence, we obtain

$$\begin{aligned}
- \int_{[Q^*(\sigma^*)]^c} e^{2\tau\varphi} B w^2 dx dt &= \mathcal{O}(\tau^3) e^{2\tau\sigma^*} \int_{[Q^*(\sigma^*)]^c} w^2 dx dt \\
&\leq C_T e^{2\tau\sigma} \int_Q w^2 dx dt. \tag{5.11}
\end{aligned}$$

Step 3. Similarly, for $Q = Q(\sigma) \cup [Q(\sigma)]^c$ with $\varphi \leq \sigma$ on $[Q(\sigma)]^c$, we obtain

$$\begin{aligned}
-2C_T \int_Q \theta^2 w^2 dQ &= -2C_T \int_Q e^{2\tau\varphi} w^2 dQ \\
&= -2C_T \int_{Q(\sigma)} e^{2\tau\varphi} w^2 dx dt - 2C_T \int_{[Q(\sigma)]^c} e^{2\tau\varphi} w^2 dx dt \\
&\geq -2C_T \int_{Q(\sigma)} e^{2\tau\varphi} w^2 dx dt - 2C_T e^{2\tau\sigma} \int_{[Q(\sigma)]^c} w^2 dx dt. \tag{5.12}
\end{aligned}$$

Moreover, since $Q^*(\sigma^*) \supset Q(\sigma)$, see (1.17), we have, via (5.12) for the two right terms of (5.10),

$$\begin{aligned}
& [2\tau^3 \beta + \mathcal{O}(\tau^2)] \int_{Q^*(\sigma^*)} e^{2\tau\varphi} w^2 dx dt - 2C_T \int_Q e^{2\tau\varphi} w^2 dQ \\
& \geq [2\tau^3 \beta + \mathcal{O}(\tau^2)] \int_{Q(\sigma)} e^{2\tau\varphi} w^2 dx dt - 2C_T \int_Q e^{2\tau\varphi} w^2 dQ \tag{5.13}
\end{aligned}$$

(by (5.12))

$$\begin{aligned}
& \geq [2\tau^3 \beta + \mathcal{O}(\tau^2) - 2C_T] \int_{Q(\sigma)} e^{2\tau\varphi} w^2 dx dt \\
& \quad - 2C_T e^{2\tau\sigma} \int_{[Q(\sigma)]^c} w^2 dx dt, \tag{5.14}
\end{aligned}$$

where in the last step we have invoked (5.12).

Finally, since $[Q(\sigma)]^c$ is a subset of $[0, T] \times \Omega$, we obtain, from (5.14),

$$\begin{aligned}
& [2\tau^3 \beta + \mathcal{O}(\tau^2)] \int_{Q^*(\sigma^*)} e^{2\tau\varphi} w^2 dx dt - 2C_T \int_Q e^{2\tau\varphi} w^2 dQ \\
& \geq [2\tau^3 \beta + \mathcal{O}(\tau^2) - 2C_T] \int_{Q(\sigma)} e^{2\tau\varphi} w^2 dx dt - 2C_T e^{2\tau\sigma} \int_Q w^2 dx dt. \tag{5.15}
\end{aligned}$$

We return to (5.10): on its left side we use the identity in (5.11), while on its right side we use inequality (5.15). We thus obtain

$$\begin{aligned}
& BT|_{\Sigma} + 2 \int_Q e^{2\tau\varphi} f^2 dQ + \mathcal{O}(\tau^3) e^{2\tau\sigma^*} \int_{[Q^*(\sigma^*)]^c} w^2 dx dt + 2C_T e^{2\tau\sigma} \int_{[Q(\sigma)]^c} w^2 dx dt \\
& \geq [\tau\varepsilon\rho - 2C_T] \int_0^T \int_{\Omega} e^{2\tau\varphi} [w_t^2 + |Dw|^2] dQ \\
& \quad + [2\tau^3\beta + \mathcal{O}(\tau^2) - 2C_T] \int_{Q(\sigma)} e^{2\tau\varphi} w^2 dx dt \\
& \quad - 2 \left[\int_{\Omega} (f_1 \ell_t + f_2 v_t) d\Omega \right]_0^T. \tag{5.16}
\end{aligned}$$

On the other hand, if this time we use the inequality in (5.11), as well as the fact that $[Q(\sigma)]^c$ is a subset of Q , we obtain

$$\begin{aligned}
& BT|_{\Sigma} + 2 \int_Q e^{2\tau\varphi} f^2 dQ + C_{1T} e^{2\tau\sigma} \int_Q w^2 dQ \\
& \geq \text{LHS of (5.16)} \geq \text{RHS of (5.16)}. \tag{5.17}
\end{aligned}$$

Step 4. We show in Step 5 below that

$$|f_1 \ell_t + f_2 v_t| \leq C_T \tau^3 \theta^2 [w_t^2 + |Dw|^2 + w^2]. \tag{5.18}$$

Once (5.17) is proved, it follows recalling (5.4) for $E(t)$ and (1.7) that

$$\begin{aligned}
2 \left| \left[\int_{\Omega} (f_1 \ell_t + f_2 v_t) d\Omega \right]_0^T \right| & \leq C_T \tau^3 \left[\int_{\Omega} e^{2\tau\varphi} [w_t^2 + |Dw|^2 + w^2] d\Omega \right]_0^T \\
& \text{(by (1.7))} \leq C_T \tau^3 e^{-2\tau\delta} [E(T) + E(0)], \tag{5.19}
\end{aligned}$$

where the first inequality is due to (5.17).

Inserting (5.19) into the right side of (5.17) and invoking the right-hand side of (5.16) yields the estimate

$$\begin{aligned}
& BT|_{\Sigma} + 2 \int_Q e^{2\tau\varphi} f^2 dQ + C_{1T} e^{2\tau\sigma} \int_Q w^2 dQ \\
& \geq [\tau\varepsilon\rho - 2C_T] \int_0^T \int_{\Omega} e^{2\tau\varphi} [w_t^2 + |Dw|^2] dQ + [2\tau^3\beta + \mathcal{O}(\tau^2) - 2C_T] \\
& \quad \times \int_{Q(\sigma)} e^{2\tau\varphi} w^2 dx dt - C_T \tau^3 e^{-2\tau\delta} [E(T) + E(0)], \tag{5.20}
\end{aligned}$$

which is precisely the sought-after estimate (5.1).

Step 5. It remains to show estimate (5.18). We compute

$$|f_1 \ell_t + f_2 v_t| \leq |f_1 \ell_t| + |f_2 v_t|, \tag{5.21}$$

where, recalling (4.41) for f_1 and (4.42) for f_2 , we shall show below that

$$|f_1 \ell_t| \leq C_T [\tau^3 v^2 + \tau v_t^2 + \tau |Dv|^2]; \quad (5.22)$$

$$|f_2 v_t| \leq C_T \tau [v^2 + v_t^2 + |Dv|^2]. \quad (5.23)$$

Taking momentarily (5.22) and (5.23) for granted, we next return to the relationships (4.27),

$$v_t = \tau \varphi_t \theta w + \theta w_t; \quad Dv = \tau v Dd + \theta Dw, \quad (5.24)$$

and obtain, with $v = \theta w$,

$$\begin{cases} v_t^2 \leq 2[\theta^2 w_t^2 + \tau^2 \varphi_t^2 \theta^2 w^2] \leq C_T \theta^2 [w_t^2 + \tau^2 w^2]; & (5.25) \\ |Dv|^2 \leq 2[\theta^2 |Dw|^2 + \tau^2 \theta^2 w^2 |Dd|^2] \leq C_T \theta^2 [|Dw|^2 + \tau^2 w^2]. & (5.26) \end{cases}$$

Using $v = \theta w$, (5.25), (5.26) in (5.22) and (5.23), respectively, we readily obtain

$$|f_1 \ell_t| \leq C_T \tau^3 \theta^2 [w^2 + w_t^2 + |Dw|^2]; \quad (5.27)$$

$$|f_2 v_t| \leq C_T \tau^3 \theta^2 [w^2 + w_t^2 + |Dw|^2]. \quad (5.28)$$

Using (5.27), (5.28) into (5.21) yields

$$|f_1 \ell_t + f_2 v_t| \leq C_T \tau^3 \theta^2 [w^2 + w_t^2 + |Dw|^2], \quad (5.29)$$

which is precisely the sought-after inequality (5.15). It remains to show (5.22) and (5.23).

Proof of (5.22). By (4.41) on f_1 , we compute

$$|f_1 \ell_t| = |(a + \psi)v^2 - v_t^2 + |Dv|^2| \ell_t \leq C_T [\tau^2 v^2 + v_t^2 + |Dv|^2] \tau, \quad (5.30)$$

since $a = \mathcal{O}(\tau^2)$ by (4.6), $\psi = \tau \alpha$ by (4.3), and $\ell_t = \mathcal{O}(\tau)$ by (4.4).

Then (5.28) proves (5.22), as desired.

Proof of (5.23). By (4.42) on f_2 and $\psi = \tau \alpha$ via (4.3), we compute

$$|f_2 v_t| = |2\ell_t v_t^2 - 2\tau v_t \langle Dd, Dv \rangle - \tau \alpha v v_t| \leq C_T \tau [v^2 + v_t^2 + |Dv|^2], \quad (5.31)$$

since $\ell_t = \mathcal{O}(\tau)$ by (4.4) and (5.31) proves (5.23), as desired.

The proof of part (i) of Theorem 5.1—that is, estimate (5.1)—is complete.

(ii) We take τ sufficiently large so that, since $\beta > 0$ by assumption, see (4.25), we then have that the term $[2\tau^3 \beta + \mathcal{O}(\tau^2) - 2C_T]$ is positive and we then drop the corresponding lower-order *interior* term involving w^2 in (5.1). Moreover, we invoke the critical property (1.8) for φ on the first integral term on the right side of (5.1). Finally, we majorize $\int_{\Omega} w^2(t) d\Omega$ by $E(t)$, see (5.4), thus obtaining the term $e^{2\tau\sigma} \int_0^T E(t) dt$ on the right-hand side of (5.2). This way, (5.1) readily yields (5.2). \square

To complement Theorem 5.1, we next express the boundary terms $BT|_{\Sigma}$ in (5.3) explicitly in terms of w , not v .

Proposition 5.2. *With reference to Theorem 5.1, in particular (5.3), we have the following explicit expression for the boundary terms $BT|_{\Sigma}$:*

$$BT|_{\Sigma} = -2 \int_{\Sigma} [f_1 \langle D\ell, n \rangle + f_2 \langle Dv, n \rangle] d\Sigma \quad (5.32)$$

$$= -2\tau \int_{\Sigma} (f_1 + \theta w f_2) \langle Dd, n \rangle d\Sigma - 2 \int_{\Sigma} \theta f_2 \langle Dw, n \rangle d\Sigma, \quad (5.33)$$

where $\langle Dd, n \rangle = \partial d / \partial n$ and $\langle Dw, n \rangle = \partial w / \partial n$ and where

$$f_1 + \theta w f_2 = \theta^2 \{ [|Dw|^2 - w_t^2] + [a - \tau^2(|Dd|^2 + \varphi_t^2) + 2\tau \ell_t \varphi_t] w^2 + 2[\ell_t - \tau \varphi_t] w w_t \}; \quad (5.34)$$

$$\theta f_2 = 2\theta^2 \{ \ell_t [w_t + \tau \varphi_t w] - \tau \langle Dw, Dd \rangle - \tau^2 w |Dd|^2 \} - \tau \theta \alpha w. \quad (5.35)$$

Proof. To pass from the definition (5.32) = (5.3) to (5.33), we merely use $D\ell = \tau Dd$ by (4.4) and $Dv = \theta Dw + \tau \theta w Dd$ by (4.27). Next, recalling the definition (4.41), (4.42) for f_1 and f_2 , along with the formulas in (4.27) for v_t and Dv where $\psi = \tau \alpha$, we readily obtain

$$f_1 = (a + \psi)v^2 - v_t^2 + |Dv|^2 = \theta^2 \{ (a + \psi)w^2 - (w_t + \tau \varphi_t w)^2 + |Dw + \tau w Dd|^2 \}; \quad (5.36)$$

$$f_2 = 2\ell_t v_t - 2\tau \langle Dv, Dd \rangle - \tau \alpha v = 2\theta \{ \ell_t [w_t + \tau \varphi_t w] - \tau \langle Dw, Dd \rangle - \tau^2 w |Dd|^2 \} - \tau \theta \alpha w. \quad (5.37)$$

Then (5.36) and (5.37) yield

$$f_1 + \theta w f_2 = \theta^2 \{ (a + \psi)w^2 - (w_t + \tau \varphi_t w)^2 + |Dw + \tau w Dd|^2 \} + 2\theta^2 \{ w \ell_t [w_t + \tau \varphi_t w] - \tau w \langle Dw, Dd \rangle - \tau^2 w^2 |Dd|^2 \} - \tau \theta \alpha w^2, \quad (5.38)$$

from which (5.34) follows using $\psi = \tau \alpha$ and

$$|Dw + \tau Dd|^2 - 2\tau w \langle Dw, Dd \rangle - 2\tau^2 w^2 |Dd|^2 = |Dw|^2 - \tau^2 w^2 |Dd|^2. \quad (5.39)$$

Proposition 5.2 is proved. \square

6. Carleman Estimate for Smooth Solutions of (1.1a). Second Version

A Preliminary Equivalence. Let $u \in H^1(\Omega)$ and let $\tilde{\Gamma}_1$ be any (fixed) portion of the boundary Γ with positive measure. Then the following inequality holds true: there exist positive constants $0 < \tilde{k}_1 < \tilde{k}_2 < \infty$, independent of u , such that

$$\tilde{k}_1 \int_{\Omega} [u^2 + |Du|^2] d\Omega \leq \int_{\Omega} |Du|^2 d\Omega + \int_{\tilde{\Gamma}_1} u^2 d\Gamma \leq \tilde{k}_2 \int_{\Omega} [u^2 + |Du|^2] d\Omega, \quad (6.1)$$

see (6.1) of [LTZ]. Next, for $w \in C^2(\mathbb{R}_t \times \Omega)$, or even $w \in H^{2,2}(Q)$, we introduce

$$\mathcal{E}(t) \equiv \int_{\Omega} [w_t + |Dw|^2] d\Omega + \int_{\Gamma_1} w^2 d\Gamma_1, \quad (6.2)$$

with $\Gamma_1 = \Gamma \setminus \Gamma_0$, where Γ_0 will be the uncontrolled/unobserved portion of the boundary to be defined in Section 8: in (8.2) in the Dirichlet case and in (9.2) in the Neumann case. Recalling $E(t)$ from (2.1) = (5.4), we then see that (6.1) yields the equivalence

$$k_1 E(t) \leq \mathcal{E}(t) \leq k_2 E(t), \quad (6.3)$$

for some positive constants $k_1 > 0, k_2 > 0$.

We can state the main result of the present section.

Theorem 6.1. *With $0 < c < 1$ chosen in (1.6d), let $d(x) \in C^3(\bar{\Omega})$, $\alpha(x) \in C^1(\bar{\Omega})$ be two functions such that inequalities (H.1) = (4.18), (H.2) = (4.19), (H.3) = (4.20) hold true. As seen in Section 1, (1.11)–(1.22), this is the case, in particular, if $d(x)$ is the strictly convex function provided by assumptions (A.1) and (A.2) and suitably scaled to achieve conditions (1.4b), and then $\alpha(x) = \Delta d(x) - c - 1$, as in (1.10) = (4.21). Let $\varphi(t, x)$ be the pseudo-convex function defined by (1.6) and define $\theta(t, x) = \exp(\tau\varphi(t, x))$ as in (4.3). Finally, let $w \in H^{2,2}(Q)$ be a solution of (1.1a) (and no B.C.), subject to the standing assumptions (1.2) on $F(w)$ and (1.3) on f . Then the following one-parameter family of estimates hold true, for all τ sufficiently large, and any $\varepsilon > 0$ small as in (4.22):*

$$\begin{aligned} \overline{BT}|_{\Sigma} + \int_0^T \int_{\Omega} e^{2\tau\varphi} f^2 dQ + \text{const}_{\varphi} \int_0^T \int_{\Omega} f^2 dQ \\ \geq \left\{ \left[\frac{k_1}{2} (\varepsilon\tau\rho - 2C_T)(t_1 - t_2) e^{-c_T T} - \frac{C_{1,T}k_2}{2k_1} T e^{c_T T} \right] e^{2\tau\sigma} - C_T \tau^3 e^{-2\tau\delta} \right\} \\ \times [E(0) + E(T)] \end{aligned} \quad (6.4)$$

$$\geq k_{\varphi} [E(0) + E(T)], \quad \text{for a constant } k_{\varphi} > 0, \quad (6.5)$$

since $\sigma > -\delta$, see (1.8). Here, the boundary terms $\overline{BT}|_{\Sigma}$ are given in terms of the boundary terms $BT|_{\Sigma}$ in (5.3) by

$$\begin{aligned} \overline{BT}|_{\Sigma} = BT|_{\Sigma} + C_T(\tau + 1)e^{2\tau\sigma} \left[\int_0^T \int_{\Gamma} \left| w_t \frac{\partial w}{\partial n} \right| d\Sigma + \int_0^T \int_{\Gamma_1} |ww_t| d\Sigma_1 \right. \\ \left. + \int_{t_0}^{t_1} \int_{\Gamma_1} w^2 d\Gamma_1 dt \right]. \end{aligned} \quad (6.6)$$

Proof. Step 1. We return to estimate (5.2) of Theorem 5.1(ii), add the term

$$(\tau\varepsilon\rho - 2C_T)e^{2\tau\sigma} \int_{t_0}^{t_1} \int_{\Gamma_1} w^2 d\Gamma_1 dt$$

to both sides, recall (6.2) for $\mathcal{E}(t)$, and obtain

$$\begin{aligned} & BT|_{\Sigma} + (\varepsilon\tau\rho - 2C_T)e^{2\tau\sigma} \int_{t_0}^{t_1} \int_{\Gamma_1} w^2 d\Gamma_1 dt + 2 \int_0^T \int_{\Omega} e^{2\tau\varphi} f^2 dQ \\ & \geq (\tau\varepsilon\rho - 2C_T)e^{2\tau\sigma} \int_{t_0}^{t_1} \mathcal{E}(t) dt - C_{1,T}e^{2\tau\sigma} \int_0^T E(t) dt \\ & \quad - C_T\tau^3 e^{-2\tau\delta} [E(0) + E(T)].0 \end{aligned} \quad (6.7)$$

The remaining part of the proof follows closely the proof of Theorem 6.1 of [LTZ] and [LT8] and is included for completeness.

Step 2. In a standard way, multiplying (1.1) by w_t and integrating over Ω yields, after an application of the first Green identity,

$$\begin{aligned} & \frac{1}{2} \frac{\partial}{\partial t} \left(\int_{\Omega} [w_t^2 + |Dw|^2] d\Omega + \int_{\Gamma_1} w^2 d\Gamma_1 \right) \\ & = \int_{\Gamma} \frac{\partial w}{\partial n} w_t d\Gamma + \int_{\Gamma_1} w w_t d\Gamma_1 + \int_{\Omega} [F(w) + f] w_t d\Omega. \end{aligned} \quad (6.8)$$

Notice that on both sides of (6.8) we have added the term $\frac{1}{2}(\partial/\partial t) \int_{\Gamma_1} w^2 d\Gamma_1 = \int_{\Gamma_1} w w_t d\Gamma_1$. Recalling $\mathcal{E}(t)$ in (6.2), we integrate (6.8) over (s, t) and obtain

$$\begin{aligned} \mathcal{E}(t) & = \mathcal{E}(s) + 2 \int_s^t \left[\int_{\Gamma} \frac{\partial w}{\partial n} w_t d\Gamma + \int_{\Gamma_1} w w_t d\Gamma_1 \right] dr \\ & \quad + 2 \int_s^t [F(w) + f] w_t d\Omega dr. \end{aligned} \quad (6.9)$$

We apply Schwarz inequality on $[F(w) + f] w_t$, recall estimate (1.2) for $F(w)$, invoke the left side $E(t) \leq (1/k_1)\mathcal{E}(t)$ of equivalence (6.3), and obtain

$$\mathcal{E}(t) \leq [\mathcal{E}(s) + N(T)] + C_T \int_s^t \mathcal{E}(r) dr; \quad (6.10)$$

$$\mathcal{E}(s) \leq [\mathcal{E}(t) + N(T)] + C_T \int_s^t \mathcal{E}(r) dr \quad (6.11)$$

(C_T includes the constant $1/k_1$ of equivalence), where we have set

$$N(T) = \int_0^T \int_{\Omega} f^2 dQ + 2 \int_0^T \int_{\Gamma} \left| \frac{\partial w}{\partial n} w_t \right| d\Sigma + 2 \int_0^T \int_{\Gamma_1} |w w_t| d\Sigma_1. \quad (6.12)$$

Gronwall's inequality applied on (6.10), (6.11) then yields, for $0 \leq s \leq t \leq T$,

$$\mathcal{E}(t) \leq [\mathcal{E}(s) + N(T)]e^{C_T(t-s)}; \quad \mathcal{E}(s) \leq [\mathcal{E}(t) + N(T)]e^{C_T(t-s)}. \quad (6.13)$$

Set $t = T$ and $s = t$ in the first (left) inequality of (6.13); and set $s = 0$ in the second (right) inequality of (6.13), to obtain

$$\mathcal{E}(T) \leq [\mathcal{E}(t) + N(T)]e^{C_T T}; \quad \mathcal{E}(0) \leq [\mathcal{E}(t) + N(T)]e^{C_T T}. \quad (6.14)$$

Summing up these two inequalities in (6.14) yields, for $0 \leq t \leq T$,

$$\mathcal{E}(t) \geq \frac{\mathcal{E}(T) + \mathcal{E}(0)}{2} e^{-C_\tau T} - N(T) \tag{6.15}$$

$$\geq \frac{k_1}{2} [E(T) + E(0)] e^{-C_\tau T} - N(T), \tag{6.16}$$

after recalling the left side of the equivalence in (6.3). Similarly, summing up the left inequality of (6.13) for $s = 0$ and the right inequality of (6.13) for $s = t$ and $t = T$, and using the equivalence (6.3) yields, for $0 \leq t \leq T$,

$$E(t) \leq \frac{1}{k_1} \left[\frac{k_2(E(0) + E(T))}{2} + N(T) \right] e^{C_\tau T}, \tag{6.17}$$

and hence, by (6.17),

$$\begin{aligned} & -C_{1,T} e^{2\tau\sigma} \int_0^T E(t) dt \\ & \geq -\frac{C_{1,T} k_2}{2k_1} T e^{C_\tau T} e^{2\tau\sigma} [E(0) + E(T)] - \frac{C_{1,T} T}{k_1} e^{C_\tau T} e^{2\tau\sigma} N(T). \end{aligned} \tag{6.18}$$

Step 3. We insert (6.16) into the first integral on the right side of (6.7) and use (6.18) and readily obtain (6.4), (6.6), by invoking (6.12) for $N(T)$.

Finally, we recall the critical relation $\sigma > 0$, $\delta > 0$, $\sigma > -\delta$ from (1.7), (1.18), so that $[\varepsilon \tau e^{2\tau\sigma} - \tau^3 e^{-2\tau\delta}]$ is positive for all τ large enough. Thus, (6.4) yields (6.5). \square

7. Extension of Estimates to Finite Energy Solutions

So far our estimates have been stated and proved for $C^2(\mathbb{R}_t \times \Omega)$ -solutions, hence $H^{2,2}(Q)$ -solutions (Theorems 5.1(iii) and 6.1) of (1.1) with $f \in L_2(Q)$ as in (1.3). In this section we point out that it is possible to extend all our previous estimates to finite energy solutions of (1.1) in the following class:

$$\begin{cases} w \in H^{1,1}(Q) \equiv L_2(0, T; H^1(\Omega)) \cap H^1(0, T; L_2(\Omega)); \\ w_t, \frac{\partial w}{\partial n} \in L_2(0, T; L_2(\Gamma)). \end{cases} \tag{7.1}$$

Thus, the present section is the counterpart of Section 8 in [LTZ], from the Euclidean to the Riemannian setting. As in this reference, we notice that in order to achieve this goal, it suffices to extend the validity of estimate (5.1) of Theorem 5.1(i) from $H^{2,2}(Q)$ -solutions to finite energy solutions defined by the class in (7.1). Here, the main difficulty is the fact that finite energy solutions subject to homogeneous Neumann B.C. do not produce (in dimension ≥ 2) H^1 -traces on the boundary [LT6]. (By contrast, finite energy solutions subject to homogeneous Dirichlet B.C. do produce H^1 -traces: the $\partial w / \partial n$ in $L_2(\Sigma)$ is dominated by the $(H^1(\Omega) \times L_2(\Omega))$ -energy of the initial data, and the $L_1(0, T; L_2(\Omega))$ -norm of f , thanks to sharp results on the regularity of general second-order hyperbolic equations with Dirichlet B.C. given in [LLT] and [LT1], which hold true also in the

Riemannian case. To overcome this difficulty, Section 8 of [LTZ] employed a regularizing procedure inspired by [LaTa]. The same proof of Theorem 8.2 of [LTZ] works in the Riemannian setting and yields:

Theorem 7.1. *Let $f \in L_2(Q)$. Let $w \in H^{2,2}(Q)$ be a solution of (1.1) for which inequality (5.1) of Theorem 5.1(i) holds true, at least as guaranteed by Theorem 5.1(iii). Let u be a solution of (1.1) in the class defined by (7.1). Then estimate (5.1) is satisfied by such a solution u as well. Accordingly, estimate (5.2) of Theorem 5.1(ii), as well as estimate (6.5) of Theorem 6.1, can be extended from $H^{2,2}(Q)$ -solutions of (1.1) to finite energy solutions in the class (7.1).*

8. Continuous Observability Inequality—a Fortiori, a Global Uniqueness Theorem—with Pure Homogeneous Dirichlet B.C. on Σ

In this section we consider the following problem:

$$\begin{cases} w_{tt} = \Delta w + F(w) & \text{in } (0, T] \times \Omega \equiv Q; \\ w|_{\Sigma} \equiv 0 & \text{in } (0, T] \times \Gamma \equiv \Sigma, \end{cases} \quad (8.1a)$$

$$(8.1b)$$

where $F(w)$ satisfies assumption (1.2) and we define

$$\Gamma_{0,D} \equiv \left\{ x \in \Gamma : \langle Dd(x), n(x) \rangle \equiv \frac{\partial d(x)}{\partial n} \leq 0 \right\}; \quad \Gamma_{1,D} = \Gamma \setminus \Gamma_{0,D}. \quad (8.2)$$

The following corollary of Theorem 6.1 provides a continuous observability inequality in the Dirichlet B.C. case; a fortiori, a global uniqueness result.

Theorem 8.1. *Assume hypotheses (A.1) and (A.2): thus there exists a strictly convex function $d(x)$ (rescaled as in (1.5)), which along with the choice $\alpha(x) = \Delta d(x) - c - 1$ in (1.10) = (4.21), where $0 < c < 1$ as in (1.6d), satisfies properties $(p_1) = (1.11)$, $(p_2) = (1.22)$ so that inequalities (H.1) = (4.18), (H.2) = (4.19), (H.3) = (4.20) hold true. Let $\Gamma_{0,D}$, $\Gamma_{1,D}$ be defined by (8.2), and let $T > T_0$ in problem (8.1), where T_0 is defined in (1.6b). Let $w \in H^{1,1}(Q) = L_2(0, T; H^1(\Omega)) \cap H^1(0, T; L_2(\Omega))$ be a solution of problem (8.1a,b). Then:*

(i) *the following inequality holds true:*

$$\int_0^T \int_{\Gamma_{1,D}} \left(\frac{\partial w}{\partial n} \right)^2 d\Gamma_1 dt \geq k_\varphi [E(0) + E(T)]; \quad (8.3)$$

(ii) *a fortiori, if $w \in H^{1,1}(Q)$ is a solution of problem (8.1a,b) satisfying, in addition, $(\partial w / \partial n)|_{\Sigma_1} \equiv 0$, then $w \equiv 0$ in Q ; in fact, in $\mathbb{R}_t \times \Omega$.*

Proof. (i) We first prove the continuous observability inequality (8.3) for a solution $w \in H^{2,2}(Q)$ of problem (8.1a,b). In this case, then, such a solution w satisfies Theorem 6.1, that is estimate (6.5), under present assumptions. Moreover, by (6.6) and (5.32)–(5.35),

we obtain

$$\begin{aligned} w &\equiv 0 \quad \text{on } \Sigma \Rightarrow \overline{BT}|_{\Sigma} = BT|_{\Sigma}, \quad \text{where } [f_1 + \theta w f_2]|_{\Sigma} \equiv \theta^2 |Dw|^2; \\ \theta f_2|_{\Sigma} &= -2\tau \theta^2 \langle Dw, Dd \rangle, \end{aligned} \quad (8.4)$$

so that

$$\begin{aligned} \overline{BT}|_{\Sigma} &= BT|_{\Sigma} \\ &= -2\tau \int_{\Sigma} \theta^2 |Dw|^2 \langle Dw, n \rangle d\Sigma + 4\tau \int_{\Sigma} \theta^2 \langle Dw, Dd \rangle \langle Dw, n \rangle d\Sigma. \end{aligned} \quad (8.5)$$

However, $w \equiv 0$ on Σ implies also $Dw = \langle Dw, n \rangle n$, hence

$$|Dw|^2 = |\langle Dw, n \rangle|^2, \quad \text{as well as } \langle Dw, Dd \rangle = \langle Dw, n \rangle \langle Dd, n \rangle, \quad (8.6)$$

so that (8.5) becomes

$$\begin{aligned} \overline{BT}|_{\Sigma} &= BT|_{\Sigma} = 2\tau \int_{\Sigma} \theta^2 |\langle Dw, n \rangle|^2 \langle Dd, n \rangle d\Sigma \\ &= 2\tau \int_{\Sigma} \theta^2 \left(\frac{\partial w}{\partial n} \right)^2 \langle Dd, n \rangle d\Sigma. \end{aligned} \quad (8.7)$$

Finally, since $\langle Dd, n \rangle \leq 0$ on $\Gamma_{0,D}$ by definition (8.2), estimate (6.5) of Theorem 6.1 becomes, by (8.7),

$$\begin{aligned} 2\tau \int_0^T \int_{\Gamma_{1,D}} \theta^2 \left(\frac{\partial w}{\partial n} \right)^2 \langle Dd, n \rangle d\Gamma_1 dt &\geq 2\tau \int_0^T \int_{\Gamma} \theta^2 \left(\frac{\partial w}{\partial n} \right)^2 \langle Dd, n \rangle d\Gamma dt \\ &= \overline{BT}|_{\Sigma} \geq k_{\varphi} [E(T) + E(0)], \end{aligned} \quad (8.8)$$

and (8.8) shows (8.3), as desired, at least for $H^{2,2}(Q)$ -solutions w . A density argument based on the regularity trace inequality [LT1], [LLT] then extends the validity of (8.3) to $H^{1,1}(Q)$ -solutions w .

(ii) If a solution $w \in H^{1,1}(Q)$ of (8.1a,b) also satisfies $\partial w / \partial n = 0$ on Σ_1 , then (8.3) implies $E(0) = 0$. Since problem (8.1a,b) is well posed—forward and backward—it follows then that $w \equiv 0$ on $\mathbb{R}_t \times \Omega$. \square

9. Global Uniqueness and Continuous Observability with Pure Homogeneous Neumann B.C. A Global Uniqueness Theorem

In this section we consider the following overdetermined problem:

$$\begin{cases} w_t = \Delta w + F(w) & \text{in } (0, T] \times \Omega = Q; & (9.1a) \\ \frac{\partial w}{\partial n} \Big|_{\Sigma} = 0 & \text{in } (0, T] \times \Gamma = \Sigma; & (9.1b) \\ w|_{\Sigma_{1,N}} = 0 & \text{in } (0, T] \times \Gamma_{1,N} = \Sigma_{1,N}, & (9.1c) \end{cases}$$

where $F(w)$ satisfies assumption (1.2), and we define

$$\Gamma_{0,N} \equiv \left\{ x \in \Gamma : \langle Dd(x), n(x) \rangle = \frac{\partial d(x)}{\partial n} = 0 \right\}, \quad \Gamma_{1,N} = \Gamma \setminus \Gamma_{0,N}. \quad (9.2)$$

The following corollary of Theorem 6.1 provides a global uniqueness result.

Theorem 9.1. *Assume hypotheses (A.1) and (A.2); thus, there exists a strictly convex function $d(x)$ (rescaled as in (1.5)), which along with the choice $\alpha(x) = \Delta d(x) - c - 1$ in (1.10) = (4.21), where $0 < c < 1$ as in (1.6d), satisfies properties $(p_1) = (1.11)$, $(p'_2) = (1.22)$, so that inequalities (H.1) = (4.18), (H.2) = (4.19), (H.3) = (4.20) hold true. Let $\Gamma_{0,N}, \Gamma_{1,N}$ be defined by (9.2), and let $T > T_0$ in problem (9.1), where T_0 is defined by (1.6b). Let $w \in H^{1,1}(Q) = L_2(0, T; H^1(\Omega)) \cap H^1(0, T; L_2(\Omega))$ be a solution of problem (9.1a–c). Then $w \equiv 0$ in Q ; in fact, in $\mathbb{R}_t \times \Omega$.*

Proof. We first prove the desired uniqueness for a solution w of problem (9.1a–c) of class $H^{2,2}(Q)$. In this case, such a solution satisfies Theorem 6.1, that is estimate (6.5). Our present goal is to show that, for such a solution, the boundary terms $\overline{BT}|_\Sigma$ given by (6.6) do vanish: $\overline{BT}|_\Sigma = 0$. In fact, by (9.1b), we see via (6.6) that $\overline{BT}|_\Sigma = BT|_\Sigma$, the latter being given by (5.32)–(5.35). Indeed, these same identities show at once that because of (9.1a–c), (9.2), we then have

$$\overline{BT}|_\Sigma = BT|_\Sigma = -2\tau \int_{\Sigma_1} (f_1 + \theta w f_2) \langle Dd, n \rangle d\Sigma_1 \quad (9.3a)$$

$$\text{on } \Sigma_1: \quad (f_1 + \theta w f_2)|_{\Sigma_1} = f_1|_{\Sigma_1} = \theta^2 |Dw|^2. \quad (9.3b)$$

Finally, on Σ_1 , $w = 0$ and $\partial w / \partial n = \langle Dw, n \rangle$ imply $Dw = \langle Dw, n \rangle n + \langle Dw, \bar{\tau} \rangle \bar{\tau} = 0$, where $\bar{\tau}$ is a unit tangential vector. Thus $Dw = 0$ on Σ , used in (9.3a,b), yields $\overline{BT}|_\Sigma = 0$, as desired. However, such an $H^{2,2}(Q)$ -solution satisfies Theorem 6.1, hence inequality (6.5) holds true and yields $E(0) = 0$. Then $w \equiv 0$ on $\mathbb{R}_t \times \Omega$, since problem (9.1a,b) is well-posed forward and backward.

Now an extension of the result to $H^{1,1}(Q)$ -solutions employs, critically, Theorem 7.1. \square

Continuous Observability without Geometrical Conditions on Γ_1 . Non-Explicit Constant. Key to the elimination of geometrical conditions on the (controlled or observed) portion Γ_1 of the boundary Γ , is the following result from Section 7.2 of [LT8].

Lemma 9.2. *Let w be a solution of (1.1) in the class (7.1). Given $\varepsilon > 0$, $\varepsilon_0 > 0$ arbitrary, given $T > 0$, there exists a constant $C = C_{\varepsilon, \varepsilon_0, T} > 0$, such that*

$$\begin{aligned} & \int_{\varepsilon}^{T-\varepsilon} \int_{\tilde{\Gamma}_1} |\nabla_{\tan} w|^2 d\tilde{\Gamma}_1 dt \\ & \leq C \left\{ \int_0^T \int_{\tilde{\Gamma}_1} w_r^2 d\tilde{\Gamma}_1 dt + \int_0^T \int_{\Gamma} \left(\frac{\partial w}{\partial n} \right)^2 d\Gamma dt \right. \\ & \quad \left. + \|w\|_{L_2(0, T; H^{1/2+\varepsilon_0}(\Omega))}^2 + \|f\|_{H^{-1/2+\varepsilon_0}(Q)}^2 \right\}. \end{aligned} \quad (9.4)$$

Using Lemma 9.2, we shall establish as in Section 9 of [LTZ] the sought-after continuous observability inequality

Theorem 9.3. *Assume hypotheses (A.1) and (A.2). Let $w \in H^{1,1}(Q)$ be a solution of problem (9.1a,b). Then the following continuous observability inequality holds true for $T > T_0$, with T_0 given by (1.6b): there exists a constant $C_T > 0$ such that*

$$\int_0^T \int_{\Gamma_1} [w^2 + w_t^2] d\Sigma_1 \geq C_T E(0). \quad (9.5)$$

Proof. As in Section 9 of [LTZ]. (The constant C_T is not explicit in this case, as elimination of the l.o.t. in (9.4) requires a compactness/uniqueness contradiction argument, which employs Theorem 9.1 and loses control of the constant in (9.5), see the proof of Theorem 9.2 of [LTZ].) \square

10. Replacement of Assumption (A.2) = (1.5) by Virtue of Two Vector Fields

Orientation. This section is the counterpart of the presently relevant part of Section 10 of [LTZ] from the Euclidean to the Riemannian setting. The goal is to dispense with the working assumption (A.2) = (1.5). This is done by writing Ω as the overlapping union of two “nice” subdomains Ω_1 and Ω_2 , in correspondence of two functions d_i , $i = 1, 2$, each strictly convex in Ω and thus satisfying (1.4), where now, however, each d_i has *no critical point* on Ω_i , $i = 1, 2$. This way, two radial vector fields are employed.

10.1. Basic Setting Using Two Conservative Vector Fields as in (1.4). Statement of Main Results

Postulated Setting. We divide the original open bounded set Ω into two overlapping subdomains Ω_1 and Ω_2 : $\Omega = \Omega_1 \cup \Omega_2$, $\Omega_1 \cap \Omega_2 \neq \emptyset$, chosen (in infinitely many ways) to fulfill all the conditions, in particular, (a)–(c) below (following the treatment of Section 1).

We assume that there exist two functions $d_i: \bar{\Omega} \Rightarrow \mathbb{R}$, of class C^3 , $i = 1, 2$, which are strictly convex in the Riemann metric, such that the following preliminary conditions (a) and (b) are satisfied:

(a)

$$D^2 d_i(X, X) \equiv \langle D_X(Dd_i), X \rangle_g \geq 2|X|_g^2, \quad \forall x \in \Omega, \quad \forall X \in M_x, \quad (10.1.1)$$

$$\min_{\bar{\Omega}} d_i(x) \geq m > 0, \quad i = 1, 2; \quad (10.1.2)$$

(b)

$$\inf_{\bar{\Omega}_i} |Dd_i| \geq p > 0, \text{ so that we can take } \kappa_i = \inf_{\bar{\Omega}_i} \frac{|Dd_i|}{d_i} > 4, \quad (10.1.3)$$

by rescaling, see Remark 1.2. Next, following Section 1, we define the function for $i = 1, 2$:

$$\varphi_i(x, t) = d_i(x) - c \left(t - \frac{T}{2} \right)^2, \quad x \in \Omega, \quad 0 \leq t \leq T, \quad (10.1.4)$$

with $0 < c < 1$ and T selected as follow: $T > T_{0,i}$, $i = 1, 2$, where

$$T_{0,i}^2 = 4 \max_{\bar{\Omega}} d_i(x), \quad \text{so that} \quad cT^2 > 4 \max_{\bar{\Omega}} d_i(x) + 4\delta, \quad (10.1.5)$$

for some $\delta > 0$ suitably small and kept fixed henceforth. Such functions φ_i thus have the following properties:

$$(i) \quad \varphi_i(x, 0) = \varphi_i(x, T) = d_i(x) - c \frac{T^2}{4} \leq -\delta, \quad \text{uniformly in } \bar{\Omega}; \quad (10.1.6)$$

(ii) there are t_0 and t_1 , with $0 < t_0 < T/2 < t_1 < T$, such that

$$\min_{x \in \bar{\Omega}, t \in [t_0, t_1]} \varphi_i(x, t) \geq \sigma, \quad 0 < \sigma < m. \quad (10.1.7)$$

Next, we define the function

$$\varphi_i^*(x, t) = d_i(x) - c^2 \left(t - \frac{T}{2} \right)^2, \quad x \in \Omega, \quad 0 \leq t \leq T, \quad (10.1.8)$$

so that, since $0 < c < 1$, we have, by (10.1.4) and (10.1.8),

$$\varphi_i^*(x, t) \geq \varphi_i(x, t), \quad x \in \Omega, \quad 0 \leq t \leq T, \quad i = 1, 2. \quad (10.1.9)$$

Furthermore, we define the sets (subsets of $\Omega \times [0, T]$)

$$Q_i(\sigma) \equiv \{(x, t): x \in \Omega, 0 \leq t \leq T, \varphi_i(x, t) \geq \sigma > 0\}; \quad (10.1.10)$$

$$Q_i^*(\sigma^*) \equiv \{(x, t): x \in \Omega, 0 \leq t \leq T, \varphi_i^*(x, t) \geq \sigma^* > 0\}, \quad (10.1.11)$$

$$0 < \sigma^* < \sigma,$$

with constant σ^* selected to satisfy $0 < \sigma^* < \sigma < m$, see (10.1.2) and (10.1.7). By recalling also (10.1.7) and (10.1.9), we obtain

$$\Omega \times [t_0, t_1] \subset Q_i(\sigma) \subset Q_i^*(\sigma^*) \subset \Omega \times [0, T],$$

by (10.1.6), at $t = 0$ and $t = T$:

$$\text{no point of } \Omega \text{ belongs to } Q_i^*(\sigma^*). \quad (10.1.12)$$

(c) We finally require (assume) that

$\Omega_i \supset$ orthogonal projections of $Q_i^*(\sigma^*)$ onto Ω ,

so that for any $(x, t) \in Q_i^*(\sigma^*) \Rightarrow x \in \Omega_i$,

$$\text{and hence } |Dd_i(x)| \geq p > 0 \quad \text{by (10.1.3)}. \quad (10.1.13)$$

Consequences of the Above Setting. With reference to the above setting, we define functions $\alpha_i(x) \in C^1(\bar{\Omega})$, by setting

$$\alpha_i(x) \equiv \Delta d_i(x) - c - 1, \quad x \in \Omega, \quad (10.1.14)$$

so that the following properties $(p_{1;i})$, $(p_{2;i})$ hold true ((10.1.15) is immediate, while (10.1.16) is proved below):

$$(p_{1;i}) \quad \Delta d_i(x) - 2c - \alpha_i(x) \equiv 1 - c > 0, \quad \forall x \in \bar{\Omega}; \quad (10.1.15)$$

$(p_{2;i})$

$$\begin{aligned} \beta_i(x, t) &\equiv [2c + \Delta d_i - \alpha_i] |Dd_i|^2 + 2D^2 d_i(Dd_i, Dd_i) \\ &\quad - 4c^2(\Delta d_i + 6c - \alpha_i) \left(t - \frac{T}{2}\right)^2 \end{aligned} \quad (10.1.16)$$

$$\geq 4(1 + 7c)\varphi_i^*(x, t), \quad \forall x \in \Omega_i, \quad \forall t \in [0, T];$$

$$\text{in particular in } Q_i^*(\sigma^*) \text{ by (10.1.13).} \quad (10.1.17)$$

Proof of (10.1.17). See the proof of (1.12). The steps up to (1.19) continue to hold true, this time by virtue of (10.1.14), (10.1.1), for $i = 1, 2$:

$$\begin{aligned} \beta_i(x, t) &\geq (5 + 3c)|Dd_i|^2 - 4c^2(1 + 7c) \left(t - \frac{T}{2}\right)^2, \\ \forall x, t \in Q &\equiv [0, T] \times \Omega. \end{aligned} \quad (10.1.18)$$

The next step from (1.19) to (1.20) requires, however, assumption (1.5), whose present counterpart is assumption (10.1.3), which is valid, however, only on $\bar{\Omega}_i$. Hence, likewise (10.1.18) implies

$$\beta_i(x, t) \geq (5 + 3c)\kappa_i d_i(x) - 4c^2(1 + 7c) \left(t - \frac{T}{2}\right)^2 \quad (10.1.19)$$

$$\begin{aligned} &\geq 4(1 + 7c) \left[d_i(x) - c^2 \left(t - \frac{T}{2}\right)^2 \right] \\ &\equiv 4(1 + 7c)\varphi_i^*(x, t), \quad \forall x \in \Omega_i, \quad t \in [0, T], \end{aligned} \quad (10.1.20)$$

this time only for $x \in \Omega_i$; and (10.1.20) establishes the first statement in (10.1.17), as desired. To claim, finally, that (10.1.17) holds true, in particular, for $(x, t) \in Q_i^*(\sigma^*)$, we invoke property (10.1.13).

Critical Property to Absorb Lower-Order Terms. As a consequence of (10.1.17), we obtain

$(p'_{2;i})$

$$\begin{aligned} \beta_i(x, t) &\equiv [2c + \Delta d_i - \alpha_i] |Dd_i|^2 + 2D^2 d_i(Dd_i, Dd_i) \\ &\quad - 4c^2(\Delta d_i + 6c - \alpha_i) \left(t - \frac{T}{2}\right)^2 \end{aligned} \quad (10.1.21)$$

(by (10.1.17))

$$\geq 4(1 + 7c)\varphi_i^*(x, t) \geq 4(1 + 7c)\sigma^* \equiv \tilde{\beta} > 0,$$

$$\forall (x, t) \in Q_i^*, \quad (10.1.22)$$

recalling the definition of the set $Q_i^*(\sigma^*)$ in (10.1.11). The term on the left-hand side of (10.1.21) is precisely the term that arises as a coefficient in front of the l.o.t., see (10.2.6)–(10.2.9) below.

The *main result of the present paper* is the following Theorem 10.1.1 which extends all the previous results of Sections 1–9 to the setting of the present Section 10, where assumption (A.2) = (1.5) is removed.

Theorem 10.1.1. *Let the setting of Section 10.1 based on assumptions (a) = (10.1.1), (b) = (10.1.3), and (c) = (10.1.13) be in force. In particular, T is given by $T > T_{0,i}$, see (10.1.5). Then:*

(a) *The Carleman estimates of Theorem 5.1, (5.1), as well as of Theorem 6.1, (6.5), continue to hold true for $H^{1,1}(Q)$ -solutions.*

(b) *The following continuous observability inequality holds true for $H^{1,1}(Q)$ -solutions of the Dirichlet problem (8.1a,b):*

$$\int_0^T \int_{\Gamma_{1,D}} \left(\frac{\partial w}{\partial n} \right)^2 d\Gamma_1 dt \geq k_\varphi [E(T) + E(0)], \quad (10.1.23)$$

with $\Gamma_{1,D} = \Gamma \setminus \Gamma_{0,D}$ and $\Gamma_{0,D}$ defined by (8.2). *A fortiori, this implies a global uniqueness theorem for $H^{1,1}(Q)$ -solutions, if, in addition, $(\partial w / \partial n)|_{\Sigma_1} \equiv 0$; i.e., it follows that, then, $w \equiv 0$.*

(c) *The following continuous observability inequality holds true for $H^{1,1}(Q)$ -solutions of the Neumann problem (9.1a,b):*

$$\int_0^T \int_{\Gamma_{1,N}} [w_t^2 + w^2] d\Gamma_1 dt \geq k_\varphi [E(0) + E(T)], \quad (10.1.24)$$

with $\Gamma_{1,N} = \Gamma \setminus \Gamma_{0,N}$ and $\Gamma_{0,N}$ defined by (9.2). *A class of triples $\{\Omega, \Gamma_{0,N}, \Gamma_{1,N}\}$ satisfying all these conditions is given in Appendix B. A fortiori, this implies a global uniqueness theorem for $H^{1,1}(Q)$ -solutions, if, in addition, $w|_{\Sigma_1} \equiv 0$; i.e., it follows that, then, $w \equiv 0$.*

10.2. Cut-Off Functions $\chi_i(t, x)$ and Corresponding Subproblems for

$$w_i = \chi_i w$$

Cut-Off Functions χ_i . Let $\chi_i(t, x)$ be a smooth cut-off function. At this stage it is not important to specify how it is constructed. Eventually, in the case of purely Neumann B.C. associated with (1.1), $\chi_i(t, x)$ will be the complicated function constructed in Section 10.2 of [LTZ], which has the important feature, among others, of being *only* time-dependent (but not space-dependent) on a *small interior layer of the boundary* Γ . This latter goal is dictated by the Neumann B.C. and would not be necessary when dealing with the Dirichlet B.C. At any rate, we only assume here that such cut-off functions fulfill the requirement

$$|\chi_i| \leq \text{const} \quad \text{and} \quad \chi_i(t, x) \equiv 1 \quad \text{on } Q_i(\sigma), \quad (10.2.1)$$

which is one of the properties satisfied by the cut-off functions in Section 10.2 of [LTZ].

Dynamical System for $w_i \equiv \chi_i w$. Let $w \in C^2(\mathbb{R}_t \times M)$ be a solution of (1.1). We introduce new variables on $[0, T] \times \Omega$:

$$w_i(t, x) \equiv \chi_i(t, x)w(t, x); \quad f_i(t, x) \equiv \chi_i(t, x)f(t, x), \quad i = 1, 2. \quad (10.2.2)$$

(The f_i here should not be confused with the f_i in Sections 4 and 5.) We then see that each term w_i satisfies the following problem:

$$\begin{cases} w_{i,tt} - \Delta w_i = F(w_i) + f_i + \mathcal{K}_i w, & i = 2, 3, \\ w_i(0, \cdot) = w_{i,0}; \quad w_{i,t}(0, \cdot) = w_{i,1}; \end{cases} \quad (10.2.3a)$$

$$\mathcal{K}_i \equiv [D_t^2 - \Delta - F, \chi_i] = \text{commutator active only on } (\text{supp } \chi_i); \quad (10.2.3c)$$

$$w_{i,0} = \chi_i(0, \cdot)w(0, \cdot); \quad w_{i,1} = \chi_{i,t}(0, \cdot)w(0, \cdot) + \chi_i(0, \cdot)w_t(0, \cdot). \quad (10.2.3d)$$

In (10.2.3c), $D_t = d/dt$, while $[\]$ denotes the corresponding commutator of order 1 in time and space. Accordingly, with χ_i smooth, we then obtain, via (10.2.3a) and (1.2b) on F ,

$$\begin{aligned} (\mathcal{A}w_i)^2 &\equiv (w_{i,tt} - \Delta w_i)^2 \\ &\leq C_T \{ |w_{i,t}^2| + |Dw_i|^2 + w_i^2 + f_i^2 + [w_t^2 + |Dw|^2 + w^2](\text{supp } \chi_i) \}, \\ &(t, x) \in [0, T] \times \Omega. \end{aligned} \quad (10.2.4)$$

Preliminary Estimate: Counterpart of Corollary 4.2. As constructed above, each problem w_i in (10.2.3), $i = 1, 2$, satisfies the setting of Section 1. As a result, each problem (10.2.3) satisfies the counterpart of Theorem 4.1/Corollary 4.2, (4.22)–(4.25); in particular, we recall (4.7), (4.8) for \tilde{B} and $\beta(x, t)$. We take this result as our present starting point.

Proposition 10.2.1. *Let $w \in C^2(\mathbb{R}_t \times \Omega)$ be a solution of (1.1). Let the setting of Section 10.1 based on assumptions (a) = (10.1.1), (b) = (10.1.3), and (c) = (10.1.13) be in force. Then each problem (10.2.3), $i = 1, 2$, satisfies the following pointwise inequality for $\varepsilon > 0$ small:*

$$\begin{aligned} \theta_i^2 (\mathcal{A}w_i)^2 + 2 \widehat{\text{div}} \{ [(a_i + \psi_i)v_i^2 - v_{i,t}^2 + |Dv_i|^2] \hat{D}\ell_i \\ + [2\hat{D}\ell_i(v_i) - \psi_i v_i] \hat{D}v_i \} \end{aligned} \quad (10.2.5)$$

$$\geq \tau \varepsilon \rho \theta_i^2 [w_{i,t}^2 + |Dw_i|^2] + B_i \theta_i^2 w_i^2, \quad \forall t \in [0, T], \quad \forall x \in \Omega, \quad (10.2.6)$$

see (4.22), where $\rho = 1 - c > 0$ is a constant, and where, for $i = 1, 2$ and (10.1.22), we have (recalling (4.7), (4.8), (4.20), (4.23)–(4.25))

$$B_i = \tilde{B}_i - 2\varepsilon \rho \tau^3 (\varphi_{i,t}^2 + |D\varphi_i|^2) \geq \tilde{B}_i - 2\varepsilon \rho \tau^3 r \quad (10.2.7)$$

$$\geq [2\tau^3 \beta + \mathcal{O}(\tau^2)], \quad \forall (x, t) \in \text{set } Q_i^*(\sigma^*); \quad (10.2.8)$$

$$\beta \equiv \beta_\varepsilon \equiv \tilde{\beta} - \varepsilon\rho r > 0, \quad r = \max_i \max_{\bar{Q}} (\varphi_{i,t}^2 + |D\varphi_i|^2),$$

$$|D\varphi_i| = |Dd_i|; \tag{10.2.9}$$

$$B_i = \mathcal{O}(\tau^3), \quad \forall (x, t) \in [0, T] \times \Omega, \tag{10.2.10}$$

where the set $Q_i^*(\sigma^*)$ is defined in (10.1.11), $\tilde{\beta} > 0$ is defined in (10.1.22), and $\beta > 0$ is a constant depending on $\varepsilon > 0$. Above, all the quantities with a subscript “i” correspond to the quantities in Sections 3 and 4 without a subscript. Thus, $\theta_i = e^{\tau\varphi_i}$, φ_i as in (10.1.4); $\ell_i = \tau d_i$, $\psi_i = \tau\alpha_i$, with d_i and α_i given by (10.1.2), (10.1.14). Thus, \tilde{B}_i on the left of (10.2.7) is the counterpart of (4.7) via (4.8), while the estimate in (10.2.18) is a consequence of (10.1.22).

10.3. Carleman Estimate for the w_i -Problem

Building up on Proposition 10.2.1, we obtain the counterpart of Theorem 5.1 (Carleman estimate, first version) for the w_i -problems.

Proposition 10.3.1. (i) Let $w \in C^2(\mathbb{R}_t \times \Omega)$ be a solution of (1.1) (and no B.C.). Let the setting of Section 10.1 based on assumptions (a) = (10.1.1), (b) = (10.1.3), and (c) = (10.1.13) be in force. Let w_i , f_i be as in (10.2.2). Let $E(t)$ be defined by (5.4). Then, for $\varepsilon > 0$ small, as in (10.2.9), and for all τ sufficiently large (with $\varepsilon\tau$ large with respect to C_T in (1.2b)), the following family of estimates holds true:

$$\begin{aligned} (BT)_{w_i}|_\Sigma + C_{1,T}e^{2\tau\sigma} \int_0^T E(t) dt + C_{2,T} \int_0^T f_i^2 d\Omega dt \\ \geq [\tau\varepsilon\rho - 2C_T] \int_0^T \int_\Omega e^{2\tau\varphi_i} [w_{i,t}^2 + |Dw_i|^2] d\Omega dt \\ + [2\tau^3\beta + \mathcal{O}(\tau^2) - 2C_T] \int_{Q_i(\sigma)} e^{2\tau\varphi_i} w^2 dx dt \\ - C_T\tau^3 e^{-2\tau\delta} [E(0) + E(T)], \end{aligned} \tag{10.3.1}$$

where $Q_i(\sigma)$ is the subset of $\Omega \times [0, T]$ defined by (10.1.10); moreover, the constants $\sigma > 0$, $\delta > 0$ are defined in (10.1.7), (10.1.5), while the critical constant $\beta > 0$ is defined by (10.2.9) by use of $\tilde{\beta}$ in (10.1.22). We do not specify the boundary terms here (counterpart of (5.3)): they could be made more specific by further specializing the cut-off functions χ_i (see (10.3.3) of [LTZ] for χ_i suitably defined to handle the case of Neumann B.C.).

(ii) The above inequality (10.3.1) can be extended to $H^{2,2}(Q)$ -solutions.

Remark 10.3.1. For τ sufficiently large to make $[2\tau^3\beta + \mathcal{O}(\tau^2) - 2C_T] > 0$ (see $\beta > 0$ in (10.2.9)), we reach one of our goals and drop the integral term involving w^2 , accordingly, from inequality (10.3.1).

Proof. The proof follows closely the one of Proposition 10.3.1 of [LTZ], and thus only a central part dealing directly with the l.o.t. will be given here explicitly.

Step 1. To begin with, the first step is the counterpart of (5.6), (5.7) for the w -problem. We likewise now return to Proposition 10.2.1, estimate (10.2.6) and integrate it over $Q = (0, T] \times \Omega$. We obtain

$$\begin{aligned} & \int_Q \theta_i^2 (Aw_i)^2 dQ + 2 \int_Q \widehat{\operatorname{div}}\{[(a_i + \psi_i)v_i^2 - v_{i,t}^2 + |Dv_i|^2] \widehat{D}\ell_i \\ & \quad + [2\widehat{D}\ell_i(v_i) - \psi_i v_i] \widehat{D}v_i\} dQ \\ & \geq \tau \varepsilon \rho \int_Q \theta_i^2 [w_{i,t}^2 + |Dw_i|^2] dQ + \int_Q \theta_i^2 B_i w_i^2 dQ. \end{aligned} \quad (10.3.2)$$

This is the exact counterpart of the estimate (10.3.4) of [LTZ].

Step 2. Precisely as in Step 2 of Theorem 10.3.1 in [LTZ], one proves that

$$\begin{aligned} \int_Q \theta_i^2 (Aw_i)^2 dQ &= \int_0^T \int_\Omega \theta_i^2 (w_{i,t} - \Delta w_i)^2 d\Omega dt \\ &\leq C_T \left\{ \int_0^T \int_\Omega e^{2\tau\varphi_i} [w_{i,t}^2 + |Dw_i|^2] d\Omega dt \right. \\ & \quad + \int_{Q_i(\sigma)} e^{2\tau\varphi_i} w_i^2 dx dt + e^{2\tau\sigma} \int_0^T E(t) dt \\ & \quad \left. + \int_0^T \int_\Omega f_i^2 d\Omega dt \right\}, \end{aligned} \quad (10.3.3)$$

where the set $Q_i(\sigma)$ is defined by (1.1.10). The proof of (10.3.3) uses both properties of χ_i in (10.2.1) and (10.2.4).

Step 3. We focus on the terms in (10.3.2). (What we do here corresponds to Step 5 in the proof of Proposition 10.3.1 in [LTZ].) We proceed as in the proof of Theorem 5.1, below (5.7). We split $\Omega \times [0, T] = Q_i^*(\sigma^*) \cup [Q_i^*(\sigma^*)]^c$, where $[]^c$ is the complement in $\Omega \times [0, T]$, and $Q_i^*(\sigma^*)$ is the set defined in (10.1.11). Because of property (10.1.13) which was assumed in the selection of Ω_i , we have that for any $(x, t) \in Q_i^*(\sigma^*)$ the space coordinate $x \in \Omega_i$, and hence $|Dd_i(x)| \geq p > 0$ by (10.1.3). As a result of property (10.1.13), we obtained the validity of estimate (10.1.17) on all of $Q_i^*(\sigma^*)$, hence of estimate (10.1.22) on all of $Q_i^*(\sigma^*)$, finally of estimate (10.2.8) on all of $Q_i^*(\sigma^*)$. Employing (10.2.8) we then estimate the last term in (10.3.2):

$$\begin{aligned} \int_0^T \int_\Omega e^{2\tau\varphi_i} B_i w_i^2 d\Omega dt &= \int_{Q_i^*(\sigma^*)} e^{2\tau\varphi_i} B_i w_i^2 dx dt \\ & \quad + \int_{[Q_i^*(\sigma^*)]^c} e^{2\tau\varphi_i} B_i w_i^2 dx dt \end{aligned} \quad (10.3.4)$$

$$\begin{aligned} \text{(by (10.2.8))} &\geq [2\tau^3 \beta + \mathcal{O}(\tau^2)] \int_{Q_i^*(\sigma^*)} e^{2\tau\varphi_i} w_i^2 dx dt \\ & \quad + \int_{[Q_i^*(\sigma^*)]^c} e^{2\tau\varphi_i} B_i w_i^2 dx dt \end{aligned} \quad (10.3.5)$$

((10.3.5) is the counterpart of (5.8)). As to the last term in (10.3.5), we proceed as in Step 2 in the proof of Theorem 5.1: first, by (10.2.10), we have $B_i = \mathcal{O}(\tau^3)$ on all of $\Omega \times [0, T]$; second, we have $\varphi_i \leq \varphi_i^* \leq \sigma^*$ on $[Q_i^*(\sigma^*)]^c$ by (10.1.9) and the very definition of $Q_i^*(\sigma^*)$ in (10.1.11). Finally, since we have chosen $0 < \sigma^* < \sigma$ in (10.1.11), we then obtain

$$-\int_{[Q_i^*(\sigma^*)]^c} e^{2\tau\varphi_i} B_i w_i^2 dx dt \leq \mathcal{O}(\tau^3) \int_{[Q_i^*(\sigma^*)]^c} e^{2\tau\varphi_i} w_i^2 dx dt \quad (10.3.6)$$

$$\begin{aligned} &\leq \mathcal{O}(\tau^3) e^{2\tau\sigma^*} \int_{[Q_i^*(\sigma^*)]^c} w_i^2 dx dt \\ &\leq \mathcal{O}(\tau^3) e^{2\tau\sigma} \int_Q w_i^2 dx dt \leq \mathcal{O}(\tau^3) e^{2\tau\sigma} \\ &\quad \times \int_Q w^2 dQ, \end{aligned} \quad (10.3.7)$$

majorizing w_i^2 by $(\text{const } w^2)$, see (10.2.1), and majorizing $[Q_i^*(\sigma^*)]^c$ by $\Omega \times [0, T]$. Equation (10.3.7) is the counterpart of (5.11), and of (10.3.22) of [LTZ].

Step 4. Next, we substitute (10.3.5) for the last term on the right side of (10.3.2), move the last term of (10.3.5) on $[Q_i^*(\sigma^*)]^c$ to the left side of (10.3.2), and apply for it estimate (10.3.7). Finally, on the first integral term on the left side of (10.3.2), we invoke estimate (10.3.3). We thus obtain

$$\begin{aligned} &\mathcal{O}(\tau^3) e^{2\tau\sigma} \int_Q w^2 dQ + C_T \int_{Q_i(\sigma)} e^{2\tau\varphi_i} w_i^2 dx dt + C_T e^{2\tau\sigma} \int_0^T E(t) dt \\ &\quad + C_T \int_0^T \int_\Omega f_i^2 d\Omega dt \\ &\quad + 2 \int_Q \widehat{\text{div}}\{[(a_i + \psi_i)v_i^2 - v_{i,t}^2 + |Dv_i|^2] \hat{D}\ell_i + [2\hat{D}\ell_i(v_i) - \psi_i v_i] \hat{D}v_i\} dQ \\ &\geq [\tau\varepsilon\rho - C_T] \int_Q e^{2\tau\varphi_i} [w_{i,t}^2 + |Dw_i|^2] dQ + [2\tau^3\beta + \mathcal{O}(\tau^2)] \\ &\quad \times \int_{Q_i^*(\sigma^*)} e^{2\tau\varphi_i} w_i^2 dx dt. \end{aligned} \quad (10.3.8)$$

Regarding the integral terms on $Q_i^*(\sigma^*)$, $Q_i(\sigma)$, we see that since $Q_i^*(\sigma^*) \supset Q_i(\sigma)$ (see (10.1.12)), we have that, for all τ sufficiently large, as in Step 3 in the proof of Theorem 5.1,

$$\begin{aligned} &[2\tau^3\beta + \mathcal{O}(\tau^2)] \int_{Q_i^*(\sigma^*)} e^{2\tau\varphi_i} w_i^2 dx dt - C_T \int_{Q_i(\sigma)} e^{2\tau\varphi_i} w_i^2 dx dt \\ &\geq [2\tau^3\beta + \mathcal{O}(\tau^2) - C_T] \int_{Q_i(\sigma)} e^{2\tau\varphi_i} w^2 dx dt, \end{aligned} \quad (10.3.9)$$

since on $Q_i(\sigma)$ we have $\chi_i \equiv 1$ by (10.2.1), hence $w_i \equiv w$ by (10.2.2). Thus, using (10.3.9) in (10.3.8) and recalling $E(t)$ in (5.4), we readily obtain the desired estimate

(10.3.1), after handling the term $\int_Q \widehat{\operatorname{div}}\{ \} dQ$, which we do not do here, since we have deliberately left undefined the cut-off functions χ_i . For the critical case of Neumann B.C., this is done in Section 10.2 of [LTZ], which admits here a faithful counterpart. \square

Final Remark. The remaining Sections 10.4–10.7 of [LTZ] admit a faithful counterpart from the Euclidean to the Riemannian setting and are not repeated. This therefore leads to Theorem 10.1.1.

Acknowledgments

The work of the first author was in part done while visiting the Scuola Normale Superiore, Pisa, on a sabbatical leave from the University of Virginia. Hospitality and support by the Scuola Normale is acknowledged and greatly appreciated. Particular thanks are extended to Professor Guiseppe Da Prato for many stimulating discussions. The content of this paper was presented at a seminar at SNS.

Appendix A. Proof of (3.43)

In this appendix we prove identity (3.43), that is,

$$\langle D\ell, D_{Dv}(Dv) \rangle = \frac{1}{2} D\ell(|Dv|^2). \quad (\text{A.0})$$

Step 1. Let M be a manifold. We denote by $\{E_1, E_2, \dots, E_n\}$ a *field normal* at $x \in M$. This means that:

- (a) E_1, \dots, E_n are vector fields on the tangent space M_x at x ;
- (b) on M_x , we have that E_1, \dots, E_n are mutually orthogonal; that is, $D_{E_j}E_i = 0$, $\forall i, j$;
- (c) E_1, \dots, E_n form an orthonormal basis on the tangent space M_x . Thus, for any function v , the vector field $Dv = \nabla_g v$ on M_x is given by

$$Dv = \sum_{i=1}^n \langle Dv, E_i \rangle E_i; \quad \langle Dv, E_i \rangle = E_i(v), \quad |Dv|^2 = \sum_{i=1}^n (E_i(v))^2. \quad (\text{A.1})$$

Step 2. We first show that

$$D_{Dv}Dv = \frac{1}{2} \sum_{i=1}^n E_i(|Dv|^2)E_i \quad \text{at } x. \quad (\text{A.2})$$

Indeed, by the axioms of a connection, if Y is any vector field, we obtain, by (A.1),

$$D_{Dv}Y = D_{\sum_{i=1}^n E_i(v)E_i}Y = \sum_{j=1}^n E_j(v)D_{E_j}Y. \quad (\text{A.3})$$

Next, we specialize to $Y = Dv$ given by (A.1) and compute

$$\begin{aligned} D_{E_j}Y &= D_{E_j}Dv = D_{E_j} \left(\sum_{i=1}^n E_i(v)E_i \right) = \sum_{i=1}^n D_{E_j}(E_i(v))E_i \\ &\quad + \sum_{i=1}^n E_i(v)D_{E_j}E_i, \end{aligned} \quad (\text{A.4})$$

where vanishing takes place by orthogonality. Inserting (A.4) in (A.3), with $D_X f = X(f)$, we arrive at

$$\begin{aligned} D_{Dv} Dv &= \sum_{j=1}^n E_j(v) \sum_{i=1}^n E_j E_i(v) E_i = \sum_{i,j=1}^n E_j(v) E_j E_i(v) E_i \\ &= \sum_{i,j=1}^n E_j(v) E_i E_j(v) E_i, \end{aligned} \quad (\text{A.5})$$

changing the order of the indices. Next, we notice that

$$E_i[(E_j(v))^2] = D_{E_i}(E_j(v))^2 = 2E_j(v)E_i E_j(v), \quad (\text{A.6})$$

which substituted into (A.5), yields, recalling (A.1),

$$\begin{aligned} D_{Dv} Dv &= \frac{1}{2} \sum_{i,j=1}^n E_i[(E_j(v))^2] E_i = \frac{1}{2} \sum_{i=1}^n E_i \left[\sum_{j=1}^n (E_j(v))^2 \right] E_i \\ &\quad (\text{by (A.1)}) \\ &= \frac{1}{2} \sum_{i=1}^n E_i(|Dv|^2) E_i, \end{aligned} \quad (\text{A.7})$$

and (A.2) is proved.

Step 3. We finally establish (A.0). Using (A.2), we obtain

$$\begin{aligned} \langle D\ell, D_{Dv} Dv \rangle &= \frac{1}{2} \left\langle D\ell, \sum_{i=1}^n E_i(|Dv|^2) E_i \right\rangle = \frac{1}{2} \sum_{i=1}^n \langle D\ell, E_i(|Dv|^2) E_i \rangle \\ &= \frac{1}{2} \sum_{i=1}^n \langle D\ell, X \rangle = \frac{1}{2} \sum_{i=1}^n X(\ell) \\ &= \frac{1}{2} \sum_{i=1}^n E_i(|Dv|^2) E_i(\ell) = \frac{1}{2} \sum_{i=1}^n E_i(\ell) E_i(|Dv|^2). \end{aligned} \quad (\text{A.8})$$

On the other hand, by (A.1) we can write

$$D\ell = \sum_{i=1}^n E_i(\ell) E_i, \quad \text{hence} \quad D\ell(|Dv|^2) = \sum_{i=1}^n E_i(\ell) E_i(|Dv|^2). \quad (\text{A.9})$$

Combining (A.8) and (A.9) yields (A.0), as desired. \square

Appendix B

In this appendix we provide an attractive class of subsets Ω in a Riemannian manifold M , for which it is possible to claim the existence of a smooth function d defined around the portion Γ_0 of $\partial\Omega$, such that (i) d is strictly convex on Γ_0 , and, moreover, (ii) its normal

derivative vanishes on Γ_0 . The present appendix, in particular Theorem B.1 below and its proof, are all the close counterparts of Appendix 4 of [LTZ], in particular Theorem A.4.1, as generalized from the Euclidean setting to the Riemannian setting.

Let $\{M, g\}$ be an n -dimensional Riemannian manifold, with Levi–Civita connection D . Let $\Omega \subset M$ be an open, connected, compact subset of M , with boundary $\partial\Omega = \overline{\Gamma_0} \cup \Gamma_1$, $\Gamma_0 \cap \Gamma_1 = \emptyset$. The portion Γ_0 of $\partial\Omega$ is defined as follows. Let $\ell: M \rightarrow \mathbb{R}$ be a function of class C^2 . Then we define

$$\Gamma_0 = \{x \in \partial\Omega: \ell(x) = 0\},$$

with the further provision that $D\ell (= \nabla_g \ell) \neq 0$ on Γ_0 . (B.0)

Theorem B.1. *In the above setting, assume that (i)*

$$D^2\ell(X, X)(x) \geq 0, \quad \forall x \in \Gamma_0, \quad \forall X \in M_x$$
 (B.1)

(“convexity of ℓ near Γ_0 ”).

(ii) *There exists a function $d_0: \bar{\Omega} \rightarrow \mathbb{R}$ of class C^2 , such that*

$$(ii_1) \quad D^2d_0(X, X)(x) \geq \rho_0 |X|_g^2, \quad \forall x \in \Gamma_0, \quad \forall X \in M_x,$$
 (B.2)

for some constant $\rho_0 > 0$ (strict convexity of d_0 near Γ_0);

$$(ii_2) \quad \left. \frac{\partial d_0}{\partial n} \right|_{\Gamma_0} = \langle Dd_0, n \rangle_g \leq 0 \quad \text{on } \Gamma_0,$$
 (B.3)

where n is the unit outward normal field to Γ_0 on M which points in the same direction as $\nabla_g \ell = D\ell$.

Then there exists a function $d: \bar{\Omega} \rightarrow \mathbb{R}$ of class C^2 (which is explicitly constructed in a layer (collar) of Γ_0 , the critical set), such that it satisfies the following two conditions:

$$(a) \quad \left. \frac{\partial d}{\partial n} \right|_{\Gamma_0} = \langle Dd, n \rangle_g = 0 \quad \text{on } \Gamma_0;$$
 (B.4)

$$(b) \quad D^2d(X, X)(x) \geq (\rho_0 - \varepsilon) |X|_g^2, \quad \forall x \in \Gamma_0, \quad \forall X \in M_x,$$
 (B.5)

where $\varepsilon > 0$ is arbitrarily small.

Definition of d . The function $d(x)$ is explicitly constructed near Γ_0 , within Ω , as a perturbation of the original function d_0 assumed in (ii) above, as follows:

$$\begin{cases} d(x) \equiv d_0(x) + z(x), & x \text{ near } \Gamma_0 \text{ in } \Omega; \end{cases}$$
 (B.6)

$$\begin{cases} z(x) \equiv -\frac{\partial d_0}{\partial n}(\ell k) + \lambda \ell^2, & k \equiv \frac{1}{|D\ell|_g}, \end{cases}$$
 (B.7)

where λ is a sufficiently large parameter, to be selected below in the proof, while

$$\frac{\partial d_0}{\partial n} \text{ denotes an extension of } \left. \frac{\partial d_0}{\partial n} \right|_{\Gamma_0} \text{ from the set } \Gamma_0$$

(defined by $\ell(x) = 0$) to a layer (collar) of Γ_0 , within Ω , (B.8a)

which is defined by

$$\frac{\partial d_0}{\partial n} \equiv \langle Dd_0, n \rangle_g = \left\langle Dd_0, \frac{D\ell}{|D\ell|_g} \right\rangle_g = \langle Dd_0, kD\ell \rangle_g. \quad (\text{B.8b})$$

$n \equiv D\ell/|D\ell|_g$ (consistently with the statement below (B.3)).

Proof. (a) First we establish property (B.4). To this end, we perform direct computations and obtain

$$\left\{ \frac{\partial(\ell k)}{\partial n} \right\}_{\Gamma_0} = \left[\ell \frac{\partial k}{\partial n} + k \frac{\partial \ell}{\partial n} \right]_{\Gamma_0} = \frac{1}{|D\ell|_g} \langle D\ell, n \rangle_g \equiv 1 \quad \text{on } \Gamma_0; \quad (\text{B.9})$$

$$\left\{ \frac{\partial(\ell^2)}{\partial n} \right\}_{\Gamma_0} = 2\ell \frac{\partial \ell}{\partial n} \Big|_{\Gamma_0} = 0, \quad (\text{B.10})$$

since $\ell = 0$ on Γ_0 , and recalling n from (B.3) and k from (B.7). Returning to $z(x)$ given by (B.7), we take its normal derivative at Γ_0 , and obtain, by virtue of (B.9), (B.10),

$$\frac{\partial z}{\partial n} \Big|_{\Gamma_0} = \left[- \left(\frac{\partial^2 d_0}{\partial n^2} \right) (\ell k) - \frac{\partial d_0}{\partial n} \frac{\partial(\ell k)}{\partial n} + \lambda \frac{\partial(\ell^2)}{\partial n} \right]_{\Gamma_0} = - \frac{\partial d_0}{\partial n} \Big|_{\Gamma_0}. \quad (\text{B.11})$$

Finally, returning to (B.6), and using (B.11), we obtain

$$\frac{\partial d}{\partial n} \Big|_{\Gamma_0} = \left[\frac{\partial d_0}{\partial n} + \frac{\partial z}{\partial n} \right]_{\Gamma_0} = \left[\frac{\partial d_0}{\partial n} - \frac{\partial d_0}{\partial n} \right]_{\Gamma_0} = 0, \quad (\text{B.12})$$

and thus (B.4) is established, as desired.

(b) We now prove property (B.5) for a suitable large λ . For convenience, we set

$$\text{near } \Gamma_0: \quad p \equiv - \frac{\partial d_0}{\partial n} k, \quad \text{so that} \quad z = p\ell + \lambda\ell^2, \quad (\text{B.13})$$

recalling (B.8) and (B.7) for z . Next, from (B.6), we compute preliminarily

$$D^2 d(X, X) = D^2 d_0(X, X) + D^2 z(X, X). \quad (\text{B.14})$$

Step 1.

Lemma B.2. *The following identities hold true:*

(b₁)

$$\begin{aligned} \text{near } \Gamma_0: \quad D^2 z(X, X) &= pD^2\ell(X, X) + \ell D^2 p(X, X) + 2\lambda\ell D^2\ell(X, X) \\ &\quad + 2\lambda|\langle D\ell, X \rangle_g|^2 + 2\langle D\ell, X \rangle_g \langle Dp, X \rangle_g; \end{aligned} \quad (\text{B.15})$$

(b₂)

$$\begin{aligned} \text{on } \Gamma_0: \quad D^2 z(X, X) &= pD^2\ell(X, X) + 2\lambda|\langle D\ell, X \rangle_g|^2 \\ &\quad + 2\langle D\ell, X \rangle_g \langle Dp, X \rangle_g. \end{aligned} \quad (\text{B.16})$$

Proof. Identity (B.16) in (b₂) follows at once from identity (B.15) in (b₁) by setting $\ell = 0$ (see (B.0) of Γ_0).

Proof of (b₁). By definition of the Hessian tensor:

$$D^2z(X, X) = \langle D_X(Dz), X \rangle_g, \quad (\text{B.17})$$

where, recalling $z = p\ell + \lambda\ell^2$ (near Γ_0) by (B.13), we find

$$\begin{cases} Dz = pD\ell + \ell Dp + 2\lambda\ell D\ell, \\ D_X(Dz) = X(p)D\ell + pD_X(D\ell) + X(\ell)Dp + \ell D_X(Dp) \\ \quad + 2\lambda[X(\ell)D\ell + \ell D_X(D\ell)]. \end{cases} \quad (\text{B.18})$$

Substituting (B.19) in (B.17) yields

$$\begin{aligned} D^2z(X, X) &= X(p)\langle D\ell, X \rangle_g + p\langle D_X(D\ell), X \rangle_g \\ &\quad + X(\ell)\langle Dp, X \rangle_g + \ell\langle D_X(Dp), X \rangle_g + 2\lambda X(\ell)\langle D\ell, X \rangle_g \\ &\quad + 2\lambda\ell\langle D_X(D\ell), X \rangle_g. \end{aligned} \quad (\text{B.20})$$

Recalling again the definition of the Hessian tensor for ℓ and p , and recalling likewise that $X(f) = \langle Df, X \rangle_g$, $f = \ell, p$, we recognize at once that (B.20) is precisely (B.15), as desired. \square

Step 2. Let $\varepsilon_0 > 0$ arbitrary. We estimate the last two terms in (B.16):

$$\begin{aligned} &2\lambda|\langle D\ell, X \rangle_g|^2 + 2\langle D\ell, X \rangle_g\langle Dp, X \rangle_g \\ &\geq \left[2\lambda - \frac{1}{\varepsilon_0}\right]|\langle D\ell, X \rangle_g|^2 - \varepsilon_0|\langle Dp, X \rangle_g|^2. \end{aligned} \quad (\text{B.21})$$

Step 3. Regarding the first term in (B.16), we obtain

$$\text{on } \Gamma_0: \quad pD^2\ell(X, X) \geq 0, \quad (\text{B.22})$$

since, as $k > 0$ by (B.7), we have

$$\begin{cases} p = -\frac{\partial d_0}{\partial n}\Big|_{\Gamma_0} & k \geq 0 & \text{on } \Gamma_0, & \text{by assumption (B.3);} \\ D^2\ell(X, X)(\Gamma_0) \geq 0 & & \text{on } \Gamma_0, & \text{by assumption (B.1).} \end{cases} \quad (\text{B.23})$$

Thus, dropping the first term in (B.16) by virtue of (B.22) and invoking (B.21) for the remaining two terms, we obtain, from (B.16),

$$\text{on } \Gamma_0: \quad D^2z(X, X) \geq \left[2\lambda - \frac{1}{\varepsilon_0}\right]|\langle D\ell, X \rangle_g|^2 - \varepsilon_0|\langle Dp, X \rangle_g|^2 \quad (\text{B.24})$$

$$\geq -\varepsilon|X|_g^2, \quad (\text{B.25})$$

where, in the last step, we have chosen

$$2\lambda - \frac{1}{\varepsilon_0} > 0 \quad \text{and} \quad \varepsilon \equiv \varepsilon_0 \max_{\Gamma_0} |Dp|_g^2. \quad (\text{B.26})$$

Finally, returning to (B.14) and invoking (B.25) and assumption (B.2), we obtain

$$\text{on } \Gamma_0: \quad D^2 d(X, X) \geq D^2 d_0(X, X) - \varepsilon |X|_g^2 \quad (\text{B.27})$$

$$\geq [\rho_0 - \varepsilon] |X|_g^2, \quad \forall x \in \Gamma_0, \quad \forall X \in M_x, \quad (\text{B.28})$$

and (B.28) establishes (B.2), as desired. The proof of Theorem B.1 is complete. \square

Remark B.1. The same proof works if we interchange the signs in assumptions (B.1) and (B.3), that is, if we assume instead:

$$(i') \quad D^2 \ell(X, X)(x) \leq 0, \quad \forall x \in \Gamma_0, \quad \forall X \in M_x \quad (\text{B.29})$$

(“concavity of ℓ near Γ_0 ”);

$$(ii') \quad \left. \frac{\partial d_0}{\partial n} \right|_{\Gamma_0} = \langle Dd_0, n \rangle_g \geq 0 \quad \text{on } \Gamma_0. \quad (\text{B.30})$$

Indeed, (B.29) and (B.30) still produce inequality (B.22), recalling (B.23).

References

- [BLR] C. Bardos, G. Lebeau, and J. Rauch, Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary, *SIAM J. Control Optim.* 30 (1992), 1024–1065.
- [BGW] S. Betelu, R. Gulliver, and W. Littman, Boundary controls of PDEs via curvature flows, this issue, DOI: 10.1007/s00245-002-0742-6.
- [D] M. P. DoCarmo, *Riemannian Geometry*, Birkhäuser, Boston, MA, 1992.
- [FI] A. Fursikov and O. Yu. Imanuvilov, *Controllability of Evolution Equations*, Lecture Notes, Vol. 34, Research Institute of Mathematics, Seoul National University, Seoul.
- [GPV] G. Gentili, F. Podestá and E. Vesentini, *Lezioni di Geometria Differenziale*, Boringhieri, Torino, Italy, 1995.
- [GL] R. Gulliver and W. Littman, Chord uniqueness and controllability, the view from the boundary, in *Differential Geometric Methods in the Control of Partial Differential Equations* (R. Gulliver, W. Littman, and R. Triggiani, eds.), Contemporary Mathematics, vol. 268, American Mathematical Society, Providence, RI, 2000, pp. 145–176.
- [He] E. Hebey, *Sobolev Spaces on Riemannian Manifolds*, Lecture Notes in Mathematics, Springer-Verlag, Berlin, 1996.
- [Ho1] L. Hörmander, *Linear Partial Differential Operators*, Springer-Verlag, Berlin, 1969.
- [Ho2] L. Hörmander, *The Analysis of Linear Partial Differential Operators*, Vol. III, Springer-Verlag, Berlin, 1985.
- [Ho3] L. Hörmander, On the uniqueness of the Cauchy problem under partial analyticity assumptions, in *Geometrical Optics and Related Topics* (F. Colombini and N. Lerner, eds.), Birkhäuser, Boston, MA, 1997.
- [I] V. Isakov, *Inverse Problem for Partial Differential Equations*, Springer-Verlag, Berlin, 1998.
- [IY] V. Isakov and M. Yamamoto, Carleman estimate with Neumann boundary condition and its application to the observability inequality and inverse hyperbolic problems, in *Differential Geometric Methods in the Control of Partial Differential Equations* (R. Gulliver, W. Littman, and R. Triggiani, eds.), Contemporary Mathematics, Vol. 268, American Mathematical Society, Providence, RI, 2000, pp. 199–226.

- [KK] M. Kazemi and M. V. Klibanov, Stability estimates for ill-posed Cauchy problems involving hyperbolic equations and inequalities, *Appl. Anal.* 50 (1993), 93–102.
- [LLT] I. Lasiecka, J. L. Lions, and R. Triggiani, Non-homogeneous boundary value problems for second-order hyperbolic operators, *J. Math. Pures Appl.* 65 (1986), 149–192.
- [LaTa] I. Lasiecka and D. Tataru, Uniform boundary stabilization of semilinear wave equations with nonlinear boundary conditions, *J. Differential Integral Equations* 6(3) (1993), 507–533.
- [LT1] I. Lasiecka and R. Triggiani, Regularity of hyperbolic partial differential equations with $L_2(0, T; L_2(\Gamma))$ -Dirichlet boundary terms, *Appl. Math. Optim.* 10 (1983), 275–286.
- [LT2] I. Lasiecka and R. Triggiani, Uniform exponential energy decay of wave equations in a bounded region with $L_2(0, \infty; L_2(\Gamma))$ -feedback control in the Dirichlet boundary conditions, *J. Differential Equations* 66 (1987), 340–390.
- [LT3] I. Lasiecka and R. Triggiani, Exact controllability of the wave equation with Neumann boundary control, *Appl. Math. Optim.* 19 (1989), 243–290. Preliminary version LNCS, Vol. 100, Springer-Verlag, Berlin, 1987, pp. 316–371.
- [LT4] I. Lasiecka and R. Triggiani, A lifting theorem for the time regularity of solutions to abstract equations with unbounded operators and applications to hyperbolic equations, *Proc. Amer. Math. Soc.* 10 (1988), 745–755.
- [LT5] I. Lasiecka and R. Triggiani, Carleman estimates and exact boundary controllability for a system of coupled, nonconservative second order hyperbolic equations, in *Partial Differential Equations Methods in Control and Shape Analysis, Lecture Notes in Pure and Applied Mathematics*, Vol. 188, Marcel Dekker, New York, 1997, pp. 215–243.
- [LT6] I. Lasiecka and R. Triggiani, Sharp regularity for mixed second-order hyperbolic equations of Neumann type, part I: the L_2 -boundary case, *Ann. Mat. Pura Appl.* (4), 157 (1990), 285–367.
- [LT7] I. Lasiecka and R. Triggiani, Regularity theory of hyperbolic equations with non-homogeneous Neumann boundary conditions. Part II: general boundary data, *J. Differential Equations* 94 (1991), 112–164.
- [LT8] I. Lasiecka and R. Triggiani, Uniform stabilization of the wave equation with Dirichlet or Neumann feedback control without geometrical conditions, *Appl. Math. Optim.* 25 (1992), 189–244.
- [LT9] I. Lasiecka and R. Triggiani, Exact controllability of semilinear abstract systems with application to waves and plates, *Appl. Math. Optim.* 23 (1991), 109–154.
- [LT10] I. Lasiecka and R. Triggiani, Recent Advances in Regularity Theory of Second-Order Hyperbolic Mixed Problems and Applications, *Dynamics Reported, Expositions in Dynamical Systems*, new series, Vol. 3, Springer-Verlag, Berlin, 1994.
- [LT11] I. Lasiecka and R. Triggiani, Control Theory for Partial Differential Equations: Continuous and Approximation Theories, *Encyclopedia of Mathematics and its Applications*, Vols. 72 and 73, Cambridge University Press, Cambridge, 2000.
- [LTY1] I. Lasiecka, R. Triggiani, and P. F. Yao, Exact controllability for second-order hyperbolic equations with variable coefficient principal part and first-order terms, *Nonlinear Anal. TMA* 30 (1997), 111–122.
- [LTY2] I. Lasiecka, R. Triggiani, and P. F. Yao, Inverse/observability estimates for second order hyperbolic equations with variable coefficients, *J. Math. Anal. Appl.* 235 (1999), 13–57.
- [LTY3] I. Lasiecka, R. Triggiani, and P. F. Yao, An observability estimate in $L_2(\Omega) \times H^{-1}(\Omega)$ for second order hyperbolic equations with variable coefficients, in *Control of Distributed Parameter and Stochastic Systems* (S. Chen, X. Li, J. Yong, and X. Y. Zhou, eds.), Kluwer, Dordrecht, 1999, pp. 71–78.
- [LTZ] I. Lasiecka, R. Triggiani, and X. Zhang, Nonconservative wave equations with unobserved Neumann B.C.: global uniqueness and observability in one shot, in *Differential Geometric Methods in the Control of Partial Differential Equations* (R. Gulliver, W. Littman, and R. Triggiani, eds.), *Contemporary Mathematics*, Vol. 263, American Mathematical Society, Providence, RI, 2000, pp. 227–326.
- [LRS] M. M. Lavrentev, V. G. Romanov, and S. P. Shishataskii, *Ill-Posed Problems of Mathematical Physics and Analysis*, Vol. 64, American Mathematical Society, Providence, RI, 1986.
- [Le] J. Lee, *Riemannian Manifolds*, Springer-Verlag, New York, 1997.
- [Lio1] J. L. Lions, Exact controllability, stabilization and perturbations for distributed systems, *SIAM Rev.* 30 (1988), 1–68.
- [Lio2] J. L. Lions, *Contrôlabilité Exacte, Perturbations et Stabilisation de Systèmes Distribués*, Vol. 1, Masson, Paris, 1988.

- [Lit1] W. Littman, Near optimal time boundary controllability for a class of hyperbolic equations, in *Control Problems for Systems Described by Partial Differential Equations and Applications* (I. Lasiecka and R. Triggiani, eds.), *Lecture Notes in Control and Information*, Vol. 97, Springer-Verlag, Berlin, 1987, pp. 307–312.
- [Lit2] W. Littman, Boundary controllability for polyhedral domains, in *Boundary Control and Boundary Variation* (J. P. Zolesio, ed.), *Lecture Notes in Control and Information*, Vol. 178, Springer-Verlag, Berlin, 1992, pp. 272–284.
- [Lit3] W. Littman, Remarks on global uniqueness for partial differential equations, in *Differential Geometric Methods in the Control of Partial Differential Equations* (R. Gulliver, W. Littman, and R. Triggiani, eds.), *Contemporary Mathematics*, Vol. 268, American Mathematical Society, Providence, RI, 2000, pp. 363–372.
- [LiTa] W. Littman and S. Taylor, Smoothing evolution equations and boundary control theory, *J. Analyse Math.* 59 (1992), 117–131.
- [T1] D. Tataru, Boundary controllability for conservative PDEs, *Appl. Math. Optim.* 31 (1995), 257–295.
- [T2] D. Tataru, Unique continuation for solutions to PDEs: between Hörmander theorem and Holmgren theorem, *Comm. Partial Differential Equations* 20(5&6) (1995), 855–884.
- [T3] D. Tataru, Carleman estimates and unique continuation for solutions to boundary value problems, *J. Math. Pures Appl.* 75 (1996), 367–408.
- [T4] D. Tataru, On the regularity of boundary traces for the wave equation, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* 26 (1998), 185–206.
- [T5] D. Tataru, Unique continuation for operators with partially analytic coefficients, *J. Math. Pures Appl.* 9 (1999), 505–521.
- [Y] P. F. Yao, On the observability inequalities for exact controllability of wave equations with variable coefficients, *SIAM J. Control Optim.* 37 (1999), 1568–1599.
- [Z1] X. Zhang, Rapid exact controllability of the semi-linear wave equation, *Chinese Ann. Math. Ser. B* 20(3) (1999), 377–384.
- [Z2] X. Zhang, Explicit observability estimate for the wave equation with potential and its application, *Proc. Roy. Soc. London Ser. A*, 456 (2000), 1101–1115.
- [Z3] X. Zhang, Explicit observability inequalities for the wave equation with lower order terms by means of Carleman inequalities, *SIAM J. Control Optim.* 39 (2001), 812–834.

Online publication 29 November 2002.