



Psychrophilic lifestyles: mechanisms of adaptation and biotechnological tools

Tony Collins¹ · Rosa Margesin²

Received: 23 November 2018 / Revised: 21 January 2019 / Accepted: 22 January 2019 / Published online: 7 February 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Cold-adapted microorganisms inhabiting permanently low-temperature environments were initially just a biological curiosity but have emerged as rich sources of numerous valuable tools for application in a broad spectrum of innovative technologies. To overcome the multiple challenges inherent to life in their cold habitats, these microorganisms have developed a diverse array of highly sophisticated synergistic adaptations at all levels within their cells: from cell envelope and enzyme adaptation, to cryoprotectant and chaperone production, and novel metabolic capabilities. Basic research has provided valuable insights into how these microorganisms can thrive in their challenging habitat conditions and into the mechanisms of action of the various adaptive features employed, and such insights have served as a foundation for the knowledge-based development of numerous novel biotechnological tools. In this review, we describe the current knowledge of the adaptation strategies of cold-adapted microorganisms and the biotechnological perspectives and commercial tools emerging from this knowledge. Adaptive features and, where possible, applications, in relation to membrane fatty acids, membrane pigments, the cell wall peptidoglycan layer, the lipopolysaccharide component of the outer cell membrane, compatible solutes, antifreeze and ice-nucleating proteins, extracellular polymeric substances, biosurfactants, chaperones, storage materials such as polyhydroxyalkanoates and cyanophycins and metabolic adjustments are presented and discussed.

Keywords Psychrophiles · Cell envelope · Cryoprotection · Enzymes · Chaperones · Metabolic adjustments

Introduction

Habitats of permanently low temperature dominate the Earth's biosphere and have been successfully colonised by a wide variety of organisms collectively termed psychrophiles or cold-adapted organisms. Microorganisms prevail in these cold habitats, especially bacteria, archaea, yeasts, cyanobacteria and protists, but microalgae and viruses are also common, and the ability of psychrophilic microorganisms to thrive in such environments reveals an adaptation to their habitat (Margesin and Collins 2019). It demonstrates a capacity to

surmount the constraints inherent to life at low temperatures, which can only be achieved via a complex range of structural and functional adaptations of all cellular components, from the level of single molecules up to whole cells and even complete ecosystems (D'Amico et al. 2006; De Maayer et al. 2014; Morgan-Kiss et al. 2006; Siddiqui et al. 2013).

Since the first reports of cold dwelling organisms (Forster 1887), various comparative physiology, microbiology, biochemistry, biophysics and molecular-based approaches have been used in identifying the physiological adaptations, biogeographical distribution, diversity and ecological roles of cold adapted organisms. More recently, various new emerging technologies, namely various 'omics' approaches including genomics, transcriptomics, proteomics and metagenomics, have broadened our understanding of these, enabling identification of novel adaptation mechanisms and biotechnological tools, detection of key functions and acquirement of a more global view of the structures and roles of microbial communities in cold ecosystems (Barauna et al. 2017; Bowman 2017; Koh et al. 2017; Raymond-Bouchard et al. 2018; Singh et al. 2014; Tribelli and Lopez 2018).

✉ Tony Collins
tcollins@bio.uminho.pt

¹ Centre of Molecular and Environmental Biology (CBMA), Department of Biology, University of Minho, 4710-057 Braga, Portugal

² Institute of Microbiology, University of Innsbruck, 6020 Innsbruck, Austria

Life in the cold is characterised by a multitude of stresses in addition to low temperatures. Indeed, besides a reduced thermal energy, low temperatures also provoke further physico-chemical constraints such as an increase in solvent viscosity and solubility of gasses, namely an increased solubility of oxygen and reactive oxygen species (ROS), a decrease in the solubility of solutes and nutrients, reduced diffusion, increased osmotic stress, desiccation and ice formation. Furthermore, many cold ecosystems are often also characterised by fluctuating environmental conditions and/or multiple additional ecological limiting factors, including low nutrient levels, high salinity, oxidative stress, freeze–thaw cycles and low water activity. In the deep sea and sub-glacial environments, for example, microbial populations are additionally subjected to high pressure stress. Extremes of light exposure are also common in many cold ecosystems, from high light and UV irradiation in high altitude cold environments to low light exposure in ice-covered lakes and deep within ice layers and permafrost. Thus, life in the cold biosphere requires a multitude of synergistic adaptations not only to respond to the low-temperature challenge but also to the multitude of other interacting stresses imposed by the particular environmental conditions.

Cold-adapted microorganisms have responded to the low-temperature challenge via the development of a cold adaptation toolkit constituted by a number of elegant physiological and structural adaptations, many of which are only beginning to be understood. Importantly, many of these tools serve overlapping functions and may be used to respond to the various different challenges or combinations of challenges encountered in a specific cold habitat. In fact, a common problem encountered in characterising cold-adapted microorganisms is in unravelling the different interacting parameters and deciphering the precise function of a specific trait, whether it is a specific response to low temperatures or to another (or other) environmental stressor(s) common in the particular habitat. Importantly also, microorganisms do not always make use of all tools in their ‘cold adaptation’ toolbox, and in fact, each specific organism will use its own strategy, or combination of strategies, depending on its own specific requirements and on the environmental parameters and microbial community structure.

In the present review, we will discuss various adaptation strategies used by cold-adapted microorganisms to enable life in the harsh environmental conditions of their habitat and show how an understanding of these different strategies is leading to the development of various novel tools of commercial interest. See Table 1 for an overview of cold-adaptation tools with commercial potential. In this review, the various stresses to which microorganisms in cold habitats are exposed will be presented, the adaptation strategies developed by psychophiles to cope with the

challenges and their underlying mechanism of action discussed, and examples of the biotechnological tools being developed from these will be given.

Cell envelope

The cell envelope and its various components serve the critical functions of providing shape, support and protection as well as regulating the movement of substances into and out of cells. It protects cells from the surroundings and against turgor pressure; acts as a semipermeable membrane functioning in nutrient uptake, product export and solute transport; and participates in cell division, sensing, signalling and adhesion. Low temperatures adversely affect cell envelope properties and functions by leading to reduced membrane fluidity, permeability and diffusion rates, in addition to decreasing mobility and function of embedded proteins, increasing turgor pressure and even leading to physical cell rupture by ice formation and/or freeze–thaw cycles. While the cold adaptation traits of the cell membrane have been known for some time, the adaptation strategies of the other envelope components, the outer membrane, peptidoglycan layer and even exterior cell coatings, are now also beginning to be understood.

Cell membrane

Homeoviscous adaptation of cell membranes to low temperatures is accomplished in cold-adapted microorganisms via modification of the fatty acid composition of the lipid bilayer (D’Amico et al. 2001; Siddiqui et al. 2013). Cells mainly increase the content of unsaturated fatty acids, but increases in the content of short chain, methyl branched and/or cis-isomeric fatty acids are also common (Chintalapati et al. 2004; Russell 1997, 2008). These fatty acids disrupt the packing order and reduce the packing density of the phospholipid bilayer, leading to a lowering of the liquid phase to gel phase transition temperature and maintenance of functional fluid bilayers even at low temperatures. Indeed, in agreement with this cold adaptation strategy, an overrepresentation and upregulation of genes encoding various proteins involved in membrane biogenesis and fatty acid synthesis as well as in fatty acid desaturation (desaturases, which simultaneously also protect against ROS), production of branched chain fatty acids (KAS-II, KAS-III) and cis-isomerisation (fatty acid cis/trans isomerases) have been reported for numerous cold-adapted organisms (De Maayer et al. 2014; Goordial et al. 2016; He et al. 2015; Medigue et al. 2005; Methé et al. 2005). In addition, an increased genome content of genes for proteins involved in the degradation of membrane rigidifying molecules naturally present in the environment has also been observed

Table 1 The toolkit for a psychophilic lifestyle. Bioproducts believed to play a role in cold adaptation of microorganisms, their proposed function in cold-adaptation and their potential biotechnological applications are listed. Only those tools with a commercial potential and described potential role in cold adaptation are given. ‘?’ indicates that questions still remain as to the true function in cold adaptation

Bioproduct	Proposed cold-adaptation functions	Potential applications
Membrane fatty acids: unsaturated fatty acids, long-chain polyunsaturated fatty acids (LC-PUFA)	Unsaturated fatty acids: maintenance of membrane fluidity LC-PUFA: maintenance of membrane fluidity?	Nutrition and health
Membrane pigments: carotenoids	Maintenance of membrane fluidity? Cryoprotection?	Nutrition and health
Compatible solutes	Osmoprotection: against freezing induced osmotic stress Desiccation protection: against freezing induced desiccation Freezing point depression Colloidal glass transition temperature depression Protein and membrane stabilisation	Cryopreservation Stabilisation of biological materials Food stability and freshness Cosmetics and skin care Plant resistance enhancement
Antifreeze proteins	Ice growth inhibition (thermal hysteresis) Ice recrystallisation inhibition Membrane stabilisation? Ice adhesion?	Frozen foods preservation and quality enhancement Cryopreservation Cryosurgery Freeze tolerance enhancement, e.g. crops, fish Gas hydrate inhibition ‘Ice-prevention’ materials
Ice-nucleating proteins	Extracellular ice crystal nucleation - Prevention/reduction of damaging intracellular ice formation - Small ice crystals? - Release of latent heat of crystallisation?	Artificial snow production Frozen foods and beverages industries Microfluidic devices: freeze–thaw valves Cell surface display Climate control
Extracellular polymeric substances	Ice growth inhibition Osmoprotection: against freezing induced osmotic stress Desiccation protection: against freezing induced desiccation Ice-recrystallisation inhibition	Biopolymers Biofloculants Bioabsorbants Bioemulsifiers Bioadhesives Thickening agents Cryopreservation
Biosurfactants	Ice-recrystallisation inhibition? Osmoprotection?	Biosurfactants
Cold-adapted enzymes	Maintenance of adequate metabolic flux	Enzymes markets: low to moderate temperature processes, processes where simplified inactivation (e.g. by mild heat treatment) is beneficial
Chaperones	Promotion of protein folding and stability Destabilisation of RNA/DNA secondary structures	Low temperature recombinant protein production
Storage compounds: polyhydroxyalkanoates, cyanophycins	Overcoming carbon and nitrogen uptake deficiencies	Biopolymers Fine chemical synthesis Biofuels Stress tolerance enhancement: plants

(Medigue et al. 2005) and may serve as a further means for reducing membrane rigidity at low temperatures.

Finally, in relation to the other major components of the cell membrane, i.e. the proteins embedded in the lipid bilayer, an upregulation of membrane transport proteins has been observed in some psychrophiles and is believed to act in counteracting the reduced diffusion rates and transport inherent to low temperatures (Bakermans et al. 2007; De Maayer et al. 2014).

Cell membrane LC-PUFAs

While the role of unsaturated fatty acids with low numbers of double bonds in membrane cold adaptation is well understood, the precise role of long-chain polyunsaturated fatty acids (LC-PUFAs) is still under discussion. These have been considered part of the cold adaptation tactic for maintaining membrane fluidity, and indeed, some LC-PUFAs, such as eicosapentaenoic (EPA, 20:5 ω 3), docosahexaenoic (DHA, 22:6 ω 3) and arachidonic (ARA, 20:4 ω 6c) acid, are preferentially distributed in marine organisms, with higher levels being often produced at lower temperatures (Feng et al. 2014; Okuyama et al. 2008; Yoshida et al. 2016). In line with this, comparative genomics has indicated that the polyketide synthase gene cluster (*pfaA*, *B*, *C*, *D* and *E*), responsible for the synthesis of these, is largely restricted to marine organisms (Shulse and Allen 2011). Nevertheless, recent studies have indicated that, rather than serving in cold adaptation, LC-PUFAs may primarily serve antioxidative functions, protecting against reactive oxygen species naturally present at high levels in the marine environment and augmented at low temperatures (Nishida et al. 2007; Okuyama et al. 2008). It is believed that LC-PUFAs act as membrane shields, forming more hydrophobic interfaces between the lipid bilayers and thereby preventing entry of ROS such as H₂O₂ into the cells. Therefore, their presence in low-temperature environments may not be a direct response to the cold, per se, but a response to other stress(es) inherent to low temperatures, in this case, oxidative stress. Interestingly, in addition to their antioxidative function, functions as chaperones for membrane proteins, in efflux processes and in cell division, have also been proposed (Okuyama et al. 2008; Yoshida et al. 2016).

PUFAs have considerable nutritional and pharmaceutical value (Yoshida et al. 2016). They are components of neuronal and thrombocytes cells, neutrophils and monocytes and are found at high concentrations in the brain and retina. Importantly also, they are precursors of eicosanoid signalling molecules and endocannabinoid neurotransmitters (Ochsenreither et al. 2016). As such, they have a wide array of functions in human health: from regulating the cardiovascular system, immune system and inflammation to participating in the development and proper functioning of the brain, eyes and central nervous system. Some PUFAs such a linoleic

acid (LA) and α -linolenic acid (ALA) are essential fatty acids which have to be obtained from the diet while LC-PUFAs such as DHA and ARA are recommended in infant diets. Indeed, a balanced intake of various ω -3 (ALA, EPA and DHA) and ω -6 PUFAs (LA, ARA) is recommended as part of a normal healthy diet. Fish oils are the main source of PUFAs, especially LC-PUFAs, but problems associated with flavour, allergies, effects on global fish stocks and presence of various environmental pollutants have resulted in the search for alternative sources, including microorganisms. However, membrane levels of phospholipid PUFAs in microorganisms are in general too low for commercial production. On the other hand, oleaginous microorganisms, which produce PUFAs as components of triacylglycerol in stored oils, and which can accumulate lipids at concentrations of up to 80% of dry biomass weight, may prove suitable sources. In this sense, cold-adapted organisms, with increased levels and/or varied distributions of PUFAs may be of interest, especially for animal, and in particular aquaculture feed. Indeed, marine microalgae are potent producers of various LC-PUFA and the marine dinoflagellate *Cryptocodinium cohnii* is used in commercial production of DHA as an infant formula additive. In addition, filamentous fungi of the genus *Mortierella*, many of which are cold adapted, and various cold-adapted yeasts are also reported to accumulate high levels of PUFAs (Amaretti et al. 2010; Ochsenreither et al. 2016) and may prove potent commercial sources.

Membrane pigments

Pigments, and especially carotenoids (pigmented polyisoprenoid hydrocarbons), have also been suggested to play a role in the modulation of cell membrane fluidity. Pigment production is common in psychrophilic microorganisms, being reported in isolates from ice cores and glaciers (Shen et al. 2018), marine surface waters (Dieser et al. 2010) and high altitude soils (Pandey et al. 2018). A few reports have shown an increased production of these at low temperatures (Chattopadhyay et al. 1997; Pandey et al. 2018), with a preference for polar carotenoids (Chattopadhyay et al. 1997; Jagannadham et al. 2000), yet a recent study identified a decrease in pigmentation upon slowly reducing the temperature for a number of Arctic bacteria (Singh et al. 2017). Interestingly, polar carotenoids are believed to enhance membrane rigidity and hence increased concentrations should be counterproductive to cold-adapted organisms at low temperatures. It has been suggested that these may serve in homeoviscous adaptation by counterbalancing the fluidizing effects of the unsaturated fatty acids and stabilising the membrane (Jagannadham et al. 2000). Furthermore, these pigments have been shown to also play a number of other important roles, including, in photoprotection (acting, in conjunction with other molecules such as scytonemin and mycosporine-

like amino acids, as light screeners protecting against high light and UV radiation common to many cold habitats), as antioxidants (protecting against ROS, also commonly produced in low temperature and/or high light environments), as light harvesters (in photosynthetic microorganisms) and even as antimicrobials (Pandey et al. 2018). Importantly, a potential role as cryoprotectants, imparting resistance to freeze–thaw cycles, has also been demonstrated (Dieser et al. 2010). These various overlapping functions would variably influence pigmentation levels and could lead to the conflicting pigmentation levels observed at low temperatures. Further studies are therefore required to unravel the true role and specific functions of membrane pigments in cold-adapted organisms and their relationship to pigment type and specific structure, as well as the precise effects of the different pigments on membrane properties.

Numerous carotenoids have various beneficial health effects (Kirti et al. 2014). They have been shown to be effective antioxidants and pro-vitamins (vitamin A), and to be important for healthy growth and development, in maintenance of the immune system and in eye health. In addition, they can be used as natural food colourants, in skin care, in sunscreen products and as precursors of chemicals for fragrance products, as well as in animal/aquaculture feed to impact desired colours on products, e.g. egg yolks and salmon. Cold-adapted organisms with enhanced levels of various different carotenoids may prove interesting microbial sources of carotenoids for such commercial applications.

Cell wall: peptidoglycan layer

Cold adaptation of other cell envelope components has been much less investigated than the cell membrane, but recent studies have begun to unveil possible strategies. In relation to adaptation of the cell wall, an upregulation of peptidoglycan biosynthesis genes and a thickened peptidoglycan layer at low temperatures has been reported in some cold-adapted bacteria (Mykytczuk et al. 2013; Rodrigues et al. 2008). Similarly, *Planococcus halocryophilus* Or1 also displays a thickened outer cell surface, but this is achieved by a rather unique mechanism involving extracellular cell wall associated hydrophobic encrustations composed of peptidoglycan, calcium carbonate and choline (Mykytczuk et al. 2013). Such strategies for thickening and strengthening the cell outer surfaces would obviously lead to a reinforced physical barrier which could protect psychrophiles against cell disruption by ice formation, freezing–thawing and/or increased osmotic pressure at low temperatures.

Cell wall: outer membrane

Cold adaptation traits have also been observed in the outer membrane layer of the cell wall of gram negative bacteria

and namely in the lipopolysaccharide (LPS) structure which accounts for ~75% of this layer (Corsaro et al. 2017). Only a few cold-adapted LPS structures have been analysed to date, but it appears that most produce LPS solely as rough LPS (i.e. without the specific O-chain component) of shortened length at low temperatures (Carillo et al. 2013, 2011; Corsaro et al. 2017, 2002). Many more cold-adapted LPS structures need to be analysed to determine the extent of these observations, but it has been suggested that these alterations may increase outer membrane flexibility and stability at low temperatures (Corsaro et al. 2017). As regards to the other LPS components, a similar strategy to that used for the lipid bilayer of cell membranes is observed for the lipid A component of LPS, i.e. increased content of short chain and/or unsaturated fatty acids to increase fluidity, while a high negative charge of the core oligosaccharide component has been suggested to be important in sequestration of divalent cations common in many cold environments (Casillo et al. 2017b; Corsaro et al. 2017; Sweet et al. 2015). Finally, in common with that observed for other cell envelope components, transcriptome analyses have shown an upregulation of genes involved in biosynthesis of outer membrane components, with LPS biosynthetic genes (mainly glycosyltransferases) and outer membrane proteins being upregulated at low temperatures (De Maayer et al. 2014; Frank et al. 2011; Gao et al. 2006). In agreement with this, a recent study showed how mutation of a core LPS glycosyltransferase gene (*wapH*) impaired growth of an Antarctic bacterium at low temperatures (Benforte et al. 2018).

Cryoprotection

Sub-zero temperatures provoke ice formation which can lead to cryoinjury, osmotic stress, dehydration and even cell rupture and death. In the natural environment, onset of ice formation is usually delayed in the cell interior as compared to the cell exterior due to the lower volume and densely packed, highly crowded nature of the former. Intracellular ice crystal formation, which is normally lethal to cells, can indeed occur when temperatures decrease at high rates, but in the natural environment, cooling is usually relatively slow and ice formation is thus mostly restricted to the extracellular space (Fonseca et al. 2016). Extracellular ice formation, which can lead to physically damaging membrane fracturing, occurs with exclusion of solutes and removal of available liquid water. This leads to elevated extracellular solute concentrations and provokes intra/extracellular osmotic imbalances. In turn, this leads to stresses related to osmotic shrinkage and dehydration of the cell interior, not only negatively affecting cell function and survival but also preventing intracellular ice formation and instead leading to a non-crystalline amorphous (colloidal glassy) state with inhibition of cell metabolism

(Fonseca et al. 2016). In addition, at relatively high sub-zero temperatures, in partially frozen environments subjected to freeze–thaw cycles and during temperature fluctuations, cells can also be subjected to harmful ice recrystallization stress, a thermodynamically driven process causing ice crystal coalescence and growth of large, fatally damaging ice crystals at the expense of smaller crystals (Bar Dolev et al. 2016b). Cold-adapted microorganism respond to these multiple freezing related detrimental challenges by production of a variety of novel tools including, compatible solutes, ice-binding proteins (antifreeze and ice-nucleating proteins), extracellular polymeric substances (EPS) and/or biosurfactants.

Compatible solutes

Compatible solutes are low molecular mass, non-toxic organic osmolytes. Many cold-adapted microorganisms have an increased genome content of compatible solute biosynthesis, uptake and degradation genes and often accumulate up to molar concentrations of various different compatible solutes, with glycine betaine, trehalose, glycerol, sucrose, sarcosine, mannitol and sorbitol being commonly reported (Ghobakhlou et al. 2015; Goordial et al. 2016; Mykytczuk et al. 2013). Accumulation of these organic osmolytes contributes to restoring osmotic balance and thereby counteracts water loss and cell shrinkage during freezing. In addition, they depress the freezing point of solution and, importantly also, the intracellular colloidal glass transition temperature (T_g) (Fonseca et al. 2016). In fact, a recent study showed reductions of as much as 30 °C in the T_g of the cytoplasm of bacterial cells and improved post thaw viability upon glycerol addition (Fonseca et al. 2016). Furthermore, compatible solutes are also believed to be involved in scavenging free radicals and, due possibly to their preferential exclusion from protein surfaces and/or water entrapment effects, compacting effects on proteins and destabilising effects on the unfolded state, also play roles in preventing protein aggregation, enhancing protein folding and stabilising proteins and membranes at low temperatures.

In relation to applied aspects, various compatible solutes are commonly used in the stabilisation, preservation and cryopreservation of diverse biological materials, ranging from enzymes to whole cells and tissues. In addition, their use in increasing the freshness and stability of foods, in cosmetics and skin care products, as well as in extending the growth performance of plants in saline, dry and low-temperature environments has also been investigated (Wani et al. 2013).

Ice-binding proteins: antifreeze proteins

Antifreeze proteins (AFPs), also known as ice structuring or thermal hysteresis proteins, are noncolligative biological antifreezes which can bind to ice and inhibit ice growth and recrystallisation (for recent reviews on AFPs

see, e.g. Bar Dolev et al. (2016b); Lorv et al. (2014) and Voets (2017)). They were first identified in the blood of Antarctic fish but have since been reported in various bacteria, fungi, diatoms, plants, insects and crustaceans. Various types of AFPs of diverse structure exist in nature, being frequently glycosylated and/or lipidated and varying in size from ~2 to ~50 kDa, with a 1.5 MDa multidomain ice adhesion AFP being reported for the Antarctic bacterium *Marinomonas primoryensis* (Bar Dolev et al. 2016a). AFPs are believed to function by irreversibly binding to specific ice crystal planes, thereby blocking secondary nucleation events and shaping a unique ice morphology. Due to the Kelvin effect, ice surface expansion between the adsorbed AFPs leads to local curvature of the ice face which is energetically unfavourable for further ice growth. This leads to thermal hysteresis (TH) in which the freezing point is depressed below the equilibrium melting point to create a thermal hysteresis gap in which ice crystal growth is halted. TH activities from ~0.1 °C to as high as ~13 °C have been measured (Duman et al. 2004; Voets 2017) and these activities can be increased in the presence of other AFPs, solutes and ions. AFPs are also effective ice recrystallisation inhibitors (IRI), being frequently more effective in IRI (often requiring only sub- μ M concentrations) than in TH (often requiring mM concentrations). Indeed, IRI is believed to be the primary function of many secreted AFPs from cold-adapted organisms and AFPs from Antarctic algae and glacier ice bacteria were shown to effectively stabilise brine pockets and contribute to the preservation of a liquid environment in the cell vicinity (Raymond et al. 2008). Finally, in addition to their TH and IRI activities, AFPs are also believed to help stabilise cell membranes and protect structural integrity and, in some cases, via their adhesion domains and ice binding function, are thought to play a role in positioning cells on ice so as to enhance access to oxygen and nutrients in the phototrophic zone (Bar Dolev et al. 2016a; Lorv et al. 2014; Voets 2017).

AFPs are already being applied in the food industry for improved food preservation during freezing. They are used to preserve a smooth texture in ice creams and frozen yoghurts and can reduce drip loss in frozen meats and fish, enhance frozen dough bread quality and improve post-freezing quality of currently difficult to freeze foods such as fruit and vegetables (Muñoz et al. 2017; Regand and Goff 2006; Voets 2017). They also have obvious potential as cryoprotective agents and have been shown to improve cryopreservation efficiency and post thaw viability of various types of biological materials. Furthermore, the ability of certain AFPs at high concentrations to induce formation of needle-like structures which penetrate and destroy cells is being investigated for use in cryosurgery, e.g. in tumour ablation. They are also being developed for use

in improving freeze tolerance of crops and aquaculture fish, in gas hydrate inhibition in the petroleum industry and in anti-freeze/de-icing materials via surface coating of materials for prevention/decreasing of ice formation, e.g. on aircrafts, power lines and roads (Bar Dolev et al. 2016b; Lorv et al. 2014; Voets 2017).

Ice-binding proteins: ice-nucleating proteins

Ice-nucleating proteins (INPs) are large membrane bound proteins that facilitate ice formation. They initiate heterogeneous ice crystallisation at high sub-zero temperatures and are proposed to act by providing a template for the ordering and stabilisation of water molecules in an ice-like structure (Lorv et al. 2014; Pandey et al. 2016; Pummer et al. 2015). This lowers the activation energy barrier for freezing and nucleates ice growth at temperatures as high as $-2\text{ }^{\circ}\text{C}$ (Li et al. 1997). INPs are produced by diverse organisms, most commonly as large, extracellular, repetitive multimeric aggregates, with larger complexes enabling higher activity (Bar Dolev et al. 2016a). In cold-adapted microorganisms, INPs are believed to counteract low-temperature damage by directing ice nucleation to the extracellular space (Lorv et al. 2014) which can prevent formation of lethally damaging intracellular ice via removal of the available intracellular liquid water as discussed above. In addition, INPs favour development of small extracellular crystals which are less damaging to the cell than large crystals while the release of latent heat of crystallisation during the freezing process could also be beneficial in preventing further temperature decreases (Pummer et al. 2015). In addition to their role in cold adaptation, some INPs also play a role in nutrient mining by many plant pathogenic organisms where they are used to induce frost damage in plants and enable access to nutrients.

INPs are presently being commercialised for artificial snow production (Cochet and Widehem 2000) but are believed to have a number of other potential applications. These include reducing freezing energy costs in the food industry (Li et al. 1997), freeze-concentrating beverages, in freeze-thaw valves of microfluidic devices (Gaiteri et al. 2017), as anchoring motifs for cell surface display applications (Jung et al. 1998), as well as in cloud seeding for climate control (Pummer et al. 2015).

Extracellular polymeric substances

Extracellular polymeric substances (EPS) are multifunctional, high molecular weight biopolymer complexes secreted by various organisms into their local environment. They are large, complex, highly diverse architectural structures composed principally of carbohydrates (homo- or hetero-polysaccharides), but also proteins and lower concentrations of nucleic acids, lipids, phenols and humic substances. EPS are produced

by a wide variety of organisms and are found either attached to the cells outer surface or released into the surrounding environment. They form hydrated gels that play an important role in the formation of biofilms and in the modification of the physical, chemical and biological characteristics of the cell environment. Indeed, they are believed to have multiple functions, including in cell adhesion and nutrient scavenging, but are also thought to be important in protective functions such as osmoprotection, ROS scavenging, extracellular protein protection and even in cryoprotection. For reviews of the subject, see, e.g. Deming and Young (2017) and Ewert and Deming (2013) and references therein.

In relation to cold-adaptation, metagenomics studies have identified numerous genes for EPS biosynthesis in both Antarctic and Arctic ice shelf ponds (Varin et al. 2012) and cold-adapted bacteria have been found to produce high concentrations of EPS at low and especially sub-freezing temperatures (Caruso et al. 2018; Feng et al. 2014; Marx et al. 2009; Mykytczuk et al. 2013). The hydrated EPS gel matrix is believed to protect against low temperatures by forming a protective shell around cells which acts as a diffusion barrier to solutes and a physical-like barrier to ice formation (Caruso et al. 2018; Deming and Young 2017; Ewert and Deming 2013; Krembs et al. 2011). This diminished solute diffusion limits freezing induced osmotic stress and desiccation damage. Furthermore, the gel-like state of EPS reduces the available free energy for ice nucleation while solute exclusion to the surrounding liquid phase impedes ice crystal growth in this, thereby protecting cells from ice damage and increasing the available habitable liquid space. Interestingly also, recent studies have indicated ice binding functions and IRI activity similar to AFPs (described above) for EPSs produced by cold adapted organisms (Casillo et al. 2017a). As compared to other cold-adaptation traits, EPS are much less well studied and the important role of these in coping with low temperatures is only beginning to be unravelled. Further studies are warranted to gain a better understanding of the different inter-related cold adaptation roles and physiological and ecological functions of EPS and their relationship to composition and structure.

Microbial EPS and their various components, as biodegradable, biosustainable, non-toxic and biocompatible biopolymers, have been recognised as potential alternatives to chemical polymers in a number of applications, ranging from the pharmaceutical to the cosmetics and food industry, and environmental biotechnology. Their capacity to adsorb heavy metals and organic pollutants combined with their flocculation properties for removal of suspended solids, organic matter, etc. are advantageous for use as bioflocculants and bioabsorbants in soil and water bioremediation and decontamination as well as in wastewater and sludge treatment (More et al. 2014). Their

use, especially the polysaccharide component, as cryoprotectants is also under study and an ability to improve the freeze–thaw survival of various Antarctic bacteria has already been shown (Caruso et al. 2018). The high emulsifying activity of some EPS indicates potential as bioemulsifiers (Caruso et al. 2018) while their role as bioadhesives has also been indicated (Muralidharan and Jayachandran 2003). In the food industry, EPS polysaccharides can be used as thickening agents and emulsifiers, conferring improved texture and stability to foods and beverages while, more recently, beneficial effects of EPS on human health via potential immunomodulatory, anticoagulant, anti-inflammatory and antioxidant capacities have been suggested (Collic Jouault et al. 2001; Leroy and De Vuyst 2016). Furthermore, their use in tissue engineering has even been suggested (Kumar et al. 2018). A number of microbial EPS carbohydrates, e.g. xanthan gum, gellan gum and dextran, have already been commercialised for a number of applications, but studies of cold-adapted EPS components with high potential for identification of unique structures and functions could be expected to reveal novel products with novel applications.

Biosurfactants

Biosurfactants are surface active amphiphilic compounds of microbial origin which reduce surface and interfacial tension between liquids, solids and gasses. Generally, they are of low molecular weight, in contrast to bioemulsifiers, such as for example EPS, which tend to be of high molecular weight, and can be composed of sugars, amino acids, fatty acids and/or functional groups such as carboxylic acids. They are structurally diverse compounds but the most commonly reported are glycolipids (rhamnolipids, sophorolipids, trehalose lipids and mannosylerythritol lipids, etc.), lipopeptides (surfactin, iturin, fengycin, viscosin, etc.), phospholipids, fatty acids and neutral lipids. In nature, they are believed to play roles in enhancing the bioavailability of poorly soluble hydrophobic substrates, in regulating biofilm structure and surface attachment/detachment, in bacterial pathogenesis and quorum sensing, as well as acting as antibacterial and antifungal agents. In relation to cold adaptation, a glycolipid biosurfactant isolated from an Antarctic yeast was shown to have IRI activity (Kitamoto et al. 2001), while a role for biosurfactants as osmolytes has also been suggested (Perfumo et al. 2018). Interestingly, even though biosurfactant production appears to be widespread among cold adapted organisms (Gesheva et al. 2010; Malavenda et al. 2015; Perfumo et al. 2018; Vollú et al. 2014), studies investigating their potential role in cold adaptation are scant. Further studies are obviously required to clarify whether these compounds constitute part of the cells toolkit for adapting to cold and to elucidate their precise role and mechanism of action in this.

Biosurfactants offer non-toxic, sustainable, biodegradable and ecofriendly alternatives to the chemical surfactants currently used in a huge variety of applications as diverse as bioremediation, in cleaning agents, cosmetics, pharmaceuticals, agriculture and the petroleum industry. Interest in ‘cold’ biosurfactants has only recently been initiated, but already potential in a number of applications has been demonstrated; they have been shown to have potential for use as antiagglomerants in ice–water slurry technologies (Kitamoto et al. 2001), as low-temperature detergents (Perfumo et al. 2018), as fuel additives to improve the flow properties of biodiesel and diesel at low temperatures (Madihalli et al. 2016), and for recovery of natural gas hydrates in gas hydrate technologies (Arora et al. 2016; Perfumo et al. 2018).

Enzymes

One of the principal challenges that cold-adapted microorganisms have to contend with is the negative effect of low temperatures on reaction rates. All reactions, including enzyme catalysed reactions, are influenced by temperature according to the Arrhenius Law wherein any decrease in temperature induces an exponential decrease in the reaction rate (Arrhenius 1889). Indeed, with most non-adapted enzymes, a reduction in temperature from 37 °C to 0 °C results in a 16–80-fold reduction in activity. In contrast, most enzymes produced by psychrophilic organisms are adapted to their environment and maintain high specific activities enabling appropriate metabolic rates in their cold habitats (for recent reviews, see, e.g. Collins et al. (2002a); Collins and Gerday (2017); Fields et al. (2015); Gerday (2013); Santiago et al. (2016) and Siddiqui (2015)).

With few exceptions (e.g. Oswald et al. 2014; Roulling et al. 2016), most cold-adapted enzymes studied to date are characterised by a higher catalytic activity at low to moderate temperatures, a shift in the optimum temperature for activity towards lower temperatures and a decreased stability as compared to mesophilic and thermophilic homologous enzymes. These ‘cold enzymes’ successfully reduce the activation energy, ΔG^* , and temperature dependence of reactions through a reduced activation enthalpy, ΔH^* , thereby implying a decreased number and/or strength of enthalpic interactions that are broken during the catalytic cycle (Lonhienne et al. 2000). This is accompanied by a lower, more negative activation entropy value, ΔS^* , suggestive of greater re-ordering during activation, which, in conjunction with a more negative heat capacity change during activation, suggestive of a higher heat capacity and thus increased vibrational modes for the ground state (Arcus et al. 2016), and a lower substrate affinity (higher K_m) (Collins et al. 2002a), as compared to mesophilic and thermophilic homologues, implies a more disordered enzyme substrate (ES) ground state for psychrophilic enzymes. Such a

disordered state could give rise to an ES complex of reduced stability and higher free energy and hence lead to the observed reduced ΔG^* and enhanced catalytic rates for these enzymes.

A number of different studies using a range of techniques have investigated this disorder, or structural flexibility, in various psychrophilic enzymes. It is important to note here that protein flexibility is a complex parameter which is difficult to measure and further in-depth comparative studies characterising the specific amplitudes, time frames, regions involved and temperature dependence of protein motions are called for, with use of powerful techniques such as NMR being key to better understanding this parameter. Nonetheless, notwithstanding current limitations, it is now generally accepted that psychrophilic enzymes do indeed display an increased structural flexibility and that they have overcome the low-temperature challenge by increasing the plasticity or flexibility of specific regions (at or near the catalytic site) or of the whole protein. This flexibility enables the conformational changes necessary for activity at a low energy cost but also leads to a decreased stability. At the structural level, this increased flexibility is mainly achieved by a reduction in the number and/or strength of stabilising interactions such as H bonds, salt bridges, aromatic interactions, disulphide bonds, ion binding sites and/or a weakening of the hydrophobic core. In addition, a reduction in the proline content and increase in glycine content, especially in loop regions, and a greater exposure of hydrophobic residues have also been reported. Such modifications would allow for the increased flexibility necessary for increased low-temperature activity but would also lead to the reduced stability inherent to most psychrophilic enzymes (Collins and Gerday 2017). Importantly, it has been found that different enzymes can use different structural adaptation strategies, with each enzyme using its own specific strategy, employing any one or combination of these modifications depending on its own specific characteristics, its environment and its requirements.

The inherent characteristics of a high activity at low to moderate temperatures and a reduced stability of psychrophilic enzymes offers many advantages in a variety of applications. They can be used to enhance the efficiency and economics of low to moderate temperature processes, reduce the process temperature and hence improve process economics and environmental impact and enable more simplified enzyme inactivation. Cold enzymes are already being employed in all three sectors of the industrial enzymes markets, food, feed and technical, and are expanding into new areas in pharmaceutical and fine chemical synthesis. For a recent in-depth review, see Barroca et al. (2017a). Some of the better known commercial successes include a cold-adapted lipase used in the organic synthesis of various pharmaceutical, cosmetic and flavour compounds (Kirk and Christensen 2002); a xylanase for enhancing bread quality (Barroca et al. 2017b; Collins et al. 2012, 2006, 2002b; Dutron et al. 2012), various hydrolases

in detergents for low-temperature cleaning (Sarmiento et al. 2015) and various enzymes (alkaline phosphate, nuclease and uracil-DNA N-glycosylase) used in molecular biology for their high activity and ease of inactivation (Barroca et al. 2017a).

Chaperones

Protein and RNA/DNA chaperones, which facilitate efficient protein and RNA/DNA folding, respectively, play important roles in counteracting protein misfolding and aggregation as well as low-temperature stabilisation of RNA and DNA secondary structures. DNA and RNA chaperones are important in maintaining efficient transcription, translation and DNA replication. They are transiently produced as part of the cold-shock response in mesophilic and thermophilic microorganisms but are often continuously overexpressed as cold-acclimation proteins or upregulated at low temperatures in psychrophiles (Lim et al. 2000). In relation to protein chaperones in cold-adapted microorganisms, continuous overexpression, upregulation of production, and production of cold adapted variants, as well as, contrastingly, no overexpression and cold repression, have been reported (Ferrer et al. 2003; Godin-Roulling et al. 2015). Importantly, while protein misfolding and precipitation are believed to be strongly reduced at low temperatures, due mainly to a weakening of hydrophobic interactions, proteins at the low range of the temperature spectrum are faced with another phenomenon; cold denaturation (Collins and Gerday 2017; Romero-Romero et al. 2011). Cold denaturation is thought to be due to a preferential hydration and weakening of hydrophobic and ionic interactions at low temperatures, and intriguingly, it appears that psychrophilic enzyme may be more susceptible to this than their higher temperature adapted homologues (D'Amico et al. 2003). Nevertheless, the underlying basis for this is still not fully understood and future studies should address this.

Protein and RNA/DNA chaperones have potential for use in recombinant protein production; in the low temperature production of proteins prone to aggregation/misfolding, and an *Escherichia coli* host producing protein chaperones from an Antarctic bacterium has already been commercialised for such a use (Ferrer et al. 2003).

Metabolic adjustments

Recent studies making use of modern omics approaches such as genomics, transcriptomics and proteomics have revealed a number of additional traits common to various cold adapted microorganisms (see, e.g. Tribelli and Lopez (2018) for a recent review). In particular, studies have indicated various

metabolic adjustments at low temperatures, including a down-regulation of primary metabolism pathways and substitution with abridged or alternative secondary pathways, as well as accumulation and metabolism of reserve compounds. Oxidative metabolic processes, namely, glycolysis, the pentose phosphate pathway, the TCA cycle and/or the electron transport chain, as well as pathways involving metal ions and molybdopterin metabolism are reported to be downregulated at low temperatures in some psychrophiles (Medigue et al. 2005; Piette et al. 2011; Tribelli et al. 2015). While still poorly understood, studies have indicated their substitution with alternative/shortened pathways such as the glyoxylate, methylglyoxal or 2-methylcitrate cycles, the ethanol oxidation pathway, acetate metabolism or propionyl-CoA catabolism (Ayala-del-Río et al. 2010; Tribelli and Lopez 2018), the actual alternative pathway(s) used being dependent on the organism and its ecological niche. Such a strategy of metabolic reprogramming may alleviate oxidative stress inherent to low temperatures by avoiding ROS producing pathways but may also be important in energy conservation and long-term survival.

Screening studies have indicated a high content of polyhydroxyalkanoate (PHA) producers in cold habitats (Ciesielski et al. 2014; Goh and Tan 2012; Pärnänen et al. 2015), and many psychrophiles can accumulate and degrade PHA (López et al. 2009; Methé et al. 2005; Ting et al. 2010) and/or cyanophycin-like (Duchaud et al. 2007; Methé et al. 2005; Vollmers et al. 2013) compounds. These compounds can act as dynamic reserves of carbon, nitrogen, reducing equivalents and energy in cells and are thought to be important in overcoming low-temperature challenges to carbon and nitrogen uptake. Nevertheless, functions in cryoprotection, oxidative stress resistance, maintenance of cellular redox balance and cell motility have also been suggested for PHAs (Méthé et al. 2005; Tribelli and Lopez 2018). In relation to applied aspects, these compounds, or derivatives, are suggested as non-toxic, biodegradable, biocompatible biopolymers to replace petrobased polymers. PHAs are a family of microbial polyesters with interesting thermoplastic and elastomeric properties and potential for application in almost all areas of the conventional plastics industry. For reviews on PHA applications, see, e.g. Chen (2009) and Singh et al. (2019). A specific focus of application has been in the medical (tissue engineering, bio-implants, drug delivery, sutures, etc.), and fine chemical synthesis fields (chiral starting materials for the synthesis of antimicrobials, vitamins, fragrances etc.), but also in materials (packaging, smart materials etc.) and biofuels. Recently, novel applications such as in enhancing stress tolerance in plants have even been suggested (Stritzler et al. 2018). Cyanophycin-like compounds are polyamides composed mainly of aspartic acid and

arginine. Derivatives of these have been suggested for use in nutrition (sources of highly bioavailable amino acids/peptides), biomedicine (as polyaspartic acid hydrogels), laundry detergents (polyaspartic acid), anti-scalants (polyaspartic acid) and in bioflocculation (polyaspartic acid). For a recent review on cyanophycins and their applications, see, e.g. Frommeyer et al. (2016).

Other applications

In addition to the numerous biotechnological tools related to the cells response to low temperatures presented above, an array of other bioactive metabolites and bioproducts, including novel antimicrobials, anti-fungal agents, anti-cancer drugs, anti-tumour and anti-inflammatory agents, antioxidants, alkaloids and organic acids have also been identified in psychrophilic microorganisms. For reviews on these, see, e.g. Avila (2016), Borchert et al. (2017) and Soldatou and Baker (2017). The majority of these bioproducts used today originate from moderate temperature or warm environments, whereas cold environments have been much less investigated and hence offer an extraordinary opportunity as an underexplored source of potential novelty. Indeed, the abundance and diversity of psychrophiles, combined with the vastness, enormous diversity and severity of their habitats point to their tremendous potential as rich reservoirs of novel biomolecules and metabolites of applied interest. Recent growing interest in cold environments has led to identification of numerous new products, mainly from microbes (Soldatou and Baker 2017), and further bioprospection of these environments using modern high throughput techniques such as metagenomics-based approaches will surely lead to discovery of further novel tools with diverse bioactivities and applications (Borchert et al. 2016).

Cold-adapted microorganisms have also been shown to have potential for use in bioremediation, as probiotics, and as cell factories. Their potential for degrading a wide range of organic compounds of environmental concern, including mineral oil hydrocarbons, phenolic compounds, polyaromatic hydrocarbons, pesticides and persistent pollutants, but also proteins, carbohydrates and lipids, has found broad application in bioremediation of polluted cold soils and waters, and their use in wastewater and groundwater treatment has also been suggested (reviewed by e.g. Bajaj and Singh (2015) and Margesin (2017)). As probiotics, psychrophiles are believed to have potential for use as dietary supplements in aquaculture to improve health and nutrition of livestock (Makled et al. 2017; Sun et al. 2011; Wanka et al. 2018). While studies in this are scarce, their adaptation to low temperatures is suggested to be beneficial for a more efficient utilisation in the

marine habitat as compared to currently available terrestrial and/or moderate temperature adapted probiotic organisms. Finally, in relation to their utilisation as cell factories, cold-adapted microorganisms can be used for the production of heat-sensitive compounds and difficult to express or aggregation-prone proteins at low temperatures with reduced environmental and economic impact due to the absence of heating requirements (Miyake et al. 2007; Parrilli and Tutino 2017).

Conclusions

Cold-adapted microorganisms inhabiting permanently low-temperature environments have evolved a suite of highly sophisticated adaptations at all levels of their cells to cope with their challenging habitat conditions. Many of these adaptive features and their underlying mechanism of action are beginning to be understood but questions remain in a number of areas. These include, among others, the structure–function relationships and precise role(s) of EPS, PUFA, LPS, biosurfactants, membrane pigments and accumulated compounds such as PHA in cold adaptation; the effects of cold denaturation on proteins adapted to different temperatures and the protective measures employed by psychrophiles to counteract this; a better understanding of the dynamic motions, amplitudes and time frames of protein structures and their role in temperature adaptation and function; and a better understanding of the various metabolic adjustments employed by psychrophiles. Further studies making use of modern technologies will advance our understanding in these areas and will undoubtedly unveil further adaptive traits and novel metabolic peculiarities employed by these microorganisms, leading to identification of further novel biomolecules with novel properties for use as innovative biotechnological tools. Indeed, while a number of psychrophile derived commercial tools are already being commercialised, the adaptive features, abundance and diversity of psychrophiles and their habitats point to their high potential as rich sources of further novel biomolecules and compounds of applied interest, and further bioprospection of these organisms and their environments should lead to an improved valorisation of their commercial potential.

Acknowledgements All the technical staff at the CBMA are thanked for their skilful technical assistance. The Fundação para a Ciência e a Tecnologia (FCT), the European Social Fund (ESF), the Programa Operacional Potencial Humano (POPH) and the European Regional Development Fund (ERDF) are thanked for funding.

Funding information T.C. is supported by the FCT, the ESF, POPH, and the Investigador FCT Programme (IF/01635/2014). Funding by the

ERDF is through project EcoAgriFood (NORTE-01-0145-FEDER-000009) via the North Portugal Regional Operational Programme (NORTE 2020) under the PORTUGAL 2020 Partnership Agreement. FCT funding was through the project EngXyl (EXPL/BBB-BIO/1772/2013-FCOMP-01-0124-FEDER-041595) and the strategic programme UID/BIA/04050/2019 at the CBMA.

Compliance with ethical standards

Ethical statement This article does not contain any studies with human participants or animals performed by any of the authors.

Conflict of interest The authors declare that they have no conflict of interest.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Amaretti A, Raimondi S, Sala M, Roncaglia L, De Lucia M, Leonardi A, Rossi M (2010) Single cell oils of the cold-adapted oleaginous yeast *Rhodotorula glacialis* DBVPG 4785. *Microb Cell Factories* 9:73–76. <https://doi.org/10.1186/1475-2859-9-73>
- Arcus VL, Prentice EJ, Hobbs JK, Mulholland AJ, Van der Kamp MW, Pudney CR, Parker EJ, Schipper LA (2016) On the temperature dependence of enzyme-catalyzed rates. *Biochemistry* 55(12):1681–1688. <https://doi.org/10.1021/acs.biochem.5b01094>
- Arora A, Cameotra SS, Kumar R, Balomajumder C, Singh AK, Santhakumari B, Kumar P, Laik S (2016) Biosurfactant as a promoter of methane hydrate formation: thermodynamic and kinetic studies. *Sci Rep* 6:20893. <https://doi.org/10.1038/srep20893>
- Arrhenius S (1889) Über die Reaktionsgeschwindigkeit bei der Inversion von Rohrzucker durch Säuren. *Z Phys Chem* 4:226–248
- Avila C (2016) Biological and chemical diversity in Antarctica: from new species to new natural products. *Biodiversity* 17(1–2):5–11. <https://doi.org/10.1080/14888386.2016.1176957>
- Ayala-del-Río HL, Chain PS, Grzymalski JJ, Ponder MA, Ivanova N, Bergholz PW, Di Bartolo G, Hauser L, Land M, Bakermans C, Rodrigues D, Klappenbach J, Zarka D, Larimer F, Richardson P, Murray A, Thomashow M, Tiedje JM (2010) The genome sequence of *Psychrobacter arcticus* 273-4, a psychoactive Siberian permafrost bacterium, reveals mechanisms for adaptation to low-temperature growth. *Appl Environ Microbiol* 76(7):2304–2312. <https://doi.org/10.1128/aem.02101-09>
- Bajaj S, Singh DK (2015) Biodegradation of persistent organic pollutants in soil, water and pristine sites by cold-adapted microorganisms: mini review. *Int Biodeterior Biodegrad* 100:98–105. <https://doi.org/10.1016/j.ibiod.2015.02.023>
- Bakermans C, Tollaksen SL, Giometti CS, Wilkerson C, Tiedje JM, Thomashow MF (2007) Proteomic analysis of *Psychrobacter cryohalolentis* K5 during growth at subzero temperatures. *Extremophiles* 11(2):343–354. <https://doi.org/10.1007/s00792-006-0042-1>
- Bar Dolev M, Bernheim R, Guo S, Davies PL, Braslavsky I (2016a) Putting life on ice: bacteria that bind to frozen water. *J R Soc Interface* 13(121). <https://doi.org/10.1098/rsif.2016.0210>
- Bar Dolev M, Braslavsky I, Davies PL (2016b) Ice-binding proteins and their function. *Annu Rev Biochem* 85:515–542. <https://doi.org/10.1146/annurev-biochem-060815-014546>
- Barauna RA, Freitas DY, Pinheiro JC, Folador AR, Silva A (2017) A proteomic perspective on the bacterial adaptation to cold: integrating

- OMICs data of the psychrotrophic bacterium *Exiguobacterium antarcticum* B7. *Proteomes* 5(1). <https://doi.org/10.3390/teomes5010009>
- Barroca M, Santos G, Gerday C, Collins T (2017a) Biotechnological aspects of cold-active enzymes. In: Margesin R (ed) *Psychrophiles: from biodiversity to biotechnology*, 2nd edn. Springer, Cham, pp 461–475
- Barroca M, Santos G, Johansson B, Gillotin F, Feller G, Collins T (2017b) Deciphering the factors defining the pH-dependence of a commercial glycoside hydrolase family 8 enzyme. *Enzym Microb Technol* 96:163–169. <https://doi.org/10.1016/j.enzymtec.2016.10.011>
- Benforte FC, Colonnella MA, Ricardi MM, Solar Venero EC, Lizarraga L, Lopez NI, Tribelli PM (2018) Novel role of the LPS core glycosyltransferase WapH for cold adaptation in the Antarctic bacterium *Pseudomonas extremaustralis*. *PLoS One* 13(2):e0192559. <https://doi.org/10.1371/journal.pone.0192559>
- Borchert E, Jackson SA, O’Gara F, Dobson ADW (2016) Diversity of natural product biosynthetic genes in the microbiome of the deep sea sponges *Inflatiella pellicula*, *Poecillastra compressa*, and *Stelletta normani*. *Front Microbiol* 7(1027). <https://doi.org/10.3389/fmicb.2016.01027>
- Borchert E, Jackson SA, O’Gara F, Dobson ADW (2017) Psychrophiles as a source of novel antimicrobials. In: Margesin R (ed) *Psychrophiles: from biodiversity to biotechnology*, 2nd edn. Springer, Cham, pp 527–540
- Bowman JP (2017) Genomics of psychrophilic bacteria and archaea. In: Margesin R (ed) *Psychrophiles: from biodiversity to biotechnology*, 2nd edn. Springer, Cham, pp 345–387
- Carillo S, Pieretti G, Parrilli E, Tutino ML, Gemma S, Molteni M, Lanzetta R, Parrilli M, Corsaro MM (2011) Structural investigation and biological activity of the lipooligosaccharide from the psychrophilic bacterium *Pseudoalteromonas haloplanktis* TAB 23. *Chemistry* 17(25):7053–7060. <https://doi.org/10.1002/chem.201100579>
- Carillo S, Pieretti G, Lindner B, Parrilli E, Filomena S, Tutino ML, Lanzetta R, Parrilli M, Corsaro MM (2013) Structural characterization of the core oligosaccharide isolated from the lipopolysaccharide of the psychrophilic bacterium *Colwellia psychrerythraea* strain 34H. *Eur J Org Chem* 2013(18):3771–3779. <https://doi.org/10.1002/ejoc.201300005>
- Caruso C, Rizzo C, Mangano S, Poli A, Di Donato P, Finore I, Nicolaus B, Di Marco G, Michaud L, Lo Giudice A (2018) Production and biotechnological potential of extracellular polymeric substances from sponge-associated Antarctic bacteria. *Appl Environ Microbiol* 84(4). <https://doi.org/10.1128/AEM.01624-17>
- Casillo A, Parrilli E, Sannino F, Mitchell DE, Gibson MI, Marino G, Lanzetta R, Parrilli M, Cosconati S, Novellino E, Randazzo A, Tutino ML, Corsaro MM (2017a) Structure-activity relationship of the exopolysaccharide from a psychrophilic bacterium: a strategy for cryoprotection. *Carbohydr Polym* 156:364–371. <https://doi.org/10.1016/j.carbpol.2016.09.037>
- Casillo A, Ziacco M, Lindner B, Parrilli E, Schwudke D, Hologado A, Verstrepen L, Sannino F, Beyaert R, Lanzetta R, Tutino ML, Corsaro MM (2017b) Unusual lipid A from a cold-adapted bacterium: detailed structural characterization. *ChemBioChem* 18(18):1845–1854. <https://doi.org/10.1002/cbic.201700287>
- Chattopadhyay MK, Jagannadham MV, Vairamani M, Shivaji S (1997) Carotenoid pigments of an Antarctic psychrotrophic bacterium *Micrococcus roseus*: temperature dependent biosynthesis, structure, and interaction with synthetic membranes. *Biochem Biophys Res Commun* 239(1):85–90. <https://doi.org/10.1006/bbrc.1997.7433>
- Chen GQ (2009) A microbial polyhydroxyalkanoates (PHA) based bio-and materials industry. *Chem Soc Rev* 38(8):2434–2446. <https://doi.org/10.1039/b812677c>
- Chintalapati S, Kiran MD, Shivaji S (2004) Role of membrane lipid fatty acids in cold adaptation. *Cell Mol Biol (Noisy-le-Grand)* 50(5):631–642
- Ciesielski S, Górnjak D, Możejko J, Świątecki A, Grzesiak J, Zdanowski M (2014) The diversity of bacteria isolated from Antarctic freshwater reservoirs possessing the ability to produce polyhydroxyalkanoates. *Curr Microbiol* 69(5):594–603. <https://doi.org/10.1007/s00284-014-0629-1>
- Cochet N, Widehem P (2000) Ice crystallization by *Pseudomonas syringae*. *Appl Microbiol Biotechnol* 54(2):153–161
- Collicec Jouault S, Chevolut L, Helley D, Ratiskol J, Bros A, Sinquin C, Roger O, Fischer A-M (2001) Characterization, chemical modifications and in vitro anticoagulant properties of an exopolysaccharide produced by *Alteromonas infernus*. *Biochim Biophys Acta Gen Subj* 1528(2):141–151. [https://doi.org/10.1016/S0304-4165\(01\)00185-4](https://doi.org/10.1016/S0304-4165(01)00185-4)
- Collins T, Gerday C (2017) Enzyme catalysis in psychrophiles. In: Margesin R (ed) *Psychrophiles: from biodiversity to biotechnology*, 2nd edn. Springer, Cham, pp 209–235
- Collins T, Claverie P, D’Amico S, Georgette D, Gratia E, Hoyoux A, Meuwis MA, Poncin J, Sonan G, Feller G, Gerday C (2002a) Life in the cold: psychrophilic enzymes. In: Pandalai SG (ed) *Recent research developments in proteins*, vol 1. Transworld Research Network, Trivandrum, pp 13–26
- Collins T, Meuwis MA, Stals I, Claeysens M, Feller G, Gerday C (2002b) A novel family 8 xylanase, functional and physicochemical characterization. *J Biol Chem* 277(38):35133–35139. <https://doi.org/10.1074/jbc.M204517200>
- Collins T, Hoyoux A, Dutron A, Georis J, Genot B, Dauvrin T, Arnaut F, Gerday C, Feller G (2006) Use of glycoside hydrolase family 8 xylanases in baking. *JCS* 43:79–84
- Collins T, Feller G, Gerday C, Meuwis MA (2012) Family 8 enzymes with xylanolytic activity. US Granted Patent US8309336 B2
- Corsaro MM, Piazz FD, Lanzetta R, Parrilli M (2002) Lipid A structure of *Pseudoalteromonas haloplanktis* TAC 125: use of electrospray ionization tandem mass spectrometry for the determination of fatty acid distribution. *J Mass Spectrom* 37(5):481–488. <https://doi.org/10.1002/jms.304>
- Corsaro MM, Casillo A, Parrilli E, Tutino ML (2017) Molecular structure of lipopolysaccharides of cold-adapted bacteria. In: Margesin R (ed) *Psychrophiles: from biodiversity to biotechnology*, 2nd edn. Springer, Cham, pp 285–303
- D’Amico S, Claverie P, Collins T, Feller G, Georgette D, Gratia E, Hoyoux A, Meuwis MA, Zecchinon L, Gerday C (2001) Cold-adapted enzymes: an unachieved symphony. In: Storey KB, Storey JM (eds) *Cell and molecular responses to stress*, vol 2, protein adaptations and signal transduction, vol 3. Elsevier, Amsterdam, pp 31–42
- D’Amico S, Marx JC, Gerday C, Feller G (2003) Activity-stability relationships in extremophilic enzymes. *J Biol Chem* 278(10):7891–7896
- D’Amico S, Collins T, Marx JC, Feller G, Gerday C (2006) Psychrophilic microorganisms: challenges for life. *EMBO Rep* 7(4):385–389
- De Maayer P, Anderson D, Cary C, Cowan DA (2014) Some like it cold: understanding the survival strategies of psychrophiles. *EMBO Rep* 15(5):508–517. <https://doi.org/10.1002/embr.201338170>
- Deming JW, Young JN (2017) The role of exopolysaccharides in microbial adaptation to cold habitats. In: Margesin R (ed) *Psychrophiles: from biodiversity to biotechnology*, 2nd edn. Springer, Cham, pp 259–284
- Dieser M, Greenwood M, Foreman CM (2010) Carotenoid pigmentation in Antarctic heterotrophic bacteria as a strategy to withstand environmental stresses. *Arct Antarct Alp Res* 42(4):396–405. <https://doi.org/10.1657/1938-4246-42.4.396>
- Duchaud E, Boussaha M, Loux V, Bernardet J-F, Michel C, Kerouault B, Mondot S, Nicolas P, Bossy R, Caron C, Bessières P, Gibart J-F,

- Claverol S, Dumetz F, Hénaff ML, Benmansour A (2007) Complete genome sequence of the fish pathogen *Flavobacterium psychrophilum*. Nat Biotechnol 25:763–769. <https://doi.org/10.1038/nbt1313>
- Duman JG, Bennett V, Sformo T, Hochstrasser R, Barnes BM (2004) Antifreeze proteins in Alaskan insects and spiders. J Insect Physiol 50(4):259–266. <https://doi.org/10.1016/j.jinsphys.2003.12.003>
- Dutron A, Georis J, Genot B, Dauvrin T, Collins T, Hoyoux A, Feller G (2012) Use of family 8 enzymes with xylanolytic activity in baking. US Granted Patent US8192772
- Ewert M, Deming JW (2013) Sea ice microorganisms: environmental constraints and extracellular responses. Biology 2(2):603–628. <https://doi.org/10.3390/biology2020603>
- Feng S, Powell SM, Wilson R, Bowman JP (2014) Extensive gene acquisition in the extremely psychrophilic bacterial species *Psychroflexus torquus* and the link to sea-ice ecosystem specialism. Genome Biol Evol 6(1):133–148. <https://doi.org/10.1093/gbe/evt209>
- Ferrer M, Chernikova TN, Yakimov MM, Golyshin PN, Timmis KN (2003) Chaperonins govern growth of *Escherichia coli* at low temperatures. Nat Biotechnol 21:1266–1267. <https://doi.org/10.1038/nbt1103-1266>
- Fields PA, Dong Y, Meng X, Somero GN (2015) Adaptations of protein structure and function to temperature: there is more than one way to ‘skin a cat’. J Exp Biol 218(Pt 12):1801–1811. <https://doi.org/10.1242/jeb.114298>
- Fonseca F, Meneghel J, Cenard S, Passot S, Morris GJ (2016) Determination of intracellular vitrification temperatures for unicellular microorganisms under conditions relevant for cryopreservation. PLoS One 11(4):e0152939. <https://doi.org/10.1371/journal.pone.0152939>
- Forster J (1887) Ueber einige Eigenschaften leuchtender Bakterien. Centr Bakteriell Parasitenk 2:337–340
- Frank S, Schmidt F, Klockgether J, Davenport CF, Gesell Salazar M, Volker U, Tummler B (2011) Functional genomics of the initial phase of cold adaptation of *Pseudomonas putida* KT2440. FEMS Microbiol Lett 318(1):47–54. <https://doi.org/10.1111/j.1574-6968.2011.02237.x>
- Frommeyer M, Wiefel L, Steinbüchel A (2016) Features of the biotechnologically relevant polyamide family “cyanophycins” and their biosynthesis in prokaryotes and eukaryotes. Crit Rev Biotechnol 36(1):153–164. <https://doi.org/10.3109/07388551.2014.946467>
- Gaiteri JC, Henley WH, Siegfried NA, Linz TH, Ramsey JM (2017) Use of ice-nucleating proteins to improve the performance of freeze-thaw valves in microfluidic devices. Anal Chem 89(11):5998–6005. <https://doi.org/10.1021/acs.analchem.7b00556>
- Gao H, Yang ZK, Wu L, Thompson DK, Zhou J (2006) Global transcriptome analysis of the cold shock response of *Shewanella oneidensis* MR-1 and mutational analysis of its classical cold shock proteins. J Bacteriol 188(12):4560–4569. <https://doi.org/10.1128/JB.01908-05>
- Gerday C (2013) Psychrophily and catalysis. Biology 2(2):719–741. <https://doi.org/10.3390/biology2020719>
- Gesheva V, Stackebrandt E, Vasileva-Tonkova E (2010) Biosurfactant production by halotolerant *Rhodococcus fascians* from Casey station, Wilkes Land, Antarctica. Curr Microbiol 61(2):112–117. <https://doi.org/10.1007/s00284-010-9584-7>
- Ghobakhlu A-F, Johnston A, Harris L, Antoun H, Laberge S (2015) Microarray transcriptional profiling of Arctic *Mesorhizobium* strain N33 at low temperature provides insights into cold adaption strategies. BMC Genomics 16(1):383. <https://doi.org/10.1186/s12864-015-1611-4>
- Godin-Roulling A, Schmidpeter PAM, Schmid FX, Feller G (2015) Functional adaptations of the bacterial chaperone trigger factor to extreme environmental temperatures. Environ Microbiol 17(7):2407–2420. <https://doi.org/10.1111/1462-2920.12707>
- Goh YS, Tan IKP (2012) Polyhydroxyalkanoate production by Antarctic soil bacteria isolated from Casey Station and Signy Island. Microbiol Res 167(4):211–219. <https://doi.org/10.1016/j.micres.2011.08.002>
- Goordial J, Raymond-Bouchard I, Zolotarov Y, de Bethencourt L, Ronholm J, Shapiro N, Woyke T, Stromvik M, Greer CW, Bakermans C, Whyte L (2016) Cold adaptive traits revealed by comparative genomic analysis of the eurypsychrophile *Rhodococcus* sp. JG3 isolated from high elevation McMurdo Dry Valley permafrost, Antarctica. FEMS Microbiol Ecol 92(2). <https://doi.org/10.1093/femsec/fiv154>
- He J, Yang Z, Hu B, Ji X, Wei Y, Lin L, Zhang Q (2015) Correlation of polyunsaturated fatty acids with the cold adaptation of *Rhodotorula glutinis*. Yeast 32(11):683–690. <https://doi.org/10.1002/yea.3095>
- Jagannadham MV, Chattopadhyay MK, Subbalakshmi C, Vairamani M, Narayanan K, Rao CM, Shivaji S (2000) Carotenoids of an Antarctic psychrotolerant bacterium, *Sphingobacterium antarcticus*, and a mesophilic bacterium, *Sphingobacterium multivorum*. Arch Microbiol 173(5–6):418–424
- Jung HC, Lebeault JM, Pan JG (1998) Surface display of *Zymomonas mobilis* levansucrase by using the ice-nucleation protein of *Pseudomonas syringae*. Nat Biotechnol 16(6):576–580. <https://doi.org/10.1038/nbt0698-576>
- Kirk O, Christensen MW (2002) Lipases from *Candida antarctica*: unique biocatalysts from a unique origin. Org Process Res Dev 6(4):446–451. <https://doi.org/10.1021/op0200165>
- Kirti K, Amita S, Priti S, Mukesh Kumar A, Jyoti S (2014) Colorful world of microbes: carotenoids and their applications. Adv Biol 2014:13. <https://doi.org/10.1155/2014/837891>
- Kitamoto D, Yanagishita H, Endo A, Nakaiwa M, Nakane T, Akiya T (2001) Remarkable antiagglomeration effect of a yeast biosurfactant, diacylmannosylerythritol, on ice-water slurry for cold thermal storage. Biotechnol Prog 17(2):362–365. <https://doi.org/10.1021/bp000159f>
- Koh HY, Park H, Lee JH, Han SJ, Sohn YC, Lee SG (2017) Proteomic and transcriptomic investigations on cold-responsive properties of the psychrophilic Antarctic bacterium *Psychrobacter* sp. PAMC 21119 at subzero temperatures. Environ Microbiol 19(2):628–644. <https://doi.org/10.1111/1462-2920.13578>
- Krembs C, Eicken H, Deming JW (2011) Exopolymer alteration of physical properties of sea ice and implications for ice habitability and biogeochemistry in a warmer Arctic. Proc Natl Acad Sci U S A 108(9):3653–3658. <https://doi.org/10.1073/pnas.1100701108>
- Kumar A, Rao KM, Han SS (2018) Application of xanthan gum as polysaccharide in tissue engineering: a review. Carbohydr Polym 180:128–144. <https://doi.org/10.1016/j.carbpol.2017.10.009>
- Leroy F, De Vuyst L (2016) Advances in production and simplified methods for recovery and quantification of exopolysaccharides for applications in food and health. J Dairy Sci 99(4):3229–3238. <https://doi.org/10.3168/jds.2015-9936>
- Li J, Izquierdo MP, Lee T-C (1997) Effects of ice-nucleation active bacteria on the freezing of some model food systems. Int J Food Sci Technol 32(1):41–49. <https://doi.org/10.1046/j.1365-2621.1997.00380.x>
- Lim J, Thomas T, Cavicchioli R (2000) Low temperature regulated DEAD-box RNA helicase from the Antarctic archaeon, *Methanococcoides burtonii*. J Mol Biol 297(3):553–567
- Lonhienne T, Gerday C, Feller G (2000) Psychrophilic enzymes: revisiting the thermodynamic parameters of activation may explain local flexibility. Biochim Biophys Acta 1543(1):1–10
- López NI, Pettinari MJ, Stackebrandt E, Tribelli PM, Pötter M, Steinbüchel A, Méndez BS (2009) *Pseudomonas extremaustralis* sp. nov., a poly(3-hydroxybutyrate) producer isolated from an Antarctic environment. Curr Microbiol 59(5):514–519. <https://doi.org/10.1007/s00284-009-9469-9>

- Lorv JS, Rose DR, Glick BR (2014) Bacterial ice crystal controlling proteins. *Scientifica* 2014:976895. <https://doi.org/10.1155/2014/976895>
- Madihalli C, Sudhakar H, Doble M (2016) Mannosylerythritol lipid-A as a pour point depressant for enhancing the low-temperature fluidity of biodiesel and hydrocarbon fuels. *Energy Fuel* 30(5):4118–4125. <https://doi.org/10.1021/acs.energyfuels.6b00315>
- Makled SO, Hamdan AM, El-Sayed A-FM, Hafez EE (2017) Evaluation of marine psychrophile, *Psychrobacter namhaensis* SO89, as a probiotic in *Nile tilapia* (*Oreochromis niloticus*) diets. *Fish Shellfish Immunol* 61:194–200. <https://doi.org/10.1016/j.fsi.2017.01.001>
- Malavenda R, Rizzo C, Michaud L, Gerçe B, Bruni V, Sylдатk C, Hausmann R, Lo Giudice A (2015) Biosurfactant production by Arctic and Antarctic bacteria growing on hydrocarbons. *Polar Biol* 38(10):1565–1574. <https://doi.org/10.1007/s00300-015-1717-9>
- Margesin R (2017) Psychrophiles: from biodiversity to biotechnology, 2nd edn. Springer, Cham
- Margesin R, Collins T (2019) Microbial ecology of the cryosphere (glacial and permafrost habitats): current knowledge. *Appl Microbiol Biotechnol*. <https://doi.org/10.1007/s00253-019-09631-3>
- Marx JG, Carpenter SD, Deming JW (2009) Production of cryoprotectant extracellular polysaccharide substances (EPS) by the marine psychrophilic bacterium *Colwellia psychrerythraea* strain 34H under extreme conditions. *Can J Microbiol* 55(1):63–72. <https://doi.org/10.1139/W08-130>
- Medigue C, Krin E, Pascal G, Barbe V, Bernsel A, Bertin PN, Cheung F, Cruveiller S, D'Amico S, Duilio A, Fang G, Feller G, Ho C, Mangenot S, Marino G, Nilsson J, Parrilli E, Rocha EP, Rouy Z, Sekowska A, Tutino ML, Vallenet D, von Heijne G, Danchin A (2005) Coping with cold: the genome of the versatile marine Antarctica bacterium *Pseudoalteromonas haloplanktis* TAC125. *Genome Res* 15(10):1325–1335
- Méthé BA, Nelson KE, Deming JW, Momen B, Melamud E, Zhang X, Moulit J, Madupu R, Nelson WC, Dodson RJ, Brinkac LM, Daugherty SC, Durkin AS, DeBoy RT, Kolonay JF, Sullivan SA, Zhou L, Davidsen TM, Wu M, Huston AL, Lewis M, Weaver B, Weidman JF, Khouri H, Utterback TR, Feldblyum TV, Fraser CM (2005) The psychrophilic lifestyle as revealed by the genome sequence of *Colwellia psychrerythraea* 34H through genomic and proteomic analyses. *Proc Natl Acad Sci U S A* 102(31):10913–10918. <https://doi.org/10.1073/pnas.0504766102>
- Miyake R, Kawamoto J, Wei Y-L, Kitagawa M, Kato I, Kurihara T, Esaki N (2007) Construction of a low-temperature protein expression system using a cold-adapted bacterium, *Shewanella* sp. strain Ac10, as the host. *Appl Environ Microbiol* 73(15):4849–4856. <https://doi.org/10.1128/aem.00824-07>
- More TT, Yadav JSS, Yan S, Tyagi RD, Surampalli RY (2014) Extracellular polymeric substances of bacteria and their potential environmental applications. *J Environ Manag* 144:1–25. <https://doi.org/10.1016/j.jenvman.2014.05.010>
- Morgan-Kiss RM, Priscu JC, Pockock T, Gudynaite-Savitch L, Huner NPA (2006) Adaptation and acclimation of photosynthetic microorganisms to permanently cold environments. *Microbiol Mol Biol Rev* 70(1):222–252. <https://doi.org/10.1128/mmb.70.1.222-252.2006>
- Muñoz PA, Márquez SL, González-Nilo FD, Márquez-Miranda V, Blamey JM (2017) Structure and application of antifreeze proteins from Antarctic bacteria. *Microb Cell Factories* 16:138. <https://doi.org/10.1186/s12934-017-0737-2>
- Muralidharan J, Jayachandran S (2003) Physicochemical analyses of the exopolysaccharides produced by a marine biofouling bacterium, *Vibrio alginolyticus*. *Process Biochem* 38(6):841–847. [https://doi.org/10.1016/S0032-9592\(02\)00021-3](https://doi.org/10.1016/S0032-9592(02)00021-3)
- Mykytczuk NCS, Foote SJ, Omelon CR, Southam G, Greer CW, Whyte LG (2013) Bacterial growth at -15°C ; molecular insights from the permafrost bacterium *Planococcus halocryophilus* Or1. *ISME J* 7:1211–1226. <https://doi.org/10.1038/ismej.2013.8>
- Nishida T, Morita N, Yano Y, Orikasa Y, Okuyama H (2007) The anti-oxidative function of eicosapentaenoic acid in a marine bacterium, *Shewanella marinintestina* IK-1. *FEBS Lett* 581(22):4212–4216. <https://doi.org/10.1016/j.febslet.2007.07.065>
- Ochsenreither K, Gluck C, Stressler T, Fischer L, Sylдатk C (2016) Production strategies and applications of microbial single cell oils. *Front Microbiol* 7:1539. <https://doi.org/10.3389/fmicb.2016.01539>
- Okuyama H, Orikasa Y, Nishida T (2008) Significance of antioxidative functions of eicosapentaenoic and docosahexaenoic acids in marine microorganisms. *Appl Environ Microbiol* 74(3):570–574. <https://doi.org/10.1128/aem.02256-07>
- Oswald VF, Chen W, Harvilla PB, Magyar JS (2014) Overexpression, purification, and enthalpy of unfolding of ferricytochrome c552 from a psychrophilic microorganism. *J Inorg Biochem* 131:76–78. <https://doi.org/10.1016/j.jinorgbio.2013.11.002>
- Pandey R, Usui K, Livingstone RA, Fischer SA, Pfandtner J, Backus EHG, Nagata Y, Fröhlich-Nowoisky J, Schmäser L, Mauri S, Scheel JF, Knopf DA, Pöschl U, Bonn M, Weidner T (2016) Ice-nucleating bacteria control the order and dynamics of interfacial water. *Sci Adv* 2(4). <https://doi.org/10.1126/sciadv.1501630>
- Pandey N, Jain R, Pandey A, Tamta S (2018) Optimisation and characterisation of the orange pigment produced by a cold adapted strain of *Penicillium* sp. (GBPI_P155) isolated from mountain ecosystem. *Mycology* 9(2):81–92. <https://doi.org/10.1080/21501203.2017.1423127>
- Pärnänen K, Karkman A, Virta M, Eronen-Rasimus E, Kaartokallio H (2015) Discovery of bacterial polyhydroxyalkanoate synthase (PhaC)-encoding genes from seasonal Baltic Sea ice and cold estuarine waters. *Extremophiles* 19(1):197–206. <https://doi.org/10.1007/s00792-014-0699-9>
- Parrilli E, Tutino ML (2017) Heterologous protein expression in *Pseudoalteromonas haloplanktis* TAC125. In: M R (ed) *Psychrophiles: from biodiversity to biotechnology*, Second edition, vol 2. Springer, Cham, pp 513–525
- Perfumo A, Banat IM, Marchant R (2018) Going green and cold: biosurfactants from low-temperature environments to biotechnology applications. *Trends Biotechnol* 36(3):277–289. <https://doi.org/10.1016/j.tibtech.2017.10.016>
- Piette F, D'Amico S, Mazzucchelli G, Danchin A, Leprince P, Feller G (2011) Life in the cold: a proteomic study of cold-repressed proteins in the Antarctic bacterium *Pseudoalteromonas haloplanktis* TAC125. *Appl Environ Microbiol* 77(11):3881–3883. <https://doi.org/10.1128/aem.02757-10>
- Pummer BG, Budke C, Augustin-Bauditz S, Niedermeier D, Felgitsch L, Kampf CJ, Huber RG, Liedl KR, Loerting T, Moschen T, Schauerperl M, Tollinger M, Morris CE, Wex H, Grothe H, Pöschl U, Koop T, Fröhlich-Nowoisky J (2015) Ice nucleation by water-soluble macromolecules. *Atmos Chem Phys* 15(8):4077–4091. <https://doi.org/10.5194/acp-15-4077-2015>
- Raymond JA, Christner BC, Schuster SC (2008) A bacterial ice-binding protein from the Vostok ice core. *Extremophiles* 12(5):713–717. <https://doi.org/10.1007/s00792-008-0178-2>
- Raymond-Bouchard I, Goordial J, Zolotarov Y, Ronholm J, Stromvik M, Bakermans C, Whyte LG (2018) Conserved genomic and amino acid traits of cold adaptation in subzero-growing Arctic permafrost bacteria. *FEMS Microbiol Ecol* 94(4). <https://doi.org/10.1093/femsec/fiy023>
- Regand A, Goff HD (2006) Ice recrystallization inhibition in ice cream as affected by ice structuring proteins from winter wheat grass. *J Dairy Sci* 89(1):49–57. [https://doi.org/10.3168/jds.S0022-0302\(06\)72068-9](https://doi.org/10.3168/jds.S0022-0302(06)72068-9)
- Rodrigues DF, Ivanova N, He Z, Huebner M, Zhou J, Tiedje JM (2008) Architecture of thermal adaptation in an *Exiguobacterium sibiricum* strain isolated from 3 million year old permafrost: a genome and

- transcriptome approach. *BMC Genomics* 9:547. <https://doi.org/10.1186/1471-2164-9-547>
- Romero-Romero ML, Inglés-Prieto A, Ibarra-Molero B, Sanchez-Ruiz JM (2011) Highly anomalous energetics of protein cold denaturation linked to folding-unfolding kinetics. *PLoS One* 6(7):e23050–e23050. <https://doi.org/10.1371/journal.pone.0023050>
- Roulling F, Godin A, Cipolla A, Collins T, Miyazaki K, Feller G (2016) Activity-stability relationships revisited in blue oxidases catalyzing electron transfer at extreme temperatures. *Extremophiles* 20(5):621–629. <https://doi.org/10.1007/s00792-016-0851-9>
- Russell NJ (1997) Psychrophilic bacteria—molecular adaptations of membrane lipids. *Comp Biochem Physiol A Physiol* 118:489–493
- Russell NJ (2008) Membrane components and cold sensing. In: Margesin R, Schinner F, Marx J-C, Gerday C (eds) *Psychrophiles: from biodiversity to biotechnology*. Springer Berlin Heidelberg, Berlin, pp 177–190
- Santiago M, Ramírez-Sarmiento CA, Zamora RA, Parra LP (2016) Discovery, molecular mechanisms, and industrial applications of cold-active enzymes. *Front Microbiol* 7(1408). <https://doi.org/10.3389/fmicb.2016.01408>
- Sarmiento F, Peralta R, Blamey JM (2015) Cold and hot extremozymes: industrial relevance and current trends. *Front Bioeng Biotechnol* 3:148. <https://doi.org/10.3389/fbioe.2015.00148>
- Shen L, Liu Y, Wang N, Jiao N, Xu B, Liu X (2018) Variation with depth of the abundance, diversity and pigmentation of culturable bacteria in a deep ice core from the Yuzhufeng Glacier, Tibetan Plateau. *Extremophiles* 22(1):29–38. <https://doi.org/10.1007/s00792-017-0973-8>
- Shulse CN, Allen EE (2011) Diversity and distribution of microbial long-chain fatty acid biosynthetic genes in the marine environment. *Environ Microbiol* 13(3):684–695. <https://doi.org/10.1111/j.1462-2920.2010.02373.x>
- Siddiqui KS (2015) Some like it hot, some like it cold: temperature dependent biotechnological applications and improvements in extremophilic enzymes. *Biotechnol Adv* 33(8):1912–1922. <https://doi.org/10.1016/j.biotechadv.2015.11.001>
- Siddiqui KS, Williams TJ, Wilkins D, Yau S, Allen MA, Brown MV, Lauro FM, Cavicchioli R (2013) Psychrophiles. *Annu Rev Earth Planet Sci* 41(1):87–115. <https://doi.org/10.1146/annurev-earth-040610-133514>
- Singh AK, Sad K, Singh SK, Shivaji S (2014) Regulation of gene expression at low temperature: role of cold-inducible promoters. *Microbiology* 160(7):1291–1296. <https://doi.org/10.1099/mic.0.077594-0>
- Singh A, Krishnan KP, Prabakaran D, Sinha RK (2017) Lipid membrane modulation and pigmentation: a cryoprotection mechanism in Arctic pigmented bacteria. *J Basic Microbiol* 57(9):770–780. <https://doi.org/10.1002/jobm.201700182>
- Singh AK, Srivastava JK, Chandel AK, Sharma L, Mallick N, Singh SP (2019) Biomedical applications of microbially engineered polyhydroxyalkanoates: an insight into recent advances, bottlenecks, and solutions. *Appl Microbiol Biotechnol*. <https://doi.org/10.1007/s00253-018-09604-y>
- Soldatou S, Baker BJ (2017) Cold-water marine natural products, 2006 to 2016. *Nat Prod Rep* 34(6):585–626. <https://doi.org/10.1039/c6np00127k>
- Stritzler M, Diez Tissera A, Soto G, Ayub N (2018) Plant growth-promoting bacterium *Pseudomonas fluorescens* FR1 secretes a novel type of extracellular polyhydroxybutyrate polymerase involved in abiotic stress response in plants. *Biotechnol Lett* 40(9):1419–1423. <https://doi.org/10.1007/s10529-018-2576-6>
- Sun Y-Z, Yang H-L, Ma R-L, Zhang C-X, Lin W-Y (2011) Effect of dietary administration of *Psychrobacter* sp. on the growth, feed utilization, digestive enzymes and immune responses of grouper *Epinephelus coioides*. *Aquac Nutr* 17(3):e733–e740. <https://doi.org/10.1111/j.1365-2095.2010.00837.x>
- Sweet CR, Watson RE, Landis CA, Smith JP (2015) Temperature-dependence of lipid A acyl structure in *Psychrobacter cryohalolentis* and Arctic isolates of *Colwellia hornerae* and *Colwellia piezophila*. *Mar Drugs* 13(8):4701–4720. <https://doi.org/10.3390/md13084701>
- Ting L, Williams TJ, Cowley MJ, Lauro FM, Guilhaus M, Raftery MJ, Cavicchioli R (2010) Cold adaptation in the marine bacterium, *Sphingopyxis alaskensis*, assessed using quantitative proteomics. *Environ Microbiol* 12(10):2658–2676. <https://doi.org/10.1111/j.1462-2920.2010.02235.x>
- Tribelli PM, Lopez NI (2018) Reporting key features in cold-adapted bacteria. *Life (Basel)* 8(1). <https://doi.org/10.3390/life8010008>
- Tribelli PM, Solar Venero EC, Ricardi MM, Gómez-Lozano M, Raiger Iustman LJ, Molin S, López NI (2015) Novel essential role of ethanol oxidation genes at low temperature revealed by transcriptome analysis in the Antarctic bacterium *Pseudomonas extremaustralis*. *PLoS One* 10(12):e0145353. <https://doi.org/10.1371/journal.pone.0145353>
- Varin T, Lovejoy C, Jungblut AD, Vincent WF, Corbeil J (2012) Metagenomic analysis of stress genes in microbial mat communities from Antarctica and the High Arctic. *Appl Environ Microbiol* 78(2):549–559. <https://doi.org/10.1128/AEM.06354-11>
- Voets IK (2017) From ice-binding proteins to bio-inspired antifreeze materials. *Soft Matter* 13(28):4808–4823. <https://doi.org/10.1039/c6sm02867e>
- Vollmers J, Voget S, Dietrich S, Gollnow K, Smits M, Meyer K, Brinkhoff T, Simon M, Daniel R (2013) Poles apart: Arctic and Antarctic *Octadecabacter* strains share high genome plasticity and a new type of xanthorhodopsin. *PLoS One* 8(5):e63422. <https://doi.org/10.1371/journal.pone.0063422>
- Vollú RE, Jurelevicius D, Ramos LR, Peixoto RS, Rosado AS, Seldin L (2014) Aerobic endospore-forming bacteria isolated from Antarctic soils as producers of bioactive compounds of industrial interest. *Polar Biol* 37(8):1121–1131. <https://doi.org/10.1007/s00300-014-1505-y>
- Wani SH, Singh NB, Haribhushan A, Mir JI (2013) Compatible solute engineering in plants for abiotic stress tolerance—role of glycine betaine. *Curr Genomics* 14(3):157–165. <https://doi.org/10.2174/1389202911314030001>
- Wanka KM, Damerau T, Costas B, Krueger A, Schulz C, Wuertz S (2018) Isolation and characterization of native probiotics for fish farming. *BMC Microbiol* 18(1):119–119. <https://doi.org/10.1186/s12866-018-1260-2>
- Yoshida K, Hashimoto M, Hori R, Adachi T, Okuyama H, Orikasa Y, Nagamine T, Shimizu S, Ueno A, Morita N (2016) Bacterial long-chain polyunsaturated fatty acids: their biosynthetic genes, functions, and practical use. *Mar Drugs* 14(5):94. <https://doi.org/10.3390/md14050094>