

Reliability of striatal [^{11}C]raclopride binding in smokers wearing transdermal nicotine patches

Karmen K. Yoder · Daniel S. Albrecht · David A. Kareken · Lauren M. Federici · Kevin M. Perry · Elizabeth A. Patton · Qi-Huang Zheng · Bruce H. Mock · Sean J. O'Connor · Christine M. Herring

Received: 24 June 2011 / Accepted: 29 September 2011 / Published online: 19 October 2011
© Springer-Verlag 2011

Abstract

Purpose In studies where [^{11}C]raclopride (RAC) positron emission tomography (PET) is used to assess changes in striatal dopamine, it is important to control for cognitive states, such as drug craving, that could alter dopamine levels. In cigarette smokers, transdermal nicotine patches (TNP) can control nicotine craving, but the effects of nicotine patches on RAC binding are unknown. Thus, we sought to determine the test-retest reliability of RAC binding in the presence of nicotine patches.

Methods Eleven male smokers were scanned twice with RAC on separate days while wearing TNP.

Results Across the striatum, test-retest variability was 7.63 ± 5.88 ; percent change in binding potential was 1.11 ± 9.83 ; and the intraclass correlation coefficient was 0.91 ($p < 0.0001$).

Conclusion Baseline RAC binding is highly reproducible in smokers wearing nicotine patches. This suggests that TNP are an acceptable method for controlling cigarette craving during studies that utilize RAC to examine changes in dopamine.

K. K. Yoder · D. S. Albrecht · D. A. Kareken · L. M. Federici · K. M. Perry · E. A. Patton · Q.-H. Zheng · B. H. Mock · C. M. Herring
Indiana University Center for Neuroimaging,
Indiana University School of Medicine,
Indianapolis, IN, USA

K. K. Yoder (✉) · D. S. Albrecht · D. A. Kareken · L. M. Federici · K. M. Perry · E. A. Patton · Q.-H. Zheng · B. H. Mock · C. M. Herring
Department of Radiology and Imaging Sciences,
Indiana University School of Medicine,
R2, E124, 950 W. Walnut St.,
Indianapolis, IN 46202, USA
e-mail: kkyoder@iupui.edu

K. K. Yoder · D. S. Albrecht · D. A. Kareken · L. M. Federici
Stark Neurosciences Research Institute,
Indiana University School of Medicine,
Indianapolis, IN, USA

D. A. Kareken
Department of Neurology, Indiana University School of Medicine,
Indianapolis, IN, USA

D. A. Kareken · S. J. O'Connor
Department of Psychiatry, Indiana University School of Medicine,
Indianapolis, IN, USA

Keywords Transdermal nicotine patch · Dopamine · D_2 receptor · [^{11}C]Raclopride · Positron emission tomography · Test-retest variability

Introduction

It is possible to assess in vivo changes in relative dopamine levels with positron emission tomography (PET) and dopaminergic radioligands that are sensitive to alterations in endogenous dopamine concentration [1–4] (e.g., [^{11}C]raclopride, [^{18}F]fallypride, and [^{11}C]FLB). The goal of such studies is the comparison of dopamine D_2 receptor availability during a test condition relative to a baseline or resting condition, with the difference in D_2 signal between these two states attributed to changes in dopamine levels. There are many types of challenge paradigms (cognitive, motor, pharmacologic), methods of PET acquisition (single bolus, bolus plus infusion), and study designs (one or two scans). Regardless of the scientific question being asked, or the particulars of the methodological approach, all experiments share a critical assumption: *within* subjects, the baseline state represents a stable comparator for the challenge condition [2, 5]. Unfortunately, this assumption

can be easily violated by transient changes in cognitive states that may alter dopamine [1, 2, 5]. When possible, it would be desirable to control internal states that may contribute to confounding measurements of D₂ availability.

Several studies indicate that the internal state of drug craving is related to striatal dopamine levels [6–10]. For example, both Volkow et al. [6] and Wong et al. [7] presented cocaine users with cocaine-related cues and found that cue-induced striatal dopamine release correlated with self-reported cue-induced craving. This evidence strongly supports the concept that increases in striatal dopamine levels are related to drug craving. In 2004, Heinz and colleagues [10] reported decreased D₂ availability in the ventral striatum was negatively correlated with alcohol craving severity. It is reasonable to view lower D₂ availability as a function of higher dopamine levels, and therefore this report is consistent with the idea that higher dopamine concentration may be associated with drug craving. On the other hand, Brody et al. [8, 9] reported that smoking-induced reductions in nondisplaceable binding potential (BP_{ND}) were associated with *decreased* craving for cigarettes and proposed that increases in dopamine alleviate cigarette craving. In this case, however, the presence of chemosensory sensations and habitual motor routines of cigarette smoking may have increased dopamine independently of craving. Thus, in studies of populations with high rates of cigarette smoking, differences in nicotine withdrawal, and hence craving between scan states, present a serious challenge for data interpretation. Although transdermal nicotine patches could be used to control nicotine craving in smokers, it is not known whether stable baseline measurements of D₂ availability are possible with nicotine patches. Therefore, the purpose of this study was to determine the reliability of striatal [¹¹C] raclopride (RAC) binding in the presence of transdermal nicotine patches in nicotine-dependent cigarette smokers.

Materials and methods

All procedures were approved by the Indiana University Institutional Review Board and performed in accordance with the ethical standards of the Belmont Report (US

Department of Health and Human Services, 1979). All subjects signed informed consent statements agreeing to participate in the study. Subjects were right-handed, social drinking male smokers who were in otherwise good physical and mental health. The absence of either alcohol abuse or dependence was confirmed by the Semi-Structured Assessment for the Genetics of Alcoholism [11]. Subjects were excluded from participation if they endorsed recreational use of legal or illicit stimulants, painkillers, or sedatives, and/or consumption of >1 marijuana cigarette (or equivalent) per week. Subject demographics are presented in Table 1. Nicotine dependence was assessed with the Fagerström Test for Nicotine Dependence [12].

Study procedures

Procedures in this study are similar to those described previously [13]. Subjects underwent identical procedures on 2 separate days. Figure 1 illustrates the general time line. Briefly, subjects presented to the Indiana Clinical Research Center at approximately 8 a.m. Shortly after arrival, an IV catheter was placed in an antecubital vein, and a transdermal nicotine patch was placed on the upper arm of each subject. Patch dose was based on subjects' self-report of cigarettes smoked per day (14 mg dose with 10<20 cigarettes per day; 21 mg dose with >20 cigarettes). Two subjects were given a 14-mg patch; all others received a 21-mg patch. Subjects were given a full breakfast. As part of another study protocol, the morning of each study day, subjects received an IV alcohol infusion to a target breath alcohol concentration (BrAC) of 60 mg% using the alcohol clamp method. The rationale and implementation of the alcohol clamping technique have been described in detail elsewhere [14, 15]. The BrAC of all subjects returned to 0 mg% prior to scanning. The Cigarette Withdrawal Scale (CWS) [16], a self-report Likert rating scale, was given periodically throughout each study day. Nicotine craving was measured with the second dimension on the CWS, which specifically captures the individual's current subjective state of cigarette craving. There are four questions on this dimension, each with a 5-point scale; possible scores for cigarette craving range from 4 to 20. Ratings were taken upon arrival for the study (time 1), and before and after the

Table 1 Subject demographic, smoking, and drinking characteristics

<i>n</i>	Age	Race	Ethnicity	Education (years)	Fagerström score	D/DDay	D/Wk
11	35.0±10 (25–55)	3 AA, 8 C	0 H/L	12.6±1.92 (9–15)	4.55±1.51 (3–8)	3.27±0.99 (1.23–4.31)	5.01±3.02 (0.39–10.9)

Data expressed as mean ± SD with range in parentheses. The average Fagerström score indicates moderate nicotine dependence in this sample C Caucasian, AA African-American, H/L Hispanic/Latino, D/DDay drinks per drinking day, D/Wk drinks per week




Fig. 1 General outline of study day, starting with subject arrival at ~8 a.m. All times are approximate. *CWS* Cigarette Withdrawal Scale, *EtOH* alcohol, *MRI* magnetic resonance imaging, *RAC* [^{11}C]raclopride. Additional details are given in the text

resting (baseline) scan (times 2 and 3). Eleven subjects completed both baseline RAC scans. A 12th subject voluntarily withdrew from the study after the first RAC scan.

Scanning and reconstruction procedures

A magnetization prepared rapid acquisition gradient echo (MP-RAGE) magnetic resonance image (MRI) was acquired on all subjects using a Siemens 3T Trio for anatomic coregistration of PET data (see “[Image processing procedures](#)”).

Subjects received two baseline RAC scans in the early afternoon on 2 separate days. Time of injection was typically between 14:00 and 15:00. Breath alcohol concentration was 0 mg% prior to scanning. The time between the end of the morning alcohol infusion and the baseline RAC scan was typically ~4 h. RAC synthesis was completed as described previously [17]. RAC PET scans were acquired on a Siemens EXACT HR+(3-D mode; septa retracted). Prior to each PET scan, a 10-min transmission scan using three internal rod sources was acquired for attenuation correction. RAC PET scans were initiated with the IV infusion of 535 ± 45.4 MBq RAC (mass dose 0.13 ± 0.06 nmol/kg) over 1.5 min. Dynamic acquisition occurred for 50 min (time frames: 10×30 s, 45×60 s). Dynamic PET images were generated using Siemens Fourier rebinning (FORE) and filtered back-projection algorithms including corrections for attenuation, randoms, scatter, and dead time.

Image processing procedures

Image processing is similar to that described previously [13, 18]. MRI Digital Imaging and Communications in Medicine (DICOM) and RAC PET images were converted to Neuroimaging Informatics Technology Initiative (NIfTI) format (<http://nifti.nimh.nih.gov/>) using SPM5 (<http://www.fil.ion.ucl.ac.uk/spm/>). For each subject, dynamic PET data were coregistered to an early-time mean image to facilitate motion correction. The early mean PET image was coregistered to the MRI scan using the normalized mutual information algorithm in SPM5, with the transformation matrix from this coregistration subsequently applied to the

motion-corrected dynamic PET data. Each subject’s MRI was spatially normalized to Montreal Neurological Institute (MNI) space. The transformation matrix obtained from the spatial normalization step was then applied to the motion-corrected, MRI-registered PET data from each subject.

All regions of interest (ROIs) were drawn on an average normalized MRI from all subjects, using MRIcron (<http://www.sph.sc.edu/comd/rorden/mricron/>). Striatal ROIs consisted of the left and right ventral striatum, precommissural dorsal caudate, precommissural dorsal putamen, postcommissural caudate, and postcommissural putamen and were drawn according to specific anatomic landmarks [19, 20]. For the reference region (tissue that contains little to no D_2/D_3 receptor density), an ROI was created that contained all cerebellar gray matter except for the vermis. Time-activity curves for each ROI were generated from the dynamic RAC data using the MarsBaR toolbox for SPM5 (<http://marsbar.sourceforge.net/>). For each striatal ROI, D_2/D_3 receptor availability was indexed with BP_{ND} , the binding potential of RAC calculated as bound tracer concentration relative to nondisplaceable tracer concentration [21]. Estimations of BP_{ND} were conducted using the multilinear reference tissue model (MRTM) [22].

Metrics of test-retest reproducibility of baseline RAC binding

The relative reproducibility of striatal BP_{ND} between day 1 and day 2 was examined with three calculations, test-retest variability (TRV) to assess relative variation in BP_{ND} between days as a function of the overall average BP_{ND} across days, percent change in BP_{ND} (ΔBP_{ND}) as a qualitative descriptor of differences in BP_{ND} between day 1 and day 2, and the intraclass correlation coefficient (ICC, one-way random effects model; as implemented in the PASW statistical package [23, 24]), a metric of similarity between measurements on each day. TRV was calculated as: $|BP_{day1} - BP_{day2}| / [(BP_{day1} + BP_{day2}) / 2]$ [25, 26]. ΔBP between day 1 and day 2 was calculated as: $[(BP_{day1} - BP_{day2}) / BP_{day1}] \times 100$. Paired *t* tests were used to determine if striatal BP_{ND} was significantly different between scan days.

Other statistical tests

Independent *t* tests were used to test for differences in injected radioactivity and injected mass dose between scan days. To examine the stability of cigarette craving (CWS dimension 2 score), repeated-measures analysis of variance (ANOVA) (2 days \times 3 time points) was used to test for effects of scan day, time point, and day \times time point.

Results

RAC scan parameters

Average number of days between scans was 11.7 ± 25.2 (range 1–83 days). Injected radioactivity of RAC on day 1 and day 2 was 537 ± 54.7 and 533 ± 36.5 MBq, respectively. Corresponding mass doses were 0.13 ± 0.05 and 0.13 ± 0.06 nmol/kg. Injected radioactivity and mass doses were not significantly different between scan days.

Subject data

The demographic characteristics of the subjects are shown in Table 1. Eight subjects reported smoking a full pack of cigarettes per day; two reported a half-pack, and one reported two packs per day. Three subjects tested positive for marijuana on both scan days; these subjects endorsed sporadic recreational use of marijuana within the previous month. One subject tested positive for opiates on day 2 (subject had undergone a dental procedure and reported taking a single tablet of acetaminophen and No. 3 codeine phosphate 2 days prior). Two of these subjects had TRV for striatal RAC binding well within 1 SD of the sample average ($7.6 \pm 5.9\%$; Table 2); the other was within 1.5 SD of the mean TRV.

Exact timing of patch placement was not available for one subject on day 1. Across the remaining data points ($n=21$), the interval between patch placement and resting scan was




Fig. 2 Mean \pm SD cigarette craving ratings from day 1 (filled squares) and day 2 (open circles), and times 1, 2, and 3 (please see Fig. 1 for approximate timing of ratings). The x-axis crosses the y-axis at the value of 4 to denote that 4 is the lowest possible score on the index; for reference, 20 is the highest possible score. Craving ratings did not vary within subjects across either day or time point (see text for details)

5.9 ± 0.9 h. Ratings for the CWS were unavailable for one subject at both day 1-time 2 and day 2-time 3; for one subject at day 2-time 2; and for one subject at day 2-time 3. Cigarette craving scores are presented in Fig. 2. Repeated-measures ANOVA indicated that the CWS rating was stable within subjects, i.e., there were no main effects of day or time point, and no day \times time point interaction.

Table 2 Test-retest data from resting RAC scans in 11 cigarette smokers wearing transdermal nicotine patches

	BP _{ND} day 1	BP _{ND} day 2	% Δ	%TRV	ICC	ICC <i>p</i> value ^a
LpreDCA	2.21 \pm 0.22	2.08 \pm 0.16*	-5.51 \pm 4.85	8.43 \pm 5.67	0.41	0.09
RpreDCA	2.06 \pm 0.17	2.16 \pm 0.20	5.22 \pm 4.96	8.40 \pm 6.68	0.35	n.s.
LpreDPU	2.68 \pm 0.28	2.70 \pm 0.29	0.82 \pm 1.62	5.41 \pm 3.50	0.83	< 0.0005
RpreDPU	2.62 \pm 0.26	2.62 \pm 0.23	0.34 \pm 1.33	7.89 \pm 5.41	0.47	0.06
LVST	2.37 \pm 0.30	2.34 \pm 0.39	-0.94 \pm 0.01	11.2 \pm 6.57	0.67	< 0.01
RVST	2.25 \pm 0.26	2.35 \pm 0.33	4.30 \pm 4.70	6.75 \pm 5.84	0.77	< 0.005
LpostCA	1.70 \pm 0.19	1.64 \pm 0.22	-3.54 \pm 3.80	7.43 \pm 6.86	0.70	< 0.005
RpostCA	1.52 \pm 0.26	1.65 \pm 0.23**	9.50 \pm 9.65	9.50 \pm 7.40	0.74	< 0.005
LpostPU	2.81 \pm 0.21	2.90 \pm 0.24	3.31 \pm 3.09	5.50 \pm 4.69	0.59	< 0.05
RpostPU	2.80 \pm 0.20	2.73 \pm 0.26	-2.42 \pm 2.16	5.83 \pm 4.80	0.62	< 0.05
Striatum ^b	2.30 \pm 0.48	2.32 \pm 0.49	1.11 \pm 9.83	7.63 \pm 5.88	0.91	< 0.0001

Data expressed as mean \pm SD

BP_{ND} nondisplaceable binding potential, % Δ percent change in BP_{ND} from day 1 to day 2, %TRV percent test-retest variability, ICC intraclass correlation coefficient, L left, R right, pre precommissural, post postcommissural, DCA dorsal caudate, DPU dorsal putamen, VST ventral striatum, CA caudate, PU putamen, n.s. not significant

*Significantly different from day 1, paired *t* test, $p < 0.05$

**Significantly different from day 1, paired *t* test, $p < 0.01$

^aTrend-level significance is defined as $0.05 < p < 0.1$

^bAverage of all regions across all subjects ($n=110$)

Test-retest reproducibility of resting RAC signal

BP_{ND} values for both scan days, percent change of BP_{ND} between days (%Δ), the TRV, and ICC are presented in Table 2.

Discussion

In the current sample of 11 otherwise healthy cigarette smokers, there was good reproducibility of striatal RAC binding in the presence of transdermal nicotine patches. Overall, our test-retest metrics (Table 2) comport well with literature values for single-bolus RAC studies in healthy control subjects [25, 27–29]. The average striatal data were exceptionally stable, with slight variations in reliability between subregions. We chose to demonstrate test-retest reliability with three methods. The percent change in BP_{ND} between scans is the most commonly used parameter of effect size and corresponds to the calculation most commonly used to describe relative changes in dopamine levels between conditions. TRV quantifies the size of absolute variability across measurements. Both indices provide useful estimates of the effect sizes needed to achieve statistical significance with a two-scan dopamine challenge paradigm (although statistical significance is possible with smaller effect sizes). In this study, these assessments of RAC test-retest reliability suggest a range of variability consistent with what others have reported when scanning the same individuals across days. Overall, the ICC coefficient *r* values in this study were also quite good and indicate that estimations of BP_{ND} on day 1 correlated well with measurements on day 2. Lower ICC values were found in small regions that tend to have slightly noisier time-activity curves and hence more variable estimates of BP_{ND}.

One potential concern about the use of nicotine patches is the possibility that nicotine itself might cause measurable dopamine release [30, 31]. Two human RAC studies by Brody et al. support this view [8, 9]. However, those designs included the actual physical act of smoking cigarettes. When nicotine is delivered intranasally to humans, or intravenously to unanesthetized monkeys, there is no evidence of significant decreases in RAC binding [32, 33]. Taken together, these latter studies strongly suggest that nicotine itself does not release dopamine to levels measurable by RAC PET. It is also possible that the physical and sensory properties associated with cigarette smoking are the key components of the smoking-induced dopamine release reported by Brody et al.

In this study, the administration of IV alcohol to subjects the morning of the RAC PET study may be an unintended source of variance in baseline RAC BP_{ND}. In our previous work, we found no evidence of alcohol-induced dopamine

release in social drinkers [34, 35]. However, healthy social drinkers do not typically expose themselves to alcohol shortly after breakfast. We cannot exclude the possibility that alcohol exposure early in the morning may have caused unpredictable changes in the dopamine tone of healthy social drinkers.

Another limitation of this study was the absence of test-retest values for striatal RAC binding in smokers without nicotine patches, which could have assessed any variability in baseline striatal BP_{ND} attributable to nicotine withdrawal. However, the most important assumption in a typical dopaminergic RAC PET challenge paradigm is that the *within-subject state of basal dopamine is stable*. The present data demonstrate that it is possible to control nicotine craving (which is highly likely to alter endogenous dopamine) while keeping estimates of baseline D₂ availability stable.

Conclusion

The presence of transdermal nicotine patches does not appear to affect the stability of baseline RAC binding potential. We suggest that, in RAC challenge studies, the use of nicotine patches in smoking subjects is feasible to eliminate unwanted variance in D₂ availability caused by nicotine craving and concomitant alterations in striatal dopamine levels.

Acknowledgments This work was supported by: ABMRF/The Foundation for Alcohol Research (KKY), National Institute on Alcoholism and Alcohol Abuse P60 AA007611 Pilot 50 (KKY), and the Indiana Clinical and Translational Sciences Institute (from the National Institutes of Health, National Center for Research Resources, Clinical and Translational Sciences Award, grant #UL RR025761). The authors would like to thank Barbara Glick and Brandon Steele for assistance with radiochemistry.

Conflicts of interest None.

References

1. Egerton A, Mehta MA, Montgomery AJ, Lappin JM, Howes OD, Reeves SJ, et al. The dopaminergic basis of human behaviors: a review of molecular imaging studies. *Neurosci Biobehav Rev* 2009;33:1109–32. doi:10.1016/j.neubiorev.2009.05.005.
2. Yoder KK, Kareken DA, Morris ED. Assessing dopaminergic neurotransmission with PET: basic theory and applications in alcohol research. *Curr Med Imaging Rev* 2011;7:118–24.
3. Laruelle M. Imaging synaptic neurotransmission with in vivo binding competition techniques: a critical review. *J Cereb Blood Flow Metab* 2000;20:423–51.
4. Morris ED, Yoder KK. Positron emission tomography displacement sensitivity: predicting binding potential change for positron emission tomography tracers based on their kinetic characteristics. *J Cereb Blood Flow Metab* 2007;27:606–17. doi:10.1038/sj.jcbfm.9600359.
5. Yoder KK, Kareken DA, Morris ED. What were they thinking? Cognitive states may influence [(11)C]raclopride binding potential in the striatum. *Neurosci Lett* 2008;430:38–42.

6. Volkow ND, Wang GJ, Telang F, Fowler JS, Logan J, Childress AR, et al. Cocaine cues and dopamine in dorsal striatum: mechanism of craving in cocaine addiction. *J Neurosci* 2006;26:6583–8.
7. Wong DF, Kuwabara H, Schretlen DJ, Bonson KR, Zhou Y, Nandi A, et al. Increased occupancy of dopamine receptors in human striatum during cue-elicited cocaine craving. *Neuropsychopharmacology* 2006;31:2716–27.
8. Brody AL, Mandelkern MA, Olmstead RE, Scheibal D, Hahn E, Shiraga S, et al. Gene variants of brain dopamine pathways and smoking-induced dopamine release in the ventral caudate/nucleus accumbens. *Arch Gen Psychiatry* 2006;63:808–16.
9. Brody AL, Olmstead RE, London ED, Farahi J, Meyer JH, Grossman P, et al. Smoking-induced ventral striatum dopamine release. *Am J Psychiatry* 2004;161:1211–8.
10. Heinz A, Siessmeier T, Wrase J, Hermann D, Klein S, Grüsser-Sinopoli SM, et al. Correlation between dopamine D(2) receptors in the ventral striatum and central processing of alcohol cues and craving. *Am J Psychiatry* 2004;161:1783–9.
11. Buchholz KK, Cadoret R, Cloninger CR, Dinwiddie SH, Hesselbrock VM, Nurnberger Jr JJ, et al. A new, semi-structured psychiatric interview for use in genetic linkage studies: a report on the reliability of the SSAGA. *J Stud Alcohol* 1994;55:149–58.
12. Pomerleau CS, Carton SM, Lutzke ML, Flessland KA, Pomerleau OF. Reliability of the Fagerstrom Tolerance Questionnaire and the Fagerstrom Test for Nicotine Dependence. *Addict Behav* 1994;19:33–9.
13. Yoder KK, Albrecht DS, Kareken DA, Federici LM, Perry KM, Patton EA, et al. Test-retest variability of [(11)C]raclopride-binding potential in nontreatment-seeking alcoholics. *Synapse* 2011;65:553–61. doi:10.1002/syn.20874.
14. Ramchandani VA, Bolane J, Li TK, O'Connor S. A physiologically-based pharmacokinetic (PBPK) model for alcohol facilitates rapid BrAC clamping. *Alcohol Clin Exp Res* 1999;23:617–23.
15. O'Connor S, Morzorati S, Christian J, Li TK. Clamping breath alcohol concentration reduces experimental variance: application to the study of acute tolerance to alcohol and alcohol elimination rate. *Alcohol Clin Exp Res* 1998;22:202–10.
16. Etter JF. A self-administered questionnaire to measure cigarette withdrawal symptoms: the Cigarette Withdrawal Scale. *Nicotine Tob Res* 2005;7:47–57. doi:10.1080/14622200412331328501.
17. Fei X, Mock BH, DeGrado TR, Wang JQ, Glick-Wilson BE, Sullivan ML, et al. An improved synthesis of PET dopamine D2 receptors radioligand [(11)C]raclopride. *Synth Commun* 2004;34:1897–907.
18. Yoder KK, Morris ED, Constantinescu CC, Cheng TE, Normandin MD, O'Connor SJ, et al. When what you see isn't what you get: alcohol cues, alcohol administration, prediction error, and human striatal dopamine. *Alcohol Clin Exp Res* 2009;33:139–49. doi:10.1111/j.1530-0277.2008.00821.x.
19. Martinez D, Slifstein M, Broft A, Mawlawi O, Hwang DR, Huang Y, et al. Imaging human mesolimbic dopamine transmission with positron emission tomography. Part II: amphetamine-induced dopamine release in the functional subdivisions of the striatum. *J Cereb Blood Flow Metab* 2003;23:285–300.
20. Mai JK, Assheuer J, Paxinos G. Atlas of the human brain. San Diego: Academic; 1997.
21. Innis RB, Cunningham VJ, Delforge J, Fujita M, Gjedde A, Gunn RN, et al. Consensus nomenclature for in vivo imaging of reversibly binding radioligands. *J Cereb Blood Flow Metab* 2007;27:1533–9. doi:10.1038/sj.jcbfm.9600493.
22. Ichise M, Liow JS, Lu JQ, Takano A, Model K, Toyama H, et al. Linearized reference tissue parametric imaging methods: application to [(11)C]DASB positron emission tomography studies of the serotonin transporter in human brain. *J Cereb Blood Flow Metab* 2003;23:1096–112. doi:10.1097/01.WCB.0000085441.37552.CA.
23. McGraw KO, Wong SP. Forming inferences about some intraclass correlation coefficients. *Psychol Methods* 1996;1:30–46.
24. Shrout PE, Fleiss JL. Intraclass correlations: uses in assessing rater reliability. *Psychol Bull* 1979;86:420–8.
25. Hirvonen J, Aalto S, Lumme V, Någren K, Kajander J, Vilkmann H, et al. Measurement of striatal and thalamic dopamine D2 receptor binding with 11C-raclopride. *Nucl Med Commun* 2003;24:1207–14. doi:10.1097/01.mnm.0000104642.79626.e8.
26. Mawlawi O, Martinez D, Slifstein M, Broft A, Chatterjee R, Hwang DR, et al. Imaging human mesolimbic dopamine transmission with positron emission tomography: I. Accuracy and precision of D(2) receptor parameter measurements in ventral striatum. *J Cereb Blood Flow Metab* 2001;21:1034–57.
27. Volkow ND, Fowler JS, Wang GJ, Dewey SL, Schlyer D, MacGregor R, et al. Reproducibility of repeated measures of carbon-11-raclopride binding in the human brain. *J Nucl Med* 1993;34:609–13.
28. Hietala J, Någren K, Lehikoinen P, Ruotsalainen U, Syvälahti E. Measurement of striatal D2 dopamine receptor density and affinity with [(11)C]raclopride in vivo: a test-retest analysis. *J Cereb Blood Flow Metab* 1999;19:210–7.
29. Schlösser R, Brodie JD, Dewey SL, Alexoff D, Wang GJ, Fowler JS, et al. Long-term stability of neurotransmitter activity investigated with 11C-raclopride PET. *Synapse* 1998;28:66–70.
30. Nisell M, Nomikos GG, Svensson TH. Infusion of nicotine in the ventral tegmental area or the nucleus accumbens of the rat differentially affects accumbal dopamine release. *Pharmacol Toxicol* 1994;75:348–52.
31. Schiffer WK, Gerasimov MR, Marsteller DA, Geiger J, Barnett C, Alexoff DL, et al. Topiramate selectively attenuates nicotine-induced increases in monoamine release. *Synapse* 2001;42:196–8. doi:10.1002/syn.10000.
32. Montgomery AJ, Lingford-Hughes AR, Egerton A, Nutt DJ, Grasby PM. The effect of nicotine on striatal dopamine release in man: a [(11)C]raclopride PET study. *Synapse* 2007;61:637–45.
33. Tsukada H, Miyasato K, Kakiuchi T, Nishiyama S, Harada N, Domino EF. Comparative effects of methamphetamine and nicotine on the striatal [(11)C]raclopride binding in unanesthetized monkeys. *Synapse* 2002;45:207–12.
34. Yoder KK, Kareken DA, Seyoum RA, O'Connor SJ, Wang C, Zheng QH, et al. Dopamine D(2) receptor availability is associated with subjective responses to alcohol. *Alcohol Clin Exp Res* 2005;29:965–70.
35. Yoder KK, Constantinescu CC, Kareken DA, Normandin MD, Cheng TE, O'Connor SJ, et al. Heterogeneous effects of alcohol on dopamine release in the striatum: a PET study. *Alcohol Clin Exp Res* 2007;31:965–73.