

Clinical significance of abdominal scintigraphy using ^{99m}Tc -HMPAO-labelled leucocytes in patients with seronegative spondyloarthropathies

J.C. Alonso Farto¹, I. Almoguera Arias¹, F.J. Lopez Longo², C.M. Gonzalez Fernandez², I. Monteagudo Saez², A. Ortega Valle¹, M. Bascones², J.M. Pérez Vázquez¹, L. Carreño Perez²

¹ Department of Nuclear Medicine, "Hospital Universitario Gregorio Marañón", Universidad Complutense, Doctor Esquerdo, 46, 28007 Madrid, Spain

² Department of Rheumatology, "Hospital Universitario Gregorio Marañón", Universidad Complutense, Madrid, Spain

Published online: 9 February 2001

© Springer-Verlag 2001

Eur J Nucl Med (2001) 28:402

DOI 10.1007/s002590100493

Eur J Nucl Med (2000)

DOI 10.1007/s002590000393

The first sentence of the Introduction (p. 1768) erroneously appeared as "Spondyloarthropathies (Sp) are inflammatory multisystem diseases that present with rheumatoid factor...". In fact it should have read "Spondyloarthropathies (Sp) are inflammatory multisystem diseases that present *without* rheumatoid factor...".

The online version of the original article can be found at <http://doi.org/10.1007/s002590000393>

J.C. Alonso Farto (✉)

Department of Nuclear Medicine,
"Hospital Universitario Gregorio Marañón",
Universidad Complutense, Doctor Esquerdo, 46,
28007 Madrid, Spain

e-mail: jcaf@airtel.net or jcaf@jet.es

Tel.: +34-91-5868183, Fax: +34-91-5868469