

Erratum to: Evaluation of Pedicled Omental Flap Delivered through a Minilaparotomy for Immediate Breast Reconstruction in Obese Patients

Ashraf Khater

Published online: 19 March 2014

© Springer Science+Business Media New York and International Society of Aesthetic Plastic Surgery 2014

Erratum to: Aesth Plast Surg (2013) 37:1140–1145 DOI 10.1007/s00266-013-0217-y

In Table 2, there are five *n* (%) values in the right column—16(67) to 0—that were incorrectly aligned with the values in the left column starting with Patient satisfaction and ending with Not satisfied. These values have now been correctly aligned in this erratum.

Table 2 Operative data and postoperative outcome

Type of mastectomy	<i>n</i> (%)
NASM	17/24 (71)
SSM	7/24 (29)
Inadequate flap volume and implant usage	3/24 (13)
Total operative time, min (range)	177.92 (160–210)
Harvest time, min (range)	61.5 (50–75)
Blood loss, ml (range)	170.42 (80–300)
Blood loss with flap retrieval	–
Mean hospital stay, days (range)	3.8 (3–5)
Injury of the gastroepiploic vessels	0
Partial flap ischemia	1/24 (4)
Total flap ischemia	0
Incisional hernia	1/24 (4)
Hematoma	2/24 (8)
Wound infection	2/24 (8)
Minor deformity	5/24 (21)
Patient satisfaction	
Excellent	16 (67)
Fair	6 (25)
Good	2 (8)
Accepted	0
Not satisfied	0

NASM nipple-areola-sparing mastectomy, *SSM* skin-sparing mastectomy

The online version of the original article can be found under doi:[10.1007/s00266-013-0217-y](https://doi.org/10.1007/s00266-013-0217-y).

A. Khater (✉)
Mansoura Oncology Center (OCMU), Mansoura Faculty of
Medicine, Mansoura University, Gomhoria Street,
Mansoura 35511, Egypt
e-mail: dr.ashrafkhater@yahoo.com; ash_khater@mans.edu.eg