

Erratum to: Controlled Tissue Expansion in the Initial Management of the Short Bowel State

F. Murphy · B. A. Khalil · S. Gozzini ·
B. King · A. Bianchi · A. Morabito

Published online: 21 December 2011
© Société Internationale de Chirurgie 2011

Erratum to: World J Surg (2011) 35:1142–1145 DOI 10.1007/s00268-011-0991-0

The authors have brought to our attention the need for revision to Table 1 in their original article. While the overall conclusion of the article did not change, the correct bowel length values are represented in the table below.

Table 1 Patient diagnosis and pre and post bowel expansion lengths

Patient	Diagnosis	Pre expansion length (cm)	Post expansion length (cm)
1	Jejunal atresia	25	43
2	Gastroschisis	45	60
3	Gastroschisis	67	85
4	Vanishing Gastroschisis	30	45
5	Gastroschisis	25	25
6	Gastroschisis	25	25
7	Gastroschisis	34	34
8	Gastroschisis	10	10
9	Necrotising enterocolitis	5	8
10	Necrotising enterocolitis	30	35
Mean		29.6	37

Overall percentage increase in bowel length: 20%. Range of actual length of increase: 0–18 cm

The online version of the original article can be found under
doi:[10.1007/s00268-011-0991-0](https://doi.org/10.1007/s00268-011-0991-0).

F. Murphy (✉) · B. A. Khalil · S. Gozzini · B. King ·
A. Bianchi · A. Morabito
Department of Paediatric Surgery, Royal Manchester Children's
Hospital, Oxford Road, Manchester M13 9WL, UK
e-mail: Fiona.murphy@doctors.org.uk