

The role of PGE₂-associated inflammatory responses in gastric cancer development

Hiroko Oshima · Masanobu Oshima

Received: 1 September 2012 / Accepted: 30 September 2012 / Published online: 11 October 2012
© Springer-Verlag Berlin Heidelberg 2012

Abstract Accumulating evidence indicates that inflammation plays a critical role in cancer development. Cyclooxygenase-2 (COX-2) is a rate-limiting enzyme for prostanoid biosynthesis, including prostaglandin E₂ (PGE₂), and plays a key role in both inflammation and cancer. It has been demonstrated that inhibition of COX-2 and PGE₂ receptor signaling results in the suppression of tumor development in a variety of animal models. However, the molecular mechanisms underlying COX-2/PGE₂-associated inflammation in carcinogenesis have not yet been fully elucidated. In order to study the role of PGE₂-associated inflammatory responses in tumorigenesis, it is important to use in vivo mouse models that recapitulate human cancer development from molecular mechanisms with construction of tumor microenvironment. We have developed a gastritis model (*K19-C2mE* mice) in which an inflammatory microenvironment is constructed in the stomach via induction of the COX-2/PGE₂ pathway. We also developed a gastric cancer mouse model (*Gan* mice) in which the mice develop inflammation-associated gastric tumors via activation of both the COX-2/PGE₂ pathway and Wnt signaling. Expression analyses using these in vivo models have revealed novel mechanisms of the inflammatory responses underlying gastric cancer development. PGE₂-associated inflammatory responses activate epidermal growth factor receptor (EGFR) signaling through the induction of EGFR ligands and ADAMs that release EGFR ligands from the cell membrane. In *Gan* mice, a combination treatment with EGFR and COX-2 inhibitors

significantly suppresses gastric tumorigenesis. Moreover, PGE₂-associated inflammation downregulates tumor suppressor microRNA, miR-7, in gastric cancer cells, which suppresses epithelial differentiation. These results indicate that PGE₂-associated inflammatory responses promote in vivo gastric tumorigenesis via several different molecular mechanisms.

Keywords Gastric cancer · Inflammation · COX-2 · PGE₂ · EGFR · MicroRNA

Introduction

It has been shown that approximately 15–20 % of malignant cancers are associated with chronic infection [1, 2]. For example, infection with *Helicobacter pylori*, hepatitis C virus, and human papilloma virus is closely associated with the development of gastric cancer, hepatocellular carcinoma, and cervical cancer, respectively. On the other hand, it has also been reported that tobacco smoke and obesity contribute to tumor development through induction of inflammatory responses in the lungs and liver, respectively [3, 4]. These results, together with those of recent genetic studies, indicate that inflammatory responses play important roles in cancer development regardless of whether they are caused by infectious or noninfectious stimuli [5–7]. Among the various inflammatory networks involved in tumor microenvironment, the cyclooxygenase-2 (COX-2)/prostaglandin E₂ (PGE₂) pathway was first identified as a key player in tumorigenesis [8, 9]. COX-2 is an inducible rate-limiting enzyme for prostaglandin biosynthesis and has been shown to play an essential role in inflammatory responses. PGE₂ is a downstream product of COX-2, and the level of PGE₂ increases significantly in tumor tissues, suggesting that PGE₂ plays a role in

This article is a contribution to the special issue on Inflammation and Cancer - Guest Editor: Takuji Tanaka

H. Oshima · M. Oshima (✉)
Division of Genetics, Cancer Research Institute,
Kanazawa University,
Kakuma-machi,
Kanazawa 920-1192, Japan
e-mail: oshimam@staff.kanazawa-u.ac.jp

tumorigenesis. Although almost 20 years have passed since the discovery of the role of COX-2 in cancer development, the mechanisms underlying the role of the COX-2/PGE₂ pathway in tumorigenesis have not yet been fully elucidated. It was recently shown that COX-2 expression and PGE₂ production are induced in mesenchymal stem cells infiltrated in cancer tissues, which leads to the expression of inflammatory cytokines and the induction of epithelial mesenchymal transition through the construction of niches for tumor-initiating cells [10]. However, it remains unclear how the COX-2/PGE₂ pathway promotes in vivo tumorigenesis. To understand the role of PGE₂-induced inflammatory responses in cancer development, it is important to construct animal models in which the animals develop cancer caused by oncogenic activation in epithelial cells together with PGE₂-associated host responses like human cancers. We developed a gastric cancer mouse model (*Gan* mice for gastric neoplasia mice) in which activation of both Wnt signaling and the COX-2/PGE₂ pathway in the stomach is achieved by the transgenic expression of *Wnt1*, *Ptgs2*, and *Ptgs1*, which encode Wnt1, COX-2, and mPGES-1, respectively [11, 12]. By conducting expression analyses of mRNA and microRNA (miRNA) using murine gastritis tissues and *Gan* mouse gastric tumors, we identified novel molecular pathways that are activated or suppressed in PGE₂-associated inflammatory microenvironment and thereby contribute to gastric cancer development. In the first part of this article, we review the role of the COX-2/PGE₂ pathway in gastrointestinal cancer development and the construction of the *Gan* mouse model. Then, we discuss the novel mechanisms underlying inflammation in gastric tumorigenesis, which we found using the *Gan* mouse model system.

The COX-2/PGE₂ pathway in gastrointestinal tumor development

Approximately 20 years ago, epidemiological studies reported that the regular use of non-steroidal anti-inflammatory drugs (NSAIDs) lowers the mortality rate of patients with gastrointestinal cancer [13, 14]. The target molecules of NSAIDs are COX-1 and COX-2, which are rate-limiting enzymes for prostaglandin biosynthesis. COX-1 is constitutively expressed and plays a physiological house-keeping role, while COX-2 expression is induced in both inflammation and cancer. It has also been reported that treatment of patients with familial adenomatous polyposis (FAP) using NSAIDs results in significant decreases in the number and size of colon polyps [15]. These results strongly suggested that COX-2 plays an important role in intestinal tumorigenesis. Moreover, a large number of animal experiments have confirmed that NSAID treatment suppresses chemical carcinogen-induced colon tumor development, indicating that induction of the COX pathway is required for tumor development [16]. Mouse genetic studies clearly demonstrate the role of COX-2 in intestinal tumorigenesis. Disruption of the *Ptgs2* gene in *Apc*^{Δ716} and *Apc*^{Min} mice, mouse models for FAP, significantly suppresses the development of intestinal polyposis, thus indicating an essential role for COX-2 in tumor development [17, 18]. Interestingly, disruption of *Ptgs1* (COX-1 gene) also suppresses intestinal polyposis in *Apc*^{Min} mice. It is therefore possible that the COX-1-derived basal level of PGE₂ is also involved in tumor formation before COX-2 expression is induced [16] (Fig. 1).




Fig. 1 Schematic diagram showing the role of PGE₂ in tumorigenesis. Wnt signaling activation triggers initiation of tumorigenesis. Expression of COX-2 and mPGES-1 is induced in microadenomas, which results in an increase of PGE₂ level. Activated COX-2/PGE₂ pathway contributes to tumor development through a variety of mechanisms,


including induction of angiogenesis and Wnt activation, or construction of inflammatory microenvironment as indicated in Figs. 4 and 5. COX-1-derived PGE₂ may play a role in tumorigenesis at the early stage of tumorigenesis when COX-2 expression is not yet induced

Microsomal PGE synthase-1 (mPGES-1) is an inducible PGE-converting enzyme that functionally couples with COX-2 for PGE₂ biosynthesis, and the expression of mPGES-1 is induced in cancer tissues together with that of COX-2 [19, 20]. Accordingly, the PGE₂ level is increased in cancer tissues due to the expression of both COX-2 and mPGES-1. Importantly, a disruption of the *Ptges* gene encoding mPGES-1 in *Apc*^{Δ14} mice, another FAP model, and in a chemical carcinogen-treated colon cancer mouse model results in significant decreases in the PGE₂ level in the intestinal mucosa, which causes further suppression of intestinal tumor development [21, 22]. There are four G-protein coupled receptors for PGE₂, namely, EP1, EP2, EP3, and EP4. Signaling through EP2 and EP4 increases intracellular cyclic AMP and has been shown to be important for tumor development. Disruption of the *Ptger2* gene encoding EP2 in *Apc*^{Δ716} mice causes significant suppression of intestinal polyp development, whereas disruption of *Ptger1* and *Ptger3* does not affect the tumor phenotype [23]. EP2 signaling accelerates angiogenesis in intestinal polyps through the induction of the vascular endothelial growth factor (VEGF) and basic fibroblast growth factor expression [24]. Moreover, EP2 signaling has been shown to activate Wnt signaling in colon cancer cells through the activation of the PI3K/Akt pathway and the direct association between G protein and Axin, thus resulting in the stabilization of β-catenin [25]. These results indicate that the COX-2/mPGES-1/PGE₂/EP2 pathway is important for tumor development

through the involvement of several different mechanisms (Fig. 1). Although several studies have identified possible mechanisms underlying the COX-2/PGE₂ pathway in cancer development [8, 9], the key function of the COX-2/PGE₂ pathway in in vivo cancer development has not yet been fully elucidated.

The expression of proinflammatory cytokines, including TNF-α and IL-6, is induced in the inflammatory microenvironment of cancer, and mouse genetic studies have shown that these cytokines also promote tumorigenesis. TNF-α was originally recognized as a tumor necrosis factor; however, it is now established that TNF-α exhibits important tumor-promoting functions [26]. Inhibition of TNF-α signaling by disruption of TNF-α or its receptor gene results in the suppression of skin carcinogenesis and inflammation-associated colon cancer development [27–29]. TNF-α activates transcription factor NF-κB, which further induces the expression of inflammatory factors, including COX-2, IL-6, IL-8, and TNF-α itself, forming an inflammatory network in tumor microenvironment [30] (Fig. 2). Genetic inhibition of NF-κB in colitis-associated colon cancer model mice results in significant suppression of tumor development [31]. On the other hand, IL-6 expression is induced in colon cancer [32]. IL-6 signaling activates Stat3 through the gp130 receptor. Notably, disruption of IL-6 or Stat3 genes in colitis-associated colon cancer model mice results in significant suppression of colonic tumors [33, 34]. Taken together, these results indicate that the COX-2/PGE₂, TNF-α/NF-κB, and

Fig. 2 Schematic diagram showing inflammatory network in tumor microenvironment. Induction of COX-2/PGE₂ pathway is responsible for the construction of an inflammatory microenvironment where macrophages are recruited. NF-κB is activated in tumor-associated macrophages, resulting in the induction of inflammatory factors including COX-2, TNF-α, and IL-6. TNF-α and IL-6 activate the transcription factors NF-κB and Stat3, respectively, forming an inflammatory network in the tumor microenvironment. Such network plays a critical role in tumor promotion



IL-6/Stat3 pathways construct the inflammatory tumor microenvironment required for the promotion of tumorigenesis [30] (Fig. 2).

Inflammation-associated gastric cancer mouse model

Preneoplastic changes induced by oncogenic Wnt signaling

Based on the results of the role of the COX-2/PGE₂ pathway in cancer development, we developed a PGE₂-associated gastritis and gastric cancer mouse models as follows: Canonical Wnt signaling (Wnt/ β -catenin signaling) is important for the undifferentiated status of epithelial cells and thus induces tumorigenesis [35, 36]. Mutations in APC or β -catenin genes result in constitutive activation of canonical Wnt signaling, which causes tumor development in the entire intestinal tract [37, 38]. Nuclear or cytoplasmic accumulation of β -catenin, a hallmark of Wnt signaling activation, is detected in 30–50 % of gastric cancer tissues [39–42]. Moreover, mutations in the β -catenin gene have been found in a subpopulation of gastric cancers [39, 40, 43, 44]. These results indicate that Wnt signaling activation is an important oncogenic pathway for gastric cancer development.

To examine the role of Wnt activation in gastric tumorigenesis, we created *K19-Wnt1* transgenic mice that express Wnt1, a ligand for canonical Wnt signaling, using the *K19* gene promoter that is transcriptionally active in gastric epithelial cells [42]. Although *K19-Wnt1* mice do not develop gastric tumors, they do develop small aberrant cryptic foci on the surface of the gastric mucosa (Fig. 3a). Histologically, these lesions consist of dysplastic epithelial cells with increased cell proliferation and cytoplasmic β -catenin accumulation and are thus diagnosed as preneoplastic lesions [11, 12]. Importantly, macrophages infiltrate the stroma of preneoplastic lesions, while tissue macrophages are rarely detected in normal gastric mucosa (Fig. 3a). These results suggest that inflammatory microenvironments consisting of macrophages are constructed in the early stage of tumorigenesis and may be required for tumorous changes in Wnt-activated epithelial cells. To support this hypothesis, we demonstrated that macrophage-derived TNF- α promotes Wnt signaling activity in epithelial cells, which contributes to the development of preneoplastic lesions in *K19-Wnt1* mice [45].

Gastritis induced by the inflammatory COX-2/PGE₂ pathway

On the other hand, the COX-2/PGE₂ pathway is widely induced in gastric cancer, as found in colon cancer tissues [46]. *H. pylori* infection is tightly associated with gastric

cancer, and the levels of the COX-2 expression and PGE₂ production are related to the status of *H. pylori* infection [47, 48]. Accordingly, it is possible that the induction of the COX-2/PGE₂ pathway is an important tumor-promoting mechanism of *H. pylori* infection. To examine the role of the COX-2/PGE₂ pathway in gastric tumorigenesis, we created another group of transgenic mice, *K19-C2mE* mice, that expresses *Ptgs2* and *Ptges* encoding COX-2 and mPGES-1, respectively, in the gastric mucosa [49]. *K19-C2mE* mice develop gastric hyperplasia and mucous cell metaplasia, which are histologically similar to spasmodic polypeptide/trefoil factor 2 (TFF2)-expressing metaplasia (SPEM) (Fig. 3b) [49, 50]. SPEM is associated with human gastric cancers [51, 52], thus suggesting that SPEM is a possible precursor of gastric cancer [53]. SPEM development has also been observed to occur in the stomachs of several different gastric tumor model mice, including *Helicobacter*-infected mice, gastrin gene knockout mice, and gp130 receptor-active mutant mice [54–56], suggesting that the COX-2/PGE₂ pathway is responsible for SPEM formation in human gastric cancer as well as in these mouse models.

Hyperplastic lesions in *K19-C2mE* mice are associated with submucosal infiltration of lymphocytes, granulocytes, and mononuclear cells. Moreover, macrophage accumulation is also observed in the gastric mucosa (Fig. 3b). Notably, α -smooth muscle-expressing myofibroblasts are detected in *K19-C2mE* mouse inflamed gastric mucosa, and approximately 10 % of myofibroblasts is derived from bone marrow [57]. Myofibroblasts are also important components of the tumor microenvironment, together with tumor-associated macrophages. Accordingly, it is possible that induction of the COX-2/PGE₂ pathway is responsible for the construction of inflammatory microenvironment consisting of macrophages and myofibroblasts. Importantly, disruption of the TNF- α gene in *K19-C2mE* mice results in suppression of gastritis and SPEM development, while IL-1 receptor or Rag2 gene disruption does not affect these phenotypes [50]. Accordingly, it is possible that TNF- α induced in the COX-2/PGE₂-associated inflammatory microenvironment plays a key role in SPEM formation.

Gastric cancer induced by Wnt and the COX-2/PGE₂ pathway

By crossing *K19-Wnt1* and *K19-C2mE* transgenic mice, compound *K19-Wnt1/C2mE* transgenic mice were created that express *Wnt1*, *Ptgs2*, and *Ptges* simultaneously in the gastric epithelial cells, resulting in activation of Wnt signaling and the COX-2/PGE₂ pathway simultaneously in the stomach [42]. Importantly, *K19-Wnt1/C2mE* mice develop large gastric tumors at around 40 to 50 weeks of age with 100 % incidence (Fig. 3c). We thus named the *K19-Wnt1/*




Fig. 3 Gastric phenotypes by Wnt activation, COX-2/PGE₂ induction, or both in combination. **a** *K19-Wnt1* mice expressing *Wnt1* in gastric epithelial cells develop preneoplastic lesions by Wnt activation (*left, right top*). Expression of *Wnt1* gene in *K19-Wnt1* mouse stomach was confirmed by RT-PCR (*left, bottom*). F4/80-positive macrophages are infiltrated in the stroma of preneoplastic lesion (*right, bottom*). **b** *K19-C2mE* mice express *Ptgs2* and *Ptges* in the stomach, resulting in the induction of the COX-2/PGE₂ pathway, and they develop SPEM lesion consisting of metaplastic mucous cells (*left*). F4/80-positive macrophages are accumulated in metaplastic and hyperplastic gastric mucosa

(*right*). Expression of *Ptgs2* and *Ptges* in the glandular stomach was confirmed by Northern blotting (*bottom*). **c** *Gan* mice expressing *Wnt1*, *Ptgs2*, and *Ptges* in the stomach develop large gastric tumors (*left*) caused by the activation of both Wnt signaling and COX-2/PGE₂ pathway. Wild-type mouse stomach is shown as a control of the same magnification (*right*) (RT-PCR shown in **a** was reproduced from Oshima et al. [42] with permission from Elsevier, and Western blotting shown in **b** was reproduced from Oshima et al. [49] with permission from Nature Publishing)

C2mE mice as *Gan* mice, which stands for gastric neoplasia mice. Histologically, gastric tumors in *Gan* mice consist of dysplastic epithelial cells with an irregularly branching glandular structure, thus diagnosed as glandular-type gastric tumors. Moreover, the Ki-67 labeling index is significantly increased, indicating accelerated proliferation of tumor cells. Mucosal macrophage infiltration is also found in *Gan* mouse tumors, and the expression of proinflammatory cytokines and chemokines increases significantly. Accordingly, these results, taken together, indicate that Wnt signaling activation in the COX-2/PGE₂-associated inflammatory microenvironment cooperatively causes the development of gastric cancer. Importantly, the gene expression profiles of

Gan mouse tumors are similar to those found in human intestinal-type gastric cancer [58], thus indicating that *Gan* mice recapitulate the development of human intestinal-type gastric cancer from genetic changes and host inflammatory responses to tumor morphology and gene expression profiles.

In the inflamed tissues, reactive oxygen species (ROS) and nitric oxide are induced, which play a role in tumor development. In the gastric tumor tissues of *Gan* mice, ROS level is also increased [59]. However, we have recently demonstrated that the expression of CD44 variant form (CD44v) is significantly induced in gastritis of *K19-C2mE* mice [60], and CD44v plays a critical role in tumor

development in *Gan* mice through protection of tumor cells from oxidative stress [59]. Accordingly, it is possible that ROS plays a role in tumorigenesis, and at the same time protection from ROS leads to tumor promotion.

Interestingly, macrophage infiltration and inflammatory responses in gastric tumors are significantly suppressed in germ-free *Gan* mice, although the COX-2/PGE₂ pathway is still activated [61]. Moreover, the gastric tumor volume decreases significantly in germ-free *Gan* mice. In contrast, *Helicobacter* infection in the stomachs of germ-free *Gan* mice induces inflammatory responses and the development of large gastric tumors. These results indicate that increased PGE₂ signaling is not sufficient to induce inflammatory responses in the stomach; however, bacterial infection together with the COX-2/PGE₂ pathway is required for the construction of the inflammatory microenvironment. It has been shown that Toll-like receptor (TLR) signaling is important for intestinal epithelial homeostasis, and the suppression of TLR signaling suppresses regenerative proliferation in injured mucosa [62]. Moreover, a disruption of the *Myd88* gene encoding MyD88, an important adaptor molecule of the TLR signaling pathway, results in the suppression of intestinal polyposis in *Apc^{Min}* mice with a decreased expression level of COX-2 [63]. Accordingly, it is possible that innate immune signaling activated by bacterial infection and the TLR/MyD88 pathway is required for the construction of the PGE₂-associated tumor microenvironment. On the other hand, it has been shown that epithelial cell-derived PGE₂ suppresses inflammatory cytokine expression in the LPS-stimulated bone marrow-derived cells, which is important for protection from bacterial infection-associated colitis [64]. It is therefore possible that PGE₂ has distinct functions on inflammatory cytokine expression in immune tolerance and in cancer microenvironment, although the mechanism underlying such difference remains to be investigated.


It is therefore expected that complicated interactions between tumor cells and the inflammatory microenvironment in human gastric cancer are repeated in *Gan* mouse gastric tumor tissues. Accordingly, using the *Gan* mouse model, we are able to uncover the mechanisms underlying PGE₂-associated inflammatory responses in gastric tumorigenesis. For example, we have determined that gastric tumor cells educate and activate stromal fibroblasts and bone marrow-derived cells to be myofibroblasts, resulting in the induction of the VEGF expression and enhancement of angiogenesis in tumor tissues [57]. Moreover, we examined the gene expression profiles of mRNAs and microRNAs (miRNAs) in *K19-C2mE* gastritis and *Gan* mouse gastric tumor tissues and found several novel mechanisms of inflammation in tumor development, as described in the following section.

Activation of epidermal growth factor receptor signaling by PGE₂-associated inflammation

EGFR signaling is an important target for cancer prevention [65]. The genetic or pharmacological inhibition of EGFR signaling in *Apc^{Min}* mice results in significant suppression of intestinal polyposis [66–68], indicating that EGFR signaling is required for tumor cell proliferation even when Wnt signaling is constitutively activated. Importantly, the combination treatment of *Apc^{Min}* mice with EGFR inhibitors and NSAIDs or COX-2 inhibitors suppresses intestinal polyposis more effectively than simple treatment, thus suggesting a possible link between the COX-2/PGE₂ pathway and EGFR signaling [67, 68]. It has already been shown using in vitro experiments that the PGE₂ pathway transactivates EGFR signaling through the activation of cSrc or MMPs [69–71] or the induction of EGFR ligands and a disintegrin and metalloproteinase 17 (ADAM17), a shedding enzyme for EGFR ligands [72–74]. We thus examined the mechanisms underlying EGFR activation by the COX-2/PGE₂ pathway in in vivo tumors using *K19-C2mE* and *Gan* mice.

We compared the expression levels of all EGFR ligands, their receptors, and shedding enzymes for EGFR ligands in gastritis and gastric tumors with those in normal stomachs using microarray results [58]. Surprisingly, the expression of most EGFR ligands, including *Areg*, *Ereg*, *Hbgef*, and *Btc*, encoding amphiregulin, epiregulin, HB-EGF, and betacellulin, respectively, as well as *ErbB2* and *ErbB3* encoding Her2 and Her3, respectively, increased significantly in both *K19-C2mE* mouse gastritis and *Gan* mouse gastric cancer compared with that observed in wild-type mice [75] (Fig. 4a). ADAM family proteases activate EGFR signaling by shedding the ectodomain of EGFR ligands to release them from the cell membrane, and some ADAM family genes are induced in a variety of cancer tissues, suggesting that ADAMs play a role in the activation of EGFR signaling in cancer development [76]. Notably, the expression of *Adam8*, *Adam9*, *Adam10*, and *Adam28* is increased significantly in the stomach in *K19-C2mE* gastritis and *Gan* mouse tumors. Taken together, these results indicate that the expression of EGFR ligand members, their receptors, and ADAM family members is upregulated simultaneously in the PGE₂-associated inflammatory microenvironment, which causes activation of EGFR signaling in cancer cells. Consistently, EGFR is phosphorylated in the epithelial cells of *K19-C2mE* and *Gan* mouse stomachs, indicating the activation of EGFR, which is not found in the wild-type mouse gastric mucosa (Fig. 4b). Among the four PGE₂ receptors, the expression of EP4 is significantly increased in the stomachs of both *K19-C2mE* and *Gan* mice [77]. Notably, treatment of *Gan* mice with EP4 receptor inhibitors results in significant decreases in the EGFR ligand and

Fig. 4 Activation of EGFR signaling in COX-2/PGE₂ pathway-induced inflammatory microenvironment. **a** Expression of EGFR ligands and Adam family proteases is upregulated in both *K19-C2mE* mouse gastritis and *Gan* mouse gastric tumors. **b** Phosphorylated EGFR is detected by immunohistochemistry (green) in gastric epithelial cells of *K19-C2mE* mice and *Gan* mice, which is not found in the wild-type mouse stomach. Nuclei are stained with DAPI (red). **c** X ray CT photographs of a *Gan* mouse that was treated with EGFR inhibitor, ZD1839, and COX-2 inhibitor, celecoxib, in combination for 3 weeks. Note that the gastric tumor volume decreased dramatically by drug treatment. **d** Schematic diagram of EGFR activation in the COX-2/PGE₂ pathway-associated inflammatory microenvironment. Induction of EGFR ligands and Adams cooperatively activates EGFR signaling, which further activates PGE₂ signaling through downregulation of 15-PGDH by positive feedback mechanism (a, b, and c were reproduced from Oshima et al. [73] with permission from Wiley-Blackwell)



ADAM expression levels [75], thus indicating that PGE₂ signaling via the EP4 receptor is responsible for EGFR activation in gastric epithelial cells.

EGFR signaling activates the MAPK and PI3K/Akt pathways, which results in phosphorylation of Erk1/2 and Akt [78]. Notably, the phosphorylation levels of Erk1/2 and Akt in *Gan* mouse gastric tumors are decreased significantly by treatment of the mice with the COX-2 inhibitor celecoxib. Accordingly, it is possible that the COX-2/PGE₂ pathway is a major cause of activation of EGFR signaling in gastric cancer cells. On the other hand, EGFR signaling can activate the COX-2/PGE₂ pathway as follows: The cellular PGE₂ level is regulated by 15-hydroxyprostaglandin dehydrogenase (15-PGDH), which catalyzes and thus inactivates prostaglandins. Importantly, the expression of 15-PGDH is downregulated by EGFR signaling [79], resulting in the maintenance of an increased PGE₂ level. Moreover, disruption of the 15-PGDH gene in *Apc^{Min}* mice accelerates intestinal tumorigenesis [80]. Accordingly, the COX-2/PGE₂ pathway activates EGFR signaling through the induction of EGFR ligands and ADAM proteases, which in turn

downregulates 15-PGDH, resulting in an increased level of PGE₂ in the inflammatory microenvironment. Such PGE₂/EGFR signaling feedback loops therefore play an important role in gastric cancer development (Fig. 4d).

Treatment of *Gan* mice with either EGFR inhibitor, ZD1839, or celecoxib significantly reduces gastric tumor volume [75] (Fig. 4c). Importantly, combination treatment of *Gan* mice with ZD1839 and celecoxib results in the complete regression of gastric tumors, possibly through inhibition of individual pathways as well as feedback loops between PGE₂ and EGFR. These results suggest that the use of such combination treatment is an effective preventive strategy against gastric cancer development.

Downregulation of tumor suppressor microRNA, miR-7, by PGE₂-associated inflammation

miRNAs are single-strand, small, noncoding RNAs that regulate gene expression via post-transcriptional interference of target mRNAs [81, 82]. Accordingly, miRNAs can

function as oncogenes or tumor suppressors through the inhibition of tumor suppressor genes or of oncogene expression, respectively [83, 84]. Several mechanisms for dysregulation of the miRNA expression have been demonstrated, such as genetic or epigenetic alterations or transcriptional or post-transcriptional mechanisms [85]. Notably, it has also been shown that inflammation induces the expression of oncogenic miRNAs [86]. For example, miR-155 and miR-21 are oncogenic miRNAs that are induced by NF- κ B, TLR, interferon- β , or the Stat3 pathway [87–89]. NF- κ B and Stat3 are activated by TNF- α and IL-6, respectively, and are key factors in the network of tumor microenvironment [30] (Fig. 2).

Microarray analyses indicate that 50 microRNAs are upregulated (>2.0 -fold) and 42 miRNAs are downregulated (<0.5 -fold) in *Gan* mouse gastric tumors [90] (Fig. 5a). Among these miRNAs, 21 and 29 show the same upregulation and downregulation, respectively, in *K19-C2mE* mouse gastritis, thus indicating that expression changes in these miRNAs (21 up and 29 down) in gastric tumors are caused by COX-2/PGE₂ pathway-associated inflammatory responses. Notably, the oncogene miRNAs miR-21 and miR-155 [91] are upregulated both in *K19-C2mE* mouse gastritis and *Gan* gastric tumors, whereas the tumor suppressor miRNAs miR-145 and miR-7 [92, 93] are downregulated in both tissues. Accordingly, it is possible that




Fig. 5 Downregulation of tumor suppressor miRNA, miR-7, in COX-2/PGE₂ pathway-induced inflammatory microenvironment. **a** Venn diagrams of upregulated and downregulated miRNAs in *K19-C2mE* mouse gastritis and *Gan* mouse gastric tumors. Note that 21 and 29 miRNAs are upregulated and downregulated, respectively, both in gastritis and gastric tumors, indicating inflammation-dependent expression changes. Notably, oncogene miRNAs, miR-21 and miR-155, are upregulated, whereas tumor suppressor miRNAs, miR-7 and miR-145, are downregulated simultaneously both in gastritis and gastric tumors.

b Inverse relations between the expression levels of miR-7 and inflammatory cytokines, TNF- α and IL-1 β , in human gastric cancer tissues: namely, cytokine high gastric cancers show lower miR-7 level, indicating that the severity of inflammatory responses is associated with miR-7 downregulation. **c** Schematic diagram of miR-7 downregulation in PGE₂-associated inflammatory microenvironment. Downregulation of miR-7 promotes gastric tumorigenesis possibly through suppression of epithelial differentiation (**a** and **b** were reproduced from Kong et al. [88] with permission from Nature Publishing)

inflammation promotes tumorigenesis by inducing oncogene miRNAs and suppressing tumor suppressor miRNAs simultaneously, possibly through different mechanisms.

It has been shown that miR-7 functions as a tumor suppressor in glioblastoma, breast cancer, and lung cancer [93–97]. The expression level of miR-7 is downregulated both in gastritis and gastric tumors, indicating that miR-7 is an inflammation-dependent tumor suppressor miRNA (Fig. 5a). Moreover, the miR-7 level is significantly decreased in *Helicobacter*-infected inflamed mouse gastric mucosa compared with control inflamed stomachs. In a normal stomach, the basal level of miR-7 is found in undifferentiated gastric epithelial cells; however, its expression level increases during the differentiation of epithelial cells. Accordingly, it is possible that miR-7 plays a role in cell differentiation.

In human gastric cancer tissues, miR-7 expression is downregulated in 65 % of cases. Moreover, the expression levels of miR-7 in human gastric cancers are inversely correlated with those of the inflammatory cytokines IL-1 β and TNF- α (Fig. 5b). Namely, miR-7 expression is downregulated more significantly in high TNF- α or high IL-1 β gastric cancers compared with low TNF- α or low IL-1 β gastric cancers. Accordingly, it is possible that the severity of inflammatory responses is correlated with effective downregulation of miR-7.

Moreover, transfection of precursor miR-7 in gastric cancer cells results in decreased cell proliferation and suppression of soft agar colony formation. These results indicate that miR-7 functions as a tumor suppressor in gastric cancer cells as well. These results, taken together, indicate that miR-7 is an important factor that links inflammation and cancer development, namely, inflammation induces downregulation of the miR-7 expression, which may suppress the differentiation of epithelial cells and thus increase tumorigenicity (Fig. 5c). It has been shown that EGFR mRNA is a miR-7 target gene [93, 95]. Consistently, in gastric cancer cells, precursor miR-7 transfection causes downregulation of the EGFR expression. Moreover, p21-activated kinase 1, Raf1, and insulin-like growth factor I receptor have been identified as possible miR-7 target genes that are upregulated in cancer cells [94–96]. Using the expression profile data set [58], we identified novel miR-7 target genes, including *Lphn2*, *Baspl1*, and *MafG*. It is possible that upregulation of these genes by miR-7 downregulation contributes to gastric tumorigenesis, although the role of these factors in cancer remains to be investigated in the future.

Conclusions

Accumulating evidence has established that inflammatory responses are important for cancer development. As discussed in this review article, in vivo model systems such as *K19-C2mE* and *Gan* model mice are important for investigating the role of inflammatory responses in cancer development

because the tumor-promoting microenvironment consisting of residential fibroblasts and bone marrow-derived cells is constructed in tumor tissues. Using such a model system, we identified several novel mechanisms underlying PGE₂-associated inflammation in gastric tumorigenesis, i.e., EGFR activation through the induction of EGFR ligands and ADAM proteases and downregulation of tumor suppressor miRNAs, including miR-7. On the other hand, most malignant cancer cells that carry accumulated genetic alterations do not require expression of COX-2 or EGFR ligands in stromal cells because oncogenic mutations cause induction or activation of these endogenous pathways in the tumor cells. For example, *ras* mutation causes constitutive expression of COX-2 and activation of EGFR signaling pathway. It is thus possible that PGE₂-associated inflammatory responses are more important for cancer development during promotion stage rather than malignant progression stage. Accordingly, regulation of such PGE₂-associated inflammatory responses will be an effective preventive strategy against gastric cancer development.

Acknowledgments We thank Manami Watanabe for her excellent work in the series of *Gan* mouse studies.

References

1. Kuper H, Adami HO, Trichopoulos D (2000) Infections as a major preventable cause of human cancer. *J Intern Med* 248:171–183
2. Parkin DM (2006) The global health burden of infection-associated cancers in the year 2002. *Int J Cancer* 118:3030–3044
3. Takahashi H, Ogata H, Nishigaki R, Broide DH, Karin M (2010) Tobacco smoke promotes lung tumorigenesis by triggering IKK β - and JNK1-dependent inflammation. *Cancer Cell* 17:89–97
4. Park EJ, Lee JH, Yu GY, He G, Ali SR, Holzer RG, Osterreicher CH, Takahashi H, Karin M (2010) Dietary and genetic obesity promote liver inflammation and tumorigenesis by enhancing IL-6 and TNF expression. *Cell* 140:197–208
5. Coussens LM, Werb Z (2002) Inflammation and cancer. *Nature* 420:860–867
6. Mantovani A, Allavena P, Sica A, Balkwill F (2008) Cancer-related inflammation. *Nature* 454:436–444
7. Grivninkov SI, Greten FR, Karin M (2010) Immunity, inflammation, and cancer. *Cell* 140:883–899
8. Wang D, DuBois RN (2010) Eicosanoids and cancer. *Nat Rev Cancer* 10:181–193
9. Wang D, DuBois RN (2010) The role of COX-2 in intestinal inflammation and colorectal cancer. *Oncogene* 29:781–788
10. Li HJ, Reinhardt F, Herschman HR, Weinberg RA (2012) Cancer-stimulated mesenchymal stem cells create a carcinoma stem-cell niche via prostaglandin E₂ signaling. *Cancer Discov* 2:840–855
11. Oshima H, Oguma K, Du YC, Oshima M (2009) Prostaglandin E₂, Wnt, and BMP in gastric tumor mouse models. *Cancer Sci* 100:1779–1785
12. Oshima H, Oshima M (2010) Mouse models of gastric tumors: Wnt activation and PGE₂ induction. *Pathol Int* 60:599–607
13. Thun MJ, Namboodiri MM, Heath CW Jr (1991) Aspirin use and reduced risk of fatal colon cancer. *N Engl J Med* 325:1593–1596

14. Giovannucci E, Egan KM, Hunter DJ, Stampfer MJ, Colditz GA, Willett WC, Speizer FE (1995) Aspirin and the risk of colorectal cancer in women. *N Engl J Med* 333:609–614
15. Giardiello FM, Hamilton SR, Krush AJ, Piantadosi S, Hyland LM, Celano P, Booker SV, Robinson CR, Offerhaus GJ (1993) Treatment of colonic and rectal adenomas with sulindac in familial adenomatous polyposis. *N Engl J Med* 328:1313–1316
16. Oshima M, Taketo MM (2002) COX selectivity and animal models for colon cancer. *Curr Pharm Des* 8:1021–1034
17. Oshima M, Dinchuk JE, Kargman SL, Oshima H, Hancock B, Kwong E, Trzaskos JM, Evans JF, Taketo MM (1996) Suppression of intestinal polyposis in *Apc^{Δ716}* knockout mice by inhibition of cyclooxygenase 2 (COX-2). *Cell* 87:803–809
18. Chulada PC, Thompson MB, Mahler JF, Doyle CM, Gaul BW, Lee C, Tiano HF, Morham SG, Smithies O, Langenbach R (2000) Genetic disruption of *Ptgs-1*, as well as of *Ptgs-2*, reduces intestinal tumorigenesis in *Min* mice. *Cancer Res* 60:4705–4708
19. Yoshimatsu K, Altorki NK, Golijanin D, Zhang F, Jakobsson PJ, Dannenberg AJ, Subbaramaiah K (2001) Inducible prostaglandin E synthase is overexpressed in non-small cell lung cancer. *Clin Cancer Res* 7:2669–2674
20. van Rees BP, Sivula A, Thoren S, Yokozaki H, Jalobsson PJ, Offerhaus GJ, Ristimäki A (2003) Expression of microsomal prostaglandin E synthase-1 in intestinal gastric adenocarcinoma and in gastric cancer cell lines. *Int J Cancer* 107:551–556
21. Nakanishi M, Montrose DC, Clark P, Nambiar PR, Belinsky GS, Claffey KP, Xu D, Rosenberg DW (2008) Genetic deletion of *mPGES-1* suppresses intestinal tumorigenesis. *Cancer Res* 68:3251–3259
22. Nakanishi M, Menoret A, Tanaka T, Miyamoto S, Montrose DC, Vella AT, Rosenberg DW (2011) Selective PGE₂ suppression inhibits colon carcinogenesis and modifies local mucosal immunity. *Cancer Prev Res* 4:1198–1208
23. Sonoshita M, Takaku K, Sasaki N, Sugimoto Y, Ushikubi F, Natumiya S, Oshima M, Taketo MM (2001) Acceleration of intestinal polyposis through prostaglandin receptor EP2 in *Apc^{Δ716}* knockout mice. *Nat Med* 7:1048–1051
24. Seno H, Oshima M, Ishikawa TO, Oshima H, Takaku K, Chiba T, Narumiya S, Taketo MM (2002) Cyclooxygenase 2- and prostaglandin E₂ receptor EP₂-dependent angiogenesis in *Apc^{Δ716}* mouse intestinal polyps. *Cancer Res* 62:506–511
25. Castellone MD, Teramoto H, Williams BO, Druey KM, Gutkind JS (2005) Prostaglandin E₂ promotes colon cancer cell growth through a G_s-axin-β-catenin signaling axis. *Science* 310:1504–1510
26. Balkwill F (2009) Tumor necrosis factor and cancer. *Nat Rev Cancer* 9:361–371
27. Moore RJ, Owens DM, Stamp G, Arnott C, Burke F, East N, Holdsworth H, Turner L, Rollins B, Pasparakis M, Kollias G, Balkwill F (1999) Mice deficient in tumor necrosis factor-α are resistant to skin carcinogenesis. *Nat Med* 5:828–831
28. Arnott CH, Scott KA, Moore RJ, Robinson SC, Thompson RG, Balkwill FR (2004) Expression of both TNF-α receptor subtypes is essential for optimal skin tumor development. *Oncogene* 23:1902–1910
29. Popivanova BK, Kitamura K, Wu Y, Kondo T, Kagaya T, Kaneko S, Oshima M, Fujii C, Mukaida N (2008) Blocking TNF-α in mice reduces colorectal carcinogenesis associated with chronic colitis. *J Clin Invest* 118:560–570
30. Oshima H, Oshima M (2012) The inflammatory network in the gastrointestinal tumor microenvironment: lessons from mouse models. *J Gastroenterol* 47:97–106
31. Greten FR, Eckmann L, Greten TF, Park JM, Li ZW, Egan LJ, Kagnoff MF, Karin M (2004) IKKβ links inflammation and tumorigenesis in a mouse model of colitis-associated cancer. *Cell* 118:285–296
32. Heikkilä K, Ebrahim S, Lawlor DA (2008) Systematic review of the association between circulating interleukin-6 (IL-6) and cancer. *Eur J Cancer* 44:937–945
33. Bollrath J, Phesse TJ, von Burstin VA, Putoczki T, Bennecke M, Bateman T, Nebelsiek T, Lundgren-May T, Canli O, Schwitala S, Matthews V, Schmid RM, Kirchner T, Arkan MC, Ernst M, Greten FR (2009) gp130-mediated STAT3 activation in enterocytes regulates cell survival and cell-cycle progression during colitis-associated tumorigenesis. *Cancer Cell* 15:91–102
34. Grivennikov S, Karin E, Terzic J, Mucida D, Yu GY, Vallabhapurapu S, Scheller J, Rose-John S, Cheroutre H, Eckmann L, Karin M (2009) IL-6 and STAT3 are required for survival of intestinal epithelial cells and development of colitis-associated cancer. *Cancer Cell* 15:103–113
35. Gregorieff A, Clevers H (2005) Wnt signaling in the intestinal epithelium: from endoderm to cancer. *Genes Dev* 19:877–890
36. Taketo MM (2006) Wnt signaling and gastrointestinal tumorigenesis in mouse models. *Oncogene* 25:7522–7530
37. Oshima M, Oshima H, Kitagawa K, Kobayashi M, Itakura C, Taketo M (1995) Loss of *Apc* heterozygosity and abnormal tissue building in nascent intestinal polyps in mice carrying a truncated *Apc* gene. *Proc Natl Acad Sci USA* 92:4482–4486
38. Harada N, Tamai Y, Ishikawa T, Sauer B, Takaku K, Oshima M, Taketo MM (1999) Intestinal polyposis in mice with a dominant stable mutation of the β-catenin gene. *EMBO J* 18:5931–5942
39. Clements WM, Wang J, Sarnaik A, Kim OJ, MacDonald J, Fenoglio-Preiser C, Groden J, Lowy AM (2002) β-Catenin mutation is a frequent cause of Wnt pathway activation in gastric cancer. *Cancer Res* 62:3503–3506
40. Woo DK, Kim HS, Lee HS, Kang YH, Yang HK, Kim WH (2001) Altered expression and mutation of β-catenin gene in gastric carcinomas and cell lines. *Int J Cancer* 95:108–113
41. Cheng XX, Wang ZC, Chen XY, Sun Y, Kong QY, Liu J, Li H (2005) Correlation of Wnt-2 expression and β-catenin intracellular accumulation in Chinese gastric cancers: relevance with tumour dissemination. *Cancer Lett* 223:339–347
42. Oshima H, Matsunaga A, Fujimura T, Tsukamoto T, Taketo MM, Oshima M (2006) Carcinogenesis in mouse stomach by simultaneous activation of the Wnt signaling and prostaglandin E₂ pathway. *Gastroenterology* 131:1086–1095
43. Park WS, Oh RR, Park JY, Lee SH, Shin MS, Kim YS, Kim SY, Lee HK, Kim PJ, Oh ST, Yoo NJ, Lee JY (1999) Frequent somatic mutations of the β-catenin gene in intestinal-type gastric cancer. *Cancer Res* 59:4257–4260
44. Ebert MP, Fei G, Kahmann S, Muller O, Yu J, Sung JJ, Malfetherier P (2002) Increased β-catenin mRNA levels and mutational alterations of the APC and β-catenin gene are present in intestinal-type gastric cancer. *Carcinogenesis* 23:87–91
45. Oguma K, Oshima H, Aoki M, Uchio R, Naka K, Nakamura S, Hirao A, Saya H, Taketo MM, Oshima M (2008) Activated macrophages promote Wnt signalling through tumour necrosis factor-α in gastric tumour cells. *EMBO J* 27:1671–1681
46. Ristimäki A, Honkanen N, Jänkäälä H, Sipponen P, Härkönen M (1997) Expression of cyclooxygenase-2 in human gastric carcinoma. *Cancer Res* 57:1276–1280
47. Al-Marhoon MS, Nunn S, Soames RW (2004) *CagA* + *Helicobacter pylori* induces greater levels of prostaglandin E₂ than *cagA* – strains. *Prostaglandins Other Lipid Mediat* 73:181–189
48. Sun WH, Yu Q, Shen H, Qu XL, Cao DZ, Yu T, Qian C, Zhu F, Sun YL, Fu XL, Su H (2004) Roles of *Helicobacter pylori* infection and cyclooxygenase-2 expression in gastric carcinogenesis. *World J Gastroenterol* 10:2809–2813
49. Oshima H, Oshima M, Inaba K, Taketo MM (2004) Hyperplastic gastric tumors induced by activated macrophages in COX-2/*mPGES-1* transgenic mice. *EMBO J* 23:1669–1678
50. Oshima M, Oshima H, Matsunaga A, Taketo MM (2005) Hyperplastic gastric tumors with spasmodic polypeptide-expressing

- metaplasia caused by tumor necrosis factor- α -dependent inflammation in cyclooxygenase-2/microsomal prostaglandin E synthase-1 transgenic mice. *Cancer Res* 65:9147–9151
51. Yamaguchi H, Goldenring JR, Kaminishi M, Lee JR (2002) Identification of spasmolytic polypeptide-expressing metaplasia (SPEM) in remnant gastric cancer and surveillance post-gastrectomy biopsies. *Dig Dis Sci* 47:573–578
 52. Halldorsdottir AM, Sigurdardottir M, Jonasson JG, Oddsdottir M, Magnusson J, Lee JR, Goldenring JR (2003) Spasmolytic polypeptide-expressing metaplasia (SPEM) associated with gastric cancer in Iceland. *Dig Dis Sci* 48:431–441
 53. Goldenring JR, Nomura S (2009) Insight into the development of preneoplastic metaplasia: spasmolytic polypeptide-expressing metaplasia and oxyntic atrophy. In: Wang TC, Fox JG, Giraud AS (eds) *The biology of gastric cancer*. Springer, New York, pp 361–375
 54. Nomura S, Baxter T, Yamaguchi H, Leys C, Vartapetian AB, Fox JG, Lee JR, Wang TC, Goldenring JR (2004) Spasmolytic polypeptide expressing metaplasia to preneoplasia in *H. felis*-infected mice. *Gastroenterology* 127:582–594
 55. Kang W, Rathinavelu S, Samuelson LC, Merchang JL (2005) Interferon γ induction of gastric mucous neck cell hypertrophy. *Lab Invest* 85:702–715
 56. Judd LM, Alderman BM, Howlett M, Shulkes A, Dow C, Moverley J, Graill D, Jenkins BJ, Ernst M, Giraud AS (2004) Gastric cancer development in mice lacking the SHP2 binding site on the IL-6 family co-receptor gp130. *Gastroenterology* 126:196–207
 57. Guo X, Oshima H, Kitamura T, Taketo MM, Oshima M (2008) Stromal fibroblasts activated by tumor cells promote angiogenesis in mouse gastric cancer. *J Biol Chem* 283:19864–19871
 58. Itadani H, Oshima H, Oshima M, Kotani H (2009) Mouse gastric tumor models with prostaglandin E₂ pathway activation show similar gene expression profiles to intestinal-type human gastric cancer. *BMC Genomics* 10:615
 59. Ishimoto T, Nagano O, Yae T, Tamada M, Motohara T, Oshima H, Oshima M, Ikeda T, Asaba R, Yagi H, Masuko T, Shimizu T, Ishikawa T, Kai K, Takahashi E, Imamura Y, Baba Y, Ohmura M, Suematsu M, Baba E, Saya H (2011) CD44 variant regulates redox status in cancer cells by stabilizing the xCT subunit of xc⁻ and thereby promotes tumor growth. *Cancer Cell* 19:387–400
 60. Ishimoto T, Oshima H, Oshima M, Kai K, Torii R, Masuko T, Baba H, Saya H, Nagano O (2010) CD44(+) slow-cycling tumor cell expansion is triggered by cooperative actions of Wnt and prostaglandin E₂ in gastric tumorigenesis. *Cancer Sci* 101:673–678
 61. Oshima H, Hioki K, Popivanova BK, Oguma K, van Rooijen N, Ishikawa T, Oshima M (2011) Prostaglandin E₂ signaling and bacterial infection recruit tumor-promoting macrophages to mouse gastric tumors. *Gastroenterology* 140:596–607
 62. Rakoff-Nahoum S, Paglino J, Eslami-Varzaneh F, Edberg S, Medzhitov R (2004) Recognition of commensal microflora by toll-like receptors is required for intestinal homeostasis. *Cell* 118:229–241
 63. Rakoff-Nahoum S, Medzhitov R (2007) Regulation of spontaneous intestinal tumorigenesis through the adaptor protein MyD88. *Science* 317:124–127
 64. Chinen T, Komai K, Muto G, Morita R, Inoue N, Yoshida H, Sekiya T, Yoshida R, Nakamura K, Takayanagi R, Yoshimura A (2010) Prostaglandin E₂ and SOCS1 have a role in intestinal immune tolerance. *Nature Commun* 2:190
 65. Dannenberg AJ, Lippman SM, Mann JR, Subbaramaiah K, DuBois RN (2005) Cyclooxygenase-2 and epidermal growth factor receptor: pharmacologic targets for chemoprevention. *J Clin Oncol* 2:254–266
 66. Roberts RB, Min L, Washington MK, Olsen SJ, Settle SH, Coffey RJ, Threadgill DW (2002) Importance of epidermal growth factor receptor signaling in establishment of adenomas and maintenance of carcinomas during intestinal tumorigenesis. *Proc Natl Acad Sci USA* 99:1521–1526
 67. Torrance CJ, Jackson PE, Montgomery E, Kinzler KW, Vogelstein B, Wissner A, Nunes M, Frost P, Discafani CM (2006) Combinatorial chemoprevention of intestinal neoplasia. *Nat Med* 6:1024–1028
 68. Buchanan FG, Holla V, Katkuri S, Matta P, DuBois RN (2008) Targeting cyclooxygenase-2 and the epidermal growth factor receptor for the prevention and treatment of intestinal cancer. *Cancer Res* 67:9380–9388
 69. Buchanan FG, Wang D, Bargiacchi F, DuBois RN (2003) Prostaglandin E₂ regulates cell migration via the intracellular activation of the epidermal growth factor receptor. *J Biol Chem* 278:35451–35457
 70. Buchanan FG, Gorden DL, Matta P, Shi Q, Matrisian LM, DuBois RN (2006) Role of β -arrestin1 in the metastatic progression of colorectal cancer. *Proc Natl Acad Sci USA* 103:1492–1497
 71. Pai R, Soreghan B, Szabo IL, Pavelka M, Baatar D, Tarnawski AS (2002) Prostaglandin E₂ transactivates EGF receptor: a novel mechanism for promoting colon cancer growth and gastrointestinal hypertrophy. *Nat Med* 8:289–293
 72. Shao J, Lee SB, Guo H, Evers BM, Sheng H (2003) Prostaglandin E₂ stimulates the growth of colon cancer cells via induction of amphiregulin. *Cancer Res* 63:5218–5223
 73. Subbaramaiah K, Benezra R, Hudis C, Dannenberg AJ (2008) Cyclooxygenase-2-derived prostaglandin E₂ stimulates Id-1 transcription. *J Biol Chem* 283:33955–33968
 74. Al-Salih MA, Ulmer SC, Doan T, Nelson CD, Crotty T, Prescott SM, Stafforini DM, Topham MK (2007) Cyclooxygenase-2 transactivates the epidermal growth factor receptor through specific E-prostanoid receptors and tumor necrosis factor- α converting enzyme. *Cell Signal* 19:1956–1963
 75. Oshima H, Popivanova BK, Oguma K, Dong D, Ishikawa T, Oshima M (2011) Activation of epidermal growth factor receptor signaling by the prostaglandin E₂ receptor EP4 pathway during gastric tumorigenesis. *Cancer Sci* 102:713–719
 76. Mochizuki S, Okada Y (2007) ADAMs in cancer cell proliferation and progression. *Cancer Sci* 98:621–628
 77. Oshima H, Itadani H, Kotani H, Taketo MM, Oshima M (2009) Induction of prostaglandin E₂ pathway promotes gastric hamartoma development with suppression of bone morphogenetic protein signaling. *Cancer Res* 69:2729–2733
 78. Yarden Y, Sliwkowski MX (2001) Untangling the ErbB signaling network. *Nat Rev Mol Cell Biol* 2:127–137
 79. Mann JR, Backlund MG, Buchanan FG, Daikoku T, Holla VR, Rosenberg DW, Dey SK, DuBois RN (2006) Repression of prostaglandin dehydrogenase by epidermal growth factor and snail increases prostaglandin E₂ and promotes cancer progression. *Cancer Res* 66:6649–6656
 80. Myung SJ, Gerko RM, Yan M, Platzer P, Guda K, Dotson A, Lawrence E, Dannenberg AJ, Lovgren AK, Luo G, Pretlow TP, Newman RA, Willis J, Dawson D, Markowitz SD (2006) 15-Hydroxyprostaglandin dehydrogenase is an in vivo suppressor of colon tumorigenesis. *Proc Natl Acad Sci USA* 103:12098–12102
 81. Ambros V (2004) The functions of animal microRNAs. *Nature* 431:350–355
 82. Bartel DP (2004) MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116:281–297
 83. Esquela-Kerscher A, Slack FJ (2006) Oncomirs—microRNAs with a role in cancer. *Nat Rev Cancer* 6:259–269
 84. Ventura A, Jacks T (2009) MicroRNAs and cancer: short RNAs go a long way. *Cell* 136:586–591
 85. Di Leva G, Croce CM (2010) Roles of small RNAs in tumor formation. *Trends Mol Med* 16:257–267
 86. Sonkoly E, Pivarcsi A (2011) MicroRNAs in inflammation and response to injuries induced by environmental pollution. *Mutat Res* 717:46–53

87. O'Connell RM, Taganov KD, Boldin MP, Cheng G, Baltimore D (2007) MicroRNA-155 is induced during the macrophage inflammatory response. *Proc Natl Acad Sci USA* 104:1604–1609
88. Tili E, Michaille JJ, Cimino A, Costinean S, Dumitru CD, Adair B, Fabbri M, Alder H, Liu CG, Calin GA, Croce CM (2007) Modulation of miR-155 and miR-125b levels following lipopolysaccharide/TNF- α stimulation and their possible roles in regulating the response to endotoxin shock. *J Immunol* 179:5082–5089
89. Iliopoulos D, Jaeger SA, Hirsch HA, Bulyk ML, Struhl K (2010) STAT3 activation of miR-21 and miR-181b-1 via PTEN and CYLD are part of the epigenetic switch linking inflammation to cancer. *Mol Cell* 39:493–506
90. Kong D, Piao YS, Yamashita S, Oshima H, Oguma K, Fushida S, Fujimura T, Minamoto T, Seno H, Yamada Y, Satou K, Ushijima T, Ishikawa T, Oshima M (2012) Inflammation-induced repression of tumor suppressor miR-7 in gastric tumor cells. *Oncogene* 31:3949–3960
91. Volinia S, Calin GA, Liu CG, Ambs S, Cimmino A, Petrocca F, Visone R, Iorio M, Roldo C, Ferracin M, Prueitt RL, Yanaihara N, Lanza G, Scarpa A, Vecchione A, Negrini M, Harris CC, Croce CM (2006) A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci USA* 103:2257–2261
92. Sachdeva M, Zhu S, Wu F, Wu H, Walia V, Kumar S, Elble R, Watabe K, Mo YY (2009) p53 represses c-Myc through induction of the tumor suppressor miR-145. *Proc Natl Acad Sci USA* 106:3207–3212
93. Kefas B, Godlewski J, Comeau L, Li Y, Abounader R, Hawkinson M, Lee J, Fine H, Chiocca EA, Lawler S, Purow B (2008) MicroRNA-7 inhibits the epidermal growth factor receptor and the Akt pathway and is down-regulated in glioblastoma. *Cancer Res* 68:3566–3572
94. Reddy SD, Ohshiro K, Rayala SK, Kumar R (2008) MicroRNA-7, a homeobox D10 target, inhibits p21-activated kinase 1 and regulates its functions. *Cancer Res* 68:8195–8200
95. Webster RJ, Giles KM, Price KJ, Zhang PM, Mattick JS, Leedman PJ (2009) Regulation of epidermal growth factor receptor signaling in human cancer cells by microRNA-7. *J Biol Chem* 284:5731–5741
96. Jiang L, Liu X, Chen Z, Jin Y, Heidbreder CE, Kolokythas A, Wang A, Dai Y, Zhou X (2010) MicroRNA-7 targets IGF1R (insulin-like growth factor 1 receptor) in tongue squamous cell carcinoma cells. *Biochem J* 432:199–205
97. Saydam O, Senol O, Wurdinger T, Mizrak A, Ozdener GB, Stemmer-Rachamimov AO, Yi M, Stephens RM, Krichevsky AM, Saydam N, Brenner GJ, Breakefield XO (2011) miRNA-7 attenuation in schwannoma tumors stimulates growth by upregulating three oncogenic signaling pathways. *Cancer Res* 71:852–861