

The repeatability of genome-wide mutation rate and spectrum estimates

Megan G. Behringer¹ · David W. Hall²

Received: 25 January 2016 / Revised: 28 January 2016 / Accepted: 29 January 2016 / Published online: 26 February 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract Over the last decade, mutation studies have grown in popularity due to the affordability and accessibility of whole genome sequencing. As the number of species in which spontaneous mutation has been directly estimated approaches 20 across two domains of life, questions arise over the repeatability of results in such experiments. Five species were identified in which duplicate mutation studies have been performed. Across these studies the difference in estimated spontaneous mutation rate is at most, weakly significant ($p < 0.01$). However, a highly significant ($p < 10^{-5}$), threefold difference in the rate of insertions/deletions (indels) exists between two recent studies in *Schizosaccharomyces pombe*. Upon investigation of the ancestral genome sequence for both studies, a possible anti-mutator allele was identified. The observed variation in indel rate may imply that the use of indel markers, such as microsatellites, for the investigation of genetic diversity within and among populations may be inappropriate because of the assumption of uniform mutation rate within a species.

Keywords *Schizosaccharomyces pombe* · Mutation accumulation · indel · Yeast · Evolution · DNA repair · Microsatellite

Mutation is the ultimate source of all genetic variation, and has been a long-standing research focus in evolutionary genetics. Until recently, estimating mutation rates has necessitated using reporter loci (Drake 1991) and it has not been possible to obtain direct estimates of the mutation spectrum, which is the relative frequency of different nucleotide substitutions, insertions/deletions (indels), and rearrangements. However, with current methods, sequencing massive amounts of DNA to discover the relatively few newly arising, spontaneous mutations at the genome level is now affordable. This has allowed mutation accumulation (MA) studies (Lynch et al. 1999) to be used to estimate both the genome-wide mutation rate and its spectrum in a variety of organisms: *Arabidopsis thaliana* (Ossowski et al. 2010; Jiang et al. 2014), *Bacillus subtilis* (Sung et al. 2015), *Burkholderia cenocepacia* (Dillon et al. 2015), *Caenorhabditis elegans* (Denver et al. 2009; Meier et al. 2014), *Chlamydomonas reinhardtii* (Ness et al. 2012; Sung et al. 2012), *Daphnia pulex* (Keith et al. 2015), *Dictyostelium discoideum* (Saxer et al. 2012), *Dienococcus radiodurans* (Long et al. 2015), *Drosophila melanogaster* (Schridder et al. 2013; Keightley et al. 2014), *Escherichia coli* (Lee et al. 2012), *Heliconius melpomene* (Keightley et al. 2015), *Mesoplasma florum* (Sung et al. 2012), *Paramecium tetraurelia* (Sung et al. 2012), *Pristionchus pacificus* (Weller et al. 2014), *Pseudomonas aeruginosa* (Dettman et al. 2016), *Pseudomonas fluorescens* (Long et al. 2015), *Saccharomyces cerevisiae* (Lynch et al. 2008; Nishant et al. 2010; Zhu et al. 2014), *Schizosaccharomyces pombe* (Farlow et al. 2015; Behringer and Hall 2016), and *Tetrahymena*

Communicated by M. Kupiec.

Electronic supplementary material The online version of this article (doi:10.1007/s00294-016-0573-7) contains supplementary material, which is available to authorized users.

✉ Megan G. Behringer
megbehri@indiana.edu

¹ Department of Biology, Indiana University, 1001 East Third Street, Bloomington 47405, IN, USA

² Department of Genetics, University of Georgia, Athens 3060, Georgia

thermophila (Long et al. 2015). As the list of species for which genome-wide estimates or mutation rate and spectrum are available increases, the question of repeatability becomes paramount.

Genome-wide estimates of parameters of mutation have been performed in the same species only a few times, either as controls for an experimental evolution project or as a specific effort to understand the spontaneous mutation rate, so the reproducibility of results has seldom been examined. The five species with two independent genome-wide estimates reported before 2015 are included in Table 1, which indicates that they often suffer from low power due to small sample size, making it difficult to evaluate reproducibility.

Our recent mutation accumulation study in *Schizosaccharomyces pombe* (Behringer and Hall 2016) unintentionally overlapped a concurrent study of similar scope (Farlow et al. 2015) and allowed a statistically robust comparison of the repeatability of mutation rate and spectrum estimates in this species. The conditions of the two experiments differed with respect to starting strain, culture temperature, time between line transfers and growth medium, but were otherwise similar.

Genome-wide mutation rates vary across studies. In the five species with data prior to 2015, the single nucleotide mutation rate estimates vary 1.34–3.50-fold (Table 1). In comparison, the single nucleotide mutation rate estimates in the two *S. pombe* studies are only 1.18-fold different. This difference, though smaller than seen in the other five species, is statistically significant ($p < 0.05$). In previous studies, the mutation rate for small (≤ 50 bp) indels varied from 1.10–3.00-fold, though none of the differences were significant because of the low number of observed mutational events (Table 1). In comparison, the indel mutation rate estimates in the two *S. pombe* studies are 2.83-fold different, which is a highly statistically significant difference ($p < 10^{-5}$).

There are at least three possible explanations for the threefold difference observed in indel rates in *S. pombe*. First, indels are more challenging to accurately detect bioinformatically than base substitutions, not only because the resulting mismatches can make an indel-containing sequencing read more difficult to map, but also because indels commonly occur within microsatellites and highly repetitive regions which already have an increased PCR and sequencing error rate (Jiang et al. 2015; Narzisi et al. 2014). Two different pipelines were used for small indel detection in the two studies. In Farlow et al., sequencing reads were mapped with BWA (Li and Durbin 2009), and realigned with both Breakdancer (Chen et al. 2009) and Pindel (Ye et al. 2009). In our study, we also mapped sequencing reads with BWA, but realigned them with GATK's IndelRealigner (Auwera et al. 2013). Both practices are not without their issues; GATK is less sensitive

when it comes to larger indels, while Pindel has trouble with insertions (Narzisi et al. 2014). In our study, we estimated the false-positive and false-negative error rates and found them both to be less than 2.5 %. Even if error rates were an order of magnitude greater in the Farlow et al. study, they would be insufficient to explain the difference in indel mutation rate.

Second, the indel mutation rate may be altered by the different environmental conditions of the two experiments, with temperature being a likely candidate (Matsuba et al. 2013). Stressful temperatures have been demonstrated to affect microsatellite mutation rate in *Caenorhabditis elegans* (Matsuba et al. 2013), with increased temperatures leading to an increase in mutation rate. However, if *S. pombe* also exhibits increased indel mutation rate at stressful temperatures, we would expect the Farlow et al. study to have a higher rate, since they performed their mutation accumulation experiment at a higher, presumably more stressful, temperature (32° vs 30 °C). There is no obvious other difference that would suggest that *S. pombe* was more or less stressed in one versus the other experiment.

Third, differences in genetic background may explain the different indel mutation rates. To examine this hypothesis, we compared the ancestral strains in the two studies, specifically to ask whether ours exhibited a higher relative number of indels when compared to the reference. In our MA study, we found 315 indel and single nucleotide substitution differences between the ancestor and reference genome, which was a surprisingly large number given that both isolates have the same strain designation (972 h-). A comparison of our ancestor with the one used by Farlow et al., identified 208 shared differences. This high number of shared differences between the ancestors strongly suggests that they represent errors in the reference assembly. The reference genome is thought to have at least 190 errors (Hu et al. 2015; Wood et al. 2011), 183 of which are among the 208 shared mutations, and thus confirmed by our analysis. The remaining 25 have not been previously inferred (Supplemental Table). Of the remaining 107 differences between our ancestor and the reference, there are approximately 3.5-fold more indels than single nucleotide mutations. A similar analysis of Farlow et al.'s ancestor indicates 0.95-fold more indels than single nucleotide mutations, relative to the reference. Thus, our ancestor shows a 3.7-fold higher number of indels, relative to single nucleotide changes, compared to the Farlow et al. ancestor. This suggests that there may indeed be a genetic background difference between the two strains that is causing a relatively higher indel mutation rate in our ancestral strain. We note that there is no evidence for selection having played a major role in the mutational differences between either ancestor: the effects of mutational differences between the ancestors and the reference are not significantly different

Table 1 Comparison of repeated, genome-wide estimates of single nucleotide (μ_{SNM}) and small (≤ 50 bp) insertion/deletion (μ_{indel}) mutation rates across species obtained from mutation accumulation (MA) experiments

Species	Study ^a	N	Gens	SNMs	Indels	μ_{SNM} (CI ^b) ($\times 10^{-9}$)	μ_{SNM} diff	μ_{indel} (CI ^b) ($\times 10^{-9}$)	μ_{indel} diff
<i>Arabidopsis thaliana</i>	1	5	30	98	13	4.84 (3.86-5.82)	1.34	0.64 (0.28-1.00)	1.10
	2	9	10	44	7	3.62 (2.53-4.71)	NS	0.58 (0.14-1.01)	NS
<i>Chlamydomonas reinhardtii</i>	3	2	350	9	5	0.21 (0.07-0.35)	3.5	0.12 (0.01-0.22)	3.00
	4	4	1730	20	13	0.06 (0.03-0.09)	NS	0.04 (0.01-0.06)	NS
<i>Drosophila melanogaster</i>	5	8	147	732	60	5.49 (5.08-5.90)	1.96	0.45 (0.33-0.57)	2.67
	6	12	1	6	3	2.80 (0.51-5.09)	$p < 0.05$	1.20 (0.00-3.02)	NS
<i>Caenorhabditis elegans</i>	7	7	250	108	–	1.33 (1.07-1.59)	2.53	–	–
	8	6	20	60	7	3.37 (2.50-4.24)	$p < 0.01$	0.58 (0.100.69)	–
<i>Schizosaccharomyces pombe</i>	9	96	1716	398	117	0.20 (0.18-0.22)	1.18	0.06 (0.05-0.07)	2.83
	10	79	1952	327	335	0.17 (0.15-0.19)	$p < 0.05$	0.17 (0.16-0.19)	$p < 10^{-5}$
<i>Scaccharomyces cerevisiae</i>	11	2	1000	14	0	0.29 (0.14-0.45)	1.71	–	–
	12	140	2062	867	26	0.17 (0.16-0.18)	NS	0.005 (0.00-0.01)	–

Differences between the two estimates within a species are tested for significance using a Welch's *t* test, with appropriate degrees of freedom

To calculate the standard error for the *t* test, the variance among lines in the number of mutations is assumed to be equal to the mean number of mutations per line, which assumes that mutation is a Poisson process

The fold difference (diff) between estimates within a species is shown (=larger/smaller estimate)

The number of MA lines (N) and generations of accumulation (Gens), and the number of observed mutation events used for parameter estimation (SNMs and Indels) are shown

N number of mutation accumulation lines in the study, Gens generations, SNM single nucleotide mutation, indel insertion/deletion of less than 50 bp, CI confidence interval, NS not significant

^a 1 = Ossowski et al. 2010, 2 = Jiang et al. 2014, 3 = Ness et al. 2012, 4 = Sung et al. 2012a, 5 = Schrider et al. 2013, 6 = Keightley et al. 2014, 7 = Denver et al. 2012, 8 = Meier et al. 2014, 9 = Farlow et al. 2015, 10 = Behringer and Hall 2016, 11 = Nishant et al. 2010, 12 = Zhu et al. 2014

^b Confidence intervals are from standard errors of the number of mutations per line, which are estimated by assuming a Poisson distribution of mutations across lines

^c Diploid strains

from those that arose during MA, when selection is known to be ineffective (Fig. 1).

When examining differences between the ancestors for their potential to cause differences in indel rate,

two mutations were found in genes associated with DNA repair in the Farlow et al. ancestor. One of these is in *rev7* (SPBC12D12.09), which is a subunit of DNA polymerase zeta with inferred involvement in translesion synthesis

Fig. 1 Comparison of mutational effects in MA ancestor and MA lines. There is no difference between any of the three spectra (Chi-squared test: p value = 0.61)





Fig. 2 Comparison of mutation accumulation spectra for the four species in which there are sufficient numbers of mutations to make a comparison. In all four species, spectra are not significantly differ-

ent across studies (Chi-squared test: **a** *A. thaliana*, p value = 0.75, **b** *D. melanogaster*, p value = 0.58, **c** *C. elegans*, p value = 0.164, **d** *S. pombe*, p value = 0.66)

(Kosarek et al. 2008), but the mutation is synonymous and thus not likely to have an effect, unless it alters mRNA stability and thus protein levels in the cell. The second is a missense mutation in *cdc6* (SPBC336.04), also known as POLD3, which is a subunit of DNA polymerase delta. A mutation in *cdc6*, specifically *cdc6-121* has known mutator qualities, while another variant, *cdc6-23*, may reduce the mutation rate relative to wild type (Liu et al. 1999). It is possible that the *cdc6* missense mutation in the Farlow ancestor has anti-mutator qualities, which could account for the indel differences between the two ancestors and the estimated rate differences in the two MA experiments.

Regardless of whether it is environmental or due to genetic background, the substantial variation seen in indel rate has serious implications for the use of microsatellite repeats as genetic markers. In nature, populations that appear to be significantly different in their microsatellite genotypes and are thus inferred to be genetically isolated from others may simply have higher mutation rates. Microsatellite mutation models, such as the stepwise-mutation model (Slatkin 1995), assume uniformity in the indel rate within species. While warnings have been issued about the robustness of microsatellites due to differing mutation rates amongst loci (Putman and Carbone 2014), microsatellites may also miscalculate genetic distance because of differing mutation rates amongst populations.

Other parameters of the mutational process, which require large numbers of mutations to estimate with precision, including the spectrum of single nucleotide mutations, the insertion to deletion ratio for small indels, and the location of mutational hotspots do not differ between the two *S. pombe* studies. Three of the previous studies in other species have sufficient numbers of single nucleotide mutations to estimate the spectrum for this class of mutations (Fig. 2). In these three species, and in *S. pombe*, the spectra are remarkably similar to one another. The relative frequency of insertions versus deletions in small indels across studies within species can only be compared within *S. pombe*; there are insufficient numbers of indels in other studies. The ratio of insertions to deletions is not significantly different in the two *S. pombe* studies; it is 5.88 in Farlow et al. and 6.12 in our study. Further, the ratio of insertions to deletions is similar, 6.4, for those indels that differ between our ancestor and the reference, after removing those that are shared with the Farlow et al. ancestor. It is only in the Farlow ancestor, in which the ratio is 0.82, based on 20 indels, that we find a significant difference (Fisher's exact test, $p \leq 0.0003$) from the others we have observed.

In conclusion, the inadvertent overlap of our *S. pombe* MA experiment with that of Farlow et al. allowed one of the first statistically robust comparisons of estimates of parameters of mutation within a species. Generally, estimates

revealed remarkable repeatability. The single nucleotide mutation rates, though statistically significantly different, were within 20 % of one another, and the mutational spectrum for these mutations was not different. Further the relative occurrence of insertions to deletion was also not different across the two studies. The only substantial difference was the indel mutation rate, which varies by threefold across the two studies and is highly statistically significant. This suggests that the mutation rate for indels may be more sensitive to genetic background, environment, or both.

Acknowledgements We'd like to thank H. Long for helpful comments and communicating *S. pombe* ED668 ancestral sequences. This work was supported by the National Institutes of Health grants R01GM097415 and T32 GM007103.

References

- Auwerwa GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, Jordan T, Shakir K, Roazen D, Thibault J (2013). From FastQ data to high-confidence variant calls: the genome analysis toolkit best practices pipeline. *Curr Protoc Bioinformatics*, 11.10. 11–11.10. 33
- Behringer MG, Hall DW (2016) Genome-wide estimates of mutation rates and spectrum in *Schizosaccharomyces pombe* indicate CpG sites are highly mutagenic despite the absence of DNA methylation. *G3: Genes Genomes Genet* 6(1):149–160
- Chen K, Wallis JW, McLellan MD, Larson DE, Kalicki JM, Pohl CS, McGrath SD, Wendl MC, Zhang Q, Locke DP (2009) Break-Dancer: an algorithm for high-resolution mapping of genomic structural variation. *Nat Methods* 6:677–681
- Denver DR, Dolan PC, Wilhelm LJ, Sung W, Lucas-Lledó JI, Howe DK, Lewis SC, Okamoto K, Thomas WK, Lynch M (2009) A genome-wide view of *Caenorhabditis elegans* base-substitution mutation processes. *Proc Natl Acad Sci* 106:16310–16314
- Denver DR, Wilhelm LJ, Howe DK, Gafner K, Dolan PC, Baer CF (2012) Variation in base-substitution mutation in experimental and natural lineages of *Caenorhabditis* nematodes. *Genome Biol Evol* 4(4):513–522
- Dettman JR, Sztepanacz JL, Kassen R (2016) The properties of spontaneous mutations in the opportunistic pathogen *Pseudomonas aeruginosa*. *BMC Genom* 17:1
- Dillon MM, Sung W, Lynch M, Cooper VS (2015) The rate and molecular spectrum of spontaneous mutations in the GC-rich multi-chromosome genome of *Burkholderia cenocepacia*. *Genetics* 115:176834
- Drake JW (1991) A constant rate of spontaneous mutation in DNA-based microbes. *Proc Natl Acad Sci* 88:7160–7164
- Farlow A, Long H, Arnoux S, Sung W, Doak TG, Nordborg M, Lynch M (2015) The spontaneous mutation rate in the fission yeast *Schizosaccharomyces pombe*. *Genetics* 201:737–744
- Hu W, Suo F, Du L-L (2015) Bulk segregant analysis reveals the genetic basis of a natural trait variation in fission yeast. *Genome Biol Evol* 7:3496–3510
- Jiang C, Mithani A, Belfield EJ, Mott R, Hurst LD, Harberd NP (2014) Environmentally responsive genome-wide accumulation of *de novo Arabidopsis thaliana* mutations and epimutations. *Genome Res* 24:1821–1829
- Jiang Y, Turinsky AL, Brudno M (2015). The missing indels: an estimate of indel variation in a human genome and analysis of factors that impede detection. *Nucleic acids research*, gkv677

- Keightley PD, Ness RW, Halligan DL, Haddrill PR (2014) Estimation of the spontaneous mutation rate per nucleotide site in a *Drosophila melanogaster* full-sib family. *Genetics* 196:313–320
- Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, Mallet J, Davey JW, Jiggins CD (2015) Estimation of the spontaneous mutation rate in *Heliconius melpomene*. *Mol Biol Evol* 32(1):239
- Keith N, Tucker AE, Jackson CE, Sung W, Lledó JIL, Schrider DR, Schaack S, Dudycha JL, Ackerman MS, Younge AJ (2015) High mutational rates of large-scale duplication and deletion in *Daphnia pulex*. *Genome Res* 191338:191115
- Kosarek JN, Woodruff RV, Rivera-Begeman A, Guo C, D'Souza S, Koonin EV, Walker GC, Friedberg EC (2008) Comparative analysis of in vivo interactions between Rev1 protein and other Y-family DNA polymerases in animals and yeasts. *DNA Repair* 7:439–451
- Lee H, Popodi E, Tang H, Foster PL (2012) Rate and molecular spectrum of spontaneous mutations in the bacterium *Escherichia coli* as determined by whole-genome sequencing. *Proc Natl Acad Sci* 109:E2774–E2783
- Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760
- Liu VF, Bhaumik D, Wang TS-F (1999) Mutator phenotype induced by aberrant replication. *Mol Cell Biol* 19:1126–1135
- Long H, Kucukyildirim S, Sung W, Williams E, Lee H, Ackerman M, Doak TG, Tang H, Lynch M (2015). Background mutational features of the radiation-resistant bacterium *Deinococcus radiodurans*. *Molecular biology and evolution*, msv119
- Long H, Sung W, Miller SF, Ackerman MS, Doak TG, Lynch M (2015b) Mutation rate, spectrum, topology, and context-dependency in the dna mismatch repair-deficient *Pseudomonas fluorescens* ATCC948. *Gen Biol Evol* 7:262–271
- Long H, Winter DJ, Chang AYC, Sun W, Whu SH, Balboa M, Azevedo RB, Cartwright RA, Lynch M, Zufall RA (2015) Low base-substitution mutation rate in the ciliate *Tetrahymena thermophila*. [bioRxiv:025536](https://doi.org/10.1101/025536)
- Lynch M, Blanchard J, Houle D, Kibota T, Schultz S, Vassilieva L, Willis J (1999) Perspective: spontaneous deleterious mutation. *Evolution* 53:645–663
- Lynch M, Sung W, Morris K, Coffey N, Landry CR, Dopman EB, Dickinson WJ, Okamoto K, Kulkarni S, Hartl DL (2008) A genome-wide view of the spectrum of spontaneous mutations in yeast. *Proc Natl Acad Sci* 105:9272–9277
- Matsuba C, Ostrow DG, Salomon MP, Tolani A, Baer CF (2013) Temperature, stress and spontaneous mutation in *Caenorhabditis briggsae* and *Caenorhabditis elegans*. *Biol Lett* 9:20120334
- Meier B, Cooke SL, Weiss J, Bailly AP, Alexandrov LB, Marshall J, Raine K, Maddison M, Anderson E, Stratton MR (2014) *C. elegans* whole-genome sequencing reveals mutational signatures related to carcinogens and DNA repair deficiency. *Genome Res* 24:1624–1636
- Narzisi G, O'Rawe JA, Iossifov I, Fang H, Lee Y-H, Wang Z, Wu Y, Lyon GJ, Wigler M, Schatz MC (2014) Accurate *de novo* and transmitted indel detection in exome-capture data using microassembly. *Nat Methods* 11:1033–1036
- Ness RW, Morgan AD, Colegrave N, Keightley PD (2012) Estimate of the spontaneous mutation rate in *Chlamydomonas reinhardtii*. *Genetics* 192:1447–1454
- Nishant K, Wei W, Mancera E, Argueso JL, Schlattl A, Delhomme N, Ma X, Bustamante CD, Korbel JO, Gu Z (2010) The baker's yeast diploid genome is remarkably stable in vegetative growth and meiosis. *PLoS Genet* 6:e1001109
- Ossowski S, Schneeberger K, Lucas-Lledó JI, Warthmann N, Clark RM, Shaw RG, Weigel D, Lynch M (2010) The rate and molecular spectrum of spontaneous mutations in *Arabidopsis thaliana*. *Science* 327:92–94
- Putman AI, Carbone I (2014) Challenges in analysis and interpretation of microsatellite data for population genetic studies. *Ecol Evol* 4:4399–4428
- Saxer G, Havlak P, Fox SA, Quance MA, Gupta S, Fofanov Y, Strassmann JE, Queller DC (2012) Whole genome sequencing of mutation accumulation lines reveals a low mutation rate in the social amoeba *Dictyostelium discoideum*. *PloS one* 7(10):e46759
- Schrider DR, Houle D, Lynch M, Hahn MW (2013) Rates and genomic consequences of spontaneous mutational events in *Drosophila melanogaster*. *Genetics* 194:937–954
- Slatkin M (1995) A measure of population subdivision based on microsatellite allele frequencies. *Genetics* 139:457–462
- Sung W, Ackerman MS, Miller SF, Doak TG, Lynch M (2012a) Drift-barrier hypothesis and mutation-rate evolution. *Proc Natl Acad Sci* 109:18488–18492
- Sung W, Tucker AE, Doak TG, Choi E, Thomas WK, Lynch M (2012b) Extraordinary genome stability in the ciliate *Paramecium tetraurelia*. *Proc Natl Acad Sci* 109:19339–19344
- Sung W, Ackerman MS, Gout J-F, Miller SF, Williams E, Foster PL, Lynch M (2015) Asymmetric context-dependent mutation patterns revealed through mutation-accumulation experiments. *Mol Biol Evol*, msv055
- Weller AM, Rödelberger C, Eberhardt G, Molnar RI, Sommer RJ (2014) Opposing forces of A/T-biased mutations and G/C-biased gene conversions shape the genome of the nematode *Pristionchus pacificus*. *Genetics* 196:1145–1152
- Wood V, Harris MA, McDowall MD, Rutherford K, Vaughan BW, Staines DM, Aslett M, Lock A, Bähler J, Kersey PJ (2011). Pom-Base: a comprehensive online resource for fission yeast. *Nucleic Acids Res*, gkr853
- Ye K, Schulz MH, Long Q, Apweiler R, Ning Z (2009) Pindel: a pattern growth approach to detect break points of large deletions and medium sized insertions from paired-end short reads. *Bioinformatics* 25:2865–2871
- Zhu YO, Siegal ML, Hall DW, Petrov DA (2014) Precise estimates of mutation rate and spectrum in yeast. *Proc Natl Acad Sci* 111:E2310–E2318