



How to optimize recruitment strategies of patients with rheumatic and musculoskeletal diseases for online surveys: experience from an international study

Ihsane Hmamouchi^{1,2} · Antonella Abi Najm³ · Lina El Kibbi⁴ · Mona Metawee⁴ · Hussein Halabi⁵ · Nizar Abdulateef⁶ · Mervat Eissa⁷ · Manal El Rakawi⁸ · Basel Masri⁹ · Fatma Boutaiban¹⁰ · Wafa Hamdi¹¹ · Asal Adnan⁶ · Renaud Felten¹² · Laurent Arnaud¹² · Nelly Ziade¹³

Received: 7 June 2022 / Accepted: 22 August 2022 / Published online: 30 September 2022
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2022

Abstract

The use of online surveys as a recruitment tool for clinical research has recently expanded; nevertheless, optimal recruitment strategies remain poorly identified. Objectives. The study aimed to identify the most effective recruitment strategies for online research studies and to determine the optimal survey channels for obtaining patients' responses. This is a post-hoc analysis of the ARCOVAX (ArLAR COVID Vaccination) study. Multiple recruitment strategies were disseminated in Arabic, English, and French. The proportion of enrolled patients was correlated with each strategy. Channels used by patients to complete the survey were divided into three categories (social media (SoMe), doctor, and patients' associations). These channels were correlated with the patients' characteristics and the country's Gross Domestic Product (GDP). A total of 1595 patients from 19 Arab countries completed the survey. Patients' mean age was 39 years, 73.2% (1159) were females, 17.8% (284) had a university education level and 93.1% (1468) answered the survey in Arabic. The most effective recruitment strategies were personalized WhatsApp reminders to recruiters (30% of enrolled patients), technical support in response to access issues (27%) and sharing recruitment status by country on a WhatsApp group (24%). The channels used to complete the survey were: SoMe in 45% (711), doctor in 40% (647), and patients' associations in 8.5% (233), and correlated with age and GDP. To optimize recruitment, it is recommended to combine multiple strategies and channels, use the native language and be active (mobilize teams), reactive (provide prompt technical support), and proactive (share regular updates and reminders).

Keywords Recruitment strategies · Social media · Rheumatic diseases · Clinical research · Online survey

Introduction

Online surveys are currently one of the most effective tools for recruiting participants for clinical research [1] and have been largely stimulated by the Coronavirus Disease 2019 (COVID-19) pandemic. Using e-recruitment, a web-based recruitment method and a mass communication medium, researchers are now able to reach a large patient population [2]. Analysis of the most effective strategies to promote recruitment for online research studies is crucial to ensure

that the right approach is used in the right population to properly meet its needs.

One of the major channels to disseminate online surveys is social media (SoMe), such as Facebook, Twitter, or WhatsApp. In fact, SoMe has been widely used for specific health-related purposes, especially after the COVID-19 outbreak, such as medical education, awareness campaigns, health interventions, and medical surveillance [3]. Its use for recruiting participants in health surveys is growing and shows many advantages compared to traditional methods [4, 5]. It stimulates interest in participants, increases engagement and retention of subjects, and provides better interaction between clinical research demand and supply [6, 7]. Moreover, it improves participation, especially in hard-to-reach populations [8], reduces the cost, offers flexibility in time and place, and accelerates the rate of recruitment [4]. In particular, WhatsApp, which is the most used mobile instant

Renaud Felten and Laurent Arnaud contributed equally to the manuscript.

✉ Ihsane Hmamouchi
ihsane.hmamouchi@gmail.com

Extended author information available on the last page of the article

messaging, optimizes immediate communication between the researcher and the participant at an affordable price, offering flexibility in time and place [9].

However, e-recruitment might have a negative impact on the quality of data collection, on the diversity of participants, and is more prone to technical issues [10]. Indeed, online surveys create more selection bias and inadequacy of population representation since samples obtained tend to be younger, more educated, and with a female predominance [8]. Also, electricity and internet access in certain populations remain a barrier to recruitment through SoMe [4]. Other issues, such as the validity of informed consent and protection of privacy are also of concern [6].

Literature about online recruitment for research studies lacks detailed information about the profile of participants, optimal recruitment strategies and survey channels, especially in Arab countries.

The ARCOVAX study (Arab League of Associations for Rheumatology (ArLAR) Vaccination study) [11] is a cross-sectional study that used an anonymous web-based survey to assess the acceptability of the COVID-19 vaccine in patients with chronic rheumatic musculoskeletal diseases (RMDs) and healthcare professionals (HCPs) and included participants from 16 Arab countries. Being one of the largest recent online surveys in the region, the ARCOVAX population represents an excellent opportunity to analyze the optimal e-recruitment strategies.

The primary objective of this study was to identify the most effective recruitment strategies for online research studies, based on the ARCOVAX study. The secondary objective was to determine the optimal survey channels for collecting the patients' responses.

Methods

Study design

This is a post hoc analysis of the ARCOVAX study adapted from the VAXICOV study [12, 13], translated to Arabic by a steering committee composed of ten rheumatologists and one rheumatology fellow. The survey was validated by the ArLAR scientific committee composed of rheumatologists from 16 Arab countries. The scientific committee made a few minor revisions, with regard to the different names of medications in the different countries. A pilot test done in three languages (Arabic, English, and French) found the survey to be well understood and easy to complete in less than 10 min. The survey was disseminated online in the three languages between April 13 and May 11, 2021. Health-care professionals were considered as a convenience control group, expected to serve as a reference for optimal vaccine acceptability at the time of the study. The study included

1595 patients and 1517 HCPs by convenience sampling. Nevertheless, the two groups were very different in terms of socio-demographic characteristics, therefore, the current study was focused on the patient group.

Main outcome variable

The survey channels used by the participants to respond were recorded and grouped into three categories: SoMe platforms in general, the doctor (by direct contact at the clinic or through WhatsApp messaging), and patients' associations.

Procedures

At launch, the survey was disseminated online using the Google Forms platform, through multiple SoMe platforms (Facebook, Twitter, Instagram, Snapchat, Messenger, WhatsApp groups) of ArLAR, Arab Adult Arthritis Awareness group (AAAA, an ArLAR special-interest group), a network of participating rheumatologists within ArLAR and patient associations' accounts. In addition, mass emails were sent from the ArLAR and the AAAA group to available mailing lists of HCPs and patients with RMDs. Also, WhatsApp messages were shared on the participating rheumatologists' groups to be sent to their available patients' WhatsApp lists. Finally, posters with QR codes were created and displayed in the participating rheumatology clinics. During the recruitment period, additional strategies were used to boost e-recruitment: weekly reminders on SoMe and WhatsApp groups, sharing regular recruitment status by country on AAAA WhatsApp group which included the key recruiters, personalized WhatsApp messages sent to key recruiters, providing real-time technical support in case of survey access issues (such as technical assistance in connection with the survey link in the preferred languages), announcing the count-down to the last days of recruitment using a motivational video on SoMe.

Anonymous demographic data were collected: age, gender, educational level, rheumatic disease, anti-rheumatic treatments, vaccine perceptions, country, and Gross Domestic Product (GDP) level according to the World Bank Classification [14]. The details of the questionnaire were previously described elsewhere [11].

Ethical considerations

Accessing the Google Form platform and then clicking "Agree" after reading the survey description was considered as consent to participate. Confidentiality was maintained in the study by keeping all the participants' data anonymous. The study was approved by the scientific committee of the ArLAR and by the Institutional Review Boards of Saint-Joseph University, Beirut, and the Specialized Medical

Center, Riyadh (respectively, numbers CEHDF1825 and H-01-R-056, both in April 2021).

Statistical analysis

The participants' profiles (demographic and disease characteristics and chronic anti-rheumatic treatments) were presented descriptively as numbers and percentages for categorical variables and means and standard deviations for continuous variables.

The characteristics related to the survey were recorded: language used to respond to the survey, number of responses according to the country, and the GDP region [13].

For this study, we identified 5 sequential strategies to boost e-recruitment: (1) Launching the survey on a Social Media platform, (2) providing real-time technical support in case of survey access issues such as technical assistance in connection with the survey link in the preferred languages, (3) sharing regular recruitment status by country on AAAA WhatsApp group which included the key recruiters, (4) displaying reminder posters with QR codes in the participating rheumatology clinics, (5) personalized WhatsApp messages sent to key recruiters, (6) announcing the count-down to the last days of recruitment using a motivational video on SoMe.

Each strategy used to boost e-recruitment was presented descriptively in correlation with the number of daily enrolments over the time period of recruitment (from April 13 to May 11, 2021) and with the proportion of participants enrolled immediately after the instauration of the strategy.

Also, three channels used to reach the patients were identified: (1) Public messages via any SoMea platform including Facebook, Instagram, Twitter, Snapchat, Messenger, WhatsApp groups (2) Patients' associations on SoMe and (3) one on one contact with the doctor (either by direct contact at the clinic or through WhatsApp message from the doctors' office). The factors associated with each channel were identified using Chi-Square, Fisher exact test, and ANOVA as appropriate. A p value < 0.05 was used to define statistical significance. All statistical analyses were performed on JAMOVI 2021 version 1.6 [15].

Results

A total of 3176 participants from 19 Arab countries completed the survey, from which 1595 patients with RMDs were included in the current analysis.

Profile of the participants

The mean age of the patients was 39 years (SD 13 years); 72.7% were females and 17.8% had a university education (Table 1). The patients were from High-Income countries in

Table 1 Profile of the participants in the ArLAR COVID-19 vaccine perceptions (ARCOVAX) online survey

	All participants <i>N</i> = 3176	Patients group <i>N</i> = 1595
Age, mean [SD]	39.6 [12.4]	39.0 [13.0]
Women, <i>n</i> (%)	2073 (65.3)	1159 (72.7)
GDP Region, <i>n</i> (%)		
High income	808 (25.4)	577 (36.2)
Upper middle income	1286 (40.5)	546 (34.1)
Lower middle and low income	1082 (34.1)	472 (29.6)
Country, <i>n</i> (%)		
Iraq	800 (25.2)	293 (18.4)
Saudi Arabia	576 (18.1)	435 (27.3)
Egypt	416 (13.1)	236 (14.8)
Lebanon	186 (5.9)	111 (7.0)
Algeria	158 (5.0)	67 (4.2)
Libya	134 (4.2)	75 (4.7)
Morocco	129 (4.1)	28 (1.8)
Tunisia	120 (3.8)	36 (2.3)
Jordan	115 (3.6)	67 (4.2)
Bahrain	101 (3.2)	64 (4.0)
Syria	101 (3.2)	33 (2.1)
Palestine	95 (3.0)	34 (2.1)
Kuwait	56 (1.8)	41 (2.6)
Sudan	48 (1.5)	33 (2.1)
Oman	32 (1.0)	18 (1.1)
UAE	20 (0.6)	10 (0.6)
Qatar	18 (0.6)	8 (0.5)
Yemen	5 (0.2)	4 (0.2)
Comoros	1 (0.0)	1 (0.06)
Rheumatic diagnosis <i>n</i> (%) [*]		
Rheumatoid arthritis	–	745 (46.7)
Lupus	–	342 (21.4)
Axial spondyloarthritis	–	143 (9.0)
Psoriatic arthritis	–	79 (4.9)
Myositis	–	33 (2.1)
Educational level <i>n</i> (%)		
Up to primary school		1264 (79.2)
Up to university		284 (17.8)
Technical studies		47 (2.9)

^{*}The five most frequent diagnoses are reported here

GDP Region according to World Bank Data

36.2%, Upper-Middle Income countries in 34.1%, and Low-Middle and Low-Income countries in 29.6%.

Language used in the survey

The majority of patients responded to the survey in Arabic (93.1%) followed by French (4.3%) and English (2.6%).

However, significant differences among the countries were found. For instance, more responses in French were received from the Maghreb countries (Algeria, Morocco, and Tunisia countries) and Lebanon ($p < 0.001$) (Fig. 1).

Day of the week

The higher rate of response to the survey was observed on Mondays (21.9%), followed by Wednesdays (18.5%) and Tuesdays (15.2%), which are weekdays in all the surveyed countries.

The rate of participation

The recruitment of participants was highly fluctuating throughout the study period (Fig. 2). The first peak of inclusion was observed on day 3, after providing prompt technical support for participants having access issues, and a second peak was observed on day 14, after sending a personalized reminder to key recruiters. About 50% of the participants were enrolled in the first 10 days of the recruitment period which lasted around one month in total. In addition, more than 90% of the participants were

Fig. 1 Enrollment of patients for the online survey and correlation with the recruitment strategies. WA WhatsApp Messages, *sharing the status with key recruiters





Fig. 2 Language chosen by patients to respond to the ARCOVAX survey, by country and geographical region



enrolled within the first 21 days, after which there seemed to be a saturation effect.

Online recruitment strategies had a different impact on the rate of participation: 30% of participants were enrolled after sending personalized WhatsApp reminders to key recruiters in different countries, 27% after providing prompt technical support to resolve issues faced by the participants, 24% after sharing recruitment status by country on the AAAA's WhatsApp group, 12% after posting electronic posters and QR codes via SoMe and WhatsApp groups, 3% after the initial launch on SoMe and WhatsApp groups and 2% after sharing a motivational video announcing the countdown to the last week of recruiting.

Survey channels for collecting patients' responses

Among the 1595 patients, 711 (44.7%) patients were recruited via anonymous SoMe (Facebook, Instagram, Twitter, Snapchat), 647 (40.7%) were recruited through their doctor (either by direct contact at the clinic or through WhatsApp message from the doctors' office), and only 233 (14.6%) were recruited through patients' associations. Less than 1% (15 patients) were recruited through emails.

Factors associated with patient participation through social media

The use of SoMe channel to participate in the survey was highest in Lower-Middle and Low-Income countries (53.6% of participants from these countries were reached through SoMe), followed by High-Income countries (48.9%) and Upper-Middle-Income countries (32.2%) ($p < 0.001$).

Patients who used SoMe to participate were younger (mean age 38.2 years [SD 11.6]) compared to those who used other channels (39.7 years [SD 14.1]). No association was found with gender ($p = 0.10$), language ($p = 0.64$) and education ($p = 0.66$) (Table 2).

Factors associated with patient participation through the doctor

The survey reached the patients through their doctors mainly in the Upper Middle-Income countries (56.6% of the patients from that GDP region). This channel was also associated with older age (mean age was 40.3 years [SD 14.8] in patients who were reached through their doctor versus 38.2 years [SD 11.6] in the other channels ($p = 0.002$) (Table 2).

Factors associated with patient participation through the patients' associations

The proportion of patients who were reached through the patients' association was small in general (8.5%) but was slightly higher in the High-Income countries (9.1%) ($p < 0.001$) (Table 2).

Discussion

The most effective e-recruitment strategies for patients with RMDs for research purposes were personalized WhatsApp reminders to the key recruiters, especially in Middle-Income countries, and prompt technical support, mostly

Table 2 Channels through which patients were reached by the ARCOVAX online survey

	Social media ^a <i>N</i> = 711	Patients' associations <i>N</i> = 233	One on one contact with the Doctor <i>N</i> = 647	<i>p</i>
Age (years) ^b	38.2 (11.6)	37.9 (11.6) ^b	40.3 (14.8) ^b	
Gender <i>n</i> (%)	Female	183 (78.9)	473 (73.3)	0.07
Education <i>n</i> (%)	Primary school	174 (75.0)	521 (80.8)	0.31
	High school and university	129 (18.4)	104 (16.1)	
	Technical	17 (2.4)	20 (3.1)	
Region GDP <i>n</i> (%)	High	109 (47.0)	184 (28.5)	<0.001
	Upper middle	174 (24.9)	310 (48.1)	
	Lower middle and low	248 (35.4)	151 (23.4)	
Language <i>n</i> (%)	Arabic	220 (94.8)	596 (92.4)	0.23
	English	16 (2.3)	23 (3.6)	
	French	32 (4.6)	26 (4.0)	

^aSocial Media: Facebook, Instagram, Twitter, Snapchat, Messenger, WhatsApp groups.

^bMean (SD) One-Way ANOVA $p < 0.05$ for social media vs patients' association and social media vs doctor GDP Region according to World Bank Data.

in high-income countries. The superiority of personalized messages over mass-produced messages and random posts on SoMe was reported in previous studies [16], highlighting the importance of empathy among researchers and participants. Indeed, participants' compliance increases with their familiarity with the one making the request. In addition, the remarkable boost in response rate seen after the correction of technical issues suggests the importance of this action in all online surveys as mentioned in the literature [17].

Moreover, this study clearly demonstrated that an Arabic version of an international survey is mandatory to reach the participants living in Arab countries. However, French was also important in the Maghreb countries and Lebanon, where it is the second language. Thus, the current study confirmed that the native language is always the best way to reach a population since it offers a comprehensive form of a survey that gives less biased results and is more practical for the patient to finish the survey faster [18].

Regarding the optimal survey channels for reaching the patients, the current study highlights the importance of SoMe, especially in the younger population. This fact is concordant with previous literature [19, 20] which demonstrated that SoMe has become a global phenomenon and plays a central role in increasing participation levels. The importance of SoMe was observed in all countries regardless of the GDP levels but was more pronounced in Low to Low-Middle Income countries (53.6% of participants). Nevertheless, doctors still play a significant role as well to improve recruitment. The "doctor's" channel should not be underestimated especially in Upper Middle-income countries where 56.6% of patients were recruited through their doctors. Finally, an important point highlighted by our data was the urgent need to activate the role of patients' associations since only 8.5% of patients were recruited through this channel, mostly in high-income countries.

The profile of participants was consistent with the literature that showed that younger patients [21, 22] and females [34] were more likely to respond to online surveys. In comparison with the VAXICOV study [12] Arab patients who responded to the ARCOVAX survey were younger, with a mean age of 39 ± 13 years, compared to a mean age of 50 ± 10 years in VAXICOV. Also, 72.7% of participants were identified as females in the ARCOVAX study, whereas 90.5% of participants in the VAXICOV study were females. With Information Technology (IT) development, the young participants are much more reachable through SoMe [23]. Moreover, the young population tends to be more involved in health education and more motivated to participate in medical surveys, whereas older patients may not realize the importance of contributing to medical studies. As for gender, the female predominance may reflect the higher female/male ratio in some RMDs such as rheumatoid arthritis or lupus [24]. Moreover, females could be more responsive to online

surveys because they tend to be more patient, interested in health education, and available since in Arab countries, many females do not work [25].

This study has some limitations. First, a selection bias should be considered, as SoMe was used to promote the survey, which would undoubtedly impact the profile of participants. Moreover, the data about individual platforms (i.e., Facebook or Twitter alone) was not available for analysis; thus, profiling patients by platform type was not possible. Although the use of SoMe might be very heterogeneous among the Arab countries, it has a very strong and growing presence, with a predominance of Facebook use [26]. In a previous ArLAR survey including 2163 patients with RMDs, 73% reported that their main source of information about COVID-19 was obtained through SoMe, followed by television in 51% [27]. Nevertheless, a large number of participants (3176 participants, including 1595 patients) can give a reasonably accurate idea about the target population of patients with RMDs. In addition, and since the socio-demographic profiles of patients and HCPs were very different, only the patients were included in the current analysis, to minimize the risk of confounding bias in the analysis of factors associated with the use of the communication channel. Also, one might argue that there's an overlap between the channels for reaching the patient. For example, a patient might be reached by their doctors through the latter's SoMe account or by WhatsApp. However, we estimate that the patients identified in priority the channel they considered as the primary channel for receiving the survey. Moreover, although the association between one strategy and the subsequent recruitment was clear in the descriptive analysis, one might argue that the recruitment may also be boosted by previous strategies having a residual effect over time. Finally, the study did not consider the severity of the disease. In fact, a relationship could exist between disease severity and the motivation to respond to a survey related to the disease, as studies showed that knowledge of the topic increases the participation rate [28]. Despite these limitations, the study followed the current directives for survey-based studies, i.e., having an ethical committee approval, survey validation by a different group than the authors, pilot testing, maintaining anonymity, using various social media platforms, and utilization of descriptive and analytical approaches, which adds to the data quality and conformity with international guidance [1, 29, 30].

In summary, the most effective strategies to promote recruitment for online research studies were sending personalized reminders to key recruiters, followed by providing prompt technical support, and sharing enrollment results by country in real-time. Moreover, the importance of SoMe, as well as the role of doctors as a recruitment channel were highlighted crucial. Disparities and variations according to GDP classification and age were noted. Furthermore, the

study confirmed the participation of a predominantly young and female patient profile, and higher participation in the native Arabic language. Therefore, to optimize recruitment, it is recommended to combine multiple strategies and channels, use the native language, and be active (mobilize teams to participate), reactive (provide prompt technical support), and proactive (share regular updates and reminders).

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00296-022-05195-0>.

Author contributions IH, RF, LA and NZ designed the study. IH, NZ and AAN drafted the first version of the manuscript. IH, AAN, LK, MM, HH, NA, ME, MR, BM, FA, WH, and NZ recruited the participants. All the authors contributed to the critical revision of the manuscript for important intellectual content. All the authors approved the final version to be published. All the authors agreed to be accountable for all aspects of the work. All co-authors take full responsibility for the integrity of all aspects of the work.

Funding None.

Data availability Data can be made available upon request from the corresponding author.

Declarations

Conflict of interest None.

Ethical approval The study was approved by the scientific committee of the ArLAR and by the Institutional Review Boards of Saint Joseph University, Beirut, and Specialized Medical Center, Riyadh.

Consent to participate Accessing the Google form then clicking “Agree” after reading the survey description was considered as consent to participate.

References

- Ball HL (2019) Conducting online surveys. *J Hum Lact* 35(3):413–417. <https://doi.org/10.1177/0890334419848734>
- Smith AD, Manna DR (2005) E-recruitment of patients for clinical trials. *Int J Electron Healthc* 1(4):413. <https://doi.org/10.1504/IJEH.2005.006688>
- Chen J, Wang Y (2021) Social media use for health purposes: systematic review. *J Med Internet Res* 23(5):1–16. <https://doi.org/10.2196/17917>
- Whitaker C, Stevelink S, Fear N (2017) The Use of facebook in recruiting participants for health research purposes: a systematic review. *J Med Internet Res* 19(8):e290. <https://doi.org/10.2196/jmir.7071>
- Rolls K, Hansen M, Jackson D, Elliott D (2016) How health care professionals use social media to create virtual communities: an integrative review. *J Med Internet Res* 18(6):e166. <https://doi.org/10.2196/jmir.5312>
- Refolo P, Sacchini D, Minacori R, Daloiiso V, Spagnolo AG (2015) E-recruitment based clinical research: notes for research ethics committees/institutional review boards. *Eur Rev Med Pharmacol Sci* 19(5):800–804 (PMID: 25807433)
- Frandsen M, Thow M, Ferguson SG (2016) The effectiveness of social media (Facebook) compared with more traditional advertising methods for recruiting eligible participants to health research studies: a randomized controlled clinical trial. *JMIR Res Protoc* 5(3):e161. <https://doi.org/10.2196/resprot.5747>
- Topolovec-Vranic J, Natarajan K (2016) The use of social media in recruitment for medical research studies: a scoping review. *J Med Internet Res* 18(11):e286. <https://doi.org/10.2196/jmir.5698>
- Kaufmann K, Peil C (2020) The mobile instant messaging interview (MIMI): using WhatsApp to enhance self-reporting and explore media usage in situ. *Mobile Media Com* 8(2):229–246. <https://doi.org/10.1177/2050157919852392>
- Barber L (2006) e-Recruitment developments. *Inst Employ Stud*. <https://doi.org/10.2196/resprot.5747>
- El Kibbi L, Metawee M, Hmamouchi I, Abdulateef N, Halabi H, Eissa M, El Rakawi M, Masri B, Abutiban F, Hamdi W, Adnan A, Najm AA, Felten R, Arnaud L, Ziadé N (2022) Acceptability of the COVID-19 vaccine among patients with chronic rheumatic diseases and health-care professionals: a cross-sectional study in 19 Arab countries. *Lancet Rheumatol* Mar 4(3):160–163. [https://doi.org/10.1016/S2665-9913\(21\)00368-4](https://doi.org/10.1016/S2665-9913(21)00368-4)
- Felten R, Dubois M, Ugarte-Gil MF et al (2021) Vaccination against COVID-19: Expectations and concerns of patients with autoimmune and rheumatic diseases. *Lancet Rheumatol* 3(4):e243–e245. [https://doi.org/10.1016/S2665-9913\(21\)00039-4](https://doi.org/10.1016/S2665-9913(21)00039-4)
- Felten R, Dubois M, Ugarte-Gil MF et al (2021) Cluster analysis reveals three main patterns of beliefs and intention with respect to SARS-CoV-2 vaccination in patients with autoimmune and inflammatory diseases. *Rheumatology (Oxford)* 60(S1):SI68–SI76. <https://doi.org/10.1093/rheumatology/keab432>
- World Bank Country and Lending Groups–World Bank Data Help Desk (2019). The World Bank. Published online 2019:1–8. <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups%0A> Accessed 10 April 2022
- The jamovi project (2021). jamovi. (Version 1.6) [Computer Software]. Retrieved from <https://www.jamovi.org>. Accessed 10 April 2022
- Yuan P, Bare MG, Johnson MO, Saberi P (2014) Using Online Social Media for Recruitment of Human Immunodeficiency Virus-Positive Participants: A Cross-Sectional Survey. *J Med Internet Res* 16(5):e117. <https://doi.org/10.2196/jmir.3229>
- Park K, Park N, Heo W, Gustafson K (2018) What prompts college students to participate in online surveys? *Int Educ Stud* 12(1):69. <https://doi.org/10.5539/ies.v12n1p69>
- Walsh T, Nurkka P, Petrie H, Olsson J (2013) The effect of language in answering qualitative questions in user experience evaluation web-surveys. Proceedings of the 25th Australian Computer-Human Interaction Conference: Augmentation, Application, Innovation, Collaboration, OzCHI 2013. Published online 2013:73–82. <https://doi.org/10.1145/2541016.2541049>
- Cobanoglu C, Moreo PJ, Warde B (2001) A comparison of Mail, Fax and Web-based survey methods. *Int J Mark Res* 43(4):1–15. <https://doi.org/10.1177/147078530104300401>
- Greenlaw C, Brown-Welty S (2009) A comparison of Web-based and paper-based survey methods. *Eval Rev* 33(5):464–480. <https://doi.org/10.1177/0193841x09340214>
- Aerny-Perreten N, Domínguez-Berjón MF, Esteban-Vasallo MD, García-Riolobos C (2015) Participation and factors associated with late or non-response to an online survey in primary care. *J Eval Clin Pract* 21(4):688–693. <https://doi.org/10.1111/jep.12367>
- Listyowardjo TA, Nap RE, Johnson A (2011) Demographic differences between health care workers who did or did not respond to a safety and organizational culture survey. *BMC Res Notes* 4:328. <https://doi.org/10.1186/1756-0500-4-328>

23. Perrin A (2005) Social Media Usage: 2005–2015. (October):2005–2015. www.pewinternet.org/2015/10/08/social-networking-usage-2005-2015/. Accessed 10 April 2022
24. Ziade N, el Khoury B, Zoghbi M et al (2020) Prevalence and pattern of comorbidities in chronic rheumatic and musculoskeletal diseases: the COMORD study. *Sci Rep* 10(1):7683. <https://doi.org/10.1038/s41598-020-64732-8>
25. El-Swais M (2016) Despite high education levels, Arab women still don't have jobs. March 9:1–6. <https://blogs.worldbank.org/arabvoices/despite-high-education-levels-arab-women-still-dont-have-jobs>. Accessed 10 April 2022
26. <https://gs.statcounter.com/social-media-stats>. Accessed 10 April 2022
27. Ziadé N, El Kibbi L, Hmamouchi I, Abdulateef N et al (2020) Impact of the COVID-19 pandemic on patients with chronic rheumatic diseases: a study in 15 Arab countries. *Int J Rheum Dis* 23(11):1550–1557. <https://doi.org/10.1111/1756-185X.13960>
28. Brosnan K, Kemperman A, Dolnicar S (2021) Maximizing participation from online survey panel members. *Int J Mark Res* 63(4):416–435. <https://doi.org/10.1177/1470785319880704>
29. Gaur PS, Zimba O, Agarwal V, Gupta L (2020) Reporting Survey Based Studies – a Primer for Authors. *J Korean Med Sci* 35(45):e398. <https://doi.org/10.3346/jkms.2020.35.e398>
30. Jones TL, Baxter MA, Khanduja V (2013) A quick guide to survey research. *Ann R Coll Surg Engl* 95(1):5–7. <https://doi.org/10.1308/003588413X13511609956372>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Authors and Affiliations

Ihsane Hmamouchi^{1,2} · Antonella Abi Najm³ · Lina El Kibbi⁴ · Mona Metawee⁴ · Hussein Halabi⁵ · Nizar Abdulateef⁶ · Mervat Eissa⁷ · Manal El Rakawi⁸ · Basel Masri⁹ · Fatma Boutaiban¹⁰ · Wafa Hamdi¹¹ · Asal Adnan⁶ · Renaud Felten¹² · Laurent Arnaud¹² · Nelly Ziadé¹³

Antonella Abi Najm
antonella14997@gmail.com

Lina El Kibbi
linakibbe@hotmail.com

Mona Metawee
monatmetawee.93@gmail.com

Hussein Halabi
mfasel@gmail.com

Nizar Abdulateef
nazarlataef@yahoo.com

Mervat Eissa
mervateissa@kasralainy.edu.eg

Manal El Rakawi
m.elrakawi@gmail.com

Basel Masri
basel.masri@masriclinic.co

Fatma Boutaiban
thething99@hotmail.com

Wafa Hamdi
wafahamdi6@yahoo.fr

Asal Adnan
asaladnan1987@gmail.com

Renaud Felten
renaud.felten@chru-strasbourg.fr

Laurent Arnaud
laurent.arnaud@chru-strasbourg.fr

Nelly Ziadé
nellziade@yahoo.fr

¹ Laboratory of Biostatistics, Clinical Research and Epidemiology (LBRCE), Faculty of Medicine and Pharmacy, Mohammed V University, Rabat, BP 10000, Rabat, Morocco

² Rheumatology Unit, Temara Hospital, Temara, Morocco

³ Faculty of Medicine, Saint-Joseph University, Beirut, Lebanon

⁴ Division of Rheumatology, Department of Internal Medicine, Specialized Medical Center, Riyadh, Saudi Arabia

⁵ King Faisal Specialist Hospital and Research Center, Jeddah, Saudi Arabia

⁶ Rheumatology Unit, Department of Medicine, College of Medicine, University of Baghdad, Baghdad, Iraq

⁷ Rheumatology Department, Cairo University, Cairo, Egypt

⁸ Rheumatology Department, Faculty of Medicine, Douera Hospital, Saad Dahlab, Blida, Algeria

⁹ Jordan Hospital, Amman, Jordan

¹⁰ Rheumatology Unit, Department of Medicine, Jaber Alahmed Alsabab Hospital, Kuwait City, Kuwait

¹¹ Rheumatology Department, Kassab Institute of Orthopedics, UR17SP04, Faculty of Medicine of Tunis, Tunis El Manar University, Tunis, Tunisia

¹² Department of Rheumatology, French National Reference Center for Autoimmune Diseases (CRMR RESO), Hôpitaux Universitaires de Strasbourg, Strasbourg, France

¹³ Rheumatology Department, Saint-Joseph Medical University and Hotel-Dieu de France Hospital, Beirut, Lebanon