

Functional conservation of rice OsNF-YB/YC and *Arabidopsis* AtNF-YB/YC proteins in the regulation of flowering time

Yoon-Hyung Hwang¹ · Soon-Kap Kim^{2,4} · Keh Chien Lee¹ · Young Soo Chung³ · Jeong Hwan Lee² · Jeong-Kook Kim¹

Received: 5 November 2015 / Revised: 23 December 2015 / Accepted: 29 December 2015 / Published online: 11 January 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract

Key message Rice OsNF-YB and OsNF-YC complement the late flowering phenotype of *Arabidopsis nf-yb* double and *nf-yc* triple mutants, respectively. In addition, OsNF-YB and OsNF-YC interact with AtNF-YC and AtNF-YB, respectively.

Abstract Plant NUCLEAR FACTOR Y (NF-Y) transcription factors play important roles in plant development and abiotic stress. In *Arabidopsis thaliana*, two NF-YB (AtNF-YB2 and AtNF-YB3) and five NF-YC (AtNF-YC1, AtNF-YC2, AtNF-YC3, AtNF-YC4, and AtNF-YC9) genes regulate photoperiodic flowering by interacting with other

AtNF-Y subunit proteins. Three rice NF-YB (OsNF-YB8, OsNF-YB10, and OsNF-YB11) and five rice OsNF-YC (OsNF-YC1, OsNF-YC2, OsNF-YC4, OsNF-YC6, and OsNF-YC7) genes are clustered with two AtNF-YB and five AtNF-YC genes, respectively. To investigate the functional conservation of these NF-YB and NF-YC genes in rice and *Arabidopsis*, we analyzed the flowering phenotypes of transgenic plants overexpressing the respective OsNF-YB and OsNF-YC genes in *Arabidopsis* mutants. Overexpression of OsNF-YB8/10/11 and OsNF-YC2 complemented the late flowering phenotype of *Arabidopsis nf-yb2 nf-yb3* and *nf-yc3 nf-yc4 nf-yc9* mutants, respectively. The rescued phenotype of *35S::OsNF-YC2 nf-yc3 nf-yc4 nf-yc9* plants was attributed to the upregulation of *FLOWERING LOCUS T (FT)* and *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1 (SOC1)*. In vitro and in planta protein–protein analyses revealed that OsNF-YB8/10/11 and OsNF-YC1/2/4/6/7 interact with AtNF-YC3/4/9 and AtNF-YB2/3, respectively. Our data indicate that some OsNF-YB and OsNF-YC genes are functional equivalents of AtNF-YB2/3 and AtNF-YC3/4/9 genes, respectively, and suggest functional conservation of *Arabidopsis* and rice NF-Y genes in the control of flowering time.

Communicated by J. S. Shin.

Y.-H. Hwang and S.-K. Kim contributed equally to this work.

Electronic supplementary material The online version of this article (doi:10.1007/s00299-015-1927-1) contains supplementary material, which is available to authorized users.

✉ Jeong Hwan Lee
jhwlee90@sejong.ac.kr

✉ Jeong-Kook Kim
jkkim@korea.ac.kr

¹ Division of Life Sciences, Korea University, Anam-dong 5 ga, Seongbuk-gu, Seoul 136-701, Republic of Korea

² Department of Bioresource Engineering, Sejong University, 98 Gunja-dong, Gwangjin-gu, Seoul 143-747, Republic of Korea

³ Department of Genetic Engineering, Dong-A University, Busan 604-714, Republic of Korea

⁴ Center for Desert Agriculture, Division of Biological and Environmental Sciences and Engineering, King Abdullah University of Science and Technology, Thuwal 23955-6900, Saudi Arabia

Keywords AtNF-YB/YC · Flowering time · NUCLEAR FACTOR Y transcription factors · OsNF-YB/YC · Protein–protein interaction

Abbreviations

CO CONSTANS
NF-Y NUCLEAR FACTOR Y transcription factors
FT FLOWERING LOCUS T
GFP Green fluorescence protein
GST Glutathione S-transferase
HAP Heme-activated protein

LD	Long-day
ORFs	Open reading frames
SD	Short-day

Introduction

NUCLEAR FACTOR Y transcription factors [NF-Y, also known as heme-activated protein (HAP) in yeast] as important regulators control numerous genes in all eukaryotes (Edwards et al. 1998; Maity and de Crombrugghe 1998; Laloum et al.). The NF-Y transcription factors form a heterotrimeric complex composed of three unique subunits called NF-YA, NF-YB, and NF-YC (McNabb et al. 1995; Maity and de Crombrugghe 1998; Mantovani 1999). In filamentous fungi and mammals, the NF-YB and NF-YC proteins assemble as a heterodimer in the cytoplasm, which is translocated into the nucleus where it interacts with NF-YA and thereby forms active heterotrimeric complexes (Frontini et al. 2004; Steidl et al. 2004; Goda et al. 2005; Tuncher et al. 2005). The active complex then directly binds to highly conserved CCAAT motifs within the promoter of target genes. *In silico* analyses have revealed that approximately 30 % of eukaryotic promoters have the predicted binding sites of NF-Y transcription factors (Bucher 1990; Testa et al. 2005). This suggests that the regulation of gene expression by NF-Y transcription factor complex is widespread in many growth and developmental processes.

NF-Y subunits identified in plants play an important role in various aspects of developmental and stress-induced responses (Meinke et al. 1994; Li et al. 2008; Yamamoto et al. 2009; Petroni et al. 2012; Fornari et al. 2013; Laloum et al. 2013; Mu et al. 2013; Zhang et al. 2015). Many studies have revealed that they act as floral regulators in the modulation of flowering time. Individual homologs of NF-YB and NF-YC subunits are found as the interacting partners of CONSTANS (CO), an important regulator in the photoperiod pathway, in tomato and *Arabidopsis thaliana* (Ben-Naim et al. 2006; Wenkel et al. 2006). The *AtNF-YB2* and *AtNF-YB3* genes promote the onset of flowering through the induction of the *FLOWERING LOCUS T (FT)* under long-day (LD) conditions in *Arabidopsis* (Cai et al. 2007; Chen et al. 2007; Kumimoto et al. 2008). The *AtNF-YC* genes (*AtNF-Y3*, *AtNF-Y4*, and *AtNF-Y9*) also have functional redundancy in the activation of photoperiodic flowering through the activation of *FT* under LD conditions in *Arabidopsis* (Kumimoto et al. 2010). In addition, recent biochemical studies of *Arabidopsis* have shown that *AtNF-YB* proteins physically interact with *AtNF-YC* proteins to form the entire heterotrimeric complex, prior to binding to the consensus

CCAAT motif of their target genes (Calvenzani et al. 2012; Hackenberg et al. 2012b). The three *AtNF-YC* proteins (*AtNF-YC3*, *AtNF-YC4*, and *AtNF-YC9*) are also known to interact with the *AtNF-YB2*, *AtNF-YB3* and CO proteins (Kumimoto et al. 2010), suggesting that the *AtNF-YC* subunits make at least six possible complexes with *AtNF-YB2* or *AtNF-YB3* and CO to fine-tune the timing of flowering under a variety of environmental conditions (Wenkel et al. 2006; Kumimoto et al. 2010).

Monocot *NF-YB* and *NF-YC* genes homologous to *AtNF-YB* and *AtNF-YC*, respectively, have been shown to regulate flowering time. In rice, *OsNF-YB11/OsHAP3H/Ghd8/DTH8* has been revealed to regulate flowering time under LD and short-day (SD) conditions by changes in *Early heading date 1 (Ehd1)*, *RICE FLOWERING LOCUS T 1 (RFT1)*, and *Heading date 3a (Hd3a)* expression (Wei et al. 2010; Yan et al. 2011). Recently, our group reported that *OsNF-YC2* and *OsNF-YC4* inhibit flowering via decreased expression of *Ehd1*, *RFT1* and *Hd3a* under LD conditions in rice, whereas *OsNF-YC6* promotes flowering via increased expression under the same conditions (Kim et al. 2015). Also, *OsNF-YC2/4/6* proteins physically interacted with *OsNF-YB8/10/11* proteins. In addition, overexpression of *Brachypodium distachyon NF-YB3* (*BdNF-YB3*) and *BdNF-YB6* in *Arabidopsis nf-yb2 nf-yb3* double mutant results in flowering phenotypes similar to wild-type plants; *BdNF-YB3* and *BdNF-YB6* are found to physically interact with *AtNF-YC3*, *AtNF-YC4*, and *AtNF-YC9* using yeast-two hybrid analysis (Cao et al. 2011). Furthermore, *Ghd7/OsI*, a rice CCT domain-containing protein, regulates flowering under LD conditions (Xue et al. 2008). However, the protein–protein interactions between *OsNF-Y* proteins and Heading date 1 (*Hd1*), an *Arabidopsis* CO homolog, have not yet been characterized. In wheat (*Triticum aestivum*), the CCT domain of flowering promoter *CO2*, a rice *Hd1* homolog, and flowering repressor *VRN2* (*ZCCT1* and *2*) physically interact with *TaNF-YB*, and these CCT proteins compete with *TaNF-YA* for interaction with *TaNF-YB* (Li et al. 2011). These data suggest the functional conservation of monocot NF-Y transcription factors through a conserved molecular mechanism in the control of flowering time. However, our current knowledge about the role of *OsNF-Y* transcription factors in the regulation of flowering remains limited.

AtNF-YB2/3 and *AtNF-YC1/2/3/4/9* are known to control flowering time in *Arabidopsis* (Chen et al. 2007; Kumimoto et al. 2008, 2010; Hackenberg et al. 2012a). According to phylogenetic tree analysis, *OsNF-YB8/10/11*, *OsNF-YC1/4*, *OsNF-YC2* and *OsNF-YC6/7* are clustered with *AtNF-YB2/3*, *AtNF-YC2*, *AtNF-YC1/4*, and *AtNF-YC3/9*, respectively (Thirumurugan et al. 2008; Petroni et al. 2012; Laloum et al. 2013). Thus, we investigated the effects of the overexpression of *OsNF-YB* and *OsNF-YC*

genes in *Arabidopsis nf-yb* double mutants and *nf-yc* triple mutants, respectively, to understand the functional conservation of these genes. In addition, we examined the protein–protein interactions between OsNF-YB and AtNF-YC or OsNF-YC and AtNF-YB.

Materials and methods

Plant materials and growth conditions

Wild-type or transgenic *Arabidopsis* plants [ecotype Columbia (Col-0)] were grown in soil or Murashige and Skoog (MS) medium at 23 °C under LD conditions (16:8 h light:dark photoperiod) at a light intensity of 120 $\mu\text{mol m}^{-2} \text{s}^{-1}$. *nf-yb2 nf-yb3* and *nf-yc3 nf-yc4 nf-yc9* mutants (Cai et al. 2007; Kumimoto et al. 2008, 2010) were used for plant transformation to investigate the function of OsNF-YB8, OsNF-YB10, OsNF-YB11, OsNF-YC1, OsNF-YC2, OsNF-YC4, OsNF-YC6, and OsNF-YC7.

Plasmid construction

For the transgenic approach, full-length open reading frames (ORFs) for OsNF-YB8, OsNF-YB10, OsNF-YB11, OsNF-YC1, OsNF-YC2, OsNF-YC4, OsNF-YC6, and OsNF-YC7 were amplified from total RNA of wild-type rice seedlings (*Oryza sativa* L., cv. Hwaan; a gift from National Institute of Agricultural Biotechnology, Suwon, South Korea) by reverse transcription (RT)-polymerase chain reaction (PCR) using gene-specific primer sets (Supplementary Table S1). The amplicons were digested with several restriction enzymes and the resulting restriction fragments were subcloned into pCHF3, a plant transformation vector harboring the *CaMV 35S* promoter. For in vitro protein expression, the ORFs of AtNF-YB2, AtNF-YB3, AtNF-YC3, AtNF-YC4, AtNF-YC9, OsNF-YB8, OsNF-YB10, OsNF-YB11, OsNF-YC1, OsNF-YC2, OsNF-YC4, OsNF-YC6, and OsNF-YC7 were cloned into the pGEX-5X-1 (GE Healthcare, LC, UK) or pET21a-d(+) vector (GE Healthcare). Oligonucleotide sequences used for cloning are provided in Supplementary Table S1. For subcellular localization analysis, the ORFs of AtNF-YB2, AtNF-YB3, AtNF-YC3, AtNF-YC4, and AtNF-YC9 were cloned into pCAMBIA1300 or pCAMBIA2300 vectors. Oligonucleotide sequences used for cloning are provided in Supplementary Table S1. For bimolecular fluorescence complementation (BiFC) analysis, the ORFs of AtNF-YB2, AtNF-YB3, AtNF-YC3, AtNF-YC4, AtNF-YC9, OsNF-YB8, OsNF-YB10, OsNF-YB11, OsNF-YC1, OsNF-YC2, OsNF-YC4, OsNF-YC6, and OsNF-YC7 were cloned into pSPYNE-35S or pSPYCE-35S vectors (Walter et al. 2004).

Oligonucleotide sequences used for cloning are provided in Supplementary Table S1. The resulting recombinant plasmid was sequenced to verify the absence of PCR errors during amplification.

Generation of transgenic plants and measurement of flowering time

Transgenic plants were generated using the floral dip method with minor modifications (Weigel and Glazebrook 2002). Kanamycin-resistant transgenic seedlings whose genotype was confirmed via PCR were transferred to the soil. At least 20–30 T₁ seedlings were analyzed for each construct. To measure flowering time, the total number of rosette and cauline leaves of at least 6 independent transgenic lines (average 5 individual plants per independent transgenic line) was counted in the T₂ generation (Lee et al. 2012). At least 10 individual plants were used for measuring flowering time in the T₃ or T₄ generations. To determine the significant difference in flowering time of transgenic plants compared with that of *Arabidopsis* double or triple mutants, the data was analyzed using SPSS software version 12.0 (Sbaihat et al. 2015).

Expression analysis

Total RNA was isolated from *Arabidopsis* (Col-0) whole seedlings using the Plant RNA Purification Reagent (Invitrogen, Carlsbad, CA, USA). RNA quality was determined with a Nanodrop ND-2000 spectrophotometer (Nanodrop Technologies), and only high quality RNA samples ($A_{260}/A_{230} > 2.0$ and $A_{260}/A_{280} > 1.8$) were used for subsequent experiments. To remove possible genomic DNA contamination, RNA samples were treated with DNaseI for 60 min at 37 °C. RNA (1 μg) was used for cDNA synthesis, in accordance with the manufacturer's instructions (Roche Applied Science, Madison, WI, USA). Gene expression levels were analyzed via RT-PCR or real time (RT)-quantitative polymerase chain reaction (qPCR) methods, as previously described (Lee et al. 2005; Hong et al. 2010). RT-qPCR analysis was carried out in 384-well plates with a LightCycler 480 (Roche Applied Science) using LightCycler 480 SYBR green master (Roche Applied Science) to monitor the PCR amplification. *PP2AA3* (AT1G13320) was used as a reference gene, according to “The eleven golden rules for quantitative RT-PCR” (Udvardi et al. 2008; Hong et al. 2010). All RT-qPCR experiments were carried out in two biological replicates (independently harvested samples) with three technical replicates, each with similar results. Oligonucleotide sequences used for the expression analysis are provided in Supplementary Table S2.

In vitro protein–protein interaction analysis

Glutathione *S*-transferase (GST) fusion recombinant proteins were mixed with the *Escherichia coli* lysates, and the mixtures were gently rotated for 2 h at 4 °C. Subsequently, they were washed three times with the washing buffer and eluted with 10 mM reduced glutathione in 100 mM NaCl and 20 mM Tris–HCl (pH 7.2). Finally, the eluted protein samples were analyzed by 12.5 % SDS-PAGE and visualized by western blot analysis. The detailed procedure has been previously reported (Jang et al. 2009).

Subcellular localization and BiFC analyses

For transient expression in tobacco (*Nicotiana benthamiana*) leaves, the *Agrobacterium tumefaciens* strain C58C1 harboring the various combinations of constructs was infiltrated into the abaxial sides of 3-week-old tobacco plants. Subsequently, epidermal cells of infiltrated tobacco leaves were examined for fluorescence using a confocal microscope (LSM 510 META, Carl Zeiss, Germany). The detailed procedure has been previously reported (Jang et al. 2009).

Results and discussion

Overexpression of *OsNF-YB8/10/11* rescues the late flowering phenotype of *Arabidopsis nf-yb2 nf-yb3* double mutants


AtNF-YB2 and *AtNF-YB3* regulate flowering time in *Arabidopsis* (Chen et al. 2007; Kumimoto et al. 2008) and three rice *NF-YB* genes (*OsNF-YB8*, *OsNF-YB10*, and *OsNF-YB11*) are clustered with *AtNF-YB2* and *AtNF-YB3* (Thirumurugan et al. 2008; Petroni et al. 2012; Laloum et al. 2013). To assess the possible functional conservation of biological activities between rice and *Arabidopsis* homologues, we expressed the respective rice *OsNF-YB* genes (*OsNF-YB8*, *OsNF-YB10*, and *OsNF-YB11*) under the control of the *35S* promoter in *Arabidopsis nf-yb2 nf-yb3* double mutants. For analysis of flowering time, we selected six independent transgenic lines showing a Mendelian inheritance (3:1 ratio) of kanamycin resistance in the T₂ generation. In this experiment, we measured flowering time of individual plants in six independent lines overexpressing the respective *OsNF-YB8*, *OsNF-YB10*, and *OsNF-YB11* under LD conditions. *OsNF-YB8/10/11* transgenes were able to rescue the late flowering phenotype of *nf-yb2 nf-yb3* mutants, although the effect of *OsNF-YB11* on the complementation of the *nf-yb2 nf-yb3* mutation was very weak (Fig. 1a–d, Supplementary Table S3, Supplementary Fig. S1). Some *35S::OsNF-YB8 nf-yb2 nf-yb3* and *35S::OsNF-YB10 nf-yb2 nf-yb3* transgenic lines flowered

earlier than wild-type plants showing 12.8 leaves [line 7 (11.3 leaves) in *35S::OsNF-YB8 nf-yb2 nf-yb3*, and lines 17 (9.5 leaves) and 18 (8.5 leaves) in *35S::OsNF-YB10 nf-yb2 nf-yb3*]. The degree of rescue was correlated with the expression levels of *OsNF-YB8* and *OsNF-YB10* in *35S::OsNF-YB8 nf-yb2 nf-yb3* and *35S::OsNF-YB10 nf-yb2 nf-yb3* plants, respectively (Supplementary Fig. S2). Because the degree of overexpression of *OsNF-YB11* in transgenic plants was similar to that of *OsNF-YB8* or *OsNF-YB10* in transgenic plants (Supplementary Fig. S2), the comparatively weak rescue seen in *35S::OsNF-YB11 nf-yb2 nf-yb3* plants cannot be explained by the differential expression of transgenes among transgenic plants. This result suggests the stronger effect of *OsNF-YB8* and *OsNF-YB10* in rescuing the late flowering phenotype of *nf-yb2 nf-yb3* mutants, compared with that of *OsNF-YB11*. This notion was supported by the findings that *OsNF-YB8* and *OsNF-YB10* are closer to *AtNF-YB2/3* in the phylogenetic tree (Thirumurugan et al. 2008; Petroni et al. 2012; Laloum et al. 2013). These results suggest that *OsNF-YB8/10/11* genes are functional equivalents of *AtNF-YB2/3* genes in rice.

Overexpression of *OsNF-YC2* rescues the late flowering phenotype of *Arabidopsis nf-yc3 nf-yc4 nf-yc9* triple mutants

AtNF-YC1, *AtNF-YC2*, *AtNF-YC3*, *AtNF-YC4*, and *AtNF-YC9* are known to regulate flowering time in *Arabidopsis* (Kumimoto et al. 2010; Hackenberg et al. 2012a), and *OsNF-YC1/4*, *OsNF-YC2*, and *OsNF-YC6/7* are clustered with *AtNF-YC2*, *AtNF-YC1/4*, and *AtNF-YC3/9*, respectively (Thirumurugan et al. 2008; Petroni et al. 2012; Laloum et al. 2013). Therefore, to assess the possible functional conservation between rice and *Arabidopsis* homologues, we overexpressed five rice *OsNF-YC* genes (*OsNF-YC1*, *OsNF-YC2*, *OsNF-YC4*, *OsNF-YC6*, and *OsNF-YC7*) under the *35S* promoter in *Arabidopsis nf-yc3 nf-yc4 nf-yc9* triple mutants. To measure flowering time, we selected at least seven independent transgenic lines showing a single transgene insertion in the T₂ generation. We also analyzed flowering time of individual plants in transgenic lines overexpressing the respective *OsNF-YC1*, *OsNF-YC2*, *OsNF-YC4*, *OsNF-YC6*, and *OsNF-YC7* genes under LD conditions. The *OsNF-YC2* gene was able to rescue the late flowering phenotype of *nf-yc3 nf-yc4 nf-yc9* mutants, but *OsNF-YC1/4/6/7* genes were not able to significantly rescue this phenotype of these mutants (Fig. 2a, b; Supplementary Table S3; Supplementary Fig. S3). Furthermore, some *35S::OsNF-YC2 nf-yc3 nf-yc4 nf-yc9* transgenic lines flowered earlier than wild-type plants showing 9.1 leaves [lines 6 (8.6 leaves), 17 (7.7 leaves), 19 (8.0 leaves), 21 (9.0 leaves), and 22 (9.0 leaves)]. The degree of rescue was correlated with the expression levels of *OsNF-YC2* in

Fig. 1 Flowering phenotypes of transgenic plants overexpressing *OsNF-YB8* and *OsNF-YB10* in *Arabidopsis nf-yb2 nf-yb3* double mutants. Flowering time of *35S::OsNF-YB8 nf-yb2 nf-yb3* (a) and *35S::OsNF-YB10 nf-yb2 nf-yb3* (b) plants in the T₂ generation under LD conditions. Homozygous and hemizygous transgenic plants were used to measure flowering time. Asterisks indicate significant difference in flowering time of transgenic plants compared with that of double mutants (Student's *t* test, **P* < 0.05, ***P* < 0.01). Error bars indicate standard deviation (SD). Phenotypes of *35S::OsNF-YB8 nf-yb2 nf-yb3* (c) and *35S::OsNF-YB10 nf-yb2 nf-yb3* (d) plants in the T₂ generation under LD conditions. Photographs were captured when Col-0 plants flowered. Scale bars 2 cm



35S::OsNF-YC2 nf-yc3 nf-yc4 nf-yc9 plants (Supplementary Fig. S4). However, non-rescued flowering phenotypes seen in *35S::OsNF-YC4 nf-yc3 nf-yc4 nf-yc9* or *35S::OsNF-YC7 nf-yc3 nf-yc4 nf-yc9* plants cannot be explained by the differential expression of transgenes among transgenic plants (Supplementary Fig. S4), suggesting that *OsNF-YC2* exerts the strongest effect on the complementation of the *nf-yc3 nf-yc4 nf-yc9* mutation. Similarly, *OsNF-YC2* could strongly rescue the late flowering phenotype of *nf-yc3 nf-yc9* double mutants compared to other *OsNF-YC* genes (Supplementary Fig. S5, Supplementary Table S3). Although *OsNF-YC2* is closer to *AtNF-YC4* than to *AtNF-YC3/9* in the phylogenetic tree (Thirumurugan et al. 2008; Petroni et al. 2012; Laloum et al. 2013), these results indicate that *OsNF-YC2* is a functional equivalent of *AtNF-YC3/4/9* in rice.

OsNF-YC2* regulates *FT* and *SOC1* expression in *Arabidopsis nf-yc3 nf-yc4 nf-yc9* triple mutants overexpressing *OsNF-YC2

Because most *35S::OsNF-YC2 nf-yc3 nf-yc4 nf-yc9* plants showed wild-type-like flowering phenotypes (Fig. 2), we examined the expression of flowering time genes in

35S::OsNF-YC2 nf-yc3 nf-yc4 nf-yc9 plants under LD conditions. Overexpression of *OsNF-YC2* still rescued the late flowering phenotype of *nf-yc3 nf-yc4 nf-yc9* mutants in the T₃ generation (Fig. 3a), indicating that the phenotypes of *35S::OsNF-YC2 nf-yc3 nf-yc4 nf-yc9* plants are stably inherited.

Because *AtNF-YC3/4/9* regulate photoperiodic flowering time in *Arabidopsis* (Kumimoto et al. 2010), we checked the expression levels of photoperiod pathway genes [*GIGANTEA (GI)*, *CO*, *FT*, and *SOC1*] in transgenic plants in the T₄ generation under LD conditions. We found that *FT* and *SOC1* expression was significantly increased compared with *nf-yc3 nf-yc4 nf-yc9* mutants (Fig. 3b, c), indicating that *OsNF-YC2* regulates flowering time via the activation of *FT* and *SOC1*. In contrast, *GI* and *CO* expression remained unaltered in transgenic plants (Supplementary Fig. S6). This was consistent with the previous finding that *AtNF-YC3/4/9* controls *FT* expression with *CO* and other *AtNF-Y* components under LD conditions (Kumimoto et al. 2010). These results suggest that the *OsNF-YC2* promotes flowering via upregulation of *FT* and *SOC1* expression in *Arabidopsis* plants.




Fig. 2 Flowering phenotypes of transgenic plants overexpressing *OsNF-YC2* in *Arabidopsis nf-yc3 nf-yc4 nf-yc9* triple mutants. **a** Flowering time of *35S::OsNF-YC2 nf-yc3 nf-yc4 nf-yc9* plants in the T_2 generation under LD conditions. Homozygous and hemizygous transgenic plants were used to measure flowering time. Asterisks indicate significant differences in flowering time of transgenic plants compared with that of triple mutants (Student's *t* test, $*P < 0.05$, $**P < 0.01$). Error bars indicate SD. **b** Phenotypes of *35S::OsNF-YC2 nf-yc3 nf-yc4 nf-yc9* plants in the T_2 generation under LD conditions. Photographs were captured when Col-0 plants flowered. Scale bars 2 cm

OsNF-YB and OsNF-YC proteins interact with AtNF-YC and AtNF-YB proteins, respectively

AtNF-YC3, AtNF-YC4, and AtNF-YC9 regulate photoperiod-dependent flowering time via protein–protein interaction with AtNF-YB2, AtNF-YB3, and CO proteins in *Arabidopsis* (Kumimoto et al. 2010). In addition, NF-Y transcription factors control the expression of downstream target genes via direct binding to CCAAT consensus motifs within the promoter in the nucleus (Frontini et al. 2004; Steidl et al. 2004; Goda et al. 2005; Tuncher et al. 2005). Therefore, we initially examined the subcellular localization patterns of AtNF-YB and AtNF-YC proteins in tobacco leaves. The AtNF-YC3/4/9-GFP proteins were localized in both the nucleus and the cytoplasm, whereas the AtNF-YB2/3-GFP proteins were present only in the cytoplasm (Supplementary Fig. S7). In addition, the OsNF-YB11-GFP and OsNF-YC1/2/4/6/7-GFP proteins were found in both nucleus and cytoplasm, whereas the OsNF-YB8-GFP and OsNF-YB10-GFP proteins were present only in the cytoplasm and nucleus, respectively (Kim et al. 2015). Despite the different subcellular localizations of OsNF-YB8 and OsNF-YB10 proteins, the overexpression of either *OsNF-YB8* or *OsNF-YB10* rescued the late flowering phenotype of *nf-yb2 nf-yb3* mutants (Fig. 1). This suggests that the localization patterns of OsNF-YBs/AtNF-




Fig. 3 Expression analysis of flowering time genes in *35S::OsNF-YC2 Arabidopsis nf-yc3 nf-yc4 nf-yc9* plants. **a** Flowering time of *35S::OsNF-YC2 nf-yc3 nf-yc4 nf-yc9* plants in the T_3 generation under LD conditions. Homozygous transgenic plants were used to measure flowering time. Asterisks indicate significant differences in flowering time of transgenic plants compared with that of triple mutants (Student's *t* test, $**P < 0.01$). Error bars indicate SD. **b** Expression levels of *FT* and *SOC1* in 8-day-old *35S::OsNF-YC2 nf-yc3 nf-yc4 nf-yc9* plants in the T_4 generation under LD conditions. Samples for RT-qPCR were harvested at Zeitgeber time (ZT) 16. Expression levels of each gene in Col-0 plants were set to one

YCs protein complexes may be important for their function in the regulation of flowering time. This notion is supported by the observation that NF-Y proteins, which are present only in the cytoplasm, were transported into the nucleus via a “piggy-back” mechanism involving interacting NF-Y proteins (Steidl et al. 2004). This result indicates that the subcellular localization patterns of the AtNF-YB/YC and OsNF-YB/YC proteins overlap *in planta*.

The overlapping localization patterns of the AtNF-YB/YC-GFP proteins with the OsNF-YB/YC-GFP proteins raise the possibility of the protein–protein interaction between OsNF-YB and AtNF-YC proteins, or OsNF-YC and AtNF-YB proteins. We studied protein–protein interactions *in vitro* and *in planta*. For the *in vitro* pull-down assay, AtNF-YBs/YCs and OsNF-YBs/YCs were fused with GST or GFP and His proteins, respectively. For the BiFC assay in tobacco leaves, the N- or C-terminal yellow fluorescent protein (YFP) fragment was fused with AtNF-YBs,

AtNF-YCs, OsNF-YBs, and OsNF-YCs constructs (Walter et al. 2004). The GFP and His-tagged OsNF-YB10 protein significantly interacted with two AtNF-YC proteins (AtNF-YC3 and AtNF-YC9), compared with the other two OsNF-YB proteins (OsNF-YB8 and OsNF-YB11) (Fig. 4a; Supplementary Figs. S8a, b). This suggests that the strong effect of OsNF-YB10 complementing the *nf-yb2 nf-yb3* mutation (Fig. 1a–d; Supplementary Table S3; Supplementary Fig. S1) may be caused by the significant physical interaction between the OsNF-YB10 and AtNF-YC




Fig. 4 Protein–protein interaction between OsNF-YB10 and three AtNF-YC proteins in vitro and *in planta*. **a** In vitro pull-down assay. Glutathione *S*-transferase (GST) or GST-tagged AtNF-YC proteins were incubated with green fluorescent protein (GFP) and Histidine (His)-tagged OsNF-YB10 protein. The eluates were separated by 12.5 % SDS-PAGE, transferred to polyvinylidene difluoride (PVDF) membranes, and probed with anti-His antibody. About 10 % of the total sample in each reaction was loaded as an input control. Ponceau S-stained bands indicated by arrows show the amount and quality of the GST fusion proteins used in this assay. **b** Bimolecular fluorescence complementation (BiFC) analysis. A BiFC assay was performed using tobacco (*Nicotiana benthamiana*) leaf epidermal cells. AtNF-YC3/4/9 and OsNF-YB10 genes were fused to the N- and C-terminal yellow fluorescent protein (YFP) fragment genes, respectively. YFP signal between bZIP63-YFP^N and bZIP63-YFP^C was used as a positive control. There were no YFP signals between YFP^N and OsNF-YB10-YFP^C or AtNF-YC3/4/9-YFP^N and YFP^C used as negative controls. (I) indicates YFP fluorescence and (II) indicates bright field images. Scale bars 50 μ m

proteins. YFP signals of three OsNF-YB and three AtNF-YC proteins were also observed in the nucleus and cytoplasm of the transformed tobacco epidermal cells (Fig. 4b, Supplementary Fig. S8c).

In the protein–protein interaction between the OsNF-YC and AtNF-YB proteins, the GFP and His-tagged OsNF-YC2 protein significantly bound to AtNF-YB3 protein, compared with the other two OsNF-YC proteins (OsNF-YC1 and OsNF-YC4) (Fig. 5a; Supplementary Figs. S9a, b). This suggests that the significant interaction between OsNF-YC2 and AtNF-YB proteins may result in the strong effect of OsNF-YC2 in rescuing the late flowering phenotype of *nf-yc3 nf-yc4 nf-yc9* mutants (Fig. 2a, b; Supplementary Table S3; Supplementary Fig. S3). In addition, YFP signals of OsNF-YC2 and two AtNF-YB proteins were detected in the nucleus and cytoplasm of the tobacco epidermal cells (Fig. 5b). Other OsNF-YC proteins (OsNF-YC1, OsNF-YC4, OsNF-YC6, and OsNF-YC7) interacted with AtNF-YB2 and AtNF-YB3 proteins *in vitro* and *in planta* (Supplementary Fig. S9c). These results indicate that OsNF-YB and OsNF-YC proteins could physically interact with AtNF-YC and AtNF-YB proteins, respectively, *in vitro* and *in planta*.

Collectively, our results suggest that the functions of some OsNF-YB and OsNF-YC genes involved in the control of flowering time are conserved in *Arabidopsis*. In addition, the present study indicated that OsNF-YB and OsNF-YC proteins interact with AtNF-YC and AtNF-YB proteins, suggesting that the functional NF-Y complexes generated by the interaction between OsNF-YB and AtNF-YC, or OsNF-YC and AtNF-YB proteins can effectively bind the genomic regions of downstream targets, which could rescue the late flowering phenotype of *Arabidopsis* mutants. However, some OsNF-YC genes did not complement the *nf-yc3 nf-yc4 nf-yc9* mutation, although they interacted with AtNF-YB2/3 proteins. This suggests their different function among OsNF-YC genes in the regulation of flowering time. This notion is supported by the observation that overexpression of the OsNF-YC2 or OsNF-YC4 inhibited flowering in rice, whereas overexpression of the OsNF-YC6 in rice promoted flowering (Kim et al. 2015). In addition, the overexpression effect of OsNF-YB11 on the late flowering phenotype of *nf-yb2 nf-yb3* plants was weaker than those of OsNF-YB8/10, although the overexpression of OsNF-YB11/OsHAP3H/Ghd8/DTH8 known as a rice floral repressor in *Arabidopsis* led to flower about 10 days earlier than wild-type plants only under LD conditions (Yan et al. 2011). This discrepancy may be due to the different light conditions, because they transferred the seedlings grown under day-neutral conditions into the LD conditions (Yan et al. 2011).

Considering that some NF-Y components are involved in stress-induced processes such as ABA signaling and drought resistance (Li et al. 2008; Zhang et al. 2015),




Fig. 5 Protein–protein interaction between OsNF-YC2 and two AtNF-YB2/3 proteins *in vitro* and *in planta*. **a** *In vitro* pull-down assay. GST or GST-tagged AtNF-YB proteins were incubated with GFP and His-tagged OsNF-YC2 protein. The eluates were separated by 12.5 % SDS-PAGE, transferred to PVDF membranes, and probed with anti-His antibody. About 4 % of the total sample in each reaction was loaded as an input control. Ponceau S-stained bands indicated by arrows show the amount and quality of the GST fusion proteins used

in this assay. **b** BiFC analysis in tobacco leaf epidermal cells. AtNF-YB2/3 and OsNF-YC2 genes were fused to the N- and C-terminal YFP fragment genes, respectively. YFP signal between bZIP63-YFP^N and bZIP63-YFP^C shown in Fig. 4b was used as a positive control. There were no YFP signals between YFP^N and OsNF-YC2-YFP^C or AtNF-YB2/3-YFP^N and YFP^C used as negative controls. (I) indicates YFP fluorescence and (II) indicates bright field images. Scale bars 50 μ m

further investigation is needed on whether some OsNF-YC genes act in stress responses.

Author contribution statement J-KK conceived and designed the research. Y-HH, S-KK, and KCL conducted experiments. Y-HH, S-KK, KCL, YSC, JHL, and J-KK analyzed data. JHL and J-KK wrote the manuscript. All authors read and approved the manuscript.

Acknowledgments We thank Prof. Ben F. Holt III for providing *nf-yb2 nf-yb3* and *nf-yc3 nf-yc4 nf-yc9* mutants and other mutants not used in this study. This work was supported by the Next-Generation BioGreen 21 Program (PJ007978), Rural Development Administration, Republic of Korea and a Korea University Grant (to J.-K. Kim), Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2015R1D1A4A0101941 to J.H. Lee and 2014R1A1A2055796 to J.-K. Kim), and by the BK 21 program (to S.-K. Kim).

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

Ben-Naim O, Eshed R, Parnis A, Teper-Bamnolker P, Shalit A, Coupland G, Samach A, Lifschitz E (2006) The CCAAT binding

factor can mediate interactions between CONSTANS-like proteins and DNA. *Plant J* 46:462–476

Bucher P (1990) Weight matrix descriptions of four eukaryotic RNA polymerase II promoter elements derived from 502 unrelated promoter sequences. *J Mol Biol* 212:563–578

Cai X, Ballif J, Endo S, Davis E, Liang M, Chen D, DeWald D, Kreps J, Zhu T, Wu Y (2007) A putative CCAAT-binding transcription factor is a regulator of flowering timing in Arabidopsis. *Plant Physiol* 145:98–105

Calvenzani V, Testoni B, Gusmaroli G, Lorenzo M, Gnesutta N, Petroni K, Mantovani R, Tonelli C (2012) Interactions and CCAAT-binding of *Arabidopsis thaliana* NF-Y subunits. *PLoS ONE* 7:e42902

Cao S, Kumimoto RW, Siriwardana CL, Risinger JR, Holt BF 3rd (2011) Identification and characterization of NF-Y transcription factor families in the monocot model plant *Brachypodium distachyon*. *PLoS One* 6:e21805

Chen NZ, Zhang XQ, Wei PC, Chen QJ, Ren F, Chen J, Wang XC (2007) AtHAP3b plays a crucial role in the regulation of flowering time in *Arabidopsis* during osmotic stress. *J Biochem Mol Biol* 40:1083–1089

Edwards D, Murray JA, Smith AG (1998) Multiple genes encoding the conserved CCAAT-box transcription factor complex are expressed in Arabidopsis. *Plant Physiol* 117:1015–1022

Fornari M, Calvenzani V, Masiero S, Tonelli C, Petroni K (2013) The Arabidopsis *NF-YA3* and *NF-YA8* genes are functionally redundant and are required in early embryogenesis. *PLoS One* 8:e82043

Frontini M, Imbriano C, Manni I, Mantovani R (2004) Cell cycle regulation of NF-YC nuclear localization. *Cell Cycle* 3:217–222

Goda H, Nagase T, Tanoue S, Sugiyama J, Steidl S, Tuncher A, Kobayashi T, Tsukagoshi N, Brakhage AA, Kato M (2005) Nuclear translocation of the heterotrimeric CCAAT binding

- factor of *Aspergillus oryzae* is dependent on two redundant localising signals in a single subunit. Arch Microbiol 184:93–100
- Hackenberg D, Keetman U, Grimm B (2012a) Homologous NF-YC2 Subunit from *Arabidopsis* and Tobacco Is Activated by Photooxidative Stress and Induces Flowering. Int J Mol Sci 13:3458–3477
- Hackenberg D, Wu Y, Voigt A, Adams R, Schramm P, Grimm B (2012b) Studies on differential nuclear translocation mechanism and assembly of the three subunits of the *Arabidopsis thaliana* transcription factor NF-Y. Mol Plant 5:876–888
- Hong SM, Bahn SC, Lyu A, Jung HS, Ahn JH (2010) Identification and testing of superior reference genes for a starting pool of transcript normalization in *Arabidopsis*. Plant Cell Physiol 51:1606–1694
- Jang YH, Park HY, Kim SK, Lee JH, Suh MC, Chung YS, Paek KH, Kim JK (2009) Survey of rice proteins interacting with OsFCA and OsFY proteins which are homologous to the *Arabidopsis* flowering time proteins, FCA and FY. Plant Cell Physiol 50:1479–1492
- Kim SK, Park HY, Jang YH, Lee KC, Chung YS, Lee JH, Kim JK (2015) OsNF-YC2 and OsNF-YC4 proteins inhibit flowering under long-day conditions in rice. Planta. doi:10.1007/s00425-015-2426-x
- Kumimoto RW, Adam L, Hymus GJ, Repetti PP, Reuber TL, Marion CM, Hempel FD, Ratcliffe OJ (2008) The Nuclear Factor Y subunits NF-YB2 and NF-YB3 play additive roles in the promotion of flowering by inductive long-day photoperiods in *Arabidopsis*. Planta 228:709–723
- Kumimoto RW, Zhang Y, Siefers N, Holt BF 3rd (2010) NF-YC3, NF-YC4 and NF-YC9 are required for CONSTANS-mediated, photoperiod-dependent flowering in *Arabidopsis thaliana*. Plant J 63:379–391
- Laloum T, De Mita S, Gamas P, Baudin M, Niebel A (2013) CCAAT-box binding transcription factors in plants: Y so many? Trends Plant Sci 18:157–166
- Lee JH, Cho YS, Yoon HS, Suh MC, Moon J, Lee I, Weigel D, Yun CH, Kim JK (2005) Conservation and divergence of FCA function between *Arabidopsis* and rice. Plant Mol Biol 58:823–838
- Lee JH, Park SH, Ahn JH (2012) Functional conservation and diversification between rice OsMADS22/OsMADS55 and *Arabidopsis* SVP proteins. Plant Sci 185–186:97–104
- Li WX, Oono Y, Zhu J, He XJ, Wu JM, Iida K, Lu XY, Cui X, Jin H, Zhu JK (2008) The *Arabidopsis* NFYA5 transcription factor is regulated transcriptionally and posttranscriptionally to promote drought resistance. Plant Cell 20:2238–2251
- Li C, Distelfeld A, Comis A, Dubcovsky J (2011) Wheat flowering repressor VRN2 and promoter CO2 compete for interactions with NUCLEAR FACTOR-Y complexes. Plant J 67:763–773
- Maity SN, de Crombrughe B (1998) Role of the CCAAT-binding protein CBF/NF-Y in transcription. Trends Biochem Sci 23:174–178
- Mantovani R (1999) The molecular biology of the CCAAT-binding factor NF-Y. Gene 239:15–27
- McNabb DS, Xing Y, Guarente L (1995) Cloning of yeast HAP5: a novel subunit of a heterotrimeric complex required for CCAAT binding. Genes Dev 9:47–58
- Meinke DW, Franzmann LH, Nickle TC, Yeung EC (1994) Leafy cotyledon mutants of *Arabidopsis*. Plant Cell 6:1049–1064
- Mu J, Tan H, Hong S, Liang Y, Zuo J (2013) *Arabidopsis* transcription factor genes *NF-YA1*, 5, 6, and 9 play redundant roles in male gametogenesis, embryogenesis, and seed development. Mol Plant 6:188–201
- Petroni K, Kumimoto RW, Gnesutta N, Calvenzani V, Fornari M, Tonelli C, Holt BF 3rd, Mantovani R (2012) The promiscuous life of plant NUCLEAR FACTOR Y transcription factors. Plant Cell 24:4777–4792
- Sbaihat L, Takeyama K, Koga T, Takemoto D, Kawakita K (2015) Induced resistance in *Solanum lycopersicum* by algal elicitor extracted from *Sargassum fusiforme*. Sci World J 2015:870520
- Steidl S, Tuncher A, Goda H, Guder C, Papadopoulou N, Kobayashi T, Tsukagoshi N, Kato M, Brakhage AA (2004) A single subunit of a heterotrimeric CCAAT-binding complex carries a nuclear localization signal: piggy back transport of the pre-assembled complex to the nucleus. J Mol Biol 342:515–524
- Testa A, Donati G, Yan P, Romani F, Huang TH, Vigano MA, Mantovani R (2005) Chromatin immunoprecipitation (ChIP) on chip experiments uncover a widespread distribution of NF-Y binding CCAAT sites outside of core promoters. J Biol Chem 280:13606–13615
- Thirumurugan T, Ito Y, Kubo T, Serizawa A, Kurata N (2008) Identification, characterization and interaction of HAP family genes in rice. Mol Genet Genomics 279:279–289
- Tuncher A, Sprote P, Gehrke A, Brakhage AA (2005) The CCAAT-binding complex of eukaryotes: evolution of a second NLS in the HapB subunit of the filamentous fungus *Aspergillus nidulans* despite functional conservation at the molecular level between yeast, *A. nidulans* and human. J Mol Biol 352:517–533
- Udvardi MK, Czechowski T, Scheible WR (2008) Eleven golden rules of quantitative RT-PCR. Plant Cell 20:1736–1737
- Walter M, Chaban C, Schutze K, Batistic O, Weckermann K, Nake C, Blazevic D, Grefen C, Schumacher K, Oecking C, Harter K, Kudla J (2004) Visualization of protein interactions in living plant cells using bimolecular fluorescence complementation. Plant J 40:428–438
- Wei X, Xu J, Guo H, Jiang L, Chen S, Yu C, Zhou Z, Hu P, Zhai H, Wan J (2010) *DTH8* suppresses flowering in rice, influencing plant height and yield potential simultaneously. Plant Physiol 153:1747–1758
- Weigel D, Glazebrook J (2002) *Arabidopsis: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
- Wenkel S, Turck F, Singer K, Gissot L, Le Gourrierec J, Samach A, Coupland G (2006) CONSTANS and the CCAAT box binding complex share a functionally important domain and interact to regulate flowering of *Arabidopsis*. Plant Cell 18:2971–2984
- Xue W, Xing Y, Weng X, Zhao Y, Tang W, Wang L, Zhou H, Yu S, Xu C, Li X, Zhang Q (2008) Natural variation in *Ghd7* is an important regulator of heading date and yield potential in rice. Nat Genet 40:761–767
- Yamamoto A, Kagaya Y, Toyoshima R, Kagaya M, Takeda S, Hattori T (2009) *Arabidopsis* NF-YB subunits LEC1 and LEC1-LIKE activate transcription by interacting with seed-specific ABRE-binding factors. Plant J 58:843–856
- Yan WH, Wang P, Chen HX, Zhou HJ, Li QP, Wang CR, Ding ZH, Zhang YS, Yu SB, Xing YZ, Zhang QF (2011) A major QTL, *Ghd8*, plays pleiotropic roles in regulating grain productivity, plant height, and heading date in rice. Mol Plant 4:319–330
- Zhang T, Zhang D, Liu Y, Luo C, Zhou Y, Zhang L (2015) Overexpression of a NF-YB3 transcription factor from *Picea wilsonii* confers tolerance to salinity and drought stress in transformed *Arabidopsis thaliana*. Plant Physiol Biochem 94:153–164