

José Aguilera · Angelika Dummermuth
Ulf Karsten · Raimund Schriek · Christian Wiencke

Enzymatic defences against photooxidative stress induced by ultraviolet radiation in Arctic marine macroalgae

Accepted: 12 January 2002 / Published online: 14 March 2002
© Springer-Verlag 2002

Abstract The activities of the antioxidant enzymes superoxide dismutase (SOD), glutathione reductase (GR), ascorbate peroxidase (APX) and catalase (CAT), as well as the content of the antioxidant compound ascorbic acid, were determined in five green, seven red and ten brown macroalgal species from the Kongsfjord (Spitsbergen, Svalbard, Norway). In general, higher antioxidant enzyme activities and a higher content of ascorbic acid were measured in green algae in comparison to red and brown algae. Species from the eulittoral and upper sublittoral (*Acrosiphonia penicilliformis*, *Monostroma arcticum*, *Chaetomorpha linum*, *Chaetomorpha melagonium*, *Devaleraea ramentacea*, *Palmaria palmata*) showed higher antioxidant enzyme activities compared to species from the lower sublittoral, indicating a more efficient biochemical protection in algae exposed to higher stress conditions in the field. The activity of GR was stimulated by artificial ultraviolet radiation in the green alga *M. arcticum*, and in the red algae *Coccotylus truncatus* and *Phycodrys rubens* after 84 h under continuous ex-

posure. GR activity was even higher when the UV exposure was followed by incubation in darkness for 24 h, indicating a higher elimination rate of toxic oxygen radicals under these conditions. *D. ramentacea*, *P. palmata* and *A. penicilliformis* did not show any significant effect of UV radiation on CAT, APX and SOD activities after 8 days of culture under laboratory conditions. However, a significant reduction in activities of GR and SOD was observed in *A. penicilliformis* when solar UV radiation was cut off by selective filter foils in the field, indicating a lower oxidative stress in the absence of UV radiation. Overall, the ecological success of macroalgae in the eulittoral and upper sublittoral is supported by an enhanced oxygen-reactive scavenging system, allowing fast acclimation to the changes in environmental radiation conditions.

J. Aguilera (✉) · A. Dummermuth · U. Karsten · R. Schriek
C. Wiencke

Alfred Wegener Institute
for Polar and Marine Research,
Am Handelshafen 12,
27570 Bremerhaven, Germany

Present address: J. Aguilera
Universidad de Málaga,
Facultad de Ciencias,
Departamento de Ecología,
Campus Universitario de Teatinos
s/n. 29071 Málaga, Spain,
e-mail: jaguilera@uma.es,
Fax: +34-95-2132000

Present address: U. Karsten
University of Rostock,
Institute of Aquatic Ecology,
18059 Rostock, Germany

Present address: R. Schriek
University Hohenheim,
Institute 140, Fruhwirthstraße 12,
70593 Stuttgart, Germany

Introduction

Studies on the ability of living organisms to cope with enhanced levels of ultraviolet radiation become more and more important due to the increasing depletion of stratospheric ozone. Recent data show a dramatic trend of ozone depletion over the Antarctic regions, with a temporal decrease in springtime below 25–30% of the undisturbed conditions (NASA <http://toms.gsfc.nasa.gov/ozone/ozone.html>). Strong reduction of the stratospheric ozone is now also evident in the northern hemisphere (Ott and Amanatides 1994; Schulz et al. 2001), and predictions indicate a gradual increment of ultraviolet (UV) radiation in the northern polar regions similar to the southern hemisphere (Stolarski et al. 1992).

Accurate information is needed to assess the potential effects in organisms caused by UV radiation reaching the earth's surface. The effects of UV radiation on growth and other physiological features, such as damage in DNA, RNA, proteins and photosynthesis, in a range of higher and lower plants including phytoplankton are relatively well documented (Aguilera et al. 1999a, b;

Bischof et al. 2000; Buma et al. 1995; Clendennen et al. 1996; Figueroa et al. 1997; Häder and Figueroa 1997; Karentz et al. 1991; Smith et al. 1992; Strid et al. 1990; Tevini and Teramura 1989) while investigations on the effects of UV radiation on benthic marine macroalgae are scarce, although this group plays an important ecological role in the marine environment.

Photosynthesis can be damaged due to high photosynthetically active radiation (PAR) or UV radiation, as a result of an overreduction of the photosynthetic electron transport when not enough electrons are drained off by NADP^+ to NADPH from reduced ferredoxin of PSI. UV radiation can affect the draining-off system by damaging proteins of the Calvin cycle like Rubisco (Allen 1977; Bischof et al. 2000). Thus, in the absence of NADP^+ , the reduced ferredoxin can also reduce oxygen, leading to superoxide radicals (O_2^-). Consequently, photosystem II is inactivated by UV and finally damaged due to degradation of the reaction centre proteins, mainly the D1 protein (Aro et al. 1993; Ohad et al. 1984). Under such conditions, singlet oxygen ($^1\text{O}_2$) can be formed from triplets of chlorophyll of the antenna. Reactive oxygen species can produce lipid peroxidation, damage proteins and have many other harmful effects (Asada and Takahashi 1987; Fridovich 1986).

Cellular mechanisms of protection against such toxic oxygen species are essential for the maintenance of photosynthetic activity and other metabolic functions (Allen 1977; Asada and Takahashi 1987; Eltsner 1982; Halliwell 1982). Plants and algae are equipped with an array of defence mechanisms that eliminate toxic oxygen radicals produced as by-products of photosynthesis and photooxidative events. Superoxide radicals are eliminated by the enzyme superoxide dismutase (SOD), yielding H_2O_2 and oxygen. Hydrogen peroxide itself is not particularly reactive with most biologically important molecules, but it is probably an intracellular precursor for more reactive oxidants, such as hydroxyl radicals. H_2O_2 is deprotonated by the enzyme catalase and by specific scavengers such as ascorbate and glutathione, mediated by ascorbate peroxidase and glutathione peroxidase, respectively. The resulting oxidized reactants, namely monodehydroascorbate and oxidized glutathione, are regenerated via enzymatic reductions by monodehydroascorbate reductase and glutathione reductase, respectively, thereby closing the antioxidant scavenging cycle (Polle 1996).

Studies related to UV-induced photooxidation to the scavenging mechanisms for protection against oxidative damage are rare for macroalgae. Thus, the ability to resist high radiation stress may be one of the major factors controlling vertical macroalgal zonation patterns in communities (Bischof et al. 1998a; Hanelt 1998), and may be mediated by a higher biochemical potential against oxidative stress. Long-term exposure under UV radiation has been demonstrated to induce the activity of superoxide dismutase and ascorbate peroxidase in microalgae (Lesser 1996a, b; Malanga and Puntarulo 1997). Activities of antioxidant enzymes were higher in shallow-water coral zooxanthellae than in specimens

collected from deeper waters (Shick et al. 1995). In addition, analysis of the antioxidant spectrum in selected alpine plant species collected at different altitudes proved that the total amount of antioxidants is positively correlated with altitude (Wildi and Lütz 1996). In one of the few publications on macroalgae, it was postulated that the differential stress tolerance associated with the vertical zonation of different *Fucus* species is strictly related to the antioxidant status of the plant, based mainly on species-specific differences of antioxidant enzyme activities (Collen and Davison 1999a, b).

Despite the potential importance in providing an alternative sink for excessively absorbed radiation energy and their role in scavenging, little is known about the capacity or inducibility of macroalgal antioxidant enzyme systems. The present study was designed to characterize the oxidative stress tolerance in field material of different green, red and brown macroalgae from the Arctic by the analysis of a set of antioxidant enzyme activities and the ascorbic acid content, as well as by the response to UV radiation.

Materials and methods

Algal material and study site

The macroalgal species studied and their depth distribution are listed in Table 1. Plants were collected by scuba divers in summer 1998 at the study site in the Kongsfjord (Ny-Ålesund, Spitsbergen, Norway 78°55.5'N; 11°56.0'E) from depths between 0 and 20 m. Algal samples were collected in black bags to avoid exposure to high irradiance during transport. Material for enzymatic activities and for ascorbic acid determination was immediately frozen in liquid nitrogen and kept at -30°C prior to analysis. Samples were kept for at least 48 h under white fluorescent lamps ($35 \mu\text{mol m}^{-2} \text{s}^{-1}$) in running seawater pumped directly from the fjord before laboratory experiments started.

For laboratory experiments, approximately 10 g fresh weight (FW) of algae was incubated in 5-l plastic tanks in running seawater at a temperature of 2°C and exposed to $38 \mu\text{mol m}^{-2} \text{s}^{-1}$ PAR provided by one Osram daylight fluorescence tube, 8 W m^{-2} UVA (320–400 nm) and 0.36 W m^{-2} UVB (280–320 nm) provided by two Q-Panel UVA-340 fluorescence tubes (Q-Panel, Cleveland, Ohio). Total PAR + UVA + UVB (295–700 nm) radiation treatment was obtained by covering the tanks with Ultraphan cut-off filter foil (cut-off wavelength $<295 \text{ nm}$; Ultraphan, Digepra, Munich, Germany). For the PAR treatment, the tanks were covered with polyester cut-off filter foil (cut-off wavelength $<395 \text{ nm}$; Folex, Dreieich, Germany). Radiation measurements were carried out with a Li-Cor LI-190-SB cosine corrected sensor connected to a Li-Cor LI-1000 datalogger (Lambda Instruments, Lincoln, Neb.) for PAR (400–700 nm), and with an RM-21 broad-band UV radiometer (Dr. Gröbel, Ettlingen, Germany).

In a first set of experiments, the red algae, *Coccolyx truncatus*, *Phycodrys rubens* and the green alga, *Monostroma arcticum*, were exposed for 3 days under continuous PAR and PAR + UVA + UVB radiation. Samples were taken at the start of the experiment and after 24 h and 84 h of exposure. In parallel, at the same time, subsamples were taken and maintained in darkness for 24 h under otherwise identical culture conditions, in order to characterize the possible recovery processes in darkness after exposure to UV radiation.

In a second set of experiments, thalli of the red algae, *Palmaria palmata*, *Devaleraea ramentacea* and the green alga, *Acrosiphonia penicilliformis*, were exposed for 8 days under continuous PAR and

Table 1. Investigated macroalgal species from the Kongsfjord on the Arctic island of Spitsbergen (Svalbard, Norway) and their occurrence in the eulittoral, upper sublittoral (0–3 m) and lower sublittoral (3–20 m) zone according to Svendsen (1959) and Klekowski and Weslawski (1990)

Species	Habitat
Chlorophyta	
<i>Acrosiphonia penicilliformis</i> (Foslie) Kjellman	Eulittoral-upper sublittoral
<i>Monostroma arcticum</i> Wittrock	Upper-lower sublittoral
<i>Chaetomorpha linum</i> (Müller) Kützing	Upper-lower sublittoral
<i>Chaetomorpha melagonium</i> (F. Weber et Mohr) Kützing	Upper-lower sublittoral
<i>Prasiola crispa</i> (Lightfoot) Meneghini	Eulittoral
Rhodophyta	
<i>Coccotylus truncatus</i> (Pallas) M.J.Wynne & J.N.Heine	Lower sublittoral
<i>Devaleraea ramentacea</i> (L.) Guiry	Eulittoral-upper sublittoral
<i>Palmaria palmata</i> (L.) Grev.	Upper-lower sublittoral
<i>Phycodrys rubens</i> (L.) Batters	Lower sublittoral
<i>Odonthalia dentata</i> (L.) Lyngb	Lower sublittoral
<i>Polysiphonia arctica</i> J. Agardh	Lower sublittoral
<i>Ptilota gunneri</i> P.C.Silva, Maggs & L.M.Irvine	Lower sublittoral
Phaeophyta	
<i>Alaria esculenta</i> (L.) Greville	Lower sublittoral
<i>Chorda tomentosa</i> Lyngbye	Upper-lower sublittoral
<i>Chordaria flagelliformis</i> (O. F. Müller) C. Agardh	Upper sublittoral
<i>Desmarestia aculeata</i> (L.) Lamouroux	Lower sublittoral
<i>Elachista fucicola</i> (Vellay) Areschoug	Eulittoral-upper sublittoral
<i>Fucus distichus</i> L.	Eulittoral-upper sublittoral
<i>Laminaria saccharina</i> (L.) Lamouroux	Upper-lower sublittoral
<i>Laminaria solidungula</i> J. Agardh	Lower sublittoral
<i>Laminaria digitata</i> (Huds.) Lamouroux	Upper-lower sublittoral
<i>Saccorhiza dermatodea</i> (de la Pylaie) J. Agardh	Upper-lower sublittoral

PAR + UVA + UVB. Parallel to the laboratory experiments, thalli of *A. penicilliformis* were covered in situ by means of an 80×80 cm² UV transparent Plexiglass plate, wrapped with the 395 nm cut-off filter foil to avoid UVA plus UVB. These samples were compared with subsamples exposed to full solar radiation at the same part of the rocky shore.

Enzyme activities

Samples (0.2–0.3 g FW) of the studied species were ground in liquid nitrogen and extracted with 1–1.5 ml 50 mM potassium phosphate buffer (pH 7.0) containing Complete protease inhibitor cocktail (Boehringer, Mannheim, 2 tablets in 100 ml buffer). Extracts were centrifuged for 15 min at 15,000 rpm at 4°C. Catalase was analysed according to Aebi (1984); 10–40 µl extract was added to 810–840 µl potassium phosphate buffer (50 mM, pH 7). The reaction was started by the addition of 150 µl of H₂O₂ solution in phosphate buffer (15 mM final concentration in the cuvette) and followed by monitoring the decrease in absorbance at 240 nm at 20°C for 1–2 min. Catalase activity was calculated by subtracting the non-enzymatic reaction and using an extinction coefficient for H₂O₂ of 0.0398 mm⁻¹ cm⁻¹. Glutathione reductase (GR) was assayed according to Goldberg and Spooner (1983); 10–40 µl extract was added to 960–990 µl of a buffer containing 80 mM Tris buffer (pH 8), 1 mM EDTA, 0.1 mM NADPH, and 0.5 mM GSSG, and oxidation of NADPH was followed at 340 nm at 20°C. GR activity was calculated by subtracting the non-enzymatic reaction and using an extinction coefficient for NADPH of 6.22 mm⁻¹ cm⁻¹. Samples for ascorbate peroxidase (APX) activities were extracted with the same protocol as for the other enzymes with the modification that 0.5 mM of ascorbate was added to the extraction buffer for the stability of the APX (Chen and Asada 1989). Enzyme activities were assayed according to the same authors and the decrease of absorbance at 290 nm was followed for 1 min after adding 10–40 µl extract to 960–990 µl 50 mM phosphate buffer (pH 7) containing 0.1 mM of H₂O₂, and 0.5 mM ascorbate. All assays were performed at 20°C. APX activity was calculated by subtracting the non-enzymatic reaction and using an extinction coefficient for ascorbate of 2.8 mm⁻¹ cm⁻¹. Results for catalase, GR and APX are expressed as units (U) of

enzyme activity per milligram of total soluble protein [1 U = 1 µmol substratum (H₂O₂, NADPH and ascorbate, respectively) converted min⁻¹]. SOD was measured using the xanthine oxidase-cytochrome c reduction method (McCord and Fridovich 1969). In this coupled reaction, SOD inhibits the reduction of cytochrome c by superoxide anions generated from xanthine. The assay mixture (860–1,000 µl) contained 50 mM phosphate buffer (pH 7.8), 0.1 mM EDTA, 10 µM cytochrome c and 50 µM xanthine. Xanthine oxidase (Merck) was added to give an increase of absorbance at 550 nm of 0.025 ± 0.003 min⁻¹ at 20°C. Samples (10–50 µl) were added to the reaction mixture and the rate of reduction of cytochrome c was followed spectrophotometrically at 550 nm, and 1 unit of SOD was defined as the amount of enzyme required to inhibit the rate of cytochrome c reduction by 50%.

Ascorbic acid was measured according to Foyer et al. (1983). Thalli of 0.2–0.4 g fresh weight were ground in liquid nitrogen and extracted with 1–1.5 ml 100 mM potassium phosphate buffer (pH 5.6) containing 5 mM dithioerythritol (DTE). Extracts were centrifuged for 15 min at 15,000 rpm at 4°C. The ascorbate content was determined by the disappearance of absorbance at 265 nm after addition of 10 U ml⁻¹ ascorbate oxidase and 50 µl sample to 925 µl sodium phosphate buffer (100 mM, pH 5.6). Amounts were quantified using a standard curve with 1.25–12.5 µM of pure ascorbate in the reaction mixture.

Protein assay

Total soluble proteins of the crude extract for antioxidant enzyme activities were determined using a commercial Protein Assay (BioRad), based on the Bradford method (Bradford 1976). Protein content was determined spectrophotometrically at 595 nm and concentrations were calculated compared with a standard of bovine serum albumin (SIGMA).

Statistics

Mean values and their standard deviations were calculated from the different replicates per treatment. Statistical significances of means were tested with a model 1 one-way ANOVA, followed by a

multi-range test by Fisher's protected least significance difference (LSD) (Sokal and Rohlf 1995), and a $P < 0.05$ was considered to be statistically significant.

Results

Antioxidant enzyme distribution in Arctic macroalgae

To investigate the protection mechanisms against oxidative stress, a total of 22 species of green, red and brown Arctic macroalgae were analysed for the presence of the activity of superoxide dismutase, glutathione reductase, ascorbate peroxidase and catalase (Table 2). Clear differences were found between the three macroalgal groups, with green algae showing in general higher antioxidant enzyme activities than red and brown algae. Independent of the reference parameters tested to express enzyme activity (fresh weight or protein content), these taxon-specific differences remained similar. Maximum SOD activities were found in *M. arcticum* and *A. penicilliformis*, exhibiting values of 1,004 and 674 U mg TSP⁻¹, respectively.

SOD activities in all species tested were significantly higher ($P < 0.05$) compared to GR, APX and CAT. Maximum GR activities were measured again in the green algal group, with the highest value of 2.3 U mg TSP⁻¹ found in *A. penicilliformis*. Red and brown algae showed species-specific GR activities ranging from 0.07 to 0.32 U mg TSP⁻¹. Similar results were obtained in APX and CAT activities, and in both cases *M. arcticum*

and *Chaetomorpha* species exhibited highest values. *D. ramentacea* showed exceptionally high values of APX and CAT activities, being 6 times higher in APX and almost 2 times higher in CAT compared to the other investigated red algae (Table 2). The internal concentrations of ascorbate in the different species varied from traces in brown algae up to values of 1.63 mg ascorbate gFW⁻¹ in *Chaetomorpha melagonium*.

Ultraviolet radiation effects on antioxidant enzymes

Ultraviolet radiation leads to a significant enhancement ($P < 0.05$) of the enzymatic activities of GR in laboratory experiments (Fig. 1a, b, c: note the different species-specific scales). Exposure to 8 W m⁻² UVA and 0.36 W m⁻² UVB promoted an increase of 22% in GR activity in the green alga *M. arcticum* after 24 h of continuous irradiation compared to the control under PAR radiation (Fig. 1a). GR activity in subsamples kept for 24 h in darkness increased further. After 84 h exposure, GR activity rose under both radiation conditions, especially after exposure to PAR and UV radiation. After 84 h exposure, followed by 24 h of darkness, GR activity was slightly higher ($P < 0.05$) in specimens previously exposed to UV radiation.

In the red algae *Coccotylus truncatus* and *Phycodrys rubens*, 24 h of continuous UV radiation did not result in significant differences of GR activity with respect to the control (Fig. 1b, c). However, after 84 h of contin-

Table 2. Enzymatic activities of superoxide dismutase (SOD), glutathione reductase (GR), catalase (CAT), ascorbate peroxidase (APX) and the content of ascorbic acid in different green, red and brown algae from the Kongsfjord (Spitsbergen). Results are expressed as units (U) of enzyme activity per milligram of total soluble proteins (TSP) where 1 U = 1 μmol substrate converted min⁻¹. Standard deviations were less than 20% (– not measured)

Species	SOD (Umg TSP ⁻¹)	GR (Umg TSP ⁻¹)	APX (Umg TSP ⁻¹)	CAT (Umg TSP ⁻¹)	Ascorbate (mg gFW ⁻¹)
Chlorophyta					
<i>Acrosiphonia penicilliformis</i>	674	2.30	0.2	1.0	0.19
<i>Monostroma arcticum</i>	1004	1.58	0.97	27.11	1.63
<i>Chaetomorpha linum</i>	395	0.10	0.778	0.77	0.65
<i>Chaetomorpha melagonium</i>	200	1.54	0.5	30.00	1.57
<i>Prasiola crispa</i>	153	0.10	0.12	3.86	–
Rhodophyta					
<i>Coccotylus truncatus</i>	165	0.08	0.05	9.91	0.27
<i>Devaleraea ramentacea</i>	245	0.32	0.60	15.5	0.2
<i>Palmaria palmata</i>	185	0.22	0.45	4.5	0.43
<i>Phycodrys rubens</i>	94	0.08	0.04	–	–
<i>Odonthalia dentata</i>	109	0.09	0.10	6.24	0.43
<i>Polysiphonia arctica</i>	87	0.06	0.02	7.58	0.18
<i>Ptilota gunneri</i>	122	0.11	0.01	–	0.2
Phaeophyta					
<i>Alaria esculenta</i>	102	0.07	0.05	–	0.06
<i>Chorda tomentosa</i>	128	0.11	0.07	–	0.75
<i>Chordaria flagelliformis</i>	112	0.10	0.03	–	–
<i>Desmarestia viridis</i>	143	0.10	0.08	–	0.31
<i>Elachista fuciola</i>	191	0.10	0.05	–	–
<i>Fucus distichus</i>	151	0.09	0.07	3.60	0.29
<i>Laminaria saccharina</i>	181	0.18	0.04	7.97	0.17
<i>Laminaria solidungula</i>	142	0.08	0.01	–	traces
<i>Laminaria digitata</i>	68	0.10	0.05	–	traces
<i>Saccorhiza dermatodea</i>	88	0.09	0.06	–	0.58


uous UV radiation, a significant increment ($P < 0.05$) in GR activity from 0.18 to 0.31 U mgTSP⁻¹ for *C. truncatus* and from 0.25 to 0.32 U mgTSP⁻¹ for *P. rubens* was measured. Moreover, the GR activity increased in subsamples of both species cultured in darkness following the UV exposure.

In contrast to GR activity, SOD in the three algal species studied seemed not to be positively affected ($P > 0.05$) at the end of the exposure to the radiation treatments (Fig. 1d, e, f). In *M. arcticum* and *P. rubens*, UV promoted a significant decrease of SOD activity after 24 h of exposure and after 24 h in darkness ($P < 0.05$), but after 84 h exposure and the following 24 h of darkness, no significant difference ($P > 0.05$) between the two radiation conditions could

be detected. The red alga *C. truncatus* showed an unchanged SOD activity under all treatment conditions.

The red algae, *Palmaria palmata* and *Devaleraea ramentacea*, were kept for 8 days under continuous UV radiation under the same conditions as described above. For both species, no significant UV effect was observed ($P > 0.05$) in CAT, APX and SOD activities (Fig. 2a–f). Although CAT activities in both plants increased within 8 days, significant differences between the radiation treatments could not be detected (Fig. 2a, d). For APX, an inconsistent activity pattern was observed. After UV exposure, variation was high in both species, with no specific pattern or effect (Fig. 2b, e). In *P. palmata*, SOD activities markedly increased

Fig. 1a–f. Influence of exposure under artificial PAR (grey bars) and PAR + UVA + UVB (black bars) radiation on enzymic activities of glutathione reductase (GR) and superoxide dismutase (SOD) in the red algae *Coccotylus truncatus* and *Phycodrys rubens* and the green alga *Monostroma arcticum* from Spitsbergen. Plants were exposed for 84 h under continuous irradiation. Subsamples were cultured for 24 h in darkness after 24 and 84 h exposure. Data are given as mean values \pm SD ($n = 3$) and expressed as Units mg⁻¹ total soluble proteins (TSP). Mean values with different asterisks are significantly different (at $P = 0.05$) to the control



under PAR within the 8 days of exposure, whereas activities under UV radiation were lower (Fig. 2c). In *D. ramentacea*, no significant changes in SOD activities could be recorded (Fig. 2d).


In a third set of radiation experiments, the eulittoral green alga *Acrosiphonia penicilliformis* was kept for 8 days under continuous UV irradiation under the same laboratory conditions as before (Fig. 3a, b). In this case, no significant differences were observed in GR and SOD activities when the algae were maintained under PAR and PAR+UVA+UVB radiation ($P>0.05$). In contrast, in the field experiment, algal thalli were exposed in situ to the full solar spectrum and compared with thalli where UVA+UVB was filtered out of the natural solar radiation spectrum

(Fig. 3c, d). Samples exposed for 8 days to UV filtered radiation showed a significant decrease ($P<0.05$) in GR and SOD activity with respect to the fully exposed subsamples.

Discussion

The present study provides a survey of the qualitative and quantitative content and activities of different reactive-oxygen-scavenging enzymes in 22 macroalgal species from the Arctic. We detected a great variation with respect to the antioxidant enzyme activities and found strong species-specific differences. High activities of antioxidant enzymes found in green algae were

Fig. 2a–f. Influence of continuous exposure under artificial PAR (grey bars) and PAR+UVA+UVB (black bars) radiation on enzymatic activities of glutathione reductase (GR), ascorbate peroxidase (APX) and superoxide dismutase (SOD) in the Arctic red algae *Palmaria palmata* and *Devaleraea ramentacea*. Data are given as mean values \pm SD ($n=5$) and expressed as Units mg^{-1} total soluble proteins (TSP). Mean values with different asterisks are significantly different (at $P=0.05$) to the control






Fig. 3a–d. Changes of the activities of glutathione reductase (GR) and superoxide dismutase (SOD) in the Arctic green alga *Acrosiphonia penicilliformis*. **a, b** Laboratory experiment: continuous exposure under artificial PAR (grey bars) and PAR+UVA+UVB (black bars) radiation; **c, d** Field experiment: exposure under total (black bars) and UVA+UVB cut-off solar radiation by means of a selective UV cut-off filter (cut off at < 395 nm.). Data are given as mean values \pm SD ($n=3$) and expressed as Units mg^{-1} total soluble proteins (TSP). Mean values with different asterisks are significantly different (at $P=0.05$) to the control

comparable to those of higher plants and microalgae. For example, *M. arcticum*, which showed the highest activity in SOD within the investigated macroalgae, is in the same activity range as reported for pea leaves (Moran et al. 1994). CAT activity of *M. arcticum* is half as high as that of cotton fibres (Rajguru et al. 1999), whereas activities of SOD and GR were much higher. In comparison with other algae, APX activities of *M. arcticum* are comparable to those of the symbiotic zooxanthellae of the sea anemone, *Aiptasia pallida* (Lesser and Shick 1989) while CAT and SOD activities are much higher. Data with respect to other macroalgae are scarce and only the data on the Arctic brown alga *Fucus distichus* (Table 2) can be directly compared to those of similar species from temperate water (Collen and Davison 1999a). CAT activity of *F. distichus* was similar to that reported by other authors for the same species while SOD activity was much higher in the polar species and lower activities were measured for GR and APX. The ascorbic acid concentrations found in green

algae are very high, similar to those reported, in general, for lemons and oranges. *Chaetomorpha linum*, for example, contains $0.65 \text{ mg ascorbic acid g}^{-1} \text{ FW}$, a concentration similar to that of lemon, with approximately $0.69 \text{ mg g}^{-1} \text{ FW}$. *Fucus distichus* from Spitsbergen contained ascorbic acid values in the same range as in this and other *Fucus* species from temperate regions (Collen and Davison 1999a).

A closer analysis points to the relation between antioxidant activities and depth distribution. In particular, differences in SOD activity between algal groups are related to the depth distribution on the shore. Most green algae, which showed higher antioxidant activities, typically inhabit the upper part of the shore at the Kongsfjord. Similarly, the red algae, *Devaleraea ramentacea* and *Palmaria palmata*, occurring frequently in the upper sublittoral, exhibited higher SOD activities compared to red-algal species living in deeper waters, such as *Phycodrys rubens*. Additionally, green algae contain relatively high amounts of ascorbic acid. Therefore, species more exposed to drastic and rapid changes in environmental radiation conditions have developed an efficient biochemical defence system to withstand the stress.

Although directly exposed to solar radiation, the supralittoral species *Prasiola crispa* growing underneath bird colonies exhibits very low antioxidant enzyme activities in comparison to the other green algae. In this plant from such an unusual habitat, it seems that

another photoprotective strategy is developed, such as the biosynthesis of UV-absorbing compounds that are known to prevent radiative damage (Dunlap and Shick 1998). In *Prasiola crispa* ssp. *antarctica* from Antarctica, high amounts of two new, so far chemically uncharacterized, mycosporine-like amino acids (MAAs) have been reported (Hoyer et al. 2001). MAAs represent a group of compounds with a potential role as UV sunscreens, exhibiting absorption maxima between 310 and 360 nm (Karentz et al. 1991). Their accumulation is positively correlated to the extent of UV exposure, as shown in laboratory and field studies (García-Pichel and Castenholz 1991; Karsten et al. 1998; Lesser 1996b; Shick et al. 1995). *Prasiola* is the only genus within the green algae containing MAAs. At present it is unknown whether the uncharacterized MAAs in *Prasiola crispa* also exhibit antioxidative properties as described for other MAAs (Dunlap and Yamamoto 1995).

Compared to other algal taxa, antioxidant enzyme activities in brown algae are low. However, there is a strong adaptation and/or acclimation potential of photosynthesis (Bischof et al. 1998a, b, 2000) and growth (Aguilera et al. 1999a) to UV radiation in this macroalgal group. In this context we refer to the typically high content of phenolic compounds in brown algae (Ragan and Glombitza 1986; Van Alstyne and Paul 1990), since these substances can act as antioxidants by transferring hydrogen atoms to lipid peroxyl radicals (Foti et al. 1994). However, their role as antioxidants may be questionable because they are accumulated in special compartments, the physodes (Schoenwaelder 2001), rather than uniformly distributed in the protoplasm. Another explanation for the high adaptation and acclimation potential of brown algae to UV radiation may be the ability of phenolic compounds such as phlorotannins to act as UV sunscreen pigments, as suggested by Pavia et al. (1997).

The investigation of the effects of the ultraviolet waveband of the solar spectrum on polar marine ecosystems has become an important ecological issue as a result of a gradual depletion of the ozone layer in both hemispheres. Exposed organisms have developed different strategies for protection against this biologically harmful radiation. However, almost no studies on the mechanisms of production of reactive oxygen species by UV radiation, and the biochemical defence strategies against this reactive species, have been performed for macroalgae. In order to analyse the ecophysiological importance of the total UV region (UVA+UVB) of solar radiation, laboratory and field experiments have been performed. Our study has clearly shown that the activity of antioxidant enzymes is stimulated by UV radiation in several Arctic macroalgae. The combination of artificial UVA+UVB radiation increased the GR activity in *arcticum*, *Coccotylus truncatus* and *Phycodryx rubens* after 84 h under continuous exposure. GR stimulation under UV radiation indicates an active scavenging of H₂O₂ by means of the ascorbate-glutathione cycle in combination with the Mehler-peroxidase reaction, which is the major pathway for scavenging poten-

tially toxic intermediates of oxygen metabolism in photosynthesis, which at the same time enables down-regulation of electronflux (Polle 1996). Dehydroascorbate formed by oxidation of ascorbic acid for scavenging of H₂O₂ by means of ascorbate peroxidase is reduced again to ascorbate, taking electrons from reduced glutathione by means of dehydroascorbate reductase. The product of these reactions, glutathione disulphide (GSSG), is reduced by the activity of GR and consumption of NADPH. Plants have been shown to increase GR activity in response to stress (Edwards et al. 1994). Increments of GR activity in response to UV radiation have been described in *Arabidopsis* (Kubo et al. 1999; Rao et al. 1996). In Arctic macroalgae, it seemed to be a faster stimulation of GR activities after 24 h exposure, followed by incubation in darkness again for 24 h, indicating some kind of a dark-enhanced repair system after damage in light. Recovery from UV damage in low light or darkness has been extensively investigated in macroalgae, especially in studies on photoinhibition of photosynthesis (Hanelt 1996, 1998). Thus, stimulation of the biochemical system involved in the scavenging of reactive oxygen species generated in the photoinhibitory status, mediates this recovery in photosynthesis. The role of antioxidants in the partial recovery of photosynthetic performance has been studied in symbiotic cnidarians and their zooxanthellae (Lesser and Shick 1989). According to these authors, the fluxes of reduced oxygen intermediates cause damage to the photosynthetic apparatus. In contrast to GR activities, no UV effects on SOD activities were found in *Monostroma arcticum*, *Coccotylus truncatus* and *Phycodryx rubens* after this period of treatment. However, in *Palmaria palmata*, UV radiation seemed to directly affect the SOD activity and a decrease, just after the 1st day of culture, was observed in comparison to the PAR control. These results are comparable to those observed in the green microalga, *Chlorella vulgaris*, in which long-term effects of increasing UVB radiation resulted in a decrease in SOD activities (Malanga and Puntarulo 1997). The reason for this negative effect may be an inhibition of gene expression for this enzyme as observed by Strid (1993) in *Pisum sativum* or an unspecific effect on enzyme activity. In contrast, Lesser and Shick (1989) found a stimulation of SOD activities in the symbiotic zooxanthellae of *Aiptasia pallida* by UV radiation. In that species, an increase in the SOD activities was correlated with an increase in the CAT activity, while in the present work no significant differences were found in the red algae *Palmaria palmata* and *Devaleraea ramentacea* after 8 days treatment.

No effects of artificial UV radiation on GR and SOD have been observed in the green alga, *Acrosiphonia penicilliformis*. In contrast, a significant reduction in GR and SOD activities was observed when natural UV radiation was cut off by selective filter foils in the field. This means that survival of this species in the intertidal zone is mediated by an enhanced oxygen-reactive scavenging system, in combination with

morphological strategies, as reported by Aguilera et al. (1999a). While the apical region of this plant is mainly exposed to strong solar radiation, the basal cells are well protected due to self-shading. In the field, yellow-coloured tips were often observed, indicating lack of chlorophyll as a consequence of photobleaching of the apical parts, along with dark-green pigmented, healthy and unstressed basal parts.

In conclusion, different biochemical capabilities of the enzymatic defence systems against reactive-oxygen species were observed for several Arctic macroalgae of different taxa and habitats. The antioxidant enzyme activity is enhanced in species that grow in the upper part of the rocky shore, where they are exposed to drastic changes in environmental conditions, especially those related to rapid and drastic changes in the UV region of the solar spectrum.

Acknowledgements The authors would like to thank the diving team (Heike Lippert, Eva Philipp, Stefan Kremb and Tanja Michler) for collecting the plant material, as well as the Ny-Ålesund International Research and Monitoring Facility for their support. This project was financially supported by the European Union (Project ENV4-CT96-0188 (DG 12) – UV/marine macrophytes) and by the German Minister for Education and Research (BMBF-Project: “MONA”, 03FO229A). J. Aguilera is grateful to the Ministerio de Educación y Cultura of Spain and the Alexander von Humboldt Foundation for supporting his research.

References

- Aebi H (1984) Catalase in vitro. *Method Enzymol* 105:121–130
- Aguilera J, Karsten U, Lippert H, Vögele B, Philipp E, Hanelt D, Wiencke C (1999a) Effects of solar radiation on growth, photosynthesis and respiration of marine macroalgae from the Arctic. *Mar Ecol Prog Ser* 191:109–119
- Aguilera J, Jiménez C, Figueroa FL, Lebert M, Häder DP (1999b) Effect of ultraviolet radiation on thallus absorption and photosynthetic pigments in the red alga *Porphyra umbilicalis*. *J Photochem Photobiol B Biol* 48:75–82
- Allen JF (1977) Superoxide and photosynthetic reduction of oxygen. In: Michelson AM et al. (eds) *Superoxide and superoxide dismutases*. Academic Press, New York, pp 417–436
- Aro EM, Virgin I, Andersson B (1993) Photoinhibition of photosystem II. Inactivation, protein damage and turnover. *Biochim Biophys Acta* 1143:113–134
- Asada K, Takahashi M (1987) Production and scavenging of active oxygen in photosynthesis. In: Kyle DJ, Osmond CB, Arntzen CJ (eds) *Photoinhibition*. Elsevier Science, Amsterdam, pp 89–109
- Bischof K, Hanelt D, Wiencke C (1998a) UV-radiation can affect depth-zonation of Antarctic macroalgae. *Mar Biol* 131:597–605
- Bischof K, Hanelt D, Tüg H, Karsten U, Brouwer PEM, Wiencke C (1998b) Acclimation of brown algal photosynthesis to ultraviolet radiation in Arctic coastal waters (Spitsbergen, Norway). *Polar Biol* 20:388–395
- Bischof K, Hanelt D, Wiencke C (2000) Effects of ultraviolet radiation on photosynthesis and related enzyme reactions of marine macroalgae. *Planta* 211:555–562
- Bradford M (1976) A rapid and sensitive method for the quantification of micrograms quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–254
- Buma AGJ, Hanenab EJ, Roza L van, Veldhuis MJW, Gieskes WWC (1995) Monitoring ultraviolet-B induced DNA damage in individual diatom cells by immunofluorescence thymine dimer detection. *J Phycol* 31:314–321
- Chen GX, Asada K (1989) Ascorbate peroxidase in tea leaves: occurrence of two isozymes and the difference in their enzymatic and molecular properties. *Plant Cell Physiol* 30:987–998
- Clendennen SK, Zimmerman RC, Powers DA, Alberte RS (1996) Photosynthetic responses of the giant kelp *Macrocystis pyrifera* (Phaeophyceae) to ultraviolet radiation. *J Phycol* 32:614–620
- Collen J, Davison IR (1999a) Reactive oxygen metabolism in intertidal *Fucus* spp. (Phaeophyceae). *J Phycol* 35:62–69
- Collen J, Davison IR (1999b) Production and damage of reactive oxygen in intertidal *Fucus* (Phaeophyceae). *J Phycol* 35:54–61
- Dunlap WC, Shick JM (1998) Ultraviolet radiation-absorbing mycosporine-like amino acids in coral reef organisms: a biochemical and environmental perspective. *J Phycol* 34:418–430
- Dunlap WC, Yamamoto Y (1995) Small-molecule antioxidants in marine organisms: antioxidant activity of mycosporine-glycine. *Comp Biochem Physiol* 112:105–114
- Edwards EA, Enard C, Creissen GP, Mullineaux PM (1994) Synthesis and properties of glutathione reductase in stress peas. *Planta* 192:137–143
- Elstner EF (1982) Oxygen activation and oxygen toxicity. *Annu Rev Plant Physiol* 33:73–96
- Figueroa FL, Salles S, Aguilera J, Jimenez C, Mercado J, Viñeola B, Flores A, Altamirano M (1997) Effects of solar radiation on photoinhibition and pigmentation in the red alga *Porphyra leucosticta*. *Mar Ecol Prog Ser* 151:81–90
- Foti M, Piatelli M, Amico V, Ruberto G (1994) Antioxidant activity of phenolic meroditerpenoids from marine algae. *Photochem Photobiol* 26:159–164
- Foyer CH, Rowell J, Walker D (1983) Measurement of ascorbate content of spinach leaf protoplasts and chloroplasts during illumination. *Planta* 157:381–392
- Fridovich I (1986) Biological effects of the superoxide radical. *Arch Biochem Biophys* 247:1–11
- Garcia-Pichel F, Castenholz RW (1991) Characterization and biological implications of scytonemin, a cyanobacterial sheath pigment. *J Phycol* 27:395–409
- Goldberg DM, Spooner RJ (1983) Glutathione reductase. In: Bergmeyer HU (ed) *Enzymes*. 1. Oxidoreductases, transferases. VCH, Weinheim, pp 258–265
- Häder DP, Figueroa FL (1997) Photoecophysiology of macroalgae. *Photochem Photobiol* 66:1–14
- Halliwell B (1982) The toxic effects of oxygen on plant tissues. In: Oberley LW (ed) *Superoxide dismutase*, vol I. CRC Press, Boca Raton, Fla, pp 89–123
- Hanelt D (1996) Photoinhibition of photosynthesis in marine macroalgae. *Sci Mar* 60 [Suppl 1]:243–248
- Hanelt D (1998) Capability of dynamic photoinhibition in Arctic macroalgae is related to their depth distribution. *Mar Biol* 131:361–369
- Hoyer K, Karsten U, Sawall T, Wiencke C (2001) Photoprotective substances in Antarctic macroalgae and their variation with respect to depth distribution, different tissues and developmental stages. *Mar Ecol Prog Ser* 211:105–116
- Karentz D, Cleaver JE, Mitchell DL (1991) Cell survival characteristics and molecular responses of Antarctic phytoplankton to ultraviolet-B radiation. *J Phycol* 27:328–341
- Karsten U, Sawall T, Hanelt D, Bischof K, Figueroa FL, Flores-Moya A, Wiencke C (1998) An inventory of UV-absorbing mycosporine-like amino acids in macroalgae from polar to warm-temperate regions. *Bot Mar* 41:443–453
- Klekowski KR, Weslawski JM (1990) *Atlas of the marine fauna of Southern Spitsbergen*. Ossolineum, Wrocław
- Kubo A, Aono M, Nakajima N, Saji H, Tanaka K, Kondo N (1999) Differential responses in activity of antioxidant enzymes to different environmental stresses in *Arabidopsis thaliana*. *J Plant Res* 112:279–290
- Lesser MP (1996a) Elevated temperatures and ultraviolet radiation cause oxidative stress and inhibit photosynthesis in symbiotic dinoflagellates. *Limnol Oceanogr* 41:271–283
- Lesser MP (1996b) Acclimation of phytoplankton to UV-B radiation: oxidative stress and photoinhibition of photosyn-

- thesis are not prevented by UV-absorbing compounds in the dinoflagellate *Prorocentrum micans*. Mar Ecol Prog Ser 132:287–297
- Lesser MP, Shick JM (1989) Effects of irradiance and ultraviolet radiation on photoadaptation in the zooxanthellae of *Aiptasia pallida*: primary production, photoinhibition and enzymic defenses against oxygen toxicity. Mar Biol 102:243–255
- Malanga G, Puntarulo S (1997) Oxidative damage to chloroplasts from *Chorella vulgaris* exposed to ultraviolet-B radiation. Physiol Plant 101:455–462
- McCord JM, Fridovich I (1969) Superoxide dismutase: an enzymatic function for erythrocyte hemocuprein. J Biol Chem 244:6049–6055
- Moran JF, Becana M, Iturbe-Ormaetxe I, Frechilla S, Klucas RV, Aparicio-Tejo P (1994) Drought induces oxidative stress in pea plants. Planta 194:346–352
- Ohad I, Kyle DJ, Arntzen CJ (1984) Membrane protection damage and repair: removal and replacement of inactivated 32-kilodalton polypeptides in chloroplast membranes. J Cell Biol 99:481–485
- Ott H, Amanatides GT (1994) SESAME 1994–1995: a European contribution to the stratospheric ozone issue. DG XII. EC, Brussels
- Pavia H, Cervin G, Lindgren A, Åberg, P (1997) Effects of UV-B radiation and simulated herbivory on phlorotannins in the brown alga *Ascophyllum nodosum*. Mar Ecol Prog Ser 157:139–146
- Polle A (1996) Mehler reaction: friend or a foe in photosynthesis? Bot Acta 109:84–89
- Ragan MA, Glombitza KW (1986) Phlorotannins, brown alga polyphenols. In: Round FE, Chapman DJ (eds) Progress in phyiological research, vol 4. Biopress, Bristol, pp 129–241
- Rajguru SN, Banks SW, Gosset DR, Cran Lucas M, Fowler TE Jr, Millhollon EP (1999) Antioxidant response to salt stress during fiber development in cotton ovules. J Cotton Sci 3: 11–18
- Rao MV, Paliyath C, Ormrod DP (1996) Ultraviolet-B-induced and ozone-induced biochemical changes in antioxidant enzymes of *Arabidopsis thaliana*. Plant Physiol 110:125–136
- Schoenwaelder M (2001) The occurrence and cellular significance of physodes in brown algae. Phycologia (in press)
- Schulz A, Rex M, Harris NRP, Braathen GO, Reimer E, Alfier R, Kilbane-Dawe I, Eckermann S, Allaart M, Alpers M, Bojkov B, Cisneros J, Claude H, Cuevas E, Davies J, De Backer H, Dier H, Dorokhov V, Fast H, Godin S, Johnson B, Kois B, Kondo Y, Kosmidis E, Kyrö E, Litynska Z, Mikkelsen IS, Molyneux MJ, Murphy G, Nagai T, Nakane H, O'Connor F, Parrondo C, Schmidlin FJ, Skrivankova P, Varotsos C, Vialle C, Viatte P, Yushkov V, Zerefos C, Gathen P von der (2001) Arctic ozone loss in threshold conditions: Match observations in 1997/1998 and 1998/1999. J Geophys Res 106/D7:7495–7503
- Shick JM, Lesser MP, Dunlap WC, Stochaj WR, Chalker BE, Wu Won J (1995) Depth-dependent responses to solar ultraviolet radiation and oxidative stress in the zooxanthellate coral *Acropora microphthalmia*. Mar Biol 122:41–51
- Smith RC, Prézélin BB, Baler KS, Bidigare RR, Boucher NP, Coley T, Karentz D, MacIntyre S, Matlick HA, Menzies D, Ondrusek M, Wan Z, Waters KJ (1992) Ozone depletion: ultraviolet radiation and phytoplankton biology in Antarctic waters. Science 255:952–959
- Sokal RR, Rohlf FJ (1995) Biometry, 3rd edn. Freeman, New York
- Stolarski R, Bojkov R, Bishop L, Zerefos C, Staehelin J, Zawodny J (1992) Measured trends in ozone. Science 256:342–349
- Strid A (1993) Alteration in expression of defence genes in *Pisum sativum* after exposure to supplementary ultraviolet-B radiation. Plant Cell Physiol 34:949–953
- Strid A, Chow WS, Anderson JM (1990) Effect of supplementary ultraviolet-B irradiation on photosynthesis in *Pisum sativum*. Biochim Biophys Acta 1020:260–268
- Svendsen P (1959) The algal vegetation of Spitsbergen. Nor Polarinstit Skr 116:47
- Tevini M, Teramura AH (1989) UV-B effects on terrestrial plants. Photochem Photobiol 50:479–487
- Van Alstyne KL, Paul VJ (1990) The biogeography of polyphenolic compounds in marine macroalgae temperate brown algal defenses deter feeding by tropical herbivorous fishes. Oceanologia 84:158–163
- Wildi B, Lütz C (1996) Antioxidant composition on selected high alpine plant species from different altitudes. Plant Cell Environ 19:138–146