

Exploring the sublethal genotoxic effects of class II organophosphorus insecticide quinalphos on freshwater fish *Cyprinus carpio*

Devan HEMALATHA^{1,2}, Bojan NATARAJ¹, Basuvannan RANGASAMY¹,
Kannan MAHARAJAN¹, Mathan RAMESH^{1,*}

¹ Unit of Toxicology, Department of Zoology, School of Life Sciences, Bharathiar University, Coimbatore-641046, Tamil Nadu, India

² Department of Zoology, PSG College of Arts & Science, Coimbatore-641014, Tamil Nadu, India

Received Jul. 5, 2019; accepted in principle Sep. 6, 2019; accepted for publication Nov. 21, 2019

© Chinese Society for Oceanology and Limnology, Science Press and Springer-Verlag GmbH Germany, part of Springer Nature 2021

Abstract The genotoxic effects of quinalphos (QP) were evaluated in the erythrocytes of *Cyprinus carpio* using the comet, micronucleus, and nuclear abnormality assays. The 96 h LC₅₀ value of quinalphos, estimated by static renewal system, was 10.9 µL/L. On the basis of the LC₅₀ value, two sublethal concentrations (1.09 and 2.18 µL/L) of the compound were used together with cyclophosphamide (5 mg/L) as the positive control. The fish were exposed for 35 d, and sampling was done at the end of 7, 14, 21, 28, and 35 d. Our results reveal a significant dose-dependent increase in the frequencies of DNA strand breaks, micronucleus, and nuclear abnormalities. Similar responses were also observed in fish exposed to the positive genotoxin cyclophosphamide. The collective findings demonstrated that quinalphos exerts a genotoxic effect on *C. carpio*. The results further confirmed that the comet, micronucleus, and nuclear abnormality assays are useful tools in determining the potential genotoxicity of pesticides towards aquatic organisms.

Keyword: quinalphos (QP); micronucleus; nuclear abnormalities; cyclophosphamide; *Cyprinus carpio*

1 INTRODUCTION

Quinalphos (O,O-diethyl O-quinoxalin-2-yl phosphorothioate) is a synthetic organophosphate insecticide widely used in Indian agriculture to control a broad range of insect pests attacking crops such as cotton, groundnut, cereals, tea, coffee, sugarcane and vegetables (Hayes and Laws, 1991; Kokilavani et al., 2014). Common formulations of quinalphos (QP) are emulsion concentrates, dust and granules (Pant and Srivastava, 2003; http://agriculture.indiabizclub.com/info/organic_cultivation/crop_management). In India, farmers use a 4% solution of commercial-grade QP for spraying on the crops (Sharma and Mathur, 1972). QP is classified as number 2 pesticide (moderately hazardous) by the World Health Organization (WHO, 2002). In addition, the European Commission has included this insecticide in risk classification as hazardous to the environment. Like other organophosphates, the insecticidal activity of QP is

caused by the inhibition of the acetylcholinesterases (AChE) in the central nervous system (CNS) and peripheral nervous system (PNS). Inhibition of AChE results in the accumulation of acetylcholine (ACh) in the cholinergic synapses present at the neuromuscular junctions, leading to an excessive transmission of nerve impulses and causing mortality in the target pest (Ecobichon, 1991; Amitai et al., 1998; Li et al., 2012).

The half-life of QP in the soil is in the range of 10–16 d and in water it is >24 h (Racke, 1993). The actual half-life of QP mostly depends on environmental variables such as soil type, temperature and moisture. Upon degradation, QP produces various metabolites at different time intervals through hydrolysis, oxidation, de-alkylation and isomerization processes. These metabolites include quinalphosoxan, O-ethyl-

* Corresponding author: mathanramesh@yahoo.com

O-quinoxalin-2-yl phosphoric acid, quinoxaline-2-thiol and 2-hydroxy quinoxaline, which are further broken down to carbon-di-oxide. The metabolite 2-hydroxy quinoxaline and the parent compound persist in the environment for a longer period and are more toxic than the parent compound (Gupta et al., 2012). Extensive use of QP ultimately finds its way into aquatic environments, where it poses considerable toxicological risks to the resident organisms (Scott et al., 1990; Hemalatha et al., 2016). Generally, the toxic effect of the chemicals depends on the exposure period and also the size and age of the exposed animal. Hence, evaluating the toxicity of a chemical using a wide range of animal species is routinely employed in toxicological research (Oruc et al., 2004). Numerous studies have been conducted on the toxic effect of QP on plants and soil microorganisms (Vig et al., 2006), mammals (Rupa et al., 1991b; Vasilić et al., 1992; Subramanayaan et al., 2012) and aquatic organisms (Sastry et al., 1982; Durairaj and Selvarajan, 1992; Das and Mukherjee, 2000; Kavitha and Venkateswara Rao, 2008).

Quinalphos has been demonstrated to induce endocrine disruption, genotoxicity, reproductive toxicity and developmental toxicity in both male and female rats (Rupa et al., 1991a; Ray et al., 1992; Srivastava et al., 1992). In a recent investigation, Subramanayaan et al. (2012) have reported QP-induced oxidative stress and histological alterations in adult male albino rats. Dwivedi et al. (1998) have also documented the induction of hepatic antioxidant enzymes and hepatic P450 content in mice treated with QP. Moreover, QP can produce adverse effects on aquatic organisms (Sadiqul et al., 2016). In this line, among the aquatic fauna, fish is considered as a useful genetic model for the evaluation of pollution in aquatic ecosystems owing to its ability to efficiently metabolize and accumulate the pollutants (Park et al., 1989; Mitchell et al., 1992).

Fish models are widely used in toxicological research to assess the teratogenic and carcinogenic effects of chemicals on human beings (Harshbarger and Clark, 1990; Raisuddin and Jha, 2004; Matsumoto et al., 2006). The toxic effects of QP on fish have been assessed in several behavioral, respiratory (Chebbi and Davis, 2010a), biochemical (Das and Mukherjee, 2000), neurological (Chebbi and David, 2009), histological (Chamarthi et al., 2014), nucleic acid content and enzymatic studies (Venkata Rathnamma et al., 2013a, b). Nonetheless, information pertaining to the long-term genotoxic

effect of QP on fish is scarce.

Among the diverse genotoxicity tests, the comet assay (CA) or single cell gel electrophoresis (SCGE) has become the most important tool for monitoring the genotoxicity in aquatic environments, particularly in fish. The technique is rapid and sensitive in detecting and quantifying DNA damage, such as single and double-strand breaks and alkali-labile sites. Such damage can be induced directly by the contaminants or indirectly via repair processes (Tice et al., 2000). In toxicological research, SCGE/CA is often employed to ascertain the genotoxic effects of chemical compounds (Pandey et al., 2006; Ali et al., 2008; de Campos Ventura et al., 2008; Kumar et al., 2010). The micronucleus (MN) test in fish is also widely applied for estimating the genotoxicity in aquatic environments because of the ease of sampling and high sensitivity. Furthermore, water quality and the health condition of the aquatic species can be detected by these assays (Talapatra and Banerjee, 2007). Thus, the combined use of comet and micronucleus assays is preferred in aquatic toxicity research due to its sensitivity in detecting DNA damage and cytogenetic effects after acute and chronic exposure to various genotoxic agents. Along with the MN test, nuclear abnormalities (NA) have also been widely used in fish toxicology as indicators of genotoxicity (Ergene et al., 2007; Kumar, 2012). The test furnishes additional information for the scoring of micronuclei in genotoxicity surveys (Ayllon and Garcia-Vazquez, 2000; Muranli and Güner, 2011).

In such a context, we evaluated the mutagenic and genotoxic effects of QP using *C. carpio* as a laboratory animal model by means of comet assay, micronucleus and nuclear abnormalities tests. To our knowledge, this study presents the first in vivo evidence for the genotoxicity exerted by QP on *C. carpio* at sublethal concentrations. *Cyprinus carpio* is a widely cultivable and edible fish species in India. Moreover, it is a suitable animal model for ecotoxicological studies.

2 MATERIAL AND METHOD

2.1 Experimental animal and Chemicals

Fingerlings of the common carp *C. carpio* with average weight (5.0 ± 1 g) and length (6.0 ± 1 cm) were collected from the Aliyar Fish Farm, Tamil Nadu, India. Before commencing the experiments, the organisms were acclimated to laboratory conditions for three weeks in a large tank (1 000-L capacity)

containing aerated dechlorinated tap water of the following characteristics: temperature 27.0 ± 1 °C; pH 7.0 ± 1 ; dissolved oxygen 6.4 ± 0 mg/L; total hardness 18.4 ± 2 mg/L and total alkalinity 18.3 ± 2.0 mg/L. The photoperiod was 12 h:12 h. The fishes were fed once a day with commercial fish pellets and water renewal was performed once a day corresponding to 85% of the tank water to minimize contamination from metabolic wastes. The healthy organisms were randomly allocated to various treatment groups. Quinalphos (25% EC) was obtained from the Tudiyalur Cooperative Agricultural Services Limited, Tamil Nadu, India. Cyclophosphamide (CAS No. 50-18-0, purity: >99%) and other chemicals were purchased from Sigma-Aldrich, USA.

2.2 Experimental design

Acute toxicity test was conducted in a static renewal system to determine the 96-h LC_{50} value (APHA-AWWA-WEF, 2005). The stock solutions of QP were prepared in acetone and the range-finding test was carried out by exposing the fishes to five different concentrations (10.3, 10.5, 10.7, 10.9, 11.0 $\mu\text{L/L}$) of QP for 96 h. Control groups were also maintained. The experiments were repeated twice and the median lethal concentration (96 h LC_{50}) was determined by using the probit analysis described by Finney (1952). The 96 h LC_{50} value of QP in *C. carpio* was found to be 10.9 $\mu\text{L/L}$. During the acute toxicity experiment, the fishes were not fed and the dead ones were removed from the aquaria. Based on the 96 h LC_{50} value, $1/10^{\text{th}}$ (1.09 $\mu\text{L/L}$) and $1/5^{\text{th}}$ (2.18 $\mu\text{L/L}$) of the QP concentrations were taken as the sublethal concentration.

A group consisting of 25 fishes was exposed to each of the sublethal concentrations of QP (1.09 and 2.18 $\mu\text{L/L}$) separately for 35 d. Samples were withdrawn at intervals of 7, 14, 21, 28, and 35 d at the rate of five fishes per sampling. Additional sets were separately maintained for negative control (tap water), solvent control (acetone) and positive control (cyclophosphamide: 5 mg/L; concentration selection based on a previous investigation (Muranli and Güner, 2011)). In addition, two replicates for each concentration were also included for the two sublethal concentrations and the control groups.

2.3 Comet assay

For the SCGE/CA, the method of Singh et al. (1988) was adopted with slight changes. To an aliquot

of 30 μL diluted blood, 100 μL of low melting agarose (LMA) (0.65%) in PBS was added, mixed well, and pipetted on the slides (precoated with 0.65% normal agarose). Subsequently, the contents were incubated for 10 min for solidification. The slides were then dipped in cold lysing solution (2.5 mol/L NaCl, 100 mol/L EDTA-2Na, 10 mmol/L Tris, 10% DMSO, 1% sodium sarcosinate, 1% Triton-X, pH 10.0) and refrigerated for 4 h. The slides were transferred to an electrophoresis unit (15 cm \times 10 cm) containing denaturation buffer (300 mmol/L NaOH, 1 mmol/L EDTA, pH 13.0). Electrophoresis was done for 20 min at 0.6 V/cm and 300 mA. The buffer volume was adjusted to cover the slides. The slides were neutralized with Tris buffer (pH 7.5) for 2 min and stained with ethidium bromide (20 $\mu\text{g/mL}$) for microscopic examination. Approximately 100 cells per slide were scored randomly, examined under a florescent microscope, and the comet images were assayed using the Comet Assay Software Project (CASP). The DNA damage was quantified as % DNA tails, which indicate cellular damage (Imanikia et al., 2016).

2.4 Analysis of micronuclei (MN) and other nuclear abnormalities (NA)

The MN test was done according to Çavaş and Ergene-Gözükara (2003b). Peripheral blood samples were obtained from the caudal vein of the specimens and smeared onto pre-cleaned slides and dried overnight. The slides were then fixed with methanol for 10 min and stained with Giemsa (5%). All slides were coded and scored blind. Three slides were prepared for each fish, and a total of 1 000 erythrocytes per fish were examined under an Olympus optical microscope (1 000 \times magnifications).

2.5 Scoring criteria for micronuclei (MN) and other nuclear abnormalities (NA)

The MN test was carried out according to the protocol of Carrasco et al. (1990). The MN frequency was calculated as follows:

$$\%MN = \frac{\text{Number of cells containing micronuclei}}{\text{Total number of cells counted}} \times 100.$$

The NAs were classified according to Carrasco et al. (1990) into four categories: binucleated cells (BN), lobed nuclei (LB), blebbed nuclei (BL) and notched nuclei (NT). The NAs in the peripheral blood cells of *C. carpio* and their frequency was calculated in a manner similar to that of MN.




Fig.1 Percentage of tail DNA in erythrocytes of *C. carpio* exposed to QP

Bars represent means and vertical lines are the S.E. The values indicated by different letters are significantly different from one another ($P < 0.05$).




Fig.2 Comet assay in *C. carpio* erythrocytes

a. no damage; b. low; c. average; d. severe.

2.6 Statistical analysis

The data were subjected to one way analysis of variance (ANOVA) and Duncan's multiple range test (DMRT) at $P < 0.05$ level. All analyses were performed with SPSS 20.0 statistical software package, and the values were expressed as mean \pm S.E.

3 RESULT

The fishes exposed to the two different concentrations (1.09 and 2.18 $\mu\text{L/L}$) of QP exhibited higher DNA damage ($P < 0.05$) than the control groups (Fig.1). Prominent DNA damage was noticed in 2.18 $\mu\text{L/L}$ treatment on days 7, 14, and 21, and then a decline was noted at days 28 and 35. The DNA strand breaks in *C. carpio* after exposure to QP are depicted in Fig.2.




Fig.3 Frequencies of micronuclei (MN) in erythrocytes of *C. carpio* exposed to QP

Bars represent means and vertical lines are the S.E. The values indicated by different letters are significantly different from one another ($P < 0.05$).

The MN frequencies in the peripheral erythrocytes of *C. carpio* exposed to QP as well as those in the parallel positive, solvent and negative controls are summarized in Fig.3. The induction was significantly higher in the treatment groups than in the control groups at all sampling intervals. Furthermore, the induction of MN was found to be dose dependent as its highest frequency was recorded at the concentration of 2.18 $\mu\text{L/L}$ on the 35th day. In the positive control too, a concentration and duration-dependent induction of MN was noticed. In the negative and solvent control groups, the MN frequency remained almost stable and did not vary significantly from one sampling interval to the other. Cells containing normal nuclei and micronuclei are illustrated in Supplementary Fig.S1a & b.

The frequencies of NAs in the erythrocytes of fishes exposed to QP and their respective negative, solvent and positive control groups are summarized in Figs.4–7. Similar to MN, the frequencies of NAs were also significantly higher in the fishes treated with QP. The frequency of NAs in the erythrocytes of fishes exposed to QP was also concentration and duration dependent. The induction of NA was significantly ($P < 0.05$) higher in the fishes exposed to positive control when compared with the negative controls. No significant increases were observed in fishes belonging to the solvent and negative control groups. The BN, LB, BL, and NT cells are represented in Supplementary Fig.S1c–f.




Fig.4 Frequencies of binucleates in erythrocytes of *C. carpio* exposed to QP

Bars represent means and vertical lines are the S.E. The values indicated by different letters are significantly different from one another ($P < 0.05$).




Fig.6 Frequencies of blebbed nuclei in erythrocytes of *C. carpio* exposed to QP

Bars represent means and vertical lines are the S.E. The values indicated by different letters are significantly different from one another ($P < 0.05$).




Fig.5 Frequencies of lobed nuclei in erythrocytes of *C. carpio* exposed to QP

Bars represent means and vertical lines are the S.E. The values indicated by different letters are significantly different from one another ($P < 0.05$).




Fig.7 Frequencies of notched nuclei in erythrocytes of *C. carpio* exposed to QP

Bars represent means and vertical lines are the S.E. The values indicated by different letters are significantly different from one another ($P < 0.05$).

4 DISCUSSION

In general, organophosphorus insecticides have a faster rate of degradation and lower environmental persistence than the organochlorine compounds. However, the widespread use and application of QP in agriculture has resulted in the frequent detection of its residues in water, soil, sediments and aquatic biota. Fish is often used as the sentinel organism for investigating the carcinogenic and mutagenic potential of contaminants present in aquatic

environments. This choice is based on the various roles played by these organisms in the food web, such as bio-accumulation of environmental pollutants, biotransformation of xenobiotics, and response to mutagens at low concentrations (Çavaş and Ergene-Gözükara, 2005). In addition, they react to toxicants in a manner similar to that of higher vertebrates, which allows the measurement of substances that are potentially harmful to humans (de Lemos et al., 2007).

Acute toxicity data have been used to derive water

quality guidelines for regulatory purposes (Sundaram et al., 1994). For the present investigation, the 96 h LC₅₀ value (10.9 µL/L) of QP was taken based on our previous research (Hemalatha et al., 2016). The results reveal that QP is toxic to the fishes; In addition, the findings indicate that the toxicity of the pesticide is both time and dose dependent. The 96 h LC₅₀ reported in the present study for QP is slightly higher than the value (7.5 µL/L) reported by Chebbi and David (2010b). Muttappa et al. (2014) have also reported a lower 96 h LC₅₀ value (2.75×10^{-6}) for the same species. The differences in the LC₅₀ value observed within a single species for the same toxicant may depend on size, age and health of the organisms (Sprague, 1969). Furthermore, other environmental factors, such as temperature, pH, total hardness and dissolved oxygen content of the water, as well the formulation of the chemical and its exposure; greatly influence the results (Gupta et al., 1981).

The CA/SCGE is a widely used tool to detect genotoxicity in aquatic environments (Frenzilli et al., 2009). The assay has the advantage of determining the DNA damage in individual cells (Lee and Steinert, 2003; Buschini et al., 2004). The data obtained in the present study demonstrate that the frequency of % tail DNA damage for all concentrations of QP is significantly higher ($P < 0.05$) than that of the control. The percentage of DNA tail damage was found to be concentration and time dependent, which point to the genotoxicity of QP towards *C. carpio*. A similar observation has also been made in *Prochilodus lineatus* exposed to the pesticide Roundup (Cavalcante et al., 2008). In addition, time-dependent responses have been witnessed in *Channa punctatus* exposed to chlorpyrifos and malathion (Ali et al., 2008; Kumar et al., 2010). Lee and Steinert (2003) have reported that the interaction between the DNA molecules and the contaminants may occur in different ways, such as DNA damage by reactive oxygen species, inhibition of DNA repair, action of the compound directly on the DNA, interaction of the metabolites with DNA, and DNA damage through a multilateral mechanism. In the present study, the possible reason for the observed DNA strand breaks could not be elucidated due to the lack of any direct evidence. However, it has already been reported that QP can induce oxidative stress in fishes by producing reactive oxygen species (ROS) (Hemalatha et al., 2016). At supranormal levels, ROS has been shown to cause cellular and DNA lesions (Ayllon and Garcia-Vazquez, 2000; Cadet et al., 2003). Thus, it is

possible that the increased DNA damage in the erythrocytes of *C. carpio* after QP exposure might have resulted from the production of ROS during the metabolism of the toxicant. The metabolite might have interacted with the DNA and produced the lesions detected by the assay.

The MN frequency in fish erythrocytes is used to assess the toxicity of aquatic pollutants exhibiting aneugenic and clastogenic properties (Ferraro et al., 2004; Udrouiu, 2006). In this work, QP induced significantly higher number of MNs than the control groups, and the frequency escalated as the concentration and the time of exposure surged. A similar observation has been made in different fish species by using the assay to determine the genotoxic effect of xenobiotics (Nwani et al., 2011; Palanikumar et al., 2012). A dose-dependent increase in MN frequency has been reported in the peripheral blood of *Barbonymus gonionotus* in response to QP containing insecticides (convoy) (Sadiqul et al., 2016). The trend has also been observed in *Channa punctatus* in response to carbosulfan (Nwani et al., 2010), chlorpyrifos (Ali et al., 2008), PCP and 2,4-D (Farah et al., 2003). Time responses have also been documented in *Oreochromis niloticus* treated with petroleum and chromium processing plant effluents (Çavaş and Ergene-Gözükara, 2005). Our result provides the first evidence for the sublethal genotoxic effects of QP on *C. carpio* under laboratory conditions. Hence, the induction of micro-nucleated erythrocytes following exposure to QP asserts the potential clastogenic effects of the compound in fishes.

Studies have already reported the clastogenic potential of QP in rats (Rupa et al., 1991a; Ray et al., 1992; Srivastava et al., 1992). Furthermore, the findings from previous studies suggest that organophosphorus insecticides, such as dimethoate (Ali et al., 2014) profenofos (Pandey et al., 2011) and diclorvos (Shukla et al., 2010), induce genotoxic effects in fish erythrocytes mainly via oxidative damage. Similarly, QP may also cause oxidative stress. Certain chemicals are capable of causing oxidative stress through the generation of ROS, which interact with macromolecules such as DNA and result in molecular and clastogenic damages (Raisuddin and Jha, 2004; Jha, 2008). Thus, the findings of this research suggest that the observed genotoxic damage may be due to the increased oxidative stress resulting from exposure to QP.

In addition to MN, other NAs have also been reported in fish erythrocytes (Monteiro et al., 2011).

The formation of NAs was first described by Carrasco et al. (1990), but the mechanism was not fully understood. Pacheco and Santos (1997) uncovered that the changes in nuclear morphology originated from a genotoxic event as a result of exposure to xenobiotic contaminants. To investigate the genotoxic effects of chemicals, NAs in fishes such as LB, BL, NT, and BN are widely used (Çavaş and Ergene-Gözükara, 2003a; Kilemade et al., 2004; da Silva Souza and Fontanetti, 2006; Ergene et al., 2007). In the present study, the formation of the abnormalities indicates the damaging effect of QP. These defects may affect gene amplification and also cause deformed nuclei (Bolognesi et al., 2006; Cavalcante et al., 2008). Nuclear alterations such as BL and LB could also be related to the failure of tubulin polymerization (Vardavas et al., 2016). Additionally, de Campos Ventura et al. (2008) have reported that BN and NT arise from difficulties in the formation of mitotic fuse caused by the aneugenic action of the chemicals. Our findings on erythrocyte NAs are in agreement with the previous studies that have recorded increased NA frequencies in fishes following exposure to QP containing insecticides (convoy) (Sadiqul et al., 2016), deltamethrin (Ansari et al., 2009), formalin (Mert et al., 2015), carbosulfan, glyphosate and atrazine (Nwani et al., 2014). In the present study, the significant increase in NAs noted throughout the study period implies that they are better biomarkers for QP exposure than the MN alone. Therefore, we conclude that the formation of NAs in the exposed fishes clearly indicates the genotoxicity of QP. The NA and MN assays could together serve as suitable biomarkers for assessing the genotoxicity of chemicals in the aquatic environment.

5 CONCLUSION

The results demonstrate that QP is genotoxic to fishes since it induces DNA strand breaks and the formation of MN and NA in peripheral erythrocytes. This study also confirms that the comet, micronucleus and nuclear abnormality assays are useful tools in determining the potential genotoxicity of pesticides on non-target organisms. Furthermore, studies on QP-induced genotoxic effects in fishes are rare, and the present findings contribute to the limited literature on the exposure of fishes to sublethal levels of the pesticide.

6 DATA AVAILABILITY STATEMENT

The data used in the present investigation are available from the corresponding author upon request.

7 ACKNOWLEDGMENT

The author (D. HEMALATHA) wish to thank UGC, New Delhi for providing grant of Basic Science Research Fellowship (BSR) (No. F.7-24/2007).

References

- Ali D, Kumar P G, Kumar S, Ahmed M. 2014. Evaluation of genotoxic and oxidative stress response to dimethoate in freshwater fish *Channa punctatus* (Bloch). *Chem. Spec. Bioavailab.*, **26**(2): 111-118.
- Ali D, Nagpure N S, Kumar S, Kumar R, Kushwaha B. 2008. Genotoxicity assessment of acute exposure of chlorpyrifos to freshwater fish *Channa punctatus* (Bloch) using micronucleus assay and alkaline single-cell gel electrophoresis. *Chemosphere*, **71**(10): 1 823-1 831.
- Amitai G, Moorad D, Adani R, Doctor B P. 1998. Inhibition of acetylcholinesterase and butyrylcholinesterase by chlorpyrifos-oxon. *Biochem. Pharmacol.*, **56**(3): 293-299.
- Ansari R A, Kaur M, Ahmad F, Rahman S, Rashid H, Islam F, Raisuddin S. 2009. Genotoxic and oxidative stress-inducing effects of deltamethrin in the erythrocytes of a freshwater biomarker fish species, *Channa punctata* Bloch. *Environ. Toxicol.*, **24**(5): 429-436.
- APHA-AWWA-WEF. 2005. Standard Methods for the Examination of Water and Wastewater. 21st edn. American Public Health Association, American Water Works Association, Water Environment Federation, Washington, DC .
- Ayllon F, Garcia-Vazquez E. 2000. Induction of micronuclei and other nuclear abnormalities in European minnow *Phoxinus phoxinus* and mollie *Poecilia latipinna*: an assessment of the fish micronucleus test. *Mutat. Res./Genet. Toxicol. Environ. Mutag.*, **467**(2): 177-186.
- Bolognesi C, Perrone E, Roggieri P, Pampanin D M, Sciuotto A. 2006. Assessment of micronuclei induction in peripheral erythrocytes of fish exposed to xenobiotics under controlled conditions. *Aquat. Toxicol.*, **78**(Suppl. 1): S93-S98.
- Buschini A, Martino A, Gustavino B, Monfrinotti M, Poli P, Rossi C, Santoro M, Dörr A J M, Rizzoni M. 2004. Comet assay and micronucleus test in circulating erythrocytes of *Cyprinus carpio* specimens exposed in situ to lake waters treated with disinfectants for potabilization. *Mutat. Res./Genet. Toxicol. Environ. Mutagen.*, **557**(2): 119-129.
- Cadet J, Douki T, Gasparutto D, Ravanat J L. 2003. Oxidative damage to DNA: formation, measurement and biochemical features. *Mutat. Res./Fund. Mol. Mech. Mutagen.*, **531**: 5-23.
- Carrasco K R, Tilbury K L, Myers M S. 1990. Assessment of the piscine micronucleus test as an in situ biological indicator of chemical contaminant effects. *Can. J. Fish. Aquat. Sci.*, **47**(11): 2 123-2 136.
- Cavalcante D G S M, Martinez C B R, Sofia S H. 2008. Genotoxic effects of Roundup® on the fish *Prochilodus*

- lineatus*. *Mutat. Res./Genet. Toxicol. Environ. Mutagen.*, **655**(1-2): 41-46.
- Çavaş T, Ergene-Gözükara S. 2003a. Micronuclei, nuclear lesions and interphase silver-stained nucleolar organizer regions (AgNORs) as cyto-genotoxicity indicators in *Oreochromis niloticus* exposed to textile mill effluent. *Mutat. Res./Genet. Toxicol. Environ. Mutagen.*, **538**(1-2): 81-91.
- Çavaş T, Ergene-Gözükara S. 2003b. Evaluation of the genotoxic potential of lambda-cyhalothrin using nuclear and nucleolar biomarkers on fish cells. *Mutat. Res./Genet. Toxicol. Environ. Mutagen.*, **534**(1-2): 93-99.
- Çavaş T, Ergene-Gözükara S. 2005. Micronucleus test in fish cells: a bioassay for *in situ* monitoring of genotoxic pollution in the marine environment. *Environ. Mol. Mutagen.*, **46**(1): 64-70.
- Chamarthi R R, Bangeppagari M, Gooty J M, Mandala S, Tirado J O, Marigoudar S R. 2014. Histopathological alterations in the gill, liver and brain of *Cyprinus carpio* on exposure to quinalphos. *Am. J. Life Sci.*, **2**(4): 211-216.
- Chebbi S G, David M. 2009. Neurobehavioral responses of the freshwater teleost, *Cyprinus carpio* (Linnaeus.) under quinalphos intoxication. *Biotech. Anim. Husband.*, **25**(3-4): 241-249.
- Chebbi S G, David M. 2010a. Respiratory responses and behavioural anomalies of the carp *Cyprinus carpio* under quinalphos intoxication in sublethal doses. *Sci. Asia*, **36**(1): 12-17.
- Chebbi S G, David M. 2010b. Quinalphos induced alterations in the levels of ions and whole animal oxygen consumption of freshwater fish, *Cyprinus carpio* (Linnaeus, 1758). *J. Vet. Sci. Technol.*, **1**(1): 102.
- da Silva Souza T, Fontanetti C S. 2006. Micronucleus test and observation of nuclear alterations in erythrocytes of Nile tilapia exposed to waters affected by refinery effluent. *Mutat. Res./Genet. Toxicol. Environ. Mutagen.*, **605**(1-2): 87-93.
- Das B K, Mukherjee S C. 2000. Chronic toxic effects of quinalphos on some biochemical parameters in *Labeo rohita* (Ham.). *Toxicol. Lett.*, **114**(1-3): 11-18.
- de Campos Ventura B, de Angelis D D F, Marin-Morales M A. 2008. Mutagenic and genotoxic effects of the Atrazine herbicide in *Oreochromis niloticus* (Perciformes, Cichlidae) detected by the micronuclei test and the comet assay. *Pestic. Biochem. Phys.*, **90**(1): 42-51.
- de Lemos C T, Rödel P M, Terra N R, de Oliveira N C D, Erdtmann B. 2007. River water genotoxicity evaluation using micronucleus assay in fish erythrocytes. *Ecotoxicol. Environ. Saf.*, **66**(3): 391-407.
- Durairaj S, Selvarajan V R. 1992. Influence of quinalphos, an organophosphorous pesticide, on the biochemical constituents of the tissues of fish, *Oreochromis mossambicus*. *J. Environ. Biol.*, **13**(3): 181-185.
- Dwivedi P D, Das M, Khanna S K. 1998. Role of cytochrome P-450 in quinalphos toxicity: effect on hepatic and brain antioxidant enzymes in Rats ITRC communication No. 1965. *Food Chem. Toxicol.*, **36**(5): 437-444.
- Ecobichon D J. 1991. Toxic effect of pesticides. In: Amdur M O, Doull J, Klaassen CD eds. *Casarett and Doull's Toxicology: The Basic Science of Poisons*. Pergamon Press, New York. p.565-622.
- Ergene S, Çavaş T, CÇelik A, Köleli N, Kaya F, Karahan A. 2007. Monitoring of nuclear abnormalities in peripheral erythrocytes of three fish species from the Goksu Delta (Turkey): genotoxic damage in relation to water pollution. *Ecotoxicology*, **16**(4): 385-391.
- Farah M A, Ateeq B, Ali M N, Ahmad W. 2003. Evaluation of genotoxicity of PCP and 2, 4-D by micronucleus test in freshwater fish *Channa punctatus*. *Ecotoxicol. Environ. Saf.*, **54**(1): 25-29.
- Ferraro M V M, Fenocchio A S, Mantovani M S, de Oliveira Ribeiro C, Cestari M M. 2004. Mutagenic effects of tributyltin and inorganic lead (Pb II) on the fish *H. malabaricus* as evaluated using the comet assay and the piscine micronucleus and chromosome aberration tests. *Genet. Mol. Biol.*, **27**(1): 103-107.
- Finney D J. 1952. *Probit Analysis*. Cambridge University Press, New York, USA. 337p.
- Frenzilli G, Nigro M, Lyons B P. 2009. The Comet assay for the evaluation of genotoxic impact in aquatic environments. *Mutat. Res./Rev. Mutat. Res.*, **681**(1): 80-92.
- Gupta B, Rani M, Salunke R, Kumar R. 2012. *In vitro* and *in vivo* studies on degradation of quinalphos in rats. *J. Hazard. Mater.*, **213**: 285-291.
- Gupta P K, Khangarot B S, Durve V S. 1981. The temperature dependence of the acute toxicity of copper to a freshwater pond snail, *Viviparus bengalensis* L. *Hydrobiologia*, **83**(3): 461-464.
- Harshbarger J C, Clark J B. 1990. Epizootiology of neoplasms in bony fish of North America. *Sci. Total. Environ.*, **94**(1-2): 1-32.
- Hayes J W Jr, Laws E R Jr. 1991. *Handbook of Pesticide Toxicology Volume 1: General Principles*. Academic Press, San Diego. p.1-137.
- Hemalatha D, Amala A, Rangasamy B, Nataraj B, Ramesh M. 2016. Sublethal toxicity of quinalphos on oxidative stress and antioxidant responses in a freshwater fish *Cyprinus carpio*. *Environ. Toxicol.*, **31**(11): 1 399-1 406.
- Imanikia S, Galea F, Nagy E, Phillips D H, Stürzenbaum S R, Arlt V M., 2016. The application of the comet assay to assess the genotoxicity of environmental pollutants in the nematode *Caenorhabditis elegans*. *Environ. Toxicol. Pharmacol.*, **45**: 356-361.
- Jha A N. 2008. Ecotoxicological applications and significance of the comet assay. *Mutagenesis*, **23**(3): 207-221.
- Kavitha P, Rao J V. 2008. Toxic effects of chlorpyrifos on antioxidant enzymes and target enzyme acetylcholinesterase interaction in mosquito fish, *Gambusia affinis*. *Environ. Toxicol. Pharmacol.*, **26**(2): 192-198.
- Kilemade M F, Hartl M G J, Sheehan D, Mothersill C, Van Pelt F N A M, O'Halloran J, O'Brien N M. 2004. Genotoxicity of field-collected inter-tidal sediments from Cork Harbor, Ireland, to juvenile turbot (*Scophthalmus maximus* L.) as

- measured by the comet assay. *Environ. Mol. Mutagen.*, **44**(1): 56-64.
- Kokilavani P, Suriyakalaa U, Elumalai P, Abirami B, Ramachandran R, Sankarganesh A, Achiraman S. 2014. Antioxidant mediated ameliorative steroidogenesis by *Commelina benghalensis* L. and *Cissus quadrangularis* L. against quinalphos induced male reproductive toxicity. *Pestic. Biochem. Physiol.*, **109**: 18-33.
- Kumar R, Nagpure N S, Kushwaha B, Srivastava S K, Lakra W S. 2010. Investigation of the genotoxicity of malathion to freshwater teleost fish *Channa punctatus* (Bloch) using the micronucleus test and comet assay. *Arch. Environ. Contam. Toxicol.*, **58**(1): 123-130.
- Kumar S P. 2012. Micronucleus assay: a sensitive indicator for aquatic pollution. *Int. J. Res. BioSci.*, **1**(2): 32-37.
- Lee R F, Steinert S. 2003. Use of the single cell gel electrophoresis/comet assay for detecting DNA damage in aquatic (marine and freshwater) animals. *Mutat. Res./Rev. Mutat. Res.*, **544**(1): 43-64.
- Li H, Wang Y, Shi L Q, Mi J, Song D, Pan X J. 2012. Distribution and fractions of phosphorus and nitrogen in surface sediments from Dianchi Lake, China. *Int. J. Environ. Res.*, **6**(1): 195-208.
- Matsumoto S T, Mantovani M S, Malagutti M I A, Dias A L, Fonseca I C, Marin-Morales M A. 2006. Genotoxicity and mutagenicity of water contaminated with tannery effluents, as evaluated by the micronucleus test and comet assay using the fish *Oreochromis niloticus* and chromosome aberrations in onion root-tips. *Genet. Mol. Biol.*, **29**(1): 148-158.
- Mert R, Benli A Ç K, Arslan G. 2015. Determination of histological and genotoxic effects of formalin on Nile tilapia (*Oreochromis niloticus* L.). *Aquacult. Res.*, **46**(11): 2 798-2 807.
- Mitchell S H, Kennedy S. 1992. Tissue concentrations of organochlorine compounds in common seals from the Coast of Northern Ireland. *Sci. Total. Environ.*, **115**(1-2): 163-77.
- Monteiro V, Cavalcante D G S M, Viléla M B F A, Sofia S H, Martinez C B R. 2011. *In vivo* and *in vitro* exposures for the evaluation of the genotoxic effects of lead on the Neotropical freshwater fish *Prochilodus lineatus*. *Aquat. Toxicol.*, **104**(3-4): 291-298.
- Muranli F D G, Güner U. 2011. Induction of micronuclei and nuclear abnormalities in erythrocytes of mosquito fish (*Gambusia affinis*) following exposure to the pyrethroid insecticide lambda-cyhalothrin. *Mutat. Res./Genet. Toxicol. Environ. Mutagen.*, **726**(2): 104-108.
- Muttappa K, Reddy H R V, Rajesh M, Padmanabha A. 2014. Quinalphos induced alteration in respiratory rate and food consumption of freshwater fish *Cyprinus carpio*. *J. Environ. Biol.*, **35**(2): 395-398.
- Nwani C D, Nagpure N S, Kumar R, Kushwaha B, Kumar P, Lakra W S. 2010. Lethal concentration and toxicity stress of carbosulfan, glyphosate and atrazine to freshwater air breathing fish *Channa punctatus* (Bloch). *Int. Aquat. Res.*, **2**: 105-111.
- Nwani C D, Nagpure N S, Kumar R, Kushwaha B, Kumar P, Lakra W S. 2011. Mutagenic and genotoxic assessment of atrazine-based herbicide to freshwater fish *Channa punctatus* (Bloch) using micronucleus test and single cell gel electrophoresis. *Environ. Toxicol. Pharmacol.*, **31**(2): 314-322.
- Nwani C D, Nagpure N S, Kumar R, Kushwaha B, Kumar P, Lakra W S. 2014. Induction of micronuclei and nuclear lesions in *Channa punctatus* following exposure to carbosulfan, glyphosate and atrazine. *Drug. Chem. Toxicol.*, **37**(4): 370-377.
- Oruc E O, Sevgiler Y, Uner N. 2004. Tissue-specific oxidative stress responses in fish exposed to 2, 4-D and azinphosmethyl. *Comp. Biochem. Physiol. Part C Toxicol. Pharmacol.*, **137**(1): 43-51.
- Pacheco M, Santos M A. 1997. Induction of EROD activity and genotoxic effects by polycyclic aromatic hydrocarbons and resin acids on the juvenile eel (*Anguilla anguilla* L.). *Ecotoxicol. Environ. Saf.*, **38**(3): 252-259.
- Palanikumar L, Kumaraguru A K, Ramakritinan C M, Anand M. 2012. Genotoxic assessment of anthracene and benzo[a]pyrene to milkfish *Chanos chanos*. *Toxicol. Environ. Chem.*, **94**(2): 350-363.
- Pandey A K, Nagpure N S, Trivedi S P, Kumar R, Kushwaha B. 2011. Profenofos induced DNA damage in freshwater fish, *Channa punctatus* (Bloch) using alkaline single cell gel electrophoresis. *Mutat. Res./Genet. Toxicol. Environ. Mutagen.*, **726**(2): 209-214.
- Pandey S, Nagpure N S, Kumar R, Sharma S, Srivastava S K, Verma M S. 2006. Genotoxicity evaluation of acute doses of endosulfan to freshwater teleost *Channa punctatus* (Bloch) by alkaline single-cell gel electrophoresis. *Ecotoxicol. Environ. Saf.*, **65**(1): 56-61.
- Pant N, Srivastava S P. 2003. Testicular and spermatotoxic effects of quinalphos in rats. *J Appl Toxicol*, **23**(4): 271-274.
- Park E H, Lee J S, Yi A K, Etoh H. 1989. Fish cell line (ULF-23HU) derived from the fin of the central mudminnow (*Umbra limi*): suitable characteristics for clastogenicity assay. *In Vitro Cell. Dev. Biol.*, **25**(11): 987-994.
- Racke K D. 1993. Environmental fate of chlorpyrifos. *Rev. Environ. Contam. Toxicol.*, **131**: 1-150.
- Raisuddin S, Jha A N. 2004. Relative sensitivity of fish and mammalian cells to sodium arsenate and arsenite as determined by alkaline single-cell gel electrophoresis and cytokinesis-block micronucleus assay. *Environ. Mol. Mutagen.*, **44**(1): 83-89.
- Raisuddin S, Lee J S. 2008. Fish models in impact assessment of carcinogenic potential of environmental chemical pollutants: an appraisal of hermaphroditic mangrove killifish *Kryptolebias marmoratus*. In: Murakami Y, Nakayama K, Kitamura S I, Iwata H, Tanabe S eds. *Interdisciplinary Studies on Environmental Chemistry-Biological Responses to Chemical Pollutants*. TERRAPUB, Tokyo. p.7-15.
- Ray A, Chatterjee S, Ghosh S, Bhattacharya K, Pakrashi A, Deb C. 1992. Quinalphos-induced suppression of spermatogenesis, plasma gonadotrophins, testicular

- testosterone production, and secretion in adult rats. *Environ. Res.*, **57**(2): 181-189.
- Rupa D S, Reddy P P, Reddi O S. 1991a. Cytogenetic effects of quinalphos in mice. *Food Chem. Toxicol.*, **29**(2): 115-117.
- Rupa D S, Reddy P P, Reddi O S. 1991b. Frequency of sister chromatid exchange in peripheral lymphocytes of male pesticide applicators. *Environ. Mol. Mutagen.*, **18**(2): 136-138.
- Sadiqul I M, Ferdous Z, Nannu T A, Mostakim G M, Rahman K. 2016. Acute exposure to a quinalphos containing insecticide (convoy) causes genetic damage and nuclear changes in peripheral erythrocytes of silver barb, *Barbonymus gonionotus*. *Environ. Pollut.*, **219**: 949-956.
- Sastry K V, Siddiqui A A, Singh S K. 1982. Alteration in some biochemical and enzymological parameters in the snake head fish *Channa punctatus*, exposed chronically to quinalphos. *Chemosphere*, **11**(12): 1 211-1 216.
- Scott G I, Fulton M H, Moore D W. 1990. Agricultural insecticide runoff effects on estuarine organisms: correlating laboratory and field toxicity testing with ecotoxicological biomonitoring. U.S. National Marine Fisheries Service, Southeast Fisheries Science Center, Charleston Laboratory, Charleston, Report #CR813138-01-1.
- Sharma S K, Mathur B N. 1972. Toxicity data references for quinalphos, national institute of occupational safety and health, EPA IRIS database. *J. Agric. Sci.*, **3**: 108.
- Shukla D, Nagpure N S, Kumar R, Singh P J. 2010. Assessment of genotoxicity of dichlorvos to *Mystus vittatus* (Bloch) by comet assay. *Indian J. Fish.*, **57**(2): 39-44.
- Singh N P, McCoy M T, Tice R R, Schneider E L. 1988. A simple technique for quantitation of low levels of DNA damage in individual cells. *Exp. Cell Res.*, **175**(1): 184-191.
- Sprague J B. 1969. Measurement of pollutant toxicity to fish I. Bioassay methods for acute toxicity. *Water Res.*, **3**(11): 793-821.
- Srivastava M K, Raizada R B, Dikshith T S S. 1992. Fetotoxic response of technical quinalphos in rats. *Vet. Hum. Toxicol.*, **34**(2): 131-133.
- Subramanayan M, Jain S, Yadav C, Arora V K, Banerjee B D, Ahmed R S. 2012. Quinalphos induced oxidative stress and histoarchitectural alterations in adult male albino rats. *Environ. Toxicol. Pharmacol.*, **34**(3): 673-678.
- Sundaram K M S, Sundaram A, Hammock B D. 1994. Persistence of *Bacillus thuringiensis* deposits in a hardwood forest, after aerial application of a commercial formulation at two dosage rates. *J. Environ. Sci. Health Part B*, **29**(5): 999-1 052.
- Talapatra S N, Banerjee S K. 2007. Detection of micronucleus and abnormal nucleus in erythrocytes from the gill and kidney of *Labeo bata* cultivated in sewage-fed fish farms. *Food Chem. Toxicol.*, **45**(2): 210-215.
- Tice R R, Agurell E, Anderson D, Burlinson B, Hartmann A, Kobayashi H, Miyamae Y, Rojas E, Ryu J C, Sasaki Y F. 2000. Single cell gel/comet assay: guidelines for in vitro and in vivo genetic toxicology testing. *Environ. Mol. Mutagen.*, **35**(3): 206-221.
- Udroiu I. 2006. The micronucleus test in piscine erythrocytes. *Aquat. Toxicol.*, **79**(2): 201-204.
- Vardavas A I, Stivaktakis P D, Tzatzarakis M N, Fragkiadaki P, Vasilaki F, Tzardi M, Datseri G, Tsiaoussis J, Alegakis A K, Tsitsimpikou C, Rakitskii V N, Carvalho F, Tsatsakis A M. 2016. Long-term exposure to cypermethrin and piperonyl butoxide cause liver and kidney inflammation and induce genotoxicity in New Zealand white male rabbits. *Food Chem. Toxicol.*, **94**: 250-259.
- Vasilić Ž, Drevenkar V, Rumenjak V, Stengl B, Frobe Z. 1992. Urinary excretion of diethylphosphorus metabolites in persons poisoned by quinalphos or chlorpyrifos. *Arch. Environ. Contam. Toxicol.*, **22**(4): 351-357.
- Venkata Rathnamma V, Nagaraju B, Aanitha A. 2013a. Impact of quinalphos on acid phosphatase and acetyl cholinesterase of the fresh-water Fish *Labeo rohita* (Hamilton). *J. Biol. Today's World*, **2**: 493-497.
- Venkata Rathnamma V, Nagaraju B. 2013b. Lethal and sublethal effects of quinalphos on nucleic acids of freshwater fish *Labeo rohita* (Hamilton). *Int. J. Res. Appl. Nat. Soc. Sci.*, **1**(3): 105-110.
- Vig K, Singh D K, Sharma P K. 2006. Endosulfan and quinalphos residues and toxicity to soil microarthropods after repeated applications in a field investigation. *J. Environ. Sci. Health Part B*, **41**(5): 681-692.
- WHO. 2002. The WHO recommended classification of pesticides by hazard and guidelines to classification 2000-2002. WHO/PCS/01.5. WHO, Geneva.

Electronic supplementary material

Supplementary material (Supplementary Fig.S1) is available in the online version of this article at <https://doi.org/10.1007/s00343-019-9104-y>.