

The minimal covering set in large tournaments

Alex Scott · Mark Fey

Received: 23 February 2010 / Accepted: 1 September 2010 / Published online: 19 September 2010
© Springer-Verlag 2010

Abstract We prove that in almost all large tournaments, the minimal covering set is the entire set of alternatives. That is, as the number of alternatives gets large, the probability that the minimal covering set of a uniformly chosen random tournament is the entire set of alternatives goes to one. In contrast, it follows from a result of (Fisher and Reeves, *Linear Algebra Appl* 217:83–85, 1995) that the bipartisan set contains about half of the alternatives in almost all large tournaments.

1 Introduction

Tournaments have long played an important role in social choice theory as a means of representing and understanding majority voting. A healthy literature has developed that is centered on defining and describing various tournament solutions, which are meant to mathematically identify which alternatives are “best” in any given tournament.¹ These tournament solutions include the top cycle set (Schwartz 1972; Miller 1977), the uncovered set (Miller 1980), the Banks set (Banks 1985), the minimal covering set (Dutta 1988), the bipartisan set (Laffond et al. 1993), the tournament equilibrium set (Schwartz 1990), and others.

A central principle in the search for new tournament solutions is that “smaller is better.” That is, tournament solutions that select small sets of “best” alternatives are preferred to those that are not as discriminating. Typically, scholars focus on the

¹ For a comprehensive survey of tournament solutions see Laslier (1997).

A. Scott
Mathematical Institute, University of Oxford, Oxford, UK
e-mail: scott@maths.ox.ac.uk

M. Fey (✉)
Department of Political Science, University of Rochester, Rochester, NY, USA
e-mail: mark.fey@rochester.edu

relative size of tournament solutions to justify these claims. For example, the Banks set is always included in the uncovered set and this is taken as indicating the superiority of the Banks set to the uncovered set. However, little attention has been paid to the *absolute* size of tournament solutions. In particular, it is important to understand what the typical size of a given tournament solution is in order to evaluate how discriminating it is.

Building on initial work by Bell (1981), this question was the focus of Fey (2008). In that article, Fey shows that in almost all tournaments, several standard tournament solutions do not offer any help in discriminating among alternatives. Specifically, Fey (2008) shows that with probability approaching one, the top cycle set, the uncovered set, and the Banks set are equal to the entire set of alternatives in a uniformly chosen large tournament.² In other words, these tournament solutions almost never narrow the set of social choices.

Fey (2008) conjectures that the same result is true for the minimal covering set. In this article, we confirm that this conjecture holds. That is, we show that in almost all tournaments, the minimal covering set is the entire set of alternatives. Thus, in large tournaments, this tournament solution fails to limit the choice of alternatives at all.

In contrast, we consider the size of the bipartisan set in large tournaments. The bipartisan set is always contained in the minimal covering set and we draw on existing results in graph theory to show that in large tournaments, the bipartisan set almost always contains close to half the alternatives.

2 Tournaments and tournament solutions

Suppose there is a finite set of *alternatives*, which we denote X . A *tournament* T is a complete and asymmetric binary relation on X . Such a tournament can arise, for example, as the majority preference relation of an odd number of voters with linear preferences. As is standard, if xTy , then we say that x *beats* y . For a subset $S \subseteq X$, we call $T \cap (S \times S)$ the *subtournament* of T generated by S .³ For a fixed alternative $x \in X$, let $T(x) = \{y \in X \mid yTx\}$ denote the set of alternatives that beat x and let $N(x) = \{y \in X \mid xTy\}$ denote the set of alternatives that x beats. For a set $Y \subseteq X$, we let $N(Y) = \bigcup_{x \in Y} N(x)$ denote the set of alternatives that is beaten by some alternative in Y .

A *tournament solution* is a way of identifying a set of “best” alternatives from a tournament T . Formally, a tournament solution is a correspondence that, for every tournament T , selects a nonempty subset of X . A number of important tournament solutions are based on the *covering* relation. We say x *covers* y if xTy and for all $z \in X$, yTz implies xTz . An alternative x is *uncovered* if there is no y that covers x . The *uncovered set* of T , $UC(T)$, is the set of alternatives that are not covered by any other alternative (Miller 1980, 1983). For a tournament T and a set $S \subseteq X$, the uncovered

² A uniformly chosen tournament is one chosen from the uniform distribution on the set of all tournaments of a given size. Fey (2008) conjectures that this result would hold for majority voting tournaments generated by randomly assigning preference orders to a set of voters.

³ When there is no chance of ambiguity, we use S to refer to both the subtournament and the set of alternatives on which the subtournament is defined.

set of S , denoted $UC(S)$ is the uncovered set of the subtournament of T generated by S . We say $S \subseteq X$ is a *covering set* for T if for every $x \in X \setminus S$, $x \notin UC(S \cup \{x\})$. The set X is trivially a covering set for T and therefore we say S is a *proper covering set* for T if S is a covering set for T and $S \neq X$. Dutta (1988) shows that there is a covering set for T that is contained in all covering sets for T . Therefore, for every tournament T , there exists a unique *minimal covering set* for T , which we denote $MC(T)$.

Our result concerns the size of the minimal covering set in a random tournament. To define the notion of a *random tournament*, let \mathcal{T}_n denote the set of possible tournaments on n alternatives, $n \geq 3$. This set contains $2^{\binom{n}{2}}$ distinct tournaments, each of which we take to be equally likely. Equivalently, a random tournament $T \in \mathcal{T}_n$ is obtained by choosing independently, for each pair of alternatives $x, y \in X$, $x \neq y$, either xTy or yTx with equal probability. For a random variable Z , we use the notation $\mathbb{E}[Z]$ and $\mathbb{P}[Z = z]$ for the expectation of Z and the probability Z takes a value z , respectively.

3 Main result

Our main result states that the minimal covering set is equal to the entire set of alternatives in almost all large tournaments. Thus, with probability approaching one, the minimal covering set is completely indiscriminating in tournaments with a large number of alternatives. As is standard in the literature of random graphs (Bollobás 2001), we say a property \mathcal{Q} holds for almost all tournaments if the probability that a random tournament has property \mathcal{Q} goes to one as the number of alternatives, n , goes to infinity.

Theorem 1 *In almost all tournaments, there are no proper covering sets of T . Therefore, in almost all tournaments, $MC(T) = X$.*

Before we prove our main result, we present several lemmas. Our first lemma records a standard fact that will be useful in the proof of our main theorem.

Lemma 1 *Let Z_n be a sequence of random variables that each takes values in the set of nonnegative integers. If $\mathbb{E}[Z_n] \rightarrow 0$ as $n \rightarrow \infty$, then $\mathbb{P}[Z_n = 0] \rightarrow 1$ as $n \rightarrow \infty$.*

Proof For each Z_n we have

$$\mathbb{E}[Z_n] = \sum_{k=0}^{\infty} k\mathbb{P}[Z_n = k] \geq \sum_{k=1}^{\infty} \mathbb{P}[Z_n = k] = 1 - \mathbb{P}[Z_n = 0].$$

It follows that $\mathbb{E}[Z_n] \rightarrow 0$ implies $\mathbb{P}[Z_n = 0] \rightarrow 1$. □

Our next two lemmas are fairly standard (Moon 1968), but for completeness we provide proofs here. Our final lemma is new and may have some independent interest.

We say a subset S of T is *dominating* if $T = S \cup N(S)$, which means that for every $y \notin S$ there is $x \in S$ such that xTy . In other words, S is a dominating set if every alternative outside of S is beaten by some alternative in S . Trivially, the entire set X is a dominating set and therefore every tournament has a dominating set. We write $dom(T)$ for the minimum cardinality of a dominating set. Our first lemma gives a maximum size for $dom(T)$.

Lemma 2 *If $|X| = n$, then $\text{dom}(T) \leq \log_2(n + 1)$.*

Proof Note that the assertion is true for $n = 1$. We continue by induction. Let T be a tournament on n vertices. Then T has $\binom{n}{2}$ edges, and so there is some alternative x with $|N(x)| \geq (n - 1)/2$. Let $S = T(x)$ and let R be a dominating set for S . Then $R \cup \{x\}$ is a dominating set for T and therefore

$$\text{dom}(T) \leq 1 + \text{dom}(S).$$

But $|S| = n - |N(x)| - 1 \leq (n - 1)/2$ and so by the induction hypothesis,

$$\text{dom}(T) \leq 1 + \log_2 \left(\frac{n + 1}{2} \right) = \log_2(n + 1).$$

□

Our second lemma gives a minimum size for $\text{dom}(T)$ in almost all tournaments.

Lemma 3 *Almost all tournaments have $\text{dom}(T) > \frac{1}{2} \log_2 n$.*

Proof From the definition of $\text{dom}(T)$, it follows that $\text{dom}(T) > k$ if and only if for every set $A \subseteq X$ with $|A| = k$ there exists an alternative x with $A \subseteq N(x)$. Define a random variable K to be the number of sets $A \subseteq X$ with $|A| = k$ in a random tournament such that there is no alternative x with $A \subseteq N(x)$. Then $\mathbb{P}[K = 0]$ is the probability that a random tournament T has $\text{dom}(T) > k$.

For a given set $A \subseteq X$ with $|A| = k$ and a given alternative $x \notin A$, the probability that $A \subseteq N(x)$ in a random tournament is 2^{-k} . Therefore, the probability that there is no x with $A \subseteq N(x)$ is

$$(1 - 2^{-k})^{n-k} \leq \exp(-(n - k)2^{-k}),$$

where the right hand side follows from the fact that $1 + z \leq \exp(z)$ for all real z . The expected number of such k -sets with no such x thus satisfies

$$\begin{aligned} \mathbb{E}[K] &\leq \binom{n}{k} \exp(-(n - k)2^{-k}) \\ &\leq n^k \exp(-(n - k)2^{-k}) \\ &\leq \exp(k \log n - (n - k)2^{-k}). \end{aligned}$$

which tends to 0 if $k \log n - (n - k)2^{-k} \rightarrow -\infty$. But it is easy to verify that this is true if $k = \lceil \frac{1}{2} \log n \rceil$. Thus, by Lemma 1, $\mathbb{P}[K = 0] \rightarrow 1$ as $n \rightarrow \infty$. □

As is clear from the proof, the constant $1/2$ is not optimal (and can be replaced by any number less than 1).

Our final lemma points out an important connection between covering sets and dominating sets.

Lemma 4 *For a tournament T and a set $S \subseteq X$, if $\text{dom}(S) < \text{dom}(T)$, then S is not a covering set of T .*

Proof Fix a tournament T and a set $S \subseteq X$ with $\text{dom}(S) < \text{dom}(T)$ and let $k = \text{dom}(S)$. Let $R \subseteq S$ be a dominating set of S with $|R| = k$. We claim that there exists an alternative $x \notin S$ such that xTr for all $r \in R$. If not, then R is a dominating set for T , which contradicts that assumption that $\text{dom}(S) < \text{dom}(T)$. Thus, xTr for all $r \in R$ and for all $y \in S \setminus R$, there exists $r \in R$ such that rTy . This means that $x \in \text{UC}(S \cup \{x\})$, so S is not a covering set of T . \square

We are now ready to prove our main result.

Proof of Theorem 1 Define a random variable C to be the number of proper covering sets of a random tournament. Then $\mathbb{P}[C = 0]$ is the probability that a random tournament has the property that $\text{MC}(T) = X$. Our argument will split the set of tournaments of order n into two groups: those that have a small dominating set, and those that do not. Formally,

$$\begin{aligned} \mathbb{P}[C > 0] &= \mathbb{P}\left[C > 0 \mid \text{dom}(T) \leq \frac{\log_2 n}{2}\right] \mathbb{P}\left[\text{dom}(T) \leq \frac{\log_2 n}{2}\right] \\ &\quad + \mathbb{P}\left[C > 0 \mid \text{dom}(T) > \frac{\log_2 n}{2}\right] \mathbb{P}\left[\text{dom}(T) > \frac{\log_2 n}{2}\right] \\ &\leq \mathbb{P}\left[\text{dom}(T) \leq \frac{\log_2 n}{2}\right] + \mathbb{P}\left[C > 0 \mid \text{dom}(T) > \frac{\log_2 n}{2}\right], \end{aligned}$$

where the last step comes from the trivial bound that a probability is at most 1. To prove our result, it suffices to show that both of these terms go to zero as n goes to infinity. But Lemma 3 implies that this is true for the first term. So all that remains is to show that $\mathbb{P}[C > 0 \mid \text{dom}(T) > (\log_2 n)/2]$ goes to zero as n goes to infinity.

Consider a tournament T with $\text{dom}(T) > (\log_2 n)/2$ and $n \geq 4$. By Lemma 4, a set $S \subseteq X$ with $\text{dom}(S) < \text{dom}(T)$ is not a covering set, therefore we need only concern ourselves with subsets S with $\text{dom}(S) \geq \text{dom}(T)$. So fix such a set S and let $x \in S$ be arbitrary. The set $T(x) \cap S$ must be non-empty, because otherwise $\{x\}$ would be a dominating set for S , which would contradict $\text{dom}(S) \geq \text{dom}(T) > (\log_2 n)/2$. So let R be a dominating set for $T(x) \cap S$. Then $R \cup \{x\}$ is a dominating set for S and therefore

$$\begin{aligned} \text{dom}(S) &\leq 1 + \text{dom}(T(x) \cap S) \\ &\leq 1 + \log_2(|T(x) \cap S| + 1). \end{aligned}$$

As $\text{dom}(S) \geq \text{dom}(T) > (\log_2 n)/2$, we have

$$\begin{aligned} (\log_2 n)/2 &< 1 + \log_2(|T(x) \cap S| + 1) \\ \log_2 n^{1/2} &< \log_2 2 + \log_2(|T(x) \cap S| + 1) \\ \log_2(n^{1/2}/2) &< \log_2(|T(x) \cap S| + 1) \\ n^{1/2}/2 &< |T(x) \cap S| + 1. \end{aligned}$$

Thus, $|T(x) \cap S| > n^{1/2}/2 - 1$, which is at least $n^{1/2}/3$ for large enough n .

We now turn to calculating $\mathbb{P}[C > 0 \mid \text{dom}(T) > (\log_2 n)/2]$. A subset S can be a proper covering set only if $\text{dom}(S) \geq \text{dom}(T)$. By Lemma 2, any such set S must have at least $\sqrt{n} - 1$ vertices and, from above, every alternative x in S must have $|T(x) \cap S| \geq n^{1/2}/3$. Furthermore, if $y \notin S$ has $|N(y) \cap T(x) \cap S| > 0$ then x does not cover y in $S \cup \{y\}$. Let us therefore write C' for the number of sets S such that

$$(P1) \quad |S| \geq \sqrt{n} - 1.$$

$$(P2) \quad \text{For all } x \in S, |T(x) \cap S| \geq n^{1/2}/3.$$

$$(P3) \quad \text{For all } y \notin S \text{ there is } x \in S \text{ such that } |N(y) \cap T(x) \cap S| = 0.$$

Since a covering set S in a tournament T with $\text{dom}(T) > (\log_2 n)/2$ must satisfy all three properties, we have $C' = C$ for all such tournaments. So

$$\begin{aligned} \mathbb{P}[C > 0 \mid \text{dom}(T) > (\log_2 n)/2] &= \mathbb{P}[C' > 0 \mid \text{dom}(T) > (\log_2 n)/2] \\ &= \frac{\mathbb{P}[C' > 0 \text{ and } \text{dom}(T) > (\log_2 n)/2]}{\mathbb{P}[\text{dom}(T) > (\log_2 n)/2]} \\ &\leq 2\mathbb{P}[C' > 0], \end{aligned}$$

since $\mathbb{P}[\text{dom}(T) > (\log_2 n)/2] > 1/2$ for sufficiently large n by Lemma 3. It will therefore be sufficient to prove that $\mathbb{P}[C' > 0] \rightarrow 0$ as $n \rightarrow \infty$. This is simpler than dealing with $\mathbb{P}[C > 0 \mid \text{dom}(T) > (\log_2 n)/2]$ in two ways: the event C' is easier to analyse than C , and we are no longer conditioning on the size of $\text{dom}(T)$.

Let us first bound $\mathbb{E}[C']$. For a fixed set $S \subset X$ with $n > |S| \geq \sqrt{n} - 1$, consider a random tournament T . Then

$$\begin{aligned} \mathbb{P}[S \text{ satisfies (P2) and (P3)}] &= \mathbb{P}[S \text{ satisfies (P3)} \mid S \text{ satisfies (P2)}]\mathbb{P}[S \text{ satisfies (P2)}] \\ &\leq \mathbb{P}[S \text{ satisfies (P3)} \mid S \text{ satisfies (P2)}]. \end{aligned}$$

Now suppose S satisfies (P2). For any $y \notin S$ and any $x \in S$,

$$\mathbb{P}[|N(y) \cap T(x) \cap S| = 0] \leq 2^{-n^{1/2}/3}.$$

So for any $y \in S$, the probability that there is $x \in S$ such that $|N(y) \cap T(x) \cap S| = 0$ is at most $|S|2^{-n^{1/2}/3}$. These events are independent for each $y \notin S$, and so

$$\mathbb{P}[S \text{ satisfies (P1)–(P3)}] \leq \left(|S|2^{-n^{1/2}/3}\right)^{n-|S|} \leq n^{n-|S|}2^{-n^{1/2}(n-|S|)/3}.$$

We now use this probability to bound $\mathbb{E}[C']$. We have

$$\mathbb{E}[C'] \leq \sum_{\sqrt{n}-1 \leq k \leq n-1} \binom{n}{k} n^{n-k} 2^{-n^{1/2}(n-k)/3}.$$

We split the sum into two pieces. First,

$$\begin{aligned} \sum_{\sqrt{n}-1 \leq k \leq n/2} \binom{n}{k} n^{n-k} 2^{-n^{1/2}(n-k)/3} &\leq \sum_{\sqrt{n}-1 \leq k \leq n/2} n^k n^{n-k} 2^{-n^{1/2}(n/2)/3} \\ &= \sum_{\sqrt{n}-1 \leq k \leq n/2} n^n 2^{-n^{3/2}/6} \\ &= \sum_{\sqrt{n}-1 \leq k \leq n/2} 2^{n \log_2 n} 2^{-n^{3/2}/6} \\ &= \sum_{\sqrt{n}-1 \leq k \leq n/2} 2^{n \log_2 n - n^{3/2}/6}. \end{aligned}$$

Now for the rest:

$$\begin{aligned} \sum_{n/2 \leq k \leq n-1} \binom{n}{k} n^{n-k} 2^{-n^{1/2}(n-k)/3} &= \sum_{n/2 \leq k \leq n-1} \binom{n}{n-k} n^{n-k} 2^{-n^{1/2}(n-k)/3} \\ &\leq \sum_{n/2 \leq k \leq n-1} n^{n-k} n^{n-k} 2^{-n^{1/2}(n-k)/3} \\ &\leq \sum_{i \geq 1} n^i n^i 2^{-n^{1/2}i/3} \\ &= \sum_{i \geq 1} \left(2^{2 \log_2 n - n^{1/2}/3} \right)^i. \end{aligned}$$

It is straightforward to show that both of these pieces go to zero as n gets large. By Lemma 1, this implies that $\mathbb{P}[C' = 0] \rightarrow 1$. We conclude then that $\mathbb{P}[C > 0] \rightarrow 0$ as n gets large and thus almost all tournaments have no proper covering sets. \square

4 The bipartisan set

Now that we have established that the minimal covering set is almost always the entire set of alternatives in large tournaments, it is natural to consider if this result continues to hold for tournament solutions that are known to always be contained in the minimal covering set. We address this question by considering the bipartisan set, due to Laffond et al. (1993).

The bipartisan set is defined in terms of the *tournament game* of a tournament T . Specifically, for a given tournament T on a set of alternatives X we associate a two-player zero-sum strategic form game G as follows. We assign to each player the strategy set X , so that each player’s action is simply a choice of an alternative from X . If player i chooses alternative x and player $j \neq i$ chooses alternative y , then the payoff to player i is equal to 1 if xTy , 0 if $x = y$, and -1 if yTx . Laffond et al. (1993) and Fisher and Ryan (1992, 1995) independently prove that a tournament game always has a unique Nash equilibrium. Based on this, Laffond et al. (1993) define the *bipartisan*

set of T , denoted $\text{BP}(T)$, to be the support of the unique Nash equilibrium of the tournament game of T . They go on to show that $\text{BP}(T)$ always consists of an odd number of alternatives and that $\text{BP}(T) \subseteq \text{MC}(T)$ for every tournament T .

We now turn to the question of the size of the bipartisan set. Is it almost always the whole set of alternatives, as with the minimal covering set? It is easy to see that the answer is negative. Because the bipartisan set must have an odd number of elements, it follows that $\text{BP}(T) \neq X$ if $|X|$ is even. So what is the size of the bipartisan set? It turns out that a precise answer has been given by [Fisher and Reeves \(1995\)](#). They prove that for a random tournament T ,

$$\mathbb{P}[|\text{BP}(T)| = k] = \begin{cases} \binom{n}{k} 2^{-(n-1)} & \text{if } k \text{ is odd} \\ 0 & \text{if } k \text{ is even.} \end{cases}$$

Thus, the size distribution of the bipartisan set is a scaled binomial distribution (with $p = 1/2$) on the odd integers in $\{1, \dots, n\}$. Using this, Fisher and Reeves show that $\mathbb{E}[|\text{BP}(T)|] = n/2$. That is, the average size of the bipartisan set in a random tournament is $n/2$. Moreover, it is easy to see from the size distribution above that the distribution of $|\text{BP}(T)|/n$ converges to $1/2$. That is,

$$\lim_{n \rightarrow \infty} \mathbb{P} \left[\left| \frac{|\text{BP}(T)|}{n} - \frac{1}{2} \right| \geq \varepsilon \right] = 0$$

for all $\varepsilon > 0$. Another way to put this is that in large tournaments, the size of the bipartisan set is almost always close to $n/2$. Therefore, the bipartisan set turns out to be about half the size of the minimal covering set in large tournaments.

5 Conclusion

This article confirms a conjecture of [Fey \(2008\)](#) that in almost all tournaments, the minimal covering set is the entire set of alternatives. Thus, we now know that the top cycle set, the uncovered set, the Banks set, and the minimal covering set are all almost always equal to the entire set of alternatives. On the other hand, the bipartisan set almost always contains close to half of the alternatives.

Given these results, an obvious open question is to establish the size of other tournament solutions. For the case of the tournament equilibrium set ([Schwartz 1990](#)), [Brandt et al. \(2010\)](#) reports on computational experiments that showed the tournament equilibrium set almost always was the entire set of alternatives in uniformly chosen large tournaments. This evidence suggests that our result on the minimal covering set of large tournaments may also hold for the tournament equilibrium set. We leave this for future study.

References

- Banks JS (1985) Sophisticated voting outcomes and covering relation. *Soc Choice Welfare* 1:295–306
 Bell CE (1981) A random voting graph almost surely has a Hamiltonian cycle when the number of alternatives is large. *Econometrica* 49:1597–1603

- Bollobás B (2001) *Random graphs*, 2nd edn. Cambridge University Press, Cambridge
- Brandt F, Fischer F, Harrenstein P, Mair M (2010) A computational analysis of the tournament equilibrium set. *Soc Choice Welfare* 34:597–609
- Dutta B (1988) Covering sets and a new Condorcet choice correspondence. *J Econ Theory* 44:63–80
- Fey M (2008) Choosing from a large tournament. *Soc Choice Welfare* 31:301–309
- Fisher DC, Reeves RB (1995) Optimal strategies for random tournament games. *Linear Algebra Appl* 217:83–85
- Fisher DC, Ryan J (1992) Optimal strategies for a generalized “scissors, paper, and stone” game. *Am Math Month* 99:935–942
- Fisher DC, Ryan J (1995) Tournament games and positive tournaments. *J Graph Theory* 19:217–236
- Laffond G, Laslier J, Le Breton M (1993) The bipartisan set of a tournament game. *Games Econ Behav* 5:182–201
- Laslier JF (1997) *Tournament solutions and majority voting*. Springer-Verlag, Berlin
- Miller NR (1977) Graph-theoretical approaches to the theory of voting. *Am J Polit Sci* 21:769–803
- Miller NR (1980) A new solution set for tournaments and majority voting. *Am J Polit Sci* 24:68–96
- Miller NR (1983) The covering relation in tournaments: two corrections. *Am J Polit Sci* 27:382–385
- Moon JW (1968) *Topics on tournaments*. Rinehart and Winston, Holt
- Schwartz T (1972) Rationality and the myth of the maximum. *Nous* 6:97–117
- Schwartz T (1990) Cyclic tournaments and cooperative majority voting: a solution. *Soc Choice Welfare* 7:19–29