


Total Domination Versus Domination in Cubic Graphs

Joanna Cyman¹  · Magda Dettlaff¹ ·
Michael A. Henning² · Magdalena Lemańska¹ ·
Joanna Raczek¹

Received: 5 June 2017 / Revised: 11 November 2017 / Published online: 19 December 2017
© The Author(s) 2017. This article is an open access publication

Abstract A dominating set in a graph G is a set S of vertices of G such that every vertex not in S has a neighbor in S . Further, if every vertex of G has a neighbor in S , then S is a total dominating set of G . The domination number, $\gamma(G)$, and total domination number, $\gamma_t(G)$, are the minimum cardinalities of a dominating set and total dominating set, respectively, in G . The upper domination number, $\Gamma(G)$, and the upper total domination number, $\Gamma_t(G)$, are the maximum cardinalities of a minimal dominating set and total dominating set, respectively, in G . It is known that $\gamma_t(G)/\gamma(G) \leq 2$ and $\Gamma_t(G)/\Gamma(G) \leq 2$ for all graphs G with no isolated vertex. In this paper we characterize the connected cubic graphs G satisfying $\gamma_t(G)/\gamma(G) = 2$, and we characterize the connected cubic graphs G satisfying $\Gamma_t(G)/\Gamma(G) = 2$.

Research supported in part by the South African National Research Foundation and the University of Johannesburg.

✉ Joanna Cyman
joanna.cyman@pg.edu.pl
Magda Dettlaff
magda.dettlaff1@pg.edu.pl
Michael A. Henning
mahenning@uj.ac.za
Magdalena Lemańska
magleman@pg.edu.pl
Joanna Raczek
joanna.raczek@pg.edu.pl

¹ Department of Mathematics, Gdańsk University of Technology, ul. Narutowicza 11/12, 80-952 Gdańsk, Poland

² Department of Pure and Applied Mathematics, University of Johannesburg, Auckland Park 2006, South Africa

Keywords Domination number · Total domination number · Upper domination number · Upper total domination number · Cubic graph

1 Introduction

In this paper we continue the study of domination and total domination in graphs. Let $G = (V, E)$ be a simple graph. If $u, v \in V(G)$, then u dominates v if u and v are adjacent. A *dominating set* of a graph G is a set S of vertices of G such that every vertex outside S is dominated by at least one vertex in S . A *total dominating set*, abbreviated TD-set, of G is a set S of vertices of G such that every vertex in $V(G)$ is adjacent to at least one vertex in S . A *neighbor* of a vertex v in G is a vertex adjacent to v . A vertex *totally dominates* its neighbors. The *domination number* of G , denoted $\gamma(G)$, is the minimum cardinality of a dominating set of G , while the *total domination number* of G , denoted by $\gamma_t(G)$, is the minimum cardinality of a TD-set of G . A TD-set S in G is *minimal* if for all vertices $v \in S$, the set $S \setminus \{v\}$ is not a TD-set of G . The *upper total domination number* of G , denoted $\Gamma_t(G)$, is the maximum cardinality of a minimal TD-set in G . We refer to a minimum total dominating set of G as a $\gamma_t(G)$ -set. A minimal TD-set of cardinality $\Gamma_t(G)$ we call a $\Gamma_t(G)$ -set.

The literature on this subject of domination and its variations has been surveyed and detailed in the two books by Haynes, Hedetniemi, and Slater [9, 10]. For a recent book on total domination in graphs we refer the reader to [15]. A survey of total domination in graphs can also be found in [14]. For a survey of known results on the upper total domination we refer the reader to [16]. Recent papers on total domination in graphs can be found, for example, in [3, 6, 17, 18] and elsewhere.

Let S be a set of vertices in a graph G , and let $v \in S$. The *open neighborhood* of v is the set $N_G(v) = \{u \in V(G) \mid uv \in E(G)\}$ and the *closed neighborhood* of v is $N_G[v] = \{v\} \cup N_G(v)$. We denote the *degree* of v in G by $d_G(v) = |N_G(v)|$, or simply by $d(v)$ if the graph G is clear from context. For a subset S of vertices of G , we let $d_S(v)$ denote the number of vertices in S that are adjacent to v . A graph G is *k-regular* if every vertex has degree k in G . A *regular graph* is a graph that is k -regular for some integer $k \geq 0$. We remark that 3-regular graphs are also called *cubic graphs* in the literature. The *open neighborhood* of S is the set $N_G(S) = \bigcup_{v \in S} N_G(v)$ and its *closed neighborhood* is the set $N_G[S] = N_G(S) \cup S$. The subgraph induced by the set S is denoted by $G[S]$.

The *private neighborhood* $\text{pn}(v, S)$ of $v \in S$ is defined by $\text{pn}(v, S) = N(v) \setminus N(S \setminus \{v\})$. Equivalently, $\text{pn}(v, S) = \{u \in V \mid N(u) \cap S = \{v\}\}$. Each vertex in $\text{pn}(v, S)$ is called a *private neighbor* of v . The *external private neighborhood* $\text{epn}(v, S)$ of v with respect to S consists of the private neighbors of v in $V \setminus S$, while the *internal private neighborhood* $\text{ipn}(v, S)$ of v with respect to S consists of the private neighbors of v in S . Thus, $\text{epn}(v, S) = \text{pn}(v, S) \cap (V \setminus S)$ and $\text{ipn}(v, S) = \text{pn}(v, S) \cap S$, while $\text{pn}(v, S) = \text{epn}(v, S) \cup \text{ipn}(v, S)$. The following property of minimal total dominating sets is established in [5].

Observation 1.1 [5] *Let S be a TD-set of a graph G with no isolated vertex. Then, S is a minimal TDS of G if and only if for each $v \in S$, $\text{epn}(v, S) \neq \emptyset$ or $\text{pn}(v, S) = \text{ipn}(v, S) \neq \emptyset$.*

A *packing* in a graph G is a set of vertices whose closed neighborhoods are pairwise disjoint. Thus, if S is a packing in G , then $N_G[u] \cap N_G[v] = \emptyset$ for all pairs $u, v \in S$; equivalently, the vertices in S are pairwise at distance at least 3 apart in G . A *perfect packing* in G is a packing whose closed neighborhoods partition $V(G)$.

We use the standard notation $[k] = \{1, 2, \dots, k\}$.

2 Domination Versus Total Domination

We shall need the following well known lower bound on the domination number of a graph (see, for example, [9]). For completeness, we provide the elementary proof of this result.

Lemma 2.1 [9] *If G is a graph of order n and $\Delta = \Delta(G)$, then $\gamma(G) \geq n/(\Delta + 1)$, with equality if and only if every minimum dominating set in G is a perfect packing that contains only vertices of degree Δ .*

Proof If D is an arbitrary minimum dominating set of G , then

$$n - |D| = |V(G) \setminus D| \leq \left| \bigcup_{v \in D} N_G(v) \right| \leq \sum_{v \in D} d_G(v) \leq |D| \cdot \Delta,$$

and so $\gamma(G) = |D| \geq n/(\Delta + 1)$. Further, if $\gamma(G) = n/(\Delta + 1)$, then we must have equality throughout the above inequality chain, implying that $d_G(v) = \Delta$ for all $v \in D$ and that the closed neighborhoods, $N_G[v]$, of the vertices $v \in D$ are pairwise disjoint. Thus, D is a perfect packing in G that contains only vertices of degree Δ . Conversely, if D is a perfect packing in G that contains only vertices of degree Δ , then every dominating set D of G contains at least one vertex from every closed neighborhood, $N_G[u]$, for all $u \in D$, and so $\gamma(G) \geq |D|$. However, the set D is a dominating set of G , and so $\gamma(G) \leq |D|$. Consequently, $\gamma(G) = |D| = n/(\Delta + 1)$. \square

The following relationship between the domination and total domination numbers of a graph with no isolated vertex was first observed by Bollobás and Cockayne [2].

Theorem 2.2 [2] *For every graph G with no isolated vertex, $\gamma_t(G) \leq 2\gamma(G)$.*

A constructive characterization of trees T satisfying $\gamma_t(T) = 2\gamma(T)$ is given in [12]. However, it remains an open problem to characterize general graphs G satisfying $\gamma_t(G) = 2\gamma(G)$. We shall need the following lemma.

Lemma 2.3 *For $k \geq 2$, let G be a k -regular graph of order n . If $\gamma_t(G) = 2\gamma(G)$, then $\gamma(G) = n/(k + 1)$.*

Proof Let G be a k -regular graph of order n satisfying $\gamma_t(G) = 2\gamma(G)$. Let D be a minimum dominating set of G , and so $|D| = \gamma(G)$. For each vertex $v \in D$, let v' be an arbitrary neighbor of v , and let

$$X = \bigcup_{v \in D} \{v, v'\}.$$

Then, X is a TD-set of G , and so $\gamma_t(G) \leq |X|$. If v and w are distinct vertices of D , and $v' = w$ or $v' = w'$, then $\gamma_t(G) \leq |X| < 2|D| = 2\gamma(G)$, a contradiction. Therefore, for every pair v and w of vertices in D , the vertices v and w are not adjacent and do not have a neighbor in common; that is, $N_G[v] \cap N_G[w] = \emptyset$. Thus, D is a packing in G . Since $|N_G[v]| = d_G(v) + 1 = k + 1$ for every vertex $v \in D$, and since D is a dominating set of G , the sets $N_G[v]$ where $v \in D$ form a partition $V(G)$, implying that $n = |D|(k + 1) = \gamma(G)(k + 1)$. \square

We wish to determine the connected, k -regular graphs achieving equality in the upper bound of Theorem 2.2 for small values of k . If $k = 1$, then $G = K_2$ and $\gamma_t(G) = 2$ and $\gamma(G) = 1$, and so $\gamma_t(G)/\gamma(G) = 2$. Hence, it is only of interest to consider the cases when $k \geq 2$.

2.1 2-Regular Graphs

We first consider the case when G is a k -regular graph with $k = 2$. In this case $G \cong C_n$ and $n \geq 3$. It is well known (see, for example, [9, 11, 15]) that $\gamma_t(C_n) = \lfloor n/2 \rfloor + \lceil n/4 \rceil - \lfloor n/4 \rfloor$ and $\gamma(C_n) = \lceil n/3 \rceil$. As an immediate consequence of these results, we note that

$$\frac{\gamma_t(G)}{\gamma(G)} = \frac{3}{2} + \Upsilon(n),$$

where

$$\Upsilon(n) = \begin{cases} 0 & \text{if } n \equiv 0, 5, 10, 11 \pmod{12} \\ -\frac{3}{2(n+2)} & \text{if } n \equiv 1, 7 \pmod{12} \\ \frac{3}{2(n+1)} & \text{if } n \equiv 2 \pmod{12} \\ \frac{3}{2n} & \text{if } n \equiv 3, 9 \pmod{12} \\ -\frac{3}{n+2} & \text{if } n \equiv 4 \pmod{12} \\ \frac{3}{n} & \text{if } n \equiv 6 \pmod{12}. \\ -\frac{3}{2(n+1)} & \text{if } n \equiv 8 \pmod{12} \end{cases}$$

Thus, we have the following observations.

Observation 2.4 *If G is a connected 2-regular graph of order $n \geq 3$, then*

$$\frac{\gamma_t(G)}{\gamma(G)} = \begin{cases} \frac{3}{2} + \frac{3}{2(n+1)} & \text{if } n \equiv 2 \pmod{12} \\ \frac{3}{2} + \frac{3}{2n} & \text{if } n \equiv 3, 9 \pmod{12} \\ \frac{3}{2} + \frac{3}{n} & \text{if } n \equiv 6 \pmod{12} \end{cases}$$

and $\frac{\gamma_t(G)}{\gamma(G)} \leq \frac{3}{2}$ otherwise.




Fig. 1 Cubic graphs $G_4 \in \mathcal{G}$ and $H_4 \in \mathcal{H}$

Observation 2.5 *If G is a connected 2-regular graph, then*

$$\frac{\gamma_t(G)}{\gamma(G)} \leq 2 \text{ with equality if and only if } G \in \{C_3, C_6\}.$$

2.2 3-Regular Graphs

We next consider the case when $k \geq 3$. For this purpose, we shall need a result on the total domination number of a cubic graph. For $\ell \geq 1$, let G_ℓ be the graph constructed in [8] as follows. (This construction is also described in Chapter 5, p. 44, of [15].) Consider two copies of the path $P_{2\ell}$ with respective vertex sequences $a_1b_1a_2b_2 \cdots a_\ell b_\ell$ and $c_1d_1c_2d_2 \cdots c_\ell d_\ell$. For each $i \in [\ell]$, join a_i to d_i and b_i to c_i . To complete the construction of the graph G_ℓ join a_1 to c_1 and b_ℓ to d_ℓ . Let $\mathcal{G} = \{G_\ell \mid \ell \geq 1\}$. For $\ell \geq 2$, let H_ℓ be obtained from G_ℓ by deleting the two edges a_1c_1 and $b_\ell d_\ell$ and adding the two edges a_1b_ℓ and c_1d_ℓ . Let $\mathcal{H} = \{H_\ell \mid \ell \geq 2\}$. We note that G_ℓ and H_ℓ are cubic graphs of order 4ℓ . Further, we note that $G_1 \cong K_4$. The graphs $G_4 \in \mathcal{G}$ and $H_4 \in \mathcal{H}$, for example, are illustrated in Fig. 1.

Let $\mathcal{H}_{\text{even}}$ be the subfamily of \mathcal{H} consisting of all graphs H_ℓ where ℓ is even; that is, $\mathcal{H}_{\text{even}} = \{H_\ell \mid \ell \geq 2 \text{ is even}\}$.

The following upper bound on the total domination number of a graph with minimum degree at least 3 was established independently by many authors (see, [1, 4, 19]).

Theorem 2.6 [1, 4, 19] *If G is a graph of order n with $\delta(G) \geq 3$, then $\gamma_t(G) \leq \frac{n}{2}$.*

The connected graphs that achieve equality in the bound of Theorem 2.6 were characterized in [13].

Theorem 2.7 [13] *If G is a connected graph of order n with $\delta(G) \geq 3$, then $\gamma_t(G) \leq \frac{n}{2}$, with equality if and only if $G \in \mathcal{G} \cup \mathcal{H}$ or G is the generalized Petersen graph shown in Fig. 2.*




Fig. 2 The generalized Petersen graph

We are now in a position to characterize the connected, cubic graphs G satisfying $\gamma_t(G)/\gamma(G) = 2$.

Theorem 2.8 *If G is a connected cubic graph, then $\gamma_t(G)/\gamma(G) \leq 2$ with equality if and only if the following holds.*

- (a) $G \in \mathcal{G} \cup \mathcal{H}_{\text{even}}$.
- (b) G is the generalized Petersen graph.

Proof Let G be a connected, cubic graph of order n . By Theorem 2.2, $\gamma_t(G)/\gamma(G) \leq 2$. Hence it suffices for us to prove that $\gamma_t(G)/\gamma(G) = 2$ if and only if (a) or (b) in the statement of the theorem hold. We first prove the sufficiency. Suppose that (a) or (b) in the statement of the theorem holds. If G is the generalized Petersen graph, then $\gamma(G) = 4$ and $\gamma_t(G) = 8$, and so $\gamma_t(G)/\gamma(G) = 2$. Suppose that $G \in \mathcal{G} \cup \mathcal{H}_{\text{even}}$. If $G = G_\ell$ for $\ell \geq 2$ even or if $G = H_\ell$ for $\ell \geq 2$ even, then we let

$$D = \bigcup_{i=1}^{\ell/2} \{b_{2i-1}, c_{2i}\}.$$

If $G = G_\ell$ for $\ell \geq 1$ odd, then we let

$$D = \{b_\ell\} \cup \left(\bigcup_{i=1}^{(\ell-1)/2} \{b_{2i-1}, c_{2i}\} \right).$$

In both cases, D is a dominating set of G , implying that $\gamma(G) \leq |D| = \ell = n/4$. By Lemma 2.1, $\gamma(G) \geq n/4$. Consequently, $\gamma(G) = n/4$. By Theorem 2.7, $\gamma_t(G) = n/2$. Thus, $\gamma_t(G)/\gamma(G) = 2$. This proves the sufficiency.

Next, we prove the necessity. Suppose that G is a connected, cubic graph satisfying $\gamma_t(G)/\gamma(G) = 2$. By Lemma 2.3, $\gamma(G) = n/4$, implying that $\gamma_t(G) = n/2$. By Theorem 2.7, $G \in \mathcal{G} \cup \mathcal{H}$ or G is the generalized Petersen graph.

It remains for us to show that if $G \in \mathcal{H}$, then $G \in \mathcal{H}_{\text{even}}$. Suppose, to the contrary, that $G = H_\ell$ for some $\ell \geq 3$, where ℓ is odd. For $i \in [\ell]$, let $V_i = \{a_i, b_i, c_i, d_i\}$. Let D be a minimum dominating set in G . Since $n = 4\ell$ and $\gamma(G) = n/4$, we note that $|D| = \gamma(G) = \ell$ and, by Lemma 2.1, that the set D is a packing in G . In order to dominate the vertex a_1 , the set D contains exactly one of the vertices a_1, b_1, b_ℓ, d_1 .

By symmetry, renaming vertices if necessary, we may assume that either $a_1 \in D$ or $b_1 \in D$.

Suppose that $b_1 \in D$. Since D is a packing, this implies that $D \cap V_1 = \{b_1\}$. Further, $D \cap \{b_\ell, d_\ell\} = \emptyset$. In order to dominate the vertices b_ℓ and d_ℓ , we note that either $a_\ell \in D$ or $c_\ell \in D$; that is, $D \cap \{a_\ell, c_\ell\} \neq \emptyset$. In order to dominate the vertex d_1 , we note that $c_2 \in D$. Since D is a packing, this in turn implies that $D \cap V_2 = \{c_2\}$ and $D \cap \{a_3, c_3\} = \emptyset$. If $\ell = 3$, we contradict our earlier observation that $D \cap \{a_\ell, c_\ell\} \neq \emptyset$. Therefore, $\ell \geq 5$. In order to dominate the vertices a_3 and c_3 , we note that either $b_3 \in D$ or $d_3 \in D$; that is, $D \cap \{b_3, d_3\} \neq \emptyset$. If $b_3 \in D$, then $c_4 \in D$. If $d_3 \in D$, then $a_4 \in D$. In both cases, since D is a packing, we deduce that $D \cap \{a_5, c_5\} = \emptyset$. Continuing in this way, it therefore holds that $D \cap \{a_i, c_i\} = \emptyset$ for all odd $i \geq 1$ where $i \in [\ell]$. In particular, noting that ℓ is odd, $D \cap \{a_\ell, c_\ell\} = \emptyset$. This contradicts our earlier observation that $D \cap \{a_\ell, c_\ell\} \neq \emptyset$.

Suppose that $a_1 \in D$. Since D is a packing, this implies that $D \cap V_1 = \{a_1\}$ and $D \cap \{a_2, c_2\} = \emptyset$, for otherwise a_1 would be within distance 2 from some other vertex of D . Further, $D \cap \{a_\ell, c_\ell\} = \emptyset$ (and $b_\ell \notin D$). In order to dominate the vertices a_2 and c_2 , we note that either $b_2 \in D$ or $d_2 \in D$; that is, $D \cap \{b_2, d_2\} \neq \emptyset$. If $b_2 \in D$, then $c_3 \in D$. If $d_2 \in D$, then $a_3 \in D$. In both cases, $D \cap \{a_3, c_3\} \neq \emptyset$. If $\ell = 3$, we contradict our earlier observation that $D \cap \{a_\ell, c_\ell\} = \emptyset$. Therefore, $\ell \geq 5$. Since D is a packing and $D \cap \{a_3, c_3\} \neq \emptyset$, we deduce that $D \cap \{a_4, c_4\} = \emptyset$. In order to dominate the vertices a_4 and c_4 , we note that either $b_4 \in D$ or $d_4 \in D$; that is, $D \cap \{b_4, d_4\} \neq \emptyset$. If $b_4 \in D$, then $c_5 \in D$. If $d_4 \in D$, then $a_5 \in D$. In both cases, $D \cap \{a_5, c_5\} \neq \emptyset$. Continuing in this way, it therefore holds that $D \cap \{a_i, c_i\} \neq \emptyset$ for all odd $i \geq 1$ where $i \in [\ell]$. In particular, noting that ℓ is odd, $D \cap \{a_\ell, c_\ell\} \neq \emptyset$. This contradicts our earlier observation that $D \cap \{a_\ell, c_\ell\} = \emptyset$.

Since both cases $a_1 \in D$ and $b_1 \in D$ produce a contradiction, we deduce that if $G \in \mathcal{H}$, then $G \in \mathcal{H}_{\text{even}}$. This proves the necessity. \square

3 Upper Domination Versus Upper Total Domination

Dorbec et al. [7] established the following relationship between the upper domination and upper total domination numbers of a graph with no isolated vertex.

Theorem 3.1 [7] *For every graph G with no isolated vertex, $\Gamma_t(G) \leq 2\Gamma(G)$.*

As remarked in [7], it remains an open problem to characterize graphs G satisfying $\Gamma_t(G) = 2\Gamma(G)$. We wish to determine the connected, k -regular graphs G achieving equality in the upper bound of Theorem 3.1 for small values of k . If $k = 1$, then $G = K_2$ and $\Gamma(G) = 1$ and $\Gamma_t(G) = 2$. Hence, it is only of interest to consider the cases when $k \geq 2$.

3.1 2-Regular Graphs

We first consider the case when $k = 2$. The following result determines the upper total domination number of a cycle.

Proposition 3.2 For $n \geq 3$,

$$\Gamma_t(C_n) = \begin{cases} 2\lfloor \frac{n}{3} \rfloor + 1 & \text{if } n \equiv 5 \pmod{6} \\ 2\lfloor \frac{n}{3} \rfloor & \text{otherwise.} \end{cases}$$

Proof We proceed by induction on $n \geq 3$. The result is straightforward to check (or use a computer) for small $n \leq 9$. Hence we may assume that $n \geq 10$. Let $G \cong C_n$ be the cycle $v_1 v_2 \cdots v_n v_1$. We show firstly that $\Gamma_t(C_n) \geq 2\lfloor \frac{n}{3} \rfloor + 1$ if $n \equiv 5 \pmod{6}$ and $\Gamma_t(C_n) \geq 2\lfloor \frac{n}{3} \rfloor$, otherwise. If $n \equiv 5 \pmod{6}$, then the set

$$A = \{v_{n-3}, v_{n-2}, v_{n-1}\} \cup \left(\bigcup_{i=0}^{\frac{n-11}{6}} \{v_{6i+2}, v_{6i+3}, v_{6i+4}, v_{6i+5}\} \right)$$

is a minimal TD-set of G , and so $\Gamma_t(G) \geq |A| = 2\lfloor \frac{n}{3} \rfloor + 1$. If $n \equiv 0, 1 \pmod{3}$, let

$$B = \bigcup_{i=0}^{\lfloor n/3 \rfloor - 1} \{v_{3i+1}, v_{3i+2}\}$$

while if $n \equiv 2 \pmod{6}$, let

$$B = \left(\bigcup_{i=0}^{\lfloor n/3 \rfloor - 3} \{v_{3i+1}, v_{3i+2}\} \right) \cup \{v_{n-6}, v_{n-5}, v_{n-2}, v_{n-1}\}.$$

In both cases, the set B is a minimal TD-set of G and $|B| = 2\lfloor \frac{n}{3} \rfloor$, and so $\Gamma_t(G) \geq 2\lfloor \frac{n}{3} \rfloor$.

We show next that $\Gamma_t(C_n) \leq 2\lfloor \frac{n}{3} \rfloor + 1$ if $n \equiv 5 \pmod{6}$ and $\Gamma_t(C_n) \leq 2\lfloor \frac{n}{3} \rfloor$, otherwise. Let S be a $\Gamma_t(G)$ -set. By the minimality of S , no five consecutive vertices of G all belong to S .

Suppose that S contains four consecutive vertices of G . Renaming vertices if necessary, we may assume that $\{v_3, v_4, v_5, v_6\} \subseteq S$. Since no five consecutive vertices all belong to S , we note that $v_2 \notin S$ and $v_7 \notin S$. If $v_1 \in S$, then $S \setminus \{v_3\}$ is a TD-set of G , contradicting the minimality of S . Hence, $v_1 \notin S$. If $v_8 \in S$, then $S \setminus \{v_6\}$ is a TD-set of G , contradicting the minimality of S . Hence, $v_8 \notin S$. This implies that $\{v_9, v_{10}\} \subset S$. We now consider the cycle G' obtained from G by deleting the six vertices v_3, v_4, \dots, v_8 and adding the edge $v_2 v_9$. Thus, $G' \cong C_{n'}$ where $n' = n - 6$. Let $S' = S \setminus \{v_3, v_4, v_5, v_6\}$. Since S is a minimal TD-set of G , the set S' is a minimal TD-set of G' , implying that $|S'| \leq \Gamma_t(G')$. If $n \equiv 5 \pmod{6}$, then $n \geq 11$ and $n' \equiv 5 \pmod{6}$, and by the inductive hypothesis, $\Gamma_t(G) = |S| = |S'| + 4 \leq \Gamma_t(G') + 4 = (2\lfloor \frac{n'}{3} \rfloor + 1) + 4 = (2\lfloor \frac{n-6}{3} \rfloor + 1) + 4 = 2\lfloor \frac{n}{3} \rfloor + 1$. If $n \not\equiv 5 \pmod{6}$, then $n' \not\equiv 5 \pmod{6}$. By the inductive hypothesis, $\Gamma_t(G) = |S| = |S'| + 4 \leq \Gamma_t(G') + 4 = (2\lfloor \frac{n'}{3} \rfloor) + 4 = 2\lfloor \frac{n}{3} \rfloor$. Hence, if S contains four consecutive vertices of G , then $\Gamma_t(C_n) \leq 2\lfloor \frac{n}{3} \rfloor + 1$ if $n \equiv 5 \pmod{6}$ and

$\Gamma_t(C_n) \leq 2\lfloor \frac{n}{3} \rfloor$, otherwise. Hence, we may assume that S contains no four consecutive vertices, for otherwise the desired result follows.

Suppose that S contains three consecutive vertices of G . Renaming vertices if necessary, we may assume that $\{v_3, v_4, v_5\} \subseteq S$. Since no four consecutive vertices all belong to S , we note that $v_2 \notin S$ and $v_6 \notin S$. By the minimality of S , $v_1 \notin S$ and $v_7 \notin S$. This implies that $\{v_n, v_{n-1}\} \subset S$ and $\{v_8, v_9\} \subset S$. We now consider the cycle G' obtained from G by deleting the five vertices v_3, v_4, \dots, v_7 and adding the edge v_2v_8 . Thus, $G' \cong C_{n'}$ where $n' = n - 5$. Let $S' = S \setminus \{v_3, v_4, v_5\}$. Since S is a minimal TD-set of G , the set S' is a minimal TD-set of G' , implying that $\Gamma_t(G) = |S| = |S'| + 3 \leq \Gamma_t(G') + 3$.

If $n \equiv 5 \pmod{6}$, then $n' \equiv 0 \pmod{6}$, and by the inductive hypothesis, $\Gamma_t(G) \leq \Gamma_t(G') + 3 = 2\lfloor \frac{n'}{3} \rfloor + 3 = 2\lfloor \frac{n-5}{3} \rfloor + 3 = 2\lfloor \frac{n}{3} \rfloor + 1$. If $n \equiv 4 \pmod{6}$, then $n' \equiv 5 \pmod{6}$, and by the inductive hypothesis, $\Gamma_t(G) \leq \Gamma_t(G') + 3 = 2\lfloor \frac{n'}{3} \rfloor + 4 = 2\lfloor \frac{n-5}{3} \rfloor + 4 = 2\lfloor \frac{n}{3} \rfloor$. If $n \equiv 0, 1, 3 \pmod{6}$, then by the inductive hypothesis, $\Gamma_t(G) \leq \Gamma_t(G') + 3 = 2\lfloor \frac{n'}{3} \rfloor + 3 = 2\lfloor \frac{n-5}{3} \rfloor + 3 \leq 2\lfloor \frac{n}{3} \rfloor$.

Suppose that $n \equiv 2 \pmod{6}$. We show that $\Gamma_t(G) \leq 2\lfloor \frac{n}{3} \rfloor$. Suppose that $v_{10} \in S$. In this case, $v_{11} \notin S$ and $v_{12} \notin S$. This implies that $\{v_{13}, v_{14}\} \subset S$. We now consider the cycle G'' obtained from G by deleting the ten vertices v_3, v_4, \dots, v_{12} and adding the edge v_2v_{13} . Thus, $G'' \cong C_{n''}$ where $n'' = n - 10$. We note that $n'' \equiv 4 \pmod{6}$. Let $S'' = S \setminus \{v_3, v_4, v_5, v_8, v_9, v_{10}\}$. Since S is a minimal TD-set of G , the set S'' is a minimal TD-set of G'' , implying by the inductive hypothesis that $\Gamma_t(G) = |S| = |S''| + 6 \leq \Gamma_t(G'') + 6 \leq 2\lfloor \frac{n''}{3} \rfloor + 6 = 2\lfloor \frac{n-10}{3} \rfloor + 6 \leq 2\lfloor \frac{n}{3} \rfloor$. Hence, we may assume that $v_{10} \notin S$, for otherwise $\Gamma_t(G) \leq 2\lfloor \frac{n}{3} \rfloor$ as desired. If $v_{11} \notin S$, then $v_{12} \in S$ and $v_{13} \in S$. In this case, we consider the cycle G^* obtained from G by deleting the four vertices v_6, v_7, v_8, v_9 and adding the edge v_5v_{10} . Thus, $G^* \cong C_{n^*}$ where $n^* = n - 4$. We note that $n^* \equiv 4 \pmod{6}$. Let $S^* = S \setminus \{v_8, v_9\}$. Since S is a minimal TD-set of G , the set S^* is a minimal TD-set of G^* , implying by the inductive hypothesis that $\Gamma_t(G) = |S| = |S^*| + 2 \leq \Gamma_t(G^*) + 2 \leq 2\lfloor \frac{n^*}{3} \rfloor + 2 = 2\lfloor \frac{n-4}{3} \rfloor + 2 \leq 2\lfloor \frac{n}{3} \rfloor$. Hence, we may assume that $v_{11} \in S$, for otherwise the desired result follows, implying that $v_{12} \in S$. Using analogous and symmetrical arguments, we may assume that $v_{n-2} \notin S$ and $\{v_{n-3}, v_{n-4}\} \subset S$, for otherwise the desired upper bound follows. If $n = 14$, then $v_{n-2} = v_{12}$, contradicting the fact that with our current assumptions $v_{12} \in S$ and $v_{n-2} \notin S$. Hence, we note that $n \geq 20$. The set $(S \setminus \{v_3, v_4, v_5\}) \cup \{v_2, v_3, v_5, v_6\}$ is a minimal TD-set of G of cardinality $|S| + 1 = \Gamma_t(G) + 1$, a contradiction.

Hence, if S contains three consecutive vertices, then we have shown that $\Gamma_t(C_n) \leq 2\lfloor \frac{n}{3} \rfloor + 1$ if $n \equiv 5 \pmod{6}$ and $\Gamma_t(C_n) \leq 2\lfloor \frac{n}{3} \rfloor$, otherwise. Hence, we may assume that S contains no three consecutive vertices, for otherwise the desired result follows. With this assumption, $\Gamma_t(C_n) \leq 2\lfloor \frac{n}{3} \rfloor$ for all $n \geq 10$. This completes the proof of the proposition. □

For $n \geq 3$, we observe that $\Gamma(C_n) = \alpha(C_n) = \lfloor n/2 \rfloor$, where $\alpha(G)$ is the size of a maximum independent vertex set of G . We state this result formally as follows.

Observation 3.3 For $n \geq 3$, $\Gamma(C_n) = \lfloor n/2 \rfloor$.

As an immediate consequence of Proposition 3.2 and Observation 3.3, we note that if $G \cong C_n$ and $n \geq 3$, then

$$\frac{\Gamma_t(G)}{\Gamma(G)} = \frac{4}{3} + \Phi(n),$$

where

$$\Phi(n) = \begin{cases} 0 & \text{if } n \equiv 0, 1 \pmod{6} \\ -\frac{8}{3n} & \text{if } n \equiv 2 \pmod{6} \\ \frac{4}{3(n-1)} & \text{if } n \equiv 3 \pmod{6} \\ -\frac{4}{3n} & \text{if } n \equiv 4 \pmod{6} \\ \frac{2}{3(n-1)} & \text{if } n \equiv 5 \pmod{6}. \end{cases}$$

This yields the following results.

Corollary 3.4 If G is a connected 2-regular graph of order $n \geq 3$, then

$$\frac{\Gamma_t(G)}{\Gamma(G)} \leq \frac{4}{3} + \frac{4}{3(n-1)}.$$

Corollary 3.5 If G is a connected 2-regular graph, then $\frac{\Gamma_t(G)}{\Gamma(G)} \leq 2$, with equality if and only if $G \cong C_3$.

3.2 3-Regular Graphs

We next consider the case when $k = 3$. In order to determine the connected 3-regular (cubic) graphs G satisfying $\Gamma_t(G)/\Gamma(G) = 2$, we analyse the proof of the upper bound of Theorem 3.1 to obtain the structure of these extremal graphs. For this purpose, we shall need the following key lemma in [7]. For completeness, we provide the proof of this lemma as given in [7].

Lemma 3.6 [7] Every $\Gamma_t(G)$ -set contains as a subset a minimal dominating set S such that $|S| \geq \frac{1}{2}\Gamma_t(G)$ and $|\text{epn}(v, S)| \geq 1$ for each $v \in S$.

Proof Let D be an arbitrary $\Gamma_t(G)$ -set, and let

$$\begin{aligned} A &= \{v \in D \mid \text{epn}(v, D) \neq \emptyset\}, \\ B &= \{v \in D \setminus A \mid d_A(v) \geq 1\}, \text{ and} \\ C &= D \setminus (A \cup B). \end{aligned}$$

Then, $D = A \cup B \cup C$. As shown in [7], if $C \neq \emptyset$, then $G[C] = \frac{|C|}{2}K_2$ and each vertex of C has degree 1 in $G[D]$. Following the notation introduced in [7], we call

two adjacent vertices in $G[C]$ partners in C and we let (X, Y) be the partite sets in the graph $G[C]$. Thus, each vertex in X (resp., in Y) is adjacent in $G[D]$ only to its partner in Y (resp., in X). For each $x \in X$, let y_x be the partner of x in C . As shown in [7], for each vertex $v \in B$, we have $\text{pn}(v, D) = \text{ipn}(v, D) \subseteq A$, implying that

$$|A| \geq \sum_{v \in B} |\text{ipn}(v, D)| \geq |B|. \tag{1}$$

Following the notation introduced in the proof of Theorem 3.1 given in [7], we let $U = V(G) \setminus (D \cup N(A) \cup N(X))$ be the set of vertices in $V(G) \setminus D$ not dominated by A or X in G . Since D is a TD-set of G , the set U is dominated by $B \cup Y$. Let B_Y be a minimum subset of $B \cup Y$ that dominates U . Thus for each $v \in B_Y$, we have that $|\text{epn}(v, B_Y) \cap U| \geq 1$.

As observed in [7], the set $S = A \cup B_Y \cup X$ is a dominating set of G , although not necessarily minimal. For each vertex $x \in X$ we systematically delete x from S if $\text{epn}(x, S) = \emptyset$ at each stage in the resulting set S , and let X^* be the resulting subset of vertices of X that belong to the set S upon the completion of this process. Thus, $S = A \cup B_Y \cup X^*$ and, by construction, $|\text{epn}(v, S)| \geq 1$ for each $v \in S$. As observed in [7], the set S is a minimal dominating set of G , and so $\Gamma(G) \geq |S|$. For every vertex $x \in X$, the set S contains at least one of x and its partner in C , noting that if the partner $y_x \in Y$ of x in C is not in S , then $y_x \in \text{epn}(x, S)$, and so x is not deleted from S . Thus, $|S \cap C| \geq |X| = \frac{1}{2}|C|$. As shown earlier, $|A| \geq |B|$. Hence,

$$\begin{aligned} \Gamma(G) &\geq |S| \\ &= |S \cap A| + |S \cap B| + |S \cap C| \\ &\geq |A| + |S \cap C| \\ &\geq \frac{1}{2}(2|A| + |C|) \\ &\geq \frac{1}{2}(|A| + |B| + |C|) \\ &= \frac{1}{2}|D| \\ &= \frac{1}{2}\Gamma_t(G), \end{aligned} \tag{2}$$

or equivalently, $\Gamma_t(G) \leq 2\Gamma(G)$. As observed earlier, $S \subset D$ and S is a minimal dominating set of G . Further, $|S| \geq \frac{1}{2}\Gamma_t(G)$ and $|\text{epn}(v, S)| \geq 1$ for each $v \in S$. \square

We note that Theorem 3.1 follows as an immediate consequence of Lemma 3.6. We proceed further with the following lemma which gives structural properties of graphs G that achieve equality in the bound of Theorem 3.1. We remark that the following properties of graphs achieving equality in the bound of Theorem 3.1 is implicitly implied in [7], albeit without proof. We provide here a complete proof since we will need these properties in order to characterize the cubic graphs achieving equality in the bound of Theorem 3.1.

Lemma 3.7 *If G is a graph with no isolated vertex satisfying $\Gamma_t(G) = 2\Gamma(G)$, then the following holds.*

- (a) $\Gamma(G) = \alpha(G)$.
- (b) Every $\Gamma_t(G)$ -set induces a subgraph that consists of disjoint copies of K_2 .
- (c) Every vertex that does not belong to an arbitrary $\Gamma_t(G)$ -set is contained in a common triangle with two vertices of the set.

Proof Suppose that G is a graph with no isolated vertex satisfying $\Gamma(G) = \frac{1}{2}\Gamma_t(G)$. Let D be an arbitrary $\Gamma_t(G)$ -set. Following exactly the notation introduced in the proof of Lemma 3.6, we must have equality throughout inequality chain (2) in this proof. Hence, $S = A \cup B_Y \cup X^*$ is a $\Gamma(G)$ -set, $|A| = |B|$, $S \cap B = \emptyset$ and $|S \cap C| = |C|/2$. Since $|A| = |B|$, we have by inequality chain (1) that $|A| = \sum_{v \in B} |\text{pn}(v, S)| = |B|$. Thus, $|\text{pn}(v, S)| = 1$ for every $v \in B$ and every vertex of A is the (internal) private neighbor of some vertex of B . In particular, $d_S(v) = 1$ for every $v \in A$ and every vertex in B is adjacent to a unique vertex of A , namely to its (internal) private neighbor in S . Hence, A is an independent set and the subgraph $G[A \cup B]$ induced by $A \cup B$ contains a perfect matching. (Note that it is possible that there may be some edges joining vertices of B .)

As observed in the proof of Lemma 3.6, for every vertex $x \in X$, the set S contains at least one of x and its partner in C , and so $|S \cap C| \geq |C|/2$. However, $|S \cap C| = |C|/2$, implying that the set S contains exactly one of x and its partner y_x in C . Renaming vertices of C if necessary, we may assume that $S \cap C = X$. Since A is an independent set and there is no edge between A and C , the set $S = A \cup X$ is an independent set, and so $\alpha(G) \geq |S| = |A| + |C|/2$. Since every maximum independent set in a graph is a minimal dominating set in the graph, we note that $|S| = \Gamma(G) \geq \alpha(G) = |S|$. Consequently, the set $S = A \cup X$ is a maximum independent set in G , and so $\Gamma(G) = \alpha(G) = |S|$. This proves Part (a).

Suppose that $A \neq \emptyset$ and consider an arbitrary vertex $u \in A$. Let v be the neighbor of u in $G[D]$, and so $v \in B$ and $\text{pn}(v, D) = \{u\}$. Further, let $w \in \text{epn}(u, S)$. The vertex w exists since $S \subseteq D$ and $u \in A$ implies that u has an external private neighbor with respect to D . Let $|C| = 2c$ and recall that $G[C] = cK_2$ and each vertex of C has degree 1 in $G[D]$. In particular, we note that v is adjacent to no vertex of $A \cup C$ different from u . Thus, the set $(S \setminus \{u\}) \cup \{v, w\}$ is an independent set in G , implying that $\alpha(G) \geq |S| + 1$, a contradiction. Hence, $A = \emptyset$, which in turn implies that $B = \emptyset$. Thus, $D = C$ and $G[D] = G[C] = cK_2$, where $|C| = 2c$. Since D is an arbitrary $\Gamma_t(G)$ -set, we deduce that every $\Gamma_t(G)$ -set induces a subgraph isomorphic to cK_2 . This proves Part (b). By our earlier observations, $\Gamma(G) = \alpha(G) = c$.

To prove Part (c), let v be an arbitrary vertex in $V(G) \setminus D$. As observed above, $D = C$ and $G[C] = cK_2$. We show that v is contained in a common triangle with two vertices of C . Suppose, to the contrary, that v does not belong to a common triangle with two vertices of C . If v is adjacent to both x and its partner y_x for some vertex $x \in C$, then $vx y_x v$ is a common triangle containing v and two vertices of C , a contradiction. Hence, v is adjacent to at most one of x and its partner y_x for every vertex $x \in C$. Let C' be a subset of C constructed as follows. Initially, let $C' = \emptyset$. For each pair x and y_x where $x \in C$ (and $x y_x$ is an edge), select one vertex that is not adjacent to v and add it to the set C' . Thus, $|C'| = c$ and C' is an independent set. By

construction, no vertex of C' is adjacent to v . Hence, $C' \cup \{v\}$ is an independent set on G , and so $\alpha(G) \geq |C'| + 1 = c + 1 > \alpha(G)$, a contradiction. Therefore, the vertex v is adjacent to both x and its partner y_x in C for at least one vertex $x \in C$. \square

We are now in a position to characterize the connected cubic graphs achieving equality in Theorem 3.1.

Theorem 3.8 *If G is a connected 3-regular graph, then*

$$\frac{\Gamma_t(G)}{\Gamma(G)} \leq 2 \text{ with equality if and only if } G \cong K_4.$$

Proof The inequality follows by Theorem 3.1. Therefore in what follows we assume $\frac{\Gamma_t(G)}{\Gamma(G)} = 2$. We adopt the notation introduced in the proof of Lemmas 3.6 and 3.7. Let D be an arbitrary $\Gamma_t(G)$ -set. By Lemma 3.7(b), $G[D] \cong cK_2$ for some integer $c \geq 1$. By Lemma 3.7(a), $\Gamma(G) = \alpha(G) = c$. Let $G[D]$ consist of the c edges x_1y_1, \dots, x_cy_c . Let $\bar{D} = V(G) \setminus D$. By Lemma 3.7(c), each vertex in \bar{D} is contained in a common triangle with two vertices from the set D . Since G is a cubic graph, there is therefore a weak partition (where some of the sets in the partition may possibly be empty) of \bar{D} into sets V_1, \dots, V_c such that every vertex in V_i is adjacent to both x_i and y_i in G for all $i \in [c]$. Since G is a cubic graph and x_i and y_i are adjacent to each other and to every vertex in V_i , we note that $|V_i| \leq 2$ for all $i \in [c]$.

Suppose that $|V_i| \leq 1$ for all $i \in [c]$. Let $[D, \bar{D}]$ denote the set of edges joining D and \bar{D} . Since each vertex in D has two neighbors in \bar{D} and $|D| = 2c$, we note that $|[D, \bar{D}]| = 2|D| = 4c$. By supposition, $|V_i| \leq 1$ for all $i \in [c]$, implying that

$$|[D, \bar{D}]| \leq \sum_{i=1}^c 3|V_i| \leq 3c,$$

noting that $\bar{D} = (V_1, \dots, V_c)$ is a partition of \bar{D} and each vertex in \bar{D} has at most three neighbors in D . Thus, $4c = |[D, \bar{D}]| \leq 3c$, a contradiction. Therefore, $|V_i| = 2$ for at least one $i \in [c]$. Renaming the sets V_1, \dots, V_c if necessary, we may assume that $|V_1| = 2$. Let $V_1 = \{u_1, u_2\}$, and so both u_1 and u_2 are adjacent to x_1 and y_1 . In what follows, let

$$D_{\geq i} = \bigcup_{j=i}^c \{x_j, y_j\}$$

where $i \in [c]$. In particular, we note that $D = D_{\geq 1}$. Suppose that u_1 and u_2 are not adjacent. Thus, $c \geq 2$. If for every $i \in [c] \setminus \{1\}$, there exists a vertex in $\{x_i, y_i\}$ that is adjacent to neither u_1 nor u_2 , then there exists a subset $D'_{\geq 2}$ of $D_{\geq 2}$ such that $|D'_{\geq 2}| = c - 1$ and $D'_{\geq 2} \cup \{u_1, u_2\}$ is an independent set, implying that $\alpha(G) \geq |D'_{\geq 2}| + 2 = c + 1$, a contradiction. Hence for some $i \in [c] \setminus \{1\}$, the vertex u_1 is adjacent to one of x_i and y_i , say y_i , and the vertex u_2 is adjacent to x_i . Renaming the sets V_2, \dots, V_c if necessary, we may assume that $i = 2$. Thus, u_1 is adjacent to y_2 and u_2 is adjacent to x_2 .




Fig. 3 A subgraph of G

Let u_3 be the neighbor of x_2 different from u_2 and y_2 . We note that $u_3 \in V_i$ for some $i \in [c] \setminus \{1\}$. Suppose that $u_3 \in V_2$. Thus, the vertex u_3 is adjacent to both x_2 and y_2 , and is adjacent to at most one vertex from $D_{\geq 3}$. Hence there exists a subset $D'_{\geq 3}$ of $D_{\geq 3}$ such that $|D'_{\geq 3}| = c - 2$ and $D'_{\geq 3} \cup \{u_1, u_2, u_3\}$ is an independent set, implying that $\alpha(G) \geq |D'_{\geq 3}| + 3 = c + 1$, a contradiction. Therefore, $u_3 \notin V_2$, and so $u_3 \in V_i$ for some $i \in [c] \setminus \{1, 2\}$. Renaming the sets V_3, \dots, V_c if necessary, we may assume that $u_3 \in V_3$. Thus, u_3 is adjacent to both x_3 and y_3 .

Let u_4 be the neighbor of x_3 different from u_3 and y_3 . We note that $u_4 \in V_i$ for some $i \in [c] \setminus \{1, 2\}$. Suppose that $u_4 \in V_3$. Thus, the vertex u_4 is adjacent to both x_3 and y_3 , and is adjacent to at most one vertex from $D_{\geq 4}$. Hence there exists a subset $D'_{\geq 4}$ of $D_{\geq 4}$ such that $|D'_{\geq 4}| = c - 3$ and $D'_{\geq 4} \cup \{u_1, u_2, u_3, u_4\}$ is an independent set, implying that $\alpha(G) \geq |D'_{\geq 4}| + 4 = c + 1$, a contradiction. Therefore, $u_4 \notin V_3$, and so $u_4 \in V_i$ for some $i \in [c] \setminus \{1, 2, 3\}$. Renaming the sets V_4, \dots, V_c if necessary, we may assume that $u_4 \in V_4$. Thus, u_4 is adjacent to both x_4 and y_4 . The graph shown in Fig. 3 is therefore a subgraph of G , where the darkened vertices belong to the set D .

Continuing in this way, there exists a sequence of distinct vertices $u_3, u_4, \dots, u_{\ell+1}$ such that the vertex u_i is adjacent to x_{i-1} and $u_i \in V_i$ for $i \in [\ell] \setminus \{1, 2\}$ and $u_{\ell+1} \in V_\ell$. Let $U = \{u_1, u_2, \dots, u_{\ell+1}\}$. If $c = \ell$, then $\alpha(G) \geq |U| = c + 1$, a contradiction. Hence, $c \geq \ell + 1$. Noting that $u_{\ell+1}$ is adjacent to at most one vertex from $D_{\geq \ell+1}$, there exists a subset $D'_{\geq \ell+1}$ of $D_{\geq \ell+1}$ such that $|D'_{\geq \ell+1}| = c - \ell$ and $U \cup D'_{\geq \ell+1}$ is an independent set, implying that $\alpha(G) \geq |U| + |D'_{\geq \ell+1}| = c + 1$, a contradiction. We deduce, therefore, that u_1 and u_2 are adjacent, implying that $G \cong K_4$. Conversely, if $G \cong K_4$, then $\Gamma_t(G) = 2$ and $\Gamma(G) = 1$, and so $\Gamma_t(G)/\Gamma(G) = 2$. This completes the proof of Theorem 3.8. \square

3.3 Closing Questions and Problems

We pose the following two questions that we have yet to settle.

Question 1 *Is it true that if $G \cong K_4$ is a connected 3-regular graph, then*

$$\frac{\Gamma_t(G)}{\Gamma(G)} \leq \frac{3}{2}?$$

Question 2 For $k \geq 2$, if G is a connected k -regular graph, then is it true that

$$\frac{\Gamma_t(G)}{\Gamma(G)} \leq 2 \text{ with equality if and only if } G \cong K_{k+1}?$$

We have shown that Question 2 is true when $k = 2$ (see Corollary 3.5) and when $k = 3$ (see Theorem 3.8). However, we have yet to answer Question 2 when $k \geq 4$.

In summary, our focus in this paper is to characterize the connected cubic graphs G satisfying $\gamma_t(G)/\gamma(G) = 2$, and to characterize the connected cubic graphs G satisfying $\Gamma_t(G)/\Gamma(G) = 2$. These characterizations are presented in Theorem 2.8 and Theorem 3.8, respectively. It would be interesting to extend these characterizations to connected subcubic graphs. We remark that there are many families of subcubic graphs G that are not cubic graphs satisfying $\gamma_t(G)/\gamma(G) = 2$. For example, the 2-corona of a path or a cycle is such a family of subcubic graphs, where we recall that the 2-corona $H \circ P_2$ of a graph H is the graph of order $3|V(H)|$ obtained from H by attaching a path of length 2 to each vertex of H so that the resulting paths are vertex-disjoint. We close with the following two open problems.

Problem 1 Characterize the subcubic isolate-free graphs G satisfying $\gamma_t(G)/\gamma(G) = 2$.

Problem 2 Characterize the subcubic isolate-free graphs G satisfying $\Gamma_t(G)/\Gamma(G) = 2$.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Archdeacon, D., Ellis-Monaghan, J., Fischer, D., Froncek, D., Lam, P.C.B., Seager, S., Wei, B., Yuster, R.: Some remarks on domination. *J. Graph Theory* **46**, 207–210 (2004)
2. Bollobás, B., Cockayne, E.J.: Graph-theoretic parameters concerning domination, independence, and irredundance. *J. Graph Theory* **3**, 241–249 (1979)
3. Brešar, B., Henning, M.A., Rall, D.F.: Total dominating sequences in graphs. *Discrete Math.* **339**, 1665–1676 (2016)
4. Chvátal, V., McDiarmid, C.: Small transversals in hypergraphs. *Combinatorica* **12**, 19–26 (1992)
5. Cockayne, E.J., Dawes, R.M., Hedetniemi, S.T.: Total domination in graphs. *Networks* **10**, 211–219 (1980)
6. Desormeaux, W.J., Haynes, T.W., Henning, M.A., Yeo, A.: Total domination numbers of graphs with diameter two. *J. Graph Theory* **75**, 91–103 (2014)
7. Dorbec, P., Henning, M.A., Rall, D.F.: On the upper total domination number of Cartesian products of graphs. *J. Combin. Optim.* **16**, 68–80 (2008)
8. Favaron, O., Henning, M.A., Mynhardt, C.M., Puech, J.: Total domination in graphs with. *J. Graph Theory* **34**(1), 9–19 (2000)
9. Haynes, T.W., Hedetniemi, S.T., Slater, P.J.: *Fundamentals of Domination in Graphs*. Marcel Dekker, New York (1998)
10. Haynes, T.W., Hedetniemi, S.T., Slater, P.J. (eds.): *Domination in Graphs: Advanced Topics*. Marcel Dekker, New York (1998)

11. Henning, M.A.: Graphs with large total domination number. *J. Graph Theory* **35**(1), 21–45 (2000)
12. Henning, M.A.: Trees with large total domination number. *Utilitas Math.* **60**, 99–106 (2001)
13. Henning, M.A., Yeo, A.: Hypergraphs with large transversal number and with edge sizes at least three. *J. Graph Theory* **59**, 326–348 (2008)
14. Henning, M.A.: Recent results on total domination in graphs: a survey. *Discrete Math.* **309**, 32–63 (2009)
15. Henning, M.A., Yeo, A.: *Total domination in graphs* (Springer Monographs in Mathematics) (2013). ISBN: 978-1-4614-6524-9 (Print) 978-1-4614-6525-6 (Online)
16. Henning, M.A., Yeo, A.: Upper total domination. In: *Total domination in graphs* (Springer Monographs in Mathematics), Chapter 16, pp. 131–139 (2013). ISBN: 978-1-4614-6524-9 (Print) 978-1-4614-6525-6 (Online)
17. Henning, M.A., Yeo, A.: A new lower bound for the total domination number in graphs proving a Graffiti conjecture. *Discrete Appl. Math.* **173**, 45–52 (2014)
18. Southey, J., Henning, M.A.: Edge weighting functions on dominating sets. *J. Graph Theory* **72**, 346–360 (2013)
19. Tuza, Z.: Covering all cliques of a graph. *Discrete Math.* **86**, 117–126 (1990)