



Alternating Sign Matrices: Extensions, König-Properties, and Primary Sum-Sequences

Richard A. Brualdi¹ · Geir Dahl² 

Received: 8 January 2019 / Revised: 2 November 2019 / Published online: 18 November 2019
© Springer Japan KK, part of Springer Nature 2019

Abstract

This paper is concerned with properties of permutation matrices and alternating sign matrices (ASMs). An ASM is a square $(0, \pm 1)$ -matrix such that, ignoring 0's, the 1's and -1 's in each row and column alternate, beginning and ending with a 1. We study extensions of permutation matrices into ASMs by changing some zeros to $+1$ or -1 . Furthermore, several properties concerning the term rank and line covering of ASMs are shown. An ASM A is determined by a sum-matrix $\Sigma(A)$ whose entries are the sums of the entries of its leading submatrices (so determined by the entries of A). We show that those sums corresponding to the nonzero entries of a permutation matrix determine all the entries of the sum-matrix and investigate some of the properties of the resulting sequence of numbers. Finally, we investigate the lattice-properties of the set of ASMs (of order n), where the partial order comes from the Bruhat order for permutation matrices.

Keywords Permutation matrix · Alternating sign matrix · Term rank · Line covering · Bruhat order

1 Introduction

An alternating sign matrix, abbreviated to ASM, is an $n \times n$ $(0, \pm 1)$ -matrix such that, ignoring 0's, the 1's and -1 's in each row and column alternate, beginning and ending with a 1. An $n \times n$ ASM A can be regarded as the adjacency matrix of a signed bipartite graph B whose vertices in each set of its bipartition have a specified order (to account for the alternating sign property). Thus, B has a vertex for each row and

✉ Geir Dahl
geird@math.uio.no

Richard A. Brualdi
brualdi@math.wisc.edu

¹ Department of Mathematics, University of Wisconsin, Madison, WI 53706, USA

² Department of Mathematics, University of Oslo, Oslo, Norway

column of the matrix $A = [a_{ij}]$, and an edge between a row vertex i and a column vertex j when the corresponding entry a_{ij} is nonzero, and associated to that edge is the sign of the entry a_{ij} . The set of $n \times n$ ASMs is denoted by \mathcal{A}_n . Permutation matrices are the ASMs without any -1 's. We denote by \mathcal{P}_n the set of $n \times n$ permutation matrices corresponding to the set \mathcal{S}_n of permutations of $\{1, 2, \dots, n\}$. Thus $\mathcal{P}_n \subseteq \mathcal{A}_n$. If $\pi = (\pi_1, \pi_2, \dots, \pi_n) \in \mathcal{S}_n$, the corresponding permutation matrix has its ones in positions (i, π_i) ($i \leq n$). We will also use the shorter notation $\pi = (\pi_1 \pi_2 \dots \pi_n)$. Some recent developments concerning ASMs, related objects and generalizations may be found in [4–6].

There is a partial order defined on the set \mathcal{S}_n (and so on \mathcal{P}_n) called the *Bruhat order* and denoted by \leq_B . If $\sigma, \tau \in \mathcal{S}_n$, then $\sigma \leq_B \tau$ (read as, σ precedes τ in the Bruhat order) provided σ can be obtained from τ by a sequence of transpositions (k, l) (interchanging k and l in a permutation) each of which decreases the number of inversions. The cover relation in this partial order results when one transposition is applied and it decreases the number of inversions by exactly one. The unique minimal permutation in the partially ordered set (\mathcal{S}_n, \leq_B) is the identity permutation, denoted as $\iota_n = (1, 2, \dots, n)$, and the unique maximal permutation is the so-called *anti-identity permutation*, denoted as $\zeta_n = (n, n-1, \dots, 1)$. If $\sigma \leq_B \tau$, and σ and τ have corresponding permutation matrices P and Q , then we also write $P \leq_B Q$. The minimal permutation matrix and maximal permutation matrix are, respectively, the $n \times n$ identity matrix I_n and $n \times n$ anti-identity matrix (with 1's on the anti-diagonal) L_n . For more on the Bruhat order, and the so-called Bruhat shadow of a permutation matrix, see [3].

There is an equivalent, and computationally efficient, way to determine whether or not two permutations are related in the Bruhat order, best described in terms of their corresponding permutation matrices. It requires only the comparison of $(n-1)^2$ pairs of integers, after having computed certain sum-matrices, as defined next. For any $m \times n$ matrix $A = [a_{ij}]$, define the *sum-matrix* of A , denoted $\Sigma(A) = [\sigma_{ij}]$ to be the $m \times n$ matrix where

$$\sigma_{ij} = \sigma_{ij}(A) = \sum_{k \leq i} \sum_{l \leq j} a_{kl} \quad (1 \leq i \leq m, 1 \leq j \leq n),$$

the sum of the entries in the leading $i \times j$ submatrix of A . If P is an $n \times n$ permutation matrix, then the entries in the last row and last column of $\Sigma(A)$ are $1, 2, \dots, n$ in that order. Note that if P is a permutation matrix, then $P^{-1} = P^T$, so $\Sigma(P^{-1}) = \Sigma(P^T) = \Sigma(P)^T$. Let the entrywise ordering of two matrices A and B be written as $A \leq B$. Let $\Sigma(\mathcal{P}_n) := \{\Sigma(P) : P \in \mathcal{P}_n\}$. The characterization of Bruhat order in terms of sum-matrices is stated next (see [1] for a proof).

Lemma 1 *Let $P, Q \in \mathcal{P}_n$. Then $P \leq_B Q$ if and only if*

$$\Sigma(P) \geq \Sigma(Q),$$

that is, $\Sigma(P)$ dominates $\Sigma(Q)$ entrywise.

The Bruhat order on \mathcal{P}_n can be extended to a Bruhat order on alternating sign matrices using sum-matrices: For $A_1, A_2 \in \mathcal{A}_n$, then $A_1 \leq_B A_2$ provided $\Sigma(A_1) \geq \Sigma(A_2)$. In the next theorem we summarize some of the results of Lascoux and Schützenberger [12] which, in particular, contain a characterization of ASMs in terms of their sum-matrices. Let $\Sigma(\mathcal{A}_n) := \{\Sigma(A) : A \in \mathcal{A}_n\}$.

Theorem 1 *The partially ordered set (\mathcal{A}_n, \leq_B) is a lattice and is the smallest lattice extending the partially ordered set (\mathcal{P}_n, \leq_B) , the MacNeille completion of (\mathcal{P}_n, \leq_B) . The set $\Sigma(\mathcal{A}_n)$ consists of the set Σ_n of all $n \times n$ nonnegative integral matrices $X = [x_{ij}]$ satisfying the following properties for each $i = 1, 2, \dots, n$:*

- (LS-a) *The integers in row i and in column i are taken from the set $\{0, 1, \dots, i\}$, beginning with a 0 or 1, and ending with i .*
- (LS-b) *The integers in row i and in column i are nondecreasing.*
- (LS-c) *Two consecutive entries in row i and in column i are equal or differ by 1.*

There is a generalization of a permutation interchange that applies to $n \times n$ ASMs, namely, adding an interchange-matrix $T_n^{i,j}$ which is all zeros except for its 2×2 matrix determined by rows i and $i + 1$ and columns j and $j + 1$ which equals

$$\begin{bmatrix} +1 & -1 \\ -1 & +1 \end{bmatrix} \quad (1 \leq i, j \leq n - 1).$$

Then it follows from [12] (see also Lemma 2 in [8]) that if $A_1, A_2 \in \mathcal{A}_n$, then $A_1 \leq_B A_2$ if and only if A_1 can be obtained from A_2 by sequentially adding matrices of the form $T_n^{i,j}$ where the result of each addition is also an ASM.

A matrix is uniquely determined by its sum-matrix since we can recover $A = [a_{ij}]$ from $\Sigma(A) = [\sigma_{ij}]$ as follows:

$$a_{ij} = \sigma_{ij} - \sigma_{i-1,j} - \sigma_{i,j-1} + \sigma_{i-1,j-1}$$

where $\sigma_{0,j} = \sigma_{i,0}$ for all i and j . Thus the mapping $\Sigma : \mathcal{A}_n \rightarrow \Sigma_n$ given by $A \rightarrow \Sigma(A)$ is a bijection. The sum-matrices of the minimal and maximal $n \times n$ ASMs have the special forms shown for $n = 6$:

$$\Sigma(I_6) = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 & 3 & 3 \\ 1 & 2 & 3 & 4 & 4 & 4 \\ 1 & 2 & 3 & 4 & 5 & 5 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{bmatrix} \quad \text{and} \quad \Sigma(L_6) = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 & 2 & 3 \\ 0 & 0 & 1 & 2 & 3 & 4 \\ 0 & 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{bmatrix}.$$

Thus the sum-matrix of every $n \times n$ ASM lies entrywise between $\Sigma(L_n)$ and $\Sigma(I_n)$, that is, each entry of $\Sigma(I_n)$, respectively, $\Sigma(L_n)$, is at least as large (at most as large) as the corresponding entry of the sum-matrix of every $A \in \mathcal{A}_n$.

Let $\widehat{\Sigma(\mathcal{A}_n)}$ be the convex hull of $\Sigma(\mathcal{A}_n)$. From a proof in connection with the notion of sum-majorization in [4] we get the following characterization of the polytope $\widehat{\Sigma(\mathcal{A}_n)}$.

Theorem 2 *The polytope $\widehat{\Sigma}(\mathcal{A}_n)$ equals the set of $n \times n$ real matrices $S = [\sigma_{ij}]$ satisfying*

$$\begin{aligned} 0 &\leq \sigma_{ij} - \sigma_{i-1,j} \leq 1 \quad (1 \leq i, j \leq n) \\ 0 &\leq \sigma_{ij} - \sigma_{i,j-1} \leq 1 \quad (1 \leq i, j \leq n) \\ \sigma_{i,n} &= i \quad (1 \leq i \leq n) \\ \sigma_{n,j} &= j \quad (1 \leq j \leq n) \end{aligned} \tag{1}$$

where we define $\sigma_{0,j} = \sigma_{i,0} = 0$ for all i and j . The set of extreme points of $\widehat{\Sigma}(\mathcal{A}_n)$ equals the set $\Sigma(\mathcal{A}_n)$ of sum-matrices of the $n \times n$ ASMs.

This paper is organized as follows. In Sect. 2 we study extensions of permutation matrices to ASMs obtained by changing some zeros (possibly none) into +1 or −1. König properties of ASMs are studied in Sect. 3 where, by this, we mean the term rank and the structure of minimum line coverings of such matrices. In Sect. 4, we investigate in more detail sum-matrices and show that in the case of $n \times n$ permutation matrices, at most n entries of the sum-matrix are needed to recover the permutation matrix from its sum-matrix. In Sect. 5 we investigate certain lattice properties of the Bruhat order.

2 ASM-Extensions

Let P be an $n \times n$ permutation matrix. Let $\alpha_n(P)$ denote the number of ASM-extensions of P , defined to be $n \times n$ ASMs obtained from P by replacing some 0's (possibly none) by +1 or −1. In such an ASM-extension of P , the number of new +1's equals the number of new −1's. It is natural to ask for

$$\alpha_n^* := \max\{\alpha_n(P) : P \in \mathcal{P}_n\}$$

and those $n \times n$ permutation matrices P that satisfy $\alpha_n(P) = \alpha_n^*$.

If $n = 3$, there is only one non-permutation ASM and it does not contain 1's in a permutation set of places. So $\alpha_3^* = 1$. The 4×4 permutation matrices

$$\begin{bmatrix} & 1 & & \\ 1 & & & \\ & & & 1 \\ & & 1 & \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} & & 1 & \\ & & & 1 \\ 1 & & & \\ & 1 & & \end{bmatrix}$$

have ASM-extensions with two −1's:

$$\begin{bmatrix} & 1 & & \\ 1 & -1 & 1 & \\ & 1 & -1 & 1 \\ & & & 1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} & & 1 & \\ & 1 & -1 & 1 \\ 1 & -1 & 1 & \\ & 1 & & \end{bmatrix}.$$

(As usual, empty positions are assumed to contain a 0.) It is straightforward to verify that $\alpha_4^* = 2$ and that the above two 4×4 permutation matrices are the only ones achieving equality.

Consider the special $n \times n$ permutation matrix

$$Q_n^* = Q_{\lceil \frac{n}{2} \rceil} \oplus Q_{\lfloor \frac{n}{2} \rfloor},$$

where Q_m is the $m \times m$ permutation matrix with 1's on its anti-diagonal running from the lower left to the upper right. For instance, for $n = 5$ and 6 we have

$$Q_5^* = \begin{bmatrix} | & | & 1 & | & | \\ | & 1 & | & | & | \\ | & | & | & | & | \\ 1 & | & | & | & | \\ | & | & | & | & 1 \\ | & | & | & 1 & | \end{bmatrix} \quad \text{and} \quad Q_6^* = \begin{bmatrix} | & | & 1 & | & | & | \\ | & 1 & | & | & | & | \\ | & | & | & | & | & | \\ 1 & | & | & | & | & | \\ | & | & | & | & | & 1 \\ | & | & | & | & 1 & | \\ | & | & | & 1 & | & | \end{bmatrix}.$$

There is a ‘canonical’ symmetric ASM-extension A_n^* of Q_n^* as illustrated below for $n = 7$ and $n = 8$:

$$\left[\begin{array}{ccc|ccc} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 & & 1 \\ 0 & 1 & -1 & & 1 & 0 \\ \hline 1 & -1 & & 1 & & 0 \\ \hline 0 & & 1 & & & -1 \\ 0 & 1 & & & -1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{array} \right], \quad \left[\begin{array}{cccc|ccc} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & & & 1 \\ 0 & 1 & -1 & & & 1 & 0 \\ \hline 1 & -1 & & & 1 & & 0 \\ \hline 0 & & & 1 & & & -1 \\ 0 & & 1 & & & -1 & 1 \\ 0 & 1 & & & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \end{array} \right],$$

where the 0's that are explicitly displayed are required in any extension of Q_n^* to an ASM. In these canonical extensions, the -1 's are in the positions of the 1 's of a Q_{n-2}^* . In addition, the middle $(n - 2) \times (n - 2)$ principal submatrix contains all 1 's on its anti-diagonal. The sum-matrices of A_7^* and A_8^* are given by

$$\Sigma(A_7^*) = \left[\begin{array}{ccc|ccc} 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 & 2 \\ 0 & 1 & 1 & 1 & 2 & 3 \\ \hline 1 & 1 & 1 & 2 & 3 & 4 \\ \hline 1 & 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 4 & 4 & 5 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{array} \right] \quad \text{and} \quad \Sigma(A_8^*) = \left[\begin{array}{cccc|cccc} 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 & 2 & 2 & 2 \\ 0 & 1 & 1 & 1 & 1 & 2 & 3 & 3 \\ \hline 1 & 1 & 1 & 1 & 2 & 3 & 4 & 4 \\ \hline 1 & 1 & 1 & 2 & 3 & 4 & 4 & 5 \\ 1 & 1 & 2 & 3 & 4 & 4 & 5 & 6 \\ 1 & 2 & 3 & 4 & 4 & 5 & 6 & 7 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \end{array} \right]. \quad (2)$$

Let $\mathcal{A}_n(Q_{2m}^*)$ denote the set of $n \times n$ ASM-extensions of Q_{2m}^* , that is, ASMs containing 1's in those positions that are 1 in Q_{2m}^* . Then a matrix in $\mathcal{A}_n(Q_{2m}^*)$ has the form

$$A = \begin{bmatrix} Q'_m & X_m \\ Y_m & Q''_m \end{bmatrix},$$

where Q'_m and Q''_m have 1's wherever Q_m has 1's. It is straightforward to check that Q'_m must have all zeros above its anti-diagonal and Q''_m must have all zeros below its anti-diagonal. We conjecture that $\alpha_n^* = \alpha_n(Q_n^*)$. We remark that in [6] completions of certain $(0, -1)$ -matrices to ASMs are considered where the matrices Q_n^* also occur, and there is a related conjecture.

Let F_{2m} be the so-called $2m \times 2m$ diamond ASM, the ASM with the largest number of nonzeros [7]. Then $F_{2m} \in \mathcal{A}_n(Q_{2m}^*)$. For instance, with $m = 4$,

$$F_8 = \begin{bmatrix} & & & & 1 & & & & \\ & & & & 1 & -1 & 1 & & & \\ & & & & 1 & -1 & 1 & & & \\ & & 1 & -1 & 1 & -1 & 1 & & & \\ 1 & -1 & 1 & -1 & 1 & -1 & 1 & & & \\ & & 1 & -1 & 1 & -1 & 1 & -1 & 1 & \\ & & & & 1 & -1 & 1 & & & \\ & & & & & & 1 & & & \\ & & & & & & & & & 1 \end{bmatrix}.$$

Let $F_{2m}^{(1)}$ be the $(0, 1)$ -matrix with $m(m - 1)$ 1's, obtained from F_{2m} by replacing all its -1 's with 0's. Then $Q_{2m}^* \leq F_{2m}^{(1)}$ and it is straightforward to check that no other permutation matrix P satisfies $P \leq F_{2m}^{(1)}$. Indeed, it is easily seen that, after permutations of rows and columns,

$$F_{2m}^{(1)} = \begin{bmatrix} T_m & O_m \\ O_m & T_m \end{bmatrix}$$

where T_m is the $m \times m$ matrix with 1's on and below its main diagonal and 0's elsewhere; the 1's on the main diagonal of T_m are those corresponding to the 1's of Q_{2m}^* . In particular, the permutation matrix Q_{2m}^* is the only permutation matrix of which F_{2m} is an ASM-extension. Since the matrix F_{2m} has the maximum number of nonzeros among all matrices in \mathcal{A}_{2m} , F_{2m} is the unique ASM-extension of Q_{2m}^* with the maximum number of nonzeros.

The matrix $\Sigma(F_{2m})$ has a special form which is illustrated for $\Sigma(F_8)$ below with $\Sigma(A_8^*) - \Sigma(F_8^*)$ also given:

$$\Sigma(F_8) = \begin{bmatrix} 0 & 0 & 0 & 1 & | & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & | & 2 & 2 & 2 & 2 \\ 0 & 1 & 1 & 2 & | & 2 & 3 & 3 & 3 \\ 1 & 1 & 2 & 2 & | & 3 & 3 & 4 & 4 \\ \hline 1 & 2 & 2 & 3 & | & 3 & 4 & 4 & 5 \\ 1 & 2 & 3 & 3 & | & 4 & 4 & 5 & 6 \\ 1 & 2 & 3 & 4 & | & 4 & 5 & 6 & 7 \\ 1 & 2 & 3 & 4 & | & 5 & 6 & 7 & 8 \end{bmatrix}, \quad \Sigma(Q_8^*) - \Sigma(F_8) = \begin{bmatrix} & & & & & & & & \\ & & & & & & & & \\ & & & 1 & & & & & \\ & & 1 & 1 & 1 & & & & \\ \hline 1 & 1 & 2 & 1 & 1 & & & & \\ \hline 1 & 1 & 1 & 1 & & & & & \\ & & & 1 & & & & & \\ & & & & & & & & \\ & & & & & & & & \\ & & & & & & & & \end{bmatrix}.$$

Thus $\Sigma(Q_8^*) \geq \Sigma(F_8^*)$, and hence $F_8 \leq_B Q_8^*$. (We remark that the odd case is different, since we have $\Sigma(A_7^*) \not\leq \Sigma(F_7)$ and $\Sigma(F_7) \not\leq \Sigma(A_7^*)$: see (2) for $\Sigma(A_7^*)$.)

Example 1 We calculate that

$$\Sigma(Q_8^*) = \begin{bmatrix} 0 & 0 & 0 & 1 & | & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 2 & | & 2 & 2 & 2 & 2 \\ 0 & 1 & 2 & 3 & | & 3 & 3 & 3 & 3 \\ 1 & 2 & 3 & 4 & | & 4 & 4 & 4 & 4 \\ \hline 1 & 2 & 3 & 4 & | & 4 & 4 & 4 & 5 \\ 1 & 2 & 3 & 4 & | & 4 & 4 & 5 & 6 \\ 1 & 2 & 3 & 4 & | & 4 & 5 & 6 & 7 \\ 1 & 2 & 3 & 4 & | & 5 & 6 & 7 & 8 \end{bmatrix}.$$

Using our calculation of $\Sigma(A_8^*)$, we get

$$\Sigma(Q_8^*) - \Sigma(A_8^*) = \begin{bmatrix} & & & & & & & & \\ & & & & & & & & \\ & & & 1 & 1 & 1 & & & \\ & & 1 & 2 & 2 & 1 & & & \\ \hline 1 & 2 & 3 & 2 & 1 & & & & \\ \hline 1 & 2 & 2 & 1 & & & & & \\ 1 & 1 & 1 & & & & & & \\ & & & & & & & & \\ & & & & & & & & \end{bmatrix}.$$

In general, we have

$$\Sigma(Q_{2m}^*) = \left[\begin{array}{c|c} C_1 & C_2 \\ \hline C_2^T & C_1 + J_m \end{array} \right],$$

where C_1 is a $m \times m$ matrix with 0's above the anti-diagonal, 1's on the anti-diagonal, and 2's, 3's, . . . on the diagonals below it, C_2 is an $m \times m$ matrix whose row i contains all i 's ($1 \leq i \leq m$), and J_m is the $m \times m$ all-ones matrix. □

Lemma 2 *Let A be an $n \times n$ ASM with $n = 2m$. Then $A \in \mathcal{A}_{2m}(Q_{2m}^*)$ if and only if*

$$Q_{2m}^* \preceq_B A \preceq_B A_{2m}^*,$$

that is, if and only if

$$\Sigma(A_{2m}^*) \leq \Sigma(A) \leq \Sigma(Q_{2m}^*),$$

a certain interval in the Bruhat order on the permutations in S_{2m} .

Proof We first show that if $A \in \mathcal{A}_{2m}(Q_{2m}^*)$, then $\Sigma(A) \leq \Sigma(Q_{2m}^*)$. But this is clear from the form of $\Sigma(Q_{2m}^*)$ given in Example 1. The reason is that any ASM-completion of Q_{2m}^* has only 0’s above the anti-diagonal in its leading $m \times m$ submatrix and the other entries in the upper $m \times 2m$ submatrix are as large as they can be in any such ASM. A similar conclusion holds for the lower $m \times 2m$ submatrix.

We next verify that if $A \in \mathcal{A}_{2m}(Q_{2m}^*)$, then $\Sigma(A_{2m}^*) \leq \Sigma(A)$. Partition A_{2m}^* into $m \times m$ matrices as

$$A_{2m}^* = \begin{bmatrix} A_1^* & A_2^* \\ A_3^* & A_4^* \end{bmatrix}.$$

The matrix A_1^* has all 0’s above the anti-diagonal, all 1’s on the anti-diagonal, and all -1 ’s on the diagonal immediately below the anti-diagonal. These -1 ’s occur as early as possible in rows $2, 3, \dots, m$ of A_{2m}^* . The matrix A_2^* has 1’s on its anti-diagonal except for a 0 in its upper right corner. These 1’s occur as late as possible in rows $2, 3, \dots, m$ of A_{2m}^* . It follows from this that $\sigma_{ij}(A_{2m}^*) \leq \sigma_{ij}(A)$ for $1 \leq i \leq m$ and $1 \leq j \leq 2m$ and, similarly, for $1 \leq i \leq 2m$ and $1 \leq j \leq m$. Since an ASM remains an ASM under a 180° rotation (that is, reading an ASM from the lower right corner from leftwards and upwards also gives an ASM), the theorem now follows. \square

Corollary 1 *Let $n = 2m$. Then the cardinality of $\mathcal{A}(Q_{2m}^*)$ equals the number of $n \times n$ nonnegative integral matrices X with $\Sigma(A_{2m}^*) \leq X \leq \Sigma(Q_{2m}^*)$ (entrywise) satisfying (LS-a), (LS-b), and (LS-c). \square*

We now consider $n \times n$ permutation matrices P satisfying $\alpha_n(P) = 1$, that is, permutation matrices having no ASM-completions other than itself. We call such permutation matrices *isolated*. For instance, the following permutation matrix P is isolated:

$$P = \begin{bmatrix} | & | & 1 & | & | \\ \hline | & | & | & | & 1 \\ \hline | & 1 & | & | & | \\ \hline 1 & | & | & | & | \\ \hline | & | & | & 1 & | \end{bmatrix}. \tag{3}$$

Let $P = [p_{ij}]$ be a permutation matrix. Assume that $i_1 < i_2 < i_3 < i_4$ and $j_1 < j_2 < j_3 < j_4$ be such that either

- (i) P has 1’s in each of the positions $(i_1, j_3), (i_2, j_4), (i_3, j_1), (i_4, j_2)$,
- or

(ii) P has 1's in each of the positions $(i_1, j_2), (i_2, j_1), (i_3, j_4), (i_4, j_3)$.

If (i) or (ii) holds, we say that P has an *open square*.

Lemma 3 *Let P be a permutation matrix. Assume that P has an open square. Then $\alpha_n(P) > 1$.*

Proof Assume that case (i) above holds. Let the matrix A be obtained from P by changing the two entries in positions (i_2, j_3) and (i_3, j_2) (from 0) into -1 , and also changing the two entries in positions (i_2, j_2) and (i_3, j_3) (from 0) into 1. Then A is an ASM, and it is an ASM completion of P . Thus, P has at least two ASM completions, so $\alpha_n(P) > 1$. The proof in case (ii) is similar. \square

Note that having an open square is equivalent to saying that either (i) the permutation σ of $\{1, 2, \dots, n\}$ corresponding to the permutation matrix P contains the pattern 3, 4, 1, 2 (corresponding to j_3, j_4, j_1, j_2) or (ii) the permutation σ of $\{1, 2, \dots, n\}$ corresponding to the permutation matrix P contains the pattern 2, 1, 4, 3 (corresponding to j_2, j_1, j_4, j_3). Thus, P not having an open square is equivalent to the corresponding permutation in S_n being a 3412-avoiding and 2143-avoiding permutation. Note that 2143 and 3412 are reverses of one another, and so the number of 3412-avoiding permutations equals the number of 2143-avoiding permutations. It follows from results on pattern-avoidance permutations (see e.g. page 154 of [2]) that the number of 2143-avoiding (resp. 3412-avoiding) permutations of $\{1, 2, \dots, n\}$ is the same as the number of 1234-avoiding permutations, that is, that do not contain an increasing subsequence of length 4. By a theorem of Gessel (see page 176 of [2]), this number equals

$$\frac{1}{(n + 1)^2(n + 2)} \sum_{k=0}^n \binom{2k}{k} \binom{n + 1}{k + 1} \binom{n + 2}{k + 1}. \tag{4}$$

Example 2 The following permutation matrix P , corresponding to the permutation $(3, 5, 1, 2, 4)$, contains an open square, shown with the corresponding ASM extension A . With 3, 5, 1, 2, it is not 3412-avoiding.

$$P = \begin{bmatrix} & & 1 & & \\ & * & * & 1 & \\ 1 & * & * & & \\ & 1 & & & \\ & & & & 1 \end{bmatrix}, \quad A = \begin{bmatrix} & & 1 & & \\ & 1 & -1 & 1 & \\ 1 & -1 & 1 & & \\ & 1 & & & \\ & & & & 1 \end{bmatrix}.$$

Note that the matrix P is obtained from the matrix in (3) by a transposition. \square

Theorem 3 *Let P be an $n \times n$ permutation matrix corresponding to a permutation $\sigma \in S_n$. Then $\alpha_n(P) = 1$, that is, P is an isolated permutation matrix, if and only if P does not have an open square, that is, if and only if σ is both a 3412-avoiding and 2143-avoiding permutation.*

Proof This follows directly from Lemma 3 and Corollary 6.2 of [7]. \square

Since a permutation can be both 3412-avoiding and 2143-avoiding, we can only say that the number of $n \times n$ permutation matrices P that have only one ASM-completion is bounded above by twice the number in (4).

Consider again the $n \times n$ permutation matrix $Q_n^* = Q_{\lceil \frac{n}{2} \rceil} \oplus Q_{\lfloor \frac{n}{2} \rfloor}$. Then Q_n^* contains $\binom{\lceil \frac{n}{2} \rceil}{2} \binom{\lfloor \frac{n}{2} \rfloor}{2}$ open squares. If e.g. $n = 2m$ is even, then Q_n^* corresponds to the permutation

$$\sigma_{2m} = (m, m - 1, \dots, 2, 1, 2m, 2m - 1, \dots, m + 2, m + 1).$$

We believe that σ_{2m} contains the largest number of patterns 2143, namely $\binom{m}{2}^2$ and thus has the largest number of completions to an ASM with two -1 's, but this does not yet mean that $\alpha_n(Q_n^*)$ is maximum, although we expect it is.

3 König Properties of a Class of ASMs

Consider again the question of extensions of a permutation matrix P to ASMs A where A has 1's where P has 1's. A “dual” viewpoint is to ask, for a given ASM A , if there exists a permutation matrix P such that P has all its 1's where A has a 1. This leads to a related question concerning term ranks. The term rank of ASMs was studied in [7]. We now consider the term rank of the nonnegative part of ASMs. For a real matrix A , let A^+ denote the matrix $\max\{A, 0\}$ in which all negative entries are replaced with 0's; we call A^+ the *nonnegative part* of A . Let $\rho(B)$ denote the term rank of a matrix B , i.e., the largest number of nonzeros in B with no two of these in the same row or column.

The maximum of $\rho(A^+)$ among all ASMs of order n is clearly n , and it is attained for permutation matrices.

The next result determines the minimum of $\rho(A^+)$ among all ASMs of order n .

Theorem 4 *The minimum of $\rho(A^+)$ among ASMs of order n is $\lceil 2\sqrt{n+1} - 2 \rceil$.*

Proof Let A be an ASM of order n and let $t = \rho(A^+)$. By König's theorem there are t lines that cover (contain) all 1s in A , say e rows and f columns, where $e + f = t$. Permute rows such that these e rows become first, and permute columns such that these f columns become first. Note that this affects the ASM property of the ± 1 's alternating. Then the permuted A has the form

$$A = \begin{bmatrix} A_1 & A_2 \\ A_3 & A_4 \end{bmatrix} \tag{5}$$

where A_1 has size $e \times f$. The matrix A_4 contains only -1 's and 0 's. Let p_i and n_i (resp.) be the number of 1 's and -1 's in the submatrix A_i ($i \leq 4$). So, $p_4 = 0$. Then, as line sums are not changed by the permutations,

$$\begin{aligned} (p_1 - n_1) + (p_2 - n_2) &= e, & (p_1 - n_1) + (p_3 - n_3) &= f. \\ p_3 - n_3 - n_4 &= n - e, & p_2 - n_2 - n_4 &= n - f. \end{aligned}$$

Thus

$$n_1 - p_1 = (p_2 - n_2) - e = n_4 + n - f - e = n_4 + n - t.$$

But $n_1 - p_1$ is at most the number of nonzeros in A_1 , so $n_1 - p_1 \leq ef \leq (t/2)^2$ as $e + f = t$. Therefore,

$$n - t \leq n_4 + n - t = n_1 - p_1 \leq t^2/4$$

so that $t^2 + 4t \geq 4n$ and $(t + 2)^2 \geq 4n + 4$. This implies that

$$t \geq \lceil 2\sqrt{n+1} - 2 \rceil.$$

It follows that

$$\min\{\rho(A^+) : A \in \mathcal{A}_n\} \geq \lceil 2\sqrt{n+1} - 2 \rceil.$$

On the other hand, Corollary 4.2 in [7] shows that

$$\min\{\rho(A) : A \in \mathcal{A}_n\} = \lceil 2\sqrt{n+1} - 2 \rceil$$

where $\rho(A)$ denotes the term rank of A . Clearly $\rho(A) \geq \rho(A^+)$ for any matrix A , so it follows that $\min\{\rho(A^+) : A \in \mathcal{A}_n\} = \lceil 2\sqrt{n+1} - 2 \rceil$. □

The previous theorem and its proof show the surprising fact that the minimum term rank among $n \times n$ ASMs is the same as the minimum term rank of the matrices obtained by replacing all negative entries with zeros. In [7], for each n , an ASM A is given which attains the minimum term rank in \mathcal{A}_n , and therefore also the minimum rank of A^+ in that class.

We now study the term rank and line covers of certain ASMs. In a matrix A a *line cover* is a set of lines (rows and columns) that contain all nonzeros of A . The minimum number of lines in a line cover of A is called the *line cover number* and it is denoted by $\tau(A)$. A classical theorem of König says that $\tau(A) = \rho(A)$, i.e., that the line cover number equals the term rank of A .

Let D_3 denote the unique ASM of order 3 with a single -1 , so

$$D_3 = \begin{bmatrix} | & | & 1 \\ 1 & -1 & 1 \\ | & & 1 \end{bmatrix}.$$

Consider an ASM A of order n . Let B be a matrix obtained from A by identifying a single $+1$ of A , say in position (i, j) , with a single $+1$ of D_3 and inserting two new rows and two new columns such that D_3 is embedded in the resulting matrix. These new rows, and the new columns, may not be consecutive, but they must retain the same relative order as in D_3 and maintain the alternating sign property. We call the mentioned position (i, j) the *insertion position*. Such a matrix is an ASM and will be

denoted by $A * D_3$ and called an D_3 -extension of A . A row of a matrix with a unique nonzero (which must be 1) will be called a *unit* row, otherwise it is called a *non-unit* row. We define a unit column and a non-unit column similarly.

We now consider two specific types of such D_3 -extensions B where, as above, the insertion position is (i, j) :

(*a) Row i is a non-unit row of A . Moreover, the -1 of the inserted D_3 is in the corresponding row of B . A similar type is obtained when column j is a non-unit row and the inserted -1 is in the corresponding column.

(*b) Row i is a unit row with its nonzero ($a + 1$) in column j . Moreover, the -1 of the inserted D_3 is in row i . A similar type is obtained by interchanging the roles of rows and columns in the previous sentence.

For instance, D_3 itself is a D_3 -extension of $J_1 = [1]$ of type (*b). Let \mathcal{A}^* denote the class of ASMs that may be obtained from J_1 using a finite number of D_3 -extensions of type (*a) or (*b). Observe that every matrix except J_1 in \mathcal{A}^* has the property that every $+1$ is in a non-unit row or column. This is seen from the structure of the extension.

Example 3 The matrix B_1 below is obtained by a D_3 -extension of type (*a) of D_3 , using column 2 and insertion position $(3, 2)$. B_2 is a D_3 -extension of type (*b) of D_3 , using column 3 and insertion position $(2, 3)$.

$$D_3 = \begin{bmatrix} & + & & \\ + & - & + & \\ & + & & \end{bmatrix}, \quad B_1 = \begin{bmatrix} & & + & & \\ & + & - & + & \\ & & + & & \\ + & & - & & + \\ & & + & & \end{bmatrix}, \quad B_2 = \begin{bmatrix} & & + & & \\ & + & - & + & \\ & & + & & \\ + & & & - & + \\ & & & + & \end{bmatrix}.$$

□

For an ASM A let $n_r^-(A)$ (resp. $n_c^-(A)$) denote the number of rows (resp. columns) of A that contain at least one negative entry, i.e., the number of non-unit rows (resp. columns).

Lemma 4 *Let A be an ASM which is not the direct sum of a permutation matrix and a smaller ASM. Then*

$$\rho(A) \leq n_r^-(A) + n_c^-(A).$$

Proof Let L be the set of lines in A that contain at least one -1 . So, $|L| = n_r^-(A) + n_c^-(A)$. Moreover, by assumption, there is no 1 in A that lies both in a unit row and a unit column (because then A would be the direct sum of $[1]$ and a smaller ASM). Therefore L covers all the ones in A , and clearly all the -1 's. So, L is a line cover, and

$$\rho(A) = \tau(A) \leq |L| = n_r^-(A) + n_c^-(A).$$

□

The next result gives a simple expression for the term rank of each matrix in \mathcal{A}^* , namely that it is equal to the upper bound established in Lemma 4.

Theorem 5 *Let A be an ASM in \mathcal{A}^* of order $n \geq 3$. Then*

$$\rho(A) = \rho(A^+) = n_r^-(A) + n_c^-(A). \tag{6}$$

Moreover, a minimum line cover of A consists of all rows and columns that contain at least one entry equal to -1 , and the corresponding set of lines in A^+ is a minimum line cover of A^+ .

Proof We prove the result by induction on the number k of D_3 -extensions. If $k = 1$, then $A = D_3$, and the result trivially holds. Assume that the theorem holds for up to k D_3 -extensions, for some k . Let A be obtained from J_1 by $k + 1$ successive D_3 -extensions, and let A' be the matrix obtained by the first k of these D_3 -extensions. Then

$$\tau(A') = \rho(A') = \rho((A')^+) = n_r^-(A') + n_c^-(A').$$

So, A is obtained by a D_3 -extension of A' . Moreover,

$$\tau(A') + 1 \leq \tau(A^+) \leq \tau(A') + 2 \tag{7}$$

as the added nonzeros (in A) can be covered by at most two lines, and at least one line is needed. There are two possible cases. Let the insertion position be (i, j) .

*Case 1 : The extension is of type (*a).* Assume the extension is as described in the first part of (*a); the other case is similar. Then

$$n_r^-(A) = n_r^-(A'), \quad \text{and} \quad n_c^-(A) = n_c^-(A') + 1.$$

Consider the minimum line cover of A' consisting of all rows and columns of A' that contain a -1 . The corresponding lines in A^+ cover all 1's of A^+ except two $+1$ entries in the same column (that were added in producing A). Therefore, by adding this column, one gets a line cover of A^+ with $\tau(A') + 1$ lines, and, by (7), that must be a minimum line cover of A^+ . The corresponding lines in A also cover all the negative entries, so it is a minimum line cover of A . Thus,

$$\rho(A^+) = \tau(A^+) = \tau(A) = \tau(A') + 1 = (n_r^-(A') + n_c^-(A')) + 1 = n_r^-(A) + n_c^-(A).$$

as desired.

*Case 2 : The extension is of type (*b).* Assume the extension is as described in the first sentence of (*b); the other situation is treated similarly. (See the matrix A_2 in Example 3 where $(i, j) = (2, 3)$.) Then

$$n_r^-(A) = n_r^-(A') + 1, \quad \text{and} \quad n_c^-(A) = n_c^-(A') + 1.$$

Let $p = \tau(A^+)$, so there is a set S of p positions of 1's in A' such that no pair of these positions in S are in the same line. Then S corresponds to a matching in the bipartite graph G corresponding to A^+ .

Assume first that $(i, j) \notin S$. Then S together with two of the $+1$'s that were added in the construction of A is a matching of size $p + 2$ in G . Otherwise, S contains (i, j) . Then A^+ contains another 1 in either row i or column j , but not both, such that this 1 is in a unit row or column. Let S' be obtained from S by adding the position of such a 1 and removing position (i, j) . Thus, S' is also a matching in G and $|S'| = |S|$. Then we can proceed as just described and add two positions and obtain a matching of size $p + 2$ in G . Therefore

$$\rho(A^+) = \tau(A') + 2 = (n_r^-(A') + n_c^-(A')) + 2 = n_r^-(A) + n_c^-(A)$$

as desired. Moreover, we obtain a line cover of size equal to $\rho(A^+)$ by taking the line cover of A' and add the two lines containing the added -1 of A . By induction, this line cover consists of all rows and columns that contain a -1 in A . The theorem now follows. □

In Example 3, a minimum line cover for B_1 consists of rows 2 and 4, and column 3. A minimum line cover for B_2 consists of rows 2 and 4, and columns 3 and 4. Matchings of corresponding sizes are easy to find.

We remark that the result above also holds for some more general D_3 -extensions than those specified in (*b).

The ASM class \mathcal{A}^* studied above has the special property that

$$\rho(A) = \rho(A^+)$$

for every $A \in \mathcal{A}^*$. In other words, covering the negative entries as well does not require any more lines than just covering the positive entries. We now give an example which shows that this property does not hold for all ASMs.

Example 4 Let k be a positive integer, and consider the matrix $B^{(k)}$ given by

$$B^{(k)} = \begin{bmatrix} & v_1 & & & \\ & & + & & v_2 \\ & & + & - & + \\ v_4 & & + & & \\ & & & & v_3 \end{bmatrix},$$

where each v_i is a vector with $p = 2k$ components; here v_2 and v_4 are row vectors while v_1 and v_3 are column vectors. The components of each v_i alternate between $+1$ and -1 such that (i) the rows in $B^{(k)}$ corresponding to v_2 and v_4 are alternating, starting and ending with $+1$, and (ii) the columns in $B^{(k)}$ corresponding to v_1 and v_3 are alternating, starting and ending with $+1$. Now we extend $B^{(k)}$ into an ASM $A^{(k)}$ by adding rows and columns containing a single nonzero, a $+1$, in suitable positions. For instance, for every -1 in v_2 we introduce two new rows, one on top and one

at the bottom of the present matrix, and containing a +1 in the same column as the mentioned -1.

Let C denote the column of $A^{(k)}$ containing v_1 . Moreover, let S be the set of positions of the nonzeros in C as well as the +1's in the rows of the -1's in column C . Let L be a line cover of $A^{(k)}$. In particular, L covers S and there are two possibilities:

Case 1: L contains C . Then the set S may be covered by column C and one row for each -1 in v_1 , so $k + 1$ lines all together, and this is the minimum number of lines to cover S .

Case 2: L does not contain C . Then the set S can not be covered by less than $2k + 1$ lines, and this minimum is attained when we use the $2k + 1$ rows that contain the nonzeros of column C .

It follows that any minimum line cover L of $A^{(k)}$ must contain the column C . Similarly, one shows that L must contain the column corresponding to v_3 and each of the rows corresponding to v_1 and v_2 . It follows from this that

$$|L| = 4(k + 1) + 1 = 4k + 5.$$

Here the additional 1 is due to the fact that we must cover the -1 in the center of $B^{(k)}$. Thus, $\rho(A^{(k)}) = 4k + 5$. It also follows from this discussion that there is a line cover of $(A^{(k)})^+$ of cardinality $4k + 4$, because we do not need to cover the -1 in the center of $B^{(k)}$. In fact,

$$\rho((A^{(k)})^+) = 4k + 4 < 4k + 5 = \rho(A^{(k)}).$$

Thus we have a class of ASMs A for which $\rho(A^+) < \rho(A)$. □

4 Sum-matrices and primary sum-sequences of permutations

Let P be an $n \times n$ permutation matrix. Then clearly each row of $\Sigma(P)$ contains exactly one increase in a column that did not contain an increase in a previous row, and this increase determines where the 1 in a row occurs (in P). For example, consider the permutation (5, 2, 7, 4, 1, 6, 3) with corresponding 7×7 permutation matrix P given below:

$$P = \begin{bmatrix} & & & 1 & & & \\ & 1 & & & & & \\ & & & & & & 1 \\ & & & & & & \\ & & & 1 & & & \\ 1 & & & & & & \\ & & & & & & 1 \\ & & & & & & \\ & & 1 & & & & \end{bmatrix} \rightarrow \Sigma(P) = \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 2 & 2 & 2 \\ 0 & 1 & 1 & 1 & 2 & 2 & 3 \\ 0 & 1 & 1 & 2 & 3 & 3 & 4 \\ 1 & 2 & 2 & 3 & 4 & 4 & 5 \\ 1 & 2 & 2 & 3 & 4 & 5 & 6 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 \end{bmatrix}. \tag{8}$$

where we have shaded the cells where an increase first occurs and then those above it. As we see from (8), the set of shaded columns form a nested sequence of subsets of $\{1, 2, 3, 4, 5, 6, 7\}$, from the top row down to the corresponding positions of the 1's of




Fig. 1 Hasse diagram of (\mathcal{S}_3, \leq_B) with primary sum-sequences in brackets

P:

$$\emptyset = X_0 \subset X_1 \subset X_2 \subset \dots \subset X_7 \text{ where } |X_k| = k \text{ for } k = 1, 2, \dots, 7 :$$

$$X_1 = \{5\}, X_2 = \{2, 5\}, X_3 = \{2, 5, 7\}, X_4 = \{2, 4, 5, 7\}, X_5 = \{1, 2, 4, 5, 7\},$$

$$X_6 = \{1, 2, 4, 5, 6, 7\}, X_7 = \{1, 2, 3, 4, 5, 6, 7\}.$$

This nested property holds in general. So the matrix $\Sigma(P)$ can be said to represent the usual way in which a permutation of $\{1, 2, \dots, n\}$ is identified as a saturated chain from \emptyset to $\{1, 2, \dots, n\}$ in the lattice of subsets of $\{1, 2, \dots, n\}$ ordered by set-inclusion. We define the *nest* $\mathcal{N}(P)$ of the permutation matrix P to be this nested sequence $X_0 \subset X_1 \subset X_2 \subset \dots \subset X_n = \{1, 2, \dots, n\}$ where $|X_k| = k$ for $k = 0, 1, \dots, n$. By the above discussion the $k \times k$ submatrix of P given by $P[\{1, 2, \dots, k\}, X_k]$ is a permutation matrix.

Let $\pi = (\pi_1, \pi_2, \dots, \pi_n)$ be a permutation of $\{1, 2, \dots, n\}$ with corresponding $n \times n$ permutation matrix P . We define the *primary sum-sequence* $\chi(P) = \chi(\pi)$ of P and π to be the sequence (c_1, c_2, \dots, c_n) where c_k equals the (k, π_k) entry of the sum-matrix $\Sigma(P)$ for $k = 1, 2, \dots, n$. Thus c_k equals the number of 1's of P in its leading $k \times \pi_k$ submatrix (which therefore contains a 1 in its lower right corner), equivalently, the rank of this submatrix. In terms of the permutation π ,

$$c_k = |\{i \leq k : \pi_i \leq \pi_k\}|.$$

Thus we have a mapping $\chi_n : \mathcal{P}_n \rightarrow \mathcal{C}_n$ where \mathcal{C}_n is the set of all sequences of length n with entries in $\{1, 2, \dots, n\}$. We have $\chi(t_n) = \chi(I_n) = (1, 2, \dots, n)$ and $\chi(\zeta_n) = \chi(L_n) = (1, 1, \dots, 1)$ where $t_n = (1, 2, \dots, n)$ and $\zeta_n = (n, n - 1, \dots, 1)$ with corresponding permutation matrix L_n . In Fig. 1 we give the Hasse diagram of (\mathcal{S}_3, \leq_B) with the primary sum-sequences given in brackets.

Example 5 Let $n = 9$ and let $\pi = (3, 7, 2, 4, 9, 1, 8, 5, 6)$. Then computation shows that the primary sum-sequence of π is $\chi(\pi) = (1, 2, 1, 3, 5, 1, 6, 5, 6)$. \square

Theorem 6 Let n be a positive integer. The mapping $\chi_n : \mathcal{P}_n \rightarrow \mathcal{C}_n$ is injective, that is, a permutation (matrix) is determined by its primary sum-sequence.

Proof Let P be an $n \times n$ permutation matrix and let $\chi(P) = (c_1, c_2, \dots, c_n)$. The following recursive algorithm shows how to reconstruct P from $\chi(P)$.

- Begin with an $n \times n$ array X in which every position is empty.
- Consider c_n . This is computed from the 1 in the last row of P . Thus c_n determines which column the 1 in row n of P is located, namely column c_n . Put a 1 in position (n, c_n) of X and a 0 in all other positions of row n .
- Now consider c_{n-1} . Ignoring column c_n and using the fact that the $(n-1)$ remaining columns of P contain a 1 in rows $1, 2, \dots, n-1$, c_{n-1} determines which column t the 1 in row $(n-1)$ is. Put a 1 in row $n-1$ of X in this column t and a 0 in all other positions of X in row $n-1$.
- Continue like this to construct a unique $(0, 1)$ -matrix, and this matrix is P .


\square

The proof above implies that when $\chi(\pi) = (c_1, c_2, \dots, c_n)$, then $\pi = (\pi_1, \pi_2, \dots, \pi_n)$ may be expressed in terms of (c_1, c_2, \dots, c_n) as follows

$$\pi_k = \min\{s : s \geq c_k + |\{l > k : c_l \leq c_k\}|, s \neq \pi_l (l > k)\} \quad (k \leq n). \tag{9}$$

Theorem 6 also follows from the facts: (i) there is a bijection between the set of $n \times n$ permutation matrices and the set $\Sigma(\mathcal{P}_n)$ of their corresponding sum-matrices, and (ii) these sum-matrices are uniquely determined by their increases in rows that are not increases in previous rows. Thus the primary sum-sequence determines the permutation matrix.

Example 6 Let $c = (1, 2, 1, 4, 2)$. Then using the procedure in Theorem 6, or the expression (9), we get



Thus $c = (1, 2, 1, 4, 2)$ is the primary sum-sequence of a permutation matrix. \square

It follows from Theorem 6 that if P is an $n \times n$ permutation matrix corresponding to a permutation $\pi = (\pi_1, \pi_2, \dots, \pi_n)$, then the sum-matrix $\Sigma(P) = [\sigma_{ij}]$ is determined by its values at the positions corresponding to the positions of the 1's of P , that is, by the sequence $(\sigma_{i, \pi_i} : 1 \leq i \leq n)$. Now consider two $n \times n$ permutation matrices P and Q corresponding to permutations π and τ with primary sum-sequences $\chi(P)$ and $\chi(Q)$, respectively.

Example 7 Let $n = 4$ and consider the two permutations $\pi_1 = (2, 4, 1, 3)$ and $\pi_2 = (2, 4, 3, 1)$ where $\pi_1 \leq_B \pi_2$. Then $\chi(\pi_1) = (1, 2, 1, 3)$ and $\chi(\pi_2) = (1, 2, 2, 1)$. \square

Lemma 5 Let P_1 and P_2 be two $n \times n$ permutation matrices such that P_1 covers P_2 in the Bruhat order. Then there exists k and l with $1 \leq k < l \leq n$ and a nonnegative integer r such that $\chi(P_2) = (b_1, b_2, \dots, b_n)$ is obtained from $\chi(P_1) = (a_1, a_2, \dots, a_n)$ by decreasing b_k by r and increasing b_l by $r + 1$.

Proof Let $\pi_1 = (i_1, i_2, \dots, i_n)$ and $\pi_2 = (j_1, j_2, \dots, j_n)$ be the permutations corresponding to P_1 and P_2 , respectively. Then π_1 covers π_2 in the Bruhat order so that π_1 has exactly one more inversion than π_2 . Hence there exists k and l with $k < l$ such that $i_k > i_l$ where for each t with $k < t < l, i_t > i_l$ or $i_t < i_k$ and π_2 is obtained from π_1 by switching j_k and j_l . It is easy to check that $\chi(P_2)$ is obtained from $\chi(P_1)$ by decreasing a_l by some nonnegative integer r and increasing a_k by $r + 1$. \square

If $a = (a_1, a_2, \dots, a_n)$ and $b = (b_1, b_2, \dots, b_n)$ are two arbitrary sequences of nonnegative integers, then b is obtained from a by a *pseudoswitch*, denoted as $a \vdash b$, provided that there exists integers $k < l$ and a nonnegative integer r such that b is obtained from a by decreasing a_k by r and increasing a_l by $r + 1$. Denote the sum of the entries of a vector x by $\sigma(x)$. If $a \vdash b, \sigma(b) = \sigma(a) + 1$. The *level* of a permutation (matrix) in the Bruhat order is its number of inversions.

Theorem 7 Let P_1 and P_2 be two $n \times n$ permutation matrices where P_1 is at level k and P_2 is at level l in the poset (\mathcal{P}_n, \leq_B) where $0 \leq k < l \leq \binom{n}{2}$. Then $P_1 \leq_B P_2$ if and only if $\chi(P_1)$ can be obtained from $\chi(P_2)$ by a sequence of $(l - k)$ pseudoswitches.

Proof This theorem is an immediate consequence of Lemma 5 and the fact that if π_1 and π_2 are two permutations of $\{1, 2, \dots, n\}$ such that $\pi_2 \leq_B \pi_1$, then π_2 can be obtained from π_1 by a sequence of $\sigma(\chi(P_1)) - \sigma(\chi(P_2))$ transforms each of which increases the number of inversions by 1, that is, increases the level by 1 in (\mathcal{P}_n, \leq_B) . \square

Example 8 In the poset (\mathcal{S}_4, \leq_B) we have the saturated chain of permutations (written compactly)

$$1234, 1324, 3124, 3142, 3412, 4312, 4321,$$

with corresponding primary sum-sequences (also written compactly)

$$1234, 1133, 1124, 1132, 1212, 1112, 1111.$$

\square

The next result characterizes the primary sum-sequences of permutation matrices.

Theorem 8 *The set $\chi(\mathcal{P}_n)$ of primary sum-sequences of $n \times n$ permutation matrices P is the set of integral vectors $c = (c_1, c_2, \dots, c_n)$ satisfying*

$$1 \leq c_i \leq i \quad (i \leq n). \tag{10}$$

Proof Assume first that $(c_1, c_2, \dots, c_n) = \chi(P)$ for some permutation matrix P with corresponding permutation (j_1, j_2, \dots, j_n) . By definition, for each $k \leq n$ we have $c_k \geq 1$. Since the first k rows of P up to column j_k contain at most k 1’s, we have that $c_k \leq k$.

We prove the converse by induction on n . Assume the result holds for smaller values than n , and let $c = (c_1, c_2, \dots, c_n)$ be an integral vector satisfying (10). Let $k = c_n$. So $1 \leq k \leq n$. By induction there exists an $(n - 1) \times (n - 1)$ permutation matrix P' such that

$$\chi(P') = (c_1, c_2, \dots, c_{n-1}).$$

Let P be the matrix obtained from P' by adding a row, as row n , and a column, as column k , and place a 1 in position (n, k) of P . Then, clearly, P is a permutation matrix of order n . Moreover, $(\chi(P))_i = (\chi(P'))_i = c_i$ for $i \leq n - 1$, and $(\chi(P))_n = k$ since the previous $k - 1$ columns each have a 1 in one of the first $n - 1$ rows. Thus, $\chi(P) = c$, as desired. □

We remark that Theorem 6 is also a consequence of Theorem 8, as the set of integral vectors satisfying (10) has cardinality $1 \cdot 2 \cdot \dots \cdot n = n!$. Thus the map $P \rightarrow \chi(P)$ is surjective between two sets of the same size, and it is therefore also injective.

Let P be an $n \times n$ permutation matrix with primary sum-sequence $\chi(P) = (c_1, c_2, \dots, c_n)$. The corresponding $n \times n$ $(0, 1)$ -matrix $C(P)$ with exactly n 1’s where these 1’s are in positions (k, c_k) ($1 \leq k \leq n$) is called the *primary sum-matrix* of P . Let \mathcal{C}_n be the set of primary sum-matrices of $n \times n$ permutation matrices. By Theorem 8, \mathcal{C}_n is the set of all $n \times n$ $(0, 1)$ -matrices with exactly n 1’s where, for each $k = 1, 2, \dots, n$, the 1 in row k is in column $i \leq k$. The primary sum-matrix of the identity ι_n is the identity matrix I_n ; the primary sum-matrix matrix of the anti-identity permutation ζ_n is the matrix whose first column is the all ones vector, and all other columns are zero. It is easy to see that no other primary sum-matrix than the one associated with ι_n is a permutation matrix.

Example 9 Let P be the 5×5 permutation matrix corresponding to the permutation $(3, 5, 2, 1, 4)$. Then its primary sum-sequence equals $(1, 2, 1, 1, 4)$ and its primary sum-matrix is

$$C(P) = \begin{bmatrix} 1 & & & & \\ & 1 & & & \\ 1 & & & & \\ 1 & & & & \\ & & & & 1 \end{bmatrix}.$$

□

The column sum vector of a matrix A is denoted by S_A . Let S be a nonnegative integral vector of length n and define

$$\begin{aligned} \mathcal{C}_n(S) &= \{C \in \mathcal{C}_n : S_C = S\}, \\ \mathcal{P}_n(S) &= \{P \in \mathcal{P}_n : C(P) \in \mathcal{C}_n(S)\} \end{aligned}$$

as the set of primary sum-matrices with column sum vector S and the corresponding set of permutation matrices, respectively.

Theorem 9 *Let $S = (s_1, s_2, \dots, s_n)$ be a nonnegative, integral vector. Then $S = S_C$ for some primary sum-matrix C if and only if*

$$\begin{aligned} s_1 &\geq 1, \\ s_i &\geq 0, & (2 \leq i \leq n) \\ \sum_{j=n-k+1}^n s_j &\leq k, & (1 \leq k < n) \\ \sum_{j=1}^n s_j &= n. \end{aligned} \tag{11}$$

Proof Let $C = [c_{ij}] \in \mathcal{C}_n$. Then $S_C = (s_1, s_2, \dots, s_n)$ is clearly nonnegative. Each row of C contains exactly one 1, so $c_{11} = 1$ and $\sum_j s_j = n$. Moreover, C is lower triangular, so the last k columns cannot contain more than k 1's (again, as each row has exactly one 1) which gives the final inequalities.

Conversely, assume that S satisfies (11). We shall construct a matrix $C \in \mathcal{C}_n(S)$. Initially, let C be the matrix whose first column is all ones, and all other entries are 0. Then, if $s_n = 1$ shift the bottommost 1 in the first column of C to column n . Next, in the updated C , shift the s_{n-1} bottommost 1's in the first column to column $n - 1$ (if $s_{n-1} = 0$, nothing is done). As $s_{n-1} + s_n \leq 2$, C will remain lower triangular. In the k 'th step, for the present C , shift the s_{n-k+1} bottommost 1's in the first column to column $n - k + 1$. As $\sum_{j=n-k+1}^n s_j \leq k$, the new C will be lower triangular, and its column sum vector is clearly S , so $C \in \mathcal{C}(S)$. □

Note that the condition of the theorem is a sort of majorization condition on S . We call the primary sum-matrix constructed in the proof of Theorem 9 the *canonical primary sum-matrix* with column sum S , and it will be denoted by $\hat{C}(S)$. The (unique) corresponding permutation matrix is denoted by $\hat{P}(S)$.

Example 10 Let $n = 7$, and $S = (2, 1, 0, 2, 0, 1, 1)$. Then

$$\hat{C}(S) = \begin{bmatrix} 1 & & & & & & \\ 1 & & & & & & \\ & 1 & & & & & \\ & & 1 & & & & \\ & & & 1 & & & \\ & & & & 1 & & \\ & & & & & 1 & \\ & & & & & & 1 \end{bmatrix}, \hat{P}(S) = \begin{bmatrix} & & & 1 & & & \\ & 1 & & & & & \\ & & 1 & & & & \\ & & & & 1 & & \\ & & & & & 1 & \\ & & & & & & 1 \\ & & & & & & & 1 \end{bmatrix}.$$

□

Let L_2 denote the matrix backward identity matrix of order 2, so

$$L_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

We say that a $(0, 1)$ -matrix has *consecutive ones* in columns if for every column its ones (if any) occur consecutively.

Theorem 10 *Assume that S satisfies the conditions of Theorem 9. Then the following holds:*

- (i) *The canonical matrix $\hat{C}(S)$ has consecutive ones in columns, and it does not contain any submatrix equal to L_2 .*
- (ii) *$\hat{C}(S) \preceq_B C$ for each matrix $C \in \mathcal{C}_n(S)$, i.e., $\hat{C}(S)$ is the least element in the poset $(\mathcal{C}(S), \preceq_B)$.*

Proof The construction of $\hat{C}(S) = [\hat{c}_{ij}]$ implies that the ones in this matrix are consecutive in each column, and that $\hat{C}(S)$ has a staircase pattern. Therefore, it has no submatrix equal to L_2 .

Let $C = [c_{ij}] \in \mathcal{C}_n(S)$ and $C \neq \hat{C}(S)$. Let \bar{C} be the $(0, 1)$ -matrix whose first column is the all ones vector, and all other entries are 0. Then C may be obtained from \bar{C} by suitable shifting of 1’s to the right in every row. If, for $i = n, n - 1, \dots, 1$, the 1 in row i is moved to the maximal column index j , then C would be equal to $\hat{C}(S)$, so this is not possible. Therefore, there is an i , and $j < k$, such that $c_{ij} = 1$ and $\hat{c}_{ik} = 1$, and we choose i largest possible with this property. Since column k of both $\hat{C}(S)$ and C have the same sum, there must exist $i' < i$, such that $c_{i'j} = 1$ and $\hat{c}_{ik} = 0$. This means that the submatrix of C consisting of rows i and i' and columns j and k equals L_2 .

Let C' be the matrix obtained from C by replacing the mentioned submatrix L_2 by I_2 . Then $C' \preceq_B C$. Now, if $C' = \hat{C}(S)$, we are done. Otherwise, we repeat the process above with C replaced by C' . It follows that $\hat{C}(S)$ is a least element in the poset $(\mathcal{C}(S), \preceq_B)$. □


One may ask if part (ii) of Theorem 10 can be extended to a similar statement for the Bruhat order on the corresponding permutation matrices. This is not the case, as the next example shows.

Example 10 cont. Let again $n = 7$ and $S = (2, 1, 0, 2, 0, 1, 1)$. Another matrix C in $\mathcal{C}(S)$ and the corresponding permutation matrix P are given by

$$C = \begin{bmatrix} 1 & & & & & & \\ & 1 & & & & & \\ 1 & & & & & & \\ & & & 1 & & & \\ & & & & 1 & & \\ & & & & & & 1 \\ & & & & & & & 1 \end{bmatrix}, P = \begin{bmatrix} 1 & & & & & & \\ & 1 & & & & & \\ 1 & & & & & & \\ & & & 1 & & & \\ & & & & 1 & & \\ & & & & & & 1 \\ & & & & & & & 1 \end{bmatrix}.$$

Then $(\Sigma(P))_{12} = 1, (\Sigma(P))_{21} = 0$ while $(\Sigma(\hat{P}(S)))_{12} = 0, (\Sigma(\hat{P}(S)))_{21} = 1$. So $\hat{P}(S) \not\stackrel{B}{\sim} P$ and $P \not\stackrel{B}{\sim} \hat{P}(S)$. \square

Remark 1 In [11] Fulton defined what he called the *essential set* of a $n \times n$ permutation matrix P corresponding to a permutation π of $\{1, 2, \dots, n\}$ as follows: Think of P as an $n \times n$ array of squares with one dot in each row and column and all other squares empty (corresponding to the 1's and 0's of P). For each dot in P shade all the squares from the dot and eastwards and from the dot and southwards leaving the *diagram* of unshaded squares of P . The *essential set* of P is the set of southeast corners of the connected components of the unshaded squares of the diagram. For example, we have



where the essential set consists of the squares with a \star which are then replaced by the number of \bullet 's in the northwest submatrix they determine (equivalently, its rank).

Fulton shows that a permutation matrix is determined by these rank numbers and their locations, the *ranked essential set* and that, in general, none of these can be omitted. Notice that these rank numbers are entries of the sum-matrix of the permutation matrix P , corresponding to certain zeros of P . Thus these entries (with their locations) determine the permutation matrix P .

In [10] an algorithm is given that determines its ranked essential set. In [10] it is shown that a permutation matrix is determined by the rank function on a subset of its essential set called its *core* where the core has size at most n . In [9] it is proved that the average size of the essential set of an $n \times n$ permutation matrix is asymptotic to $\frac{n^2}{36}$. \square

We return to the notion of primary sum-sequence of a permutation matrix, and the algorithm given in the proof of Theorem 6. The following example implies that there may be no analogue of primary sum-sequence for ASMs.

Example 11 Consider the two 6×6 ASMs

$$A_1 = \begin{bmatrix} & & & 1 & & \\ & & 1 & & & \\ & 1 & -1 & & 1 & \\ 1 & & & -1 & & 1 \\ & & 1 & & & \\ & & & & 1 & \end{bmatrix} \quad \text{and} \quad A_2 = \begin{bmatrix} & & & 1 & & \\ & & 1 & & & \\ & 1 & & -1 & 1 & \\ 1 & & -1 & & & 1 \\ & & & 1 & & \\ & & & & 1 & \end{bmatrix}.$$

Their sum-matrices are

$$\Sigma(A_1) = \begin{bmatrix} 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 2 & 2 & 2 \\ 0 & 1 & 1 & 2 & 3 & 3 \\ 1 & 2 & 2 & 2 & 3 & 4 \\ 1 & 2 & 3 & 3 & 4 & 5 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{bmatrix} \quad \text{and} \quad \Sigma(A_2) = \begin{bmatrix} 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 2 & 2 & 2 \\ 0 & 1 & 2 & 2 & 3 & 3 \\ 1 & 2 & 2 & 2 & 3 & 4 \\ 1 & 2 & 3 & 3 & 4 & 5 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{bmatrix}.$$

Thus the sum-matrices $\Sigma(A_1)$ and $\Sigma(A_2)$ differ only in the the (3, 3)-entry shaded above. Hence if we try to mimic the algorithm given to determine a permutation matrix from its primary sum-sequence (in which a row is determined by the rows below it), it will fail when we transition from rows 6, 5, and 4 to row 3. \square

The notion of nests can be generalized to ASMs but, because of the presence of -1 's, the situation is more complicated.

Example 12 Consider the 8×8 ASM

$$A = \begin{bmatrix} & & & 1 & & & & \\ & 1 & -1 & & & & 1 & \\ & & & & 1 & & & \\ 1 & -1 & 1 & & & & & \\ & 1 & -1 & & 1 & & & \\ & & 1 & -1 & & & 1 & \\ & & & & & & & 1 \\ & & & & 1 & & & \end{bmatrix}$$

where

$$\Sigma(A) = \begin{bmatrix} 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 & 2 & 2 & 2 \\ 0 & 1 & 1 & 2 & 2 & 3 & 3 & 3 \\ 1 & 1 & 2 & 3 & 3 & 4 & 4 & 4 \\ 1 & 2 & 2 & 3 & 4 & 5 & 5 & 5 \\ 1 & 2 & 3 & 3 & 4 & 5 & 6 & 6 \\ 1 & 2 & 3 & 3 & 4 & 5 & 6 & 7 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \end{bmatrix},$$

and the increases in each row (here we use rows rather than columns as we did for permutation matrices, and think of an initial column of all 0's) are indicated again by shading the corresponding cells. For any ASM A , let $\Sigma(A)^*$ be the (0, 1)-matrix obtained from $\Sigma(A)$ by replacing each positive entry with a 1. Thus each row of

$\Sigma(A)^*$ records the columns in which an increase occurs. For this example we have

$$\Sigma(A)^* = \begin{bmatrix} | & | & 1 & | & | & | & | \\ | & 1 & | & | & | & 1 & | \\ | & 1 & | & 1 & | & 1 & | \\ | & 1 & | & 1 & | & 1 & | \\ | & 1 & | & 1 & | & 1 & | \\ | & 1 & | & 1 & | & 1 & | \\ | & 1 & | & 1 & | & 1 & | \\ | & 1 & | & 1 & | & 1 & | \\ | & 1 & | & 1 & | & 1 & | \\ | & 1 & | & 1 & | & 1 & | \end{bmatrix}.$$

Let $\alpha_1, \alpha_2, \dots, \alpha_8$ be the rows of $\Sigma(A)^*$ and define $\alpha_0 = (0, 0, 0, 0, 0, 0, 0, 0)$. So for $i = 1, 2, \dots, 8$, α_i has its ones in the columns of each increase. Define $\alpha_0 = (0, 0, \dots, 0)$. Then $\alpha_i - \alpha_{i-1}$ for $i = 1, 2, \dots, 8$ are the rows of the ASM A . Moreover, $\alpha_1, \alpha_2, \dots, \alpha_8$ defines a saturated chain in the partially ordered set defined by Terwilliger [13] who shows that the saturated chains are in a one-to-one correspondence with the ASMs. This partially ordered set consists of all n -tuples of 0's and 1's with partial order defined by: $(a_1, a_2, \dots, a_n) \leq (b_1, b_2, \dots, b_n)$ if and only if the difference sequence $(b_1 - a_1, b_2 - a_2, \dots, b_n - a_n)$ is a $(0, \pm 1)$ -sequence. \square

Example 13 Consider the diamond ASM D_7 :

$$D_7 = \begin{bmatrix} | & | & | & 1 & | & | & | \\ | & | & 1 & -1 & | & 1 & | \\ | & 1 & -1 & | & 1 & -1 & | & 1 \\ | & 1 & -1 & | & 1 & -1 & | & 1 & 1 \\ | & 1 & -1 & | & 1 & -1 & | & 1 \\ | & | & | & 1 & -1 & | & 1 & | \\ | & | & | & | & | & | & 1 & | \end{bmatrix}$$

where

$$\Sigma(D_7) = \begin{bmatrix} 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 2 & 2 & 2 \\ 0 & 1 & 1 & 2 & 2 & 3 & 3 \\ 1 & 1 & 2 & 2 & 3 & 3 & 4 \\ 1 & 2 & 2 & 3 & 3 & 4 & 4 \\ 1 & 2 & 3 & 3 & 4 & 5 & 6 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 \end{bmatrix} \rightarrow \Sigma(D_7^*) = \begin{bmatrix} | & | & | & 1 & | & | & | \\ | & | & 1 & | & 1 & | & | \\ | & 1 & | & 1 & | & 1 & | \\ | & 1 & | & 1 & | & 1 & | \\ | & 1 & | & 1 & | & 1 & | \\ | & 1 & | & 1 & | & 1 & | \\ | & 1 & | & 1 & | & 1 & | \end{bmatrix}.$$

Then D_7 corresponds to the saturated chain

$$(0, 0, 0, 1, 0, 0, 0), (0, 0, 1, 0, 1, 0, 0), (0, 1, 0, 1, 0, 1, 0), (1, 0, 1, 0, 1, 0, 1), (1, 1, 0, 1, 0, 1, 0), (1, 1, 1, 0, 1, 1, 1), (1, 1, 1, 1, 1, 1, 1).$$

□

5 Bruhat Lattice Properties of ASMs

If A_1 and A_2 are two $n \times n$ ASMs, then $\text{GLB}_{\mathcal{A}_n}\{A_1, A_2\}$ denotes the *greatest lower bound* of A_1 and A_2 in the lattice $(\mathcal{A}_n, \preceq_B)$. This greatest lower bound is the ASM whose sum matrix is the $n \times n$ matrix in which each entry equals the maximum of the corresponding entries of A_1 and A_2 . The next example shows that $\text{GLB}_{\mathcal{A}_n}\{A_1, A_2\}$ may be a permutation matrix even though neither A_1 nor A_2 are permutation matrices.

Example 14 Let

$$A_1 = \begin{bmatrix} 1 & & & \\ & & 1 & \\ & 1 & -1 & 1 \\ & & & 1 \end{bmatrix} \quad \text{and} \quad A_2 = \begin{bmatrix} & 1 & & \\ 1 & -1 & 1 & \\ & 1 & & \\ & & & 1 \end{bmatrix}.$$

Then

$$\Sigma(A_1) = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 2 & 2 \\ 1 & 2 & 2 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix} \quad \text{and} \quad \Sigma(A_2) = \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 1 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}.$$

We have

$$\Sigma(\text{GLB}_{\mathcal{A}_n}(A_1, A_2)) = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix},$$

and hence the increases correspond to a nested sequence and give the permutation matrix

$$\begin{bmatrix} 1 & & & \\ & & 1 & \\ & 1 & & \\ & & & 1 \end{bmatrix}.$$

A permutation matrix and an ASM which is not a permutation matrix may have a greatest lower bound equal to a permutation matrix. For instance,

$$\begin{bmatrix} 1 & & & \\ 1 & & & \\ & & 1 & \\ & & & 1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & & & \\ & & 1 & \\ & -1 & 1 & -1 \\ & & 1 & \end{bmatrix}$$

has $\text{GLB}_{\mathcal{A}_n}$ equal to

$$\begin{bmatrix} 1 & & & \\ & 1 & & \\ & & & 1 \\ & & & 1 \end{bmatrix}.$$

□

Example 15 Let $n = 4$ and $\pi_1 = (2431)$, $\pi_2 = (3241)$. Then the $\text{GLB}\{\pi_1, \pi_2\} = \tau = (2341)$. We have

$$\Sigma(\pi_1) = \begin{bmatrix} 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 2 \\ 0 & 1 & 2 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}, \quad \Sigma(\pi_2) = \begin{bmatrix} 0 & 0 & 1 & 1 \\ 0 & 1 & 2 & 2 \\ 0 & 1 & 2 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix},$$

and

$$\Sigma(\tau) = \begin{bmatrix} 0 & 1 & 1 & 1 \\ 0 & 1 & 2 & 2 \\ 0 & 1 & 2 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}.$$

In this case, both π_1 and π_2 cover τ .

Another example is $\pi_1 = (3421)$, $\pi_2 = (4312)$ where the $\text{GLB}\{\pi_1, \pi_2\} = \tau = (3412)$. We have

$$\Sigma(\pi_1) = \begin{bmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 1 & 2 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}, \quad \Sigma(\pi_2) = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 2 \\ 1 & 1 & 2 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}$$

and

$$\Sigma(\tau) = \begin{bmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 2 \\ 1 & 1 & 2 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}.$$

In this case, both π_1 and π_2 also cover τ .

□

So, if we take two permutations π_1 and π_2 of $\{1, 2, \dots, n\}$, and form the entrywise maximum of the entries of $\Sigma(\pi_1)$ and $\Sigma(\pi_2)$ to get a matrix $M = \Sigma(\pi_1) \vee \Sigma(\pi_2)$, then M is the sum-matrix of the ASM $\text{GLB}_{\mathcal{A}_n}\{\pi_1, \pi_2\}$. Then $\text{GLB}_{\mathcal{A}_n}\{\pi_1, \pi_2\}$ is not a permutation matrix if and only if in some row of M , there are two or more increases that were not increases in the previous row, see further discussion below.

Now consider the two permutations $\pi_1 = (3241)$ and $\pi_2 = (4132)$ with $\text{GLB}_{\mathcal{A}_n}\{\pi_1, \pi_2\} = \tau = (3142)$. We have

$$\Sigma(\pi_1) = \left[\begin{array}{c|c|c|c} 0 & 0 & 1 & 1 \\ \hline 0 & 1 & 2 & 2 \\ \hline 0 & 1 & 2 & 3 \\ \hline 1 & 2 & 3 & 4 \end{array} \right], \quad \Sigma(\pi_2) = \left[\begin{array}{c|c|c|c} 0 & 0 & 0 & 1 \\ \hline 1 & 1 & 1 & 2 \\ \hline 1 & 1 & 2 & 3 \\ \hline 1 & 2 & 3 & 4 \end{array} \right],$$

and

$$\Sigma(\tau) = \left[\begin{array}{c|c|c|c} 0 & 0 & 1 & 1 \\ \hline 1 & 1 & 2 & 2 \\ \hline 1 & 1 & 2 & 3 \\ \hline 1 & 2 & 3 & 4 \end{array} \right].$$

In this case, neither π_1 nor π_2 covers τ . There is an ASM different from a permutation between π_1 and τ (namely, the ASM corresponding to 13 in the Hasse diagram of (\mathcal{A}_4, \leq_B)), and another ASM different from a permutation between π_2 and τ (the ASM corresponding to 42 in the Hasse diagram of (\mathcal{A}_4, \leq_B)). (Note that e.g., 42 in the diagram denotes a 4×4 ASM with a 1 in position $(4, 2)$ which is uniquely completable to an ASM by using the only 3×3 ASM with a -1).

For small values of n we have enumerated all permutation matrices of order n , and computed the $\text{GLB}_{\mathcal{A}_n}$ of each pair of such matrices. The following table shows how many pairs that have a $\text{GLB}_{\mathcal{A}_n}$ which is a permutation matrix, and this number is denoted by #P. The other pairs have a $\text{GLB}_{\mathcal{A}_n}$ which is an ASM different from a permutation matrix, and their number is denoted by #ASM.

n	$n!$	$n!(n! - 1)/2$	#P	#ASM
2	2	1	1	0
3	6	15	14	1
4	24	276	231	45
5	120	7140	5136	2004
6	720	258840	154385	104455

Next we characterize when the $\text{GLB}_{\mathcal{A}_n}$ of a set of permutation matrices is a permutation matrix. Let P_1, P_2, \dots, P_k be permutation matrices of order n , and let $A = [a_{ij}] = \text{GLB}_{\mathcal{A}_n}\{P_1, P_2, \dots, P_k\}$. Define $\Sigma(A) = [\hat{\sigma}_{ij}]$ and $\Sigma(P_t) = [\sigma_{ij}^{(t)}]$ ($t \leq k$). Thus, $\hat{\sigma}_{ij} = \max_t \sigma_{ij}^{(t)}$ for each i, j . Also define for $1 < i, j < n$

$$\begin{aligned} K_{ij} &= \{t \leq k : \sigma_{ij}^{(t)} = \hat{\sigma}_{ij}\}, \\ I(i, j) &= \{(i, 1), \dots, (i, j - 1)\}, \\ J(i, j) &= \{(1, j), \dots, (i - 1, j)\}. \end{aligned}$$

With this notation, we have the following result.

Proposition 1 Let P_1, P_2, \dots, P_k be permutation matrices of order n , and let $A = \text{GLB}_{\mathcal{A}_n}\{P_1, P_2, \dots, P_k\}$.

Then A is a permutation matrix if and only if there is no position (i, j) , with $1 < i, j < n$, such that (a) for some $s \in K_{i-1, j-1}$ the matrix P_s has a 1 in $I(i, j)$, but no 1 in $J(i, j)$, (b) for some $t \in K_{i-1, j-1}$ the matrix P_t has a 1 in $J(i, j)$, but no 1 in $I(i, j)$, and (c) no P_l ($1 \leq l \leq k$) has a 1 in position (i, j) , or a 1 in both I_1 and in I_2 .

Proof The matrix $A = [a_{ij}]$ is a permutation matrix if and only if there is no position (i, j) , with $1 < i, j < n$, such that $a_{ij} = -1$. Using the properties of $\Sigma(A) = [\hat{\sigma}_{ij}]$, i.e., that $\Sigma(A)$ is nondecreasing in rows and columns, and the increase is at most 1, we see that $a_{ij} = -1$ if and only if, for some integer m ,

$$\hat{\sigma}_{i-1, j-1} = m, \quad \hat{\sigma}_{i-1, j} = \hat{\sigma}_{i, j-1} = \hat{\sigma}_{i, j} = m + 1.$$

But these equations correspond precisely to the statements (a)–(c) in the proposition. □

Example 16 Let $n = 4$ and let

$$P_1 = \left[\begin{array}{c|c|c|c} & 1 & & \\ \hline & & & 1 \\ \hline & & 1 & \\ \hline 1 & & & \end{array} \right] \quad \text{and} \quad P_2 = \left[\begin{array}{c|c|c|c} & & & 1 \\ \hline 1 & & & \\ \hline & & 1 & \\ \hline & 1 & & \end{array} \right].$$

Then

$$\Sigma(P_1) = \left[\begin{array}{c|c|c|c} 0 & 1 & 1 & 1 \\ \hline 0 & 1 & 1 & 2 \\ \hline 0 & 1 & 2 & 3 \\ \hline 1 & 2 & 3 & 4 \end{array} \right] \quad \text{and} \quad \Sigma(P_2) = \left[\begin{array}{c|c|c|c} 0 & 0 & 0 & 1 \\ \hline 1 & 1 & 1 & 2 \\ \hline 1 & 1 & 2 & 3 \\ \hline 1 & 2 & 3 & 4 \end{array} \right].$$

Then

$$\Sigma(\text{GLB}_{\mathcal{A}_n}\{P_1, P_2\}) = \left[\begin{array}{c|c|c|c} 0 & 1 & 1 & 1 \\ \hline 1 & 1 & 1 & 2 \\ \hline 1 & 1 & 2 & 3 \\ \hline 1 & 2 & 3 & 4 \end{array} \right],$$

whose corresponding ASM is

$$\left[\begin{array}{c|c|c|c} 0 & 1 & 0 & 0 \\ \hline 1 & -1 & 0 & 1 \\ \hline 0 & 0 & 1 & 0 \\ \hline 0 & 1 & 0 & 0 \end{array} \right].$$

In view of Proposition 1 consider $i = j = 2$, and $k = 2$. Then $K_{1,1} = \{1, 2\}$ (as $\hat{\sigma}_{11} = 0$) and P_2 satisfies condition (a) as it has a 1 in $I(2, 2) = \{(2, 1)\}$, and P_1

satisfies condition (b) due to the 1 in $I(2, 2) = \{(1, 2)\}$. Moreover, condition (c) holds, so the proposition also shows that $\text{GLB}_{\mathcal{A}_n}\{P_1, P_2\}$ is not a permutation matrix. \square

The next result determines when the greatest lower bound of a set of permutation matrices is equal to the identity matrix I_n .

Theorem 11 *Let P_1, P_2, \dots, P_k be permutation matrices of order n , and let $A = \text{GLB}_{\mathcal{A}_n}\{P_1, P_2, \dots, P_k\}$. Then $A = I_n$ if and only if for each $k = 1, 2, \dots, n - 1$, there exists a $t \leq k$ such that the leading $k \times k$ submatrix of P_t is a permutation matrix.*

Proof Let $\Sigma(A) = [\hat{\sigma}_{ij}]$ and $\Sigma(P_t) = [\sigma_{ij}^{(t)}]$ ($t \leq k$). Assume $A = I_n$. Then $\Sigma(A) = \Sigma(I)$ is symmetric and

$$\hat{\sigma}_{ij} = i \quad (i \leq n, j = i, i + 1, \dots, n). \tag{12}$$

Therefore

$$i = \hat{\sigma}_{ii} = \max_{t \leq k} \sigma_{ii}^{(t)} = \sigma_{ii}^{(s)}$$

for some $s \leq t$. So, the permutation matrix P_s has i ones in its leading $i \times i$ submatrix, and therefore this submatrix must be a permutation matrix.

Conversely, assume that for each $k = 1, 2, \dots, n - 1$ there exists a $t \leq k$ such that the leading $k \times k$ submatrix of P_t is a permutation matrix. This implies that

$$\hat{\sigma}_{ii} = \max_{t \leq k} \sigma_{ii}^{(t)} = i \quad (i \leq n).$$

Since rows and columns of $\Sigma(A)$ are nondecreasing and the last entry in row i , and in column i , is i , it follows that $\Sigma(A)$ satisfies (12). So, $A = I$, as desired. \square

Example 17 Let $n = 4$ and let

$$P_1 = \begin{bmatrix} | & 1 & | & | \\ 1 & | & | & | \\ | & | & 1 & | \\ | & | & | & 1 \end{bmatrix} \quad \text{and} \quad P_2 = \begin{bmatrix} 1 & | & | & | \\ | & | & 1 & | \\ | & | & | & 1 \\ | & 1 & | & | \end{bmatrix}.$$

Then

$$\Sigma(P_1) = \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix} \quad \text{and} \quad \Sigma(P_2) = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 2 & 2 \\ 1 & 1 & 2 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}.$$

Then

$$\Sigma (\text{GLB}_{\mathcal{A}_n} \{P_1, P_2\}) = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix},$$

which gives $\text{GLB}_{\mathcal{A}_n} \{P_1, P_2\} = I$. The leading $k \times k$ submatrix of P_1 is a permutation matrix for $k = 2, 3$ (and 4), while the leading $k \times k$ submatrix of P_2 is a permutation matrix for $k = 1$ (and 4). \square

Another way to formulate Theorem 11 is:

Let P_1, P_2, \dots, P_k be permutation matrices of order n , and let

$$A = \text{GLB}_{\mathcal{A}_n} \{P_1, P_2, \dots, P_k\}.$$

Then $A = I_n$ if and only if for each $k = 1, 2, \dots, n - 1$ there exists a $t \leq k$ such that the k th term in $\chi(P_t)$ equals k , that is, the k th term in $\chi(I_n)$.

Acknowledgements The authors thank a referee for several useful comments.

References

1. Björner, A., Brenti, F.: Combinatorics of Coxeter Groups. Graduate Texts in Mathematics, vol. 231. Springer, New York (2005)
2. Bóna, M.: Combinatorics of Permutations, 2nd edn. CRC Press, Boca Raton (2012)
3. Brualdi, R.A., Dahl, G.: The Bruhat shadow of a permutation matrix, Mathematical Papers in Honour of Eduardo Marques de Sá, O, Azenhas, Duarte, A.L., Queiró, J.F., Santana, A.P., eds., Textos de Matemática Vol. 39, Dep. de Matemática da Univ. de Coimbra, Portugal, pp. 25–38 (2006)
4. Brualdi, R.A., Dahl, G.: Alternating sign matrices, extensions and related cones. Adv. Appl. Math. **86**, 19–49 (2017)
5. Brualdi, R.A., Dahl, G.: Alternating sign matrices and hypermatrices, and a generalization of latin squares. Adv. Appl. Math. **95**, 116–151 (2018)
6. Brualdi, R.A., Kim, H.K.: Completions of alternating sign matrices. Graphs Combin. **31**, 507–522 (2015)
7. Brualdi, R.A., Kiernan, K.P., Meyer, S.A., Schroeder, M.W.: Patterns of alternating sign matrices. Linear Algebra Appl. **438**, 3967–3990 (2013)
8. Brualdi, R.A., Schroeder, M.W.: Alternating sign matrices and their Bruhat order. Discr. Math. **340**, 1996–2019 (2017)
9. Eriksson, K., Linusson, S.: The size of Fulton’s essential set. Electr. J. Combin. **2**, R6 (1995)
10. Eriksson, K., Linusson, S.: Combinatorics of Fulton’s essential set. Duke Math. J. **85**, 61–76 (1996)
11. Fulton, W.: Flags, Schubert polynomials, degeneracy loci, and determinantal formulas. Duke Math. J. **65**, 381–420 (1992)
12. Lascoux, A., Schützenberger, M.P.: Treillis et bases des groupes de Coxeter. Electron. J. Combin. **3**, Research paper 27 (1996)
13. Terwilliger, P.: A poset Φ_n whose maximal chains are in bijection with the $n \times n$ alternating sign matrices. Linear Algebra Appl. **554**, 79–85 (2018)

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.