

Erratum to: Short- and long-term inhibition of cardiac inward-rectifier potassium channel current by an antiarrhythmic drug bepridil

Fangfang Ma^{1,2} · Hiroki Takanari¹ · Kimiko Masuda¹ · Masaki Morishima¹ · Katsushige Ono¹

Published online: 25 February 2016
© Springer Japan 2016

Erratum to: Heart Vessels DOI 10.1007/s00380-015-0762-1

Unfortunately, the caption of Fig. 4 was incorrectly published in the original publication of the article. The corrected caption of Fig. 4 is as follows:

Fig. 4 Long-term effect of bepridil and calmodulin inhibition on I_{K1} . **a** Current traces of I_{K1} in the control (vehicle for 24 h) and after the long-term (24 h) treatment with 10 μ M W-7, 1 μ M bepridil with 10 μ M W-7, and 10 μ M KN93. Outward I_{K1} traces at the potentials of +20 mV at the terminal phase indicated by *box* are shown in an *inset*. **b** I–V relationships constructed by using group data in the control and after treatment with 10 μ M W-7, 1 μ M bepridil with 10 μ M W-7, 20 μ M W-7, and 10 μ M KN93. The current density of I_{K1} at –100 mV was reduced to -8.3 ± 1.2 ,

-8.5 ± 0.7 , -6.3 ± 1.1 and -3.5 ± 0.4 mV, respectively. **c** The slope conductance was reduced to 0.44 ± 0.04 by 10 μ M W-7, 0.45 ± 0.004 by 1 μ M bepridil with 10 μ M W-7, 0.39 ± 0.08 by 20 μ M W-7, and 0.25 ± 0.05 by 10 μ M KN93 at –100 mV. **d** The slope conductance was reduced to 0.03 ± 0.005 by 10 μ M W-7, 0.02 ± 0.003 by 1 μ M bepridil with 10 μ M W-7, 0.03 ± 0.001 by 20 μ M W-7, and 0.03 ± 0.004 by 10 μ M KN93 at +20 mV. # $p < 0.1$, * $p < 0.05$, ** $p < 0.01$ compared with control

The online version of the original article can be found under doi:10.1007/s00380-015-0762-1.

✉ Katsushige Ono
ono@oita-u.ac.jp

¹ Department of Pathophysiology, Oita University School of Medicine, Yufu 879-5593, Oita, Japan

² Department of Cardiology, The 1st Hospital of Hebei Medical University, Shijiazhuang 050031, Hebei, China