

***TMEM106B* protects *C9ORF72* expansion carriers against frontotemporal dementia**

Marka van Blitterswijk · Bianca Mullen · Alexandra M. Nicholson · Kevin F. Bieniek · Michael G. Heckman · Matthew C. Baker · Mariely DeJesus-Hernandez · NiCole A. Finch · Patricia H. Brown · Melissa E. Murray · Ging-Yuek R. Hsiung · Heather Stewart · Anna M. Karydas · Elizabeth Finger · Andrew Kertesz · Eileen H. Bigio · Sandra Weintraub · Marsel Mesulam · Kimmo J. Hatanpaa · Charles L. White III · Michael J. Strong · Thomas G. Beach · Zbigniew K. Wszolek · Carol Lippa · Richard Caselli · Leonard Petrucelli · Keith A. Josephs · Joseph E. Parisi · David S. Knopman · Ronald C. Petersen · Ian R. Mackenzie · William W. Seeley · Lea T. Grinberg · Bruce L. Miller · Kevin B. Boylan · Neill R. Graff-Radford · Bradley F. Boeve · Dennis W. Dickson · Rosa Rademakers

Received: 14 October 2013 / Revised: 17 December 2013 / Accepted: 20 December 2013 / Published online: 3 January 2014
© Springer-Verlag Berlin Heidelberg 2013

Abstract Variants in transmembrane protein 106 B (*TMEM106B*) modify the disease penetrance of frontotemporal dementia (FTD) in carriers of progranulin (*GRN*) mutations. We investigated whether *TMEM106B* is also a genetic modifier of disease in carriers of chromosome 9 open reading frame 72 (*C9ORF72*) expansions. We assessed the genotype of 325 *C9ORF72* expansion carriers (cohort 1), 586 FTD patients lacking *C9ORF72* expansions [with or without motor neuron disease (MND); cohort 2], and a total of 1,302 controls for *TMEM106B* variants (rs3173615 and rs1990622) using MassArray iPLEX and

Taqman genotyping assays. For our primary analysis, we focused on functional variant rs3173615, and employed a recessive genotypic model. In cohort 1, patients with *C9ORF72* expansions showed a significantly reduced frequency of carriers homozygous for the minor allele as compared to controls [11.9 vs. 19.1 %, odds ratio (OR) 0.57, $p = 0.014$; same direction as carriers of *GRN* mutations]. The strongest evidence was provided by FTD patients (OR 0.33, $p = 0.009$) followed by FTD/MND patients (OR 0.38, $p = 0.017$), whereas no significant difference was observed in MND patients (OR 0.85, $p = 0.55$). In cohort 2, the frequency of carriers homozygous for the minor allele was not significantly reduced in patients as compared to controls (OR 0.77, $p = 0.079$); however, a significant reduction was

Electronic supplementary material The online version of this article (doi:10.1007/s00401-013-1240-4) contains supplementary material, which is available to authorized users.

M. van Blitterswijk · B. Mullen · A. M. Nicholson · K. F. Bieniek · M. C. Baker · M. DeJesus-Hernandez · N. A. Finch · P. H. Brown · M. E. Murray · L. Petrucelli · D. W. Dickson · R. Rademakers (✉)
Department of Neuroscience, Mayo Clinic, 4500 San Pablo Road, Jacksonville, FL 32224, USA
e-mail: rademakers.rosa@mayo.edu

M. van Blitterswijk
e-mail: VanBlitterswijk.Marka@mayo.edu

M. G. Heckman
Section of Biostatistics, Mayo Clinic, 4500 San Pablo Road, Jacksonville, FL 32224, USA

G.-Y. R. Hsiung · H. Stewart
Division of Neurology, University of British Columbia, 2211 Wesbrook Mall, Vancouver, BC V6T 2B5, Canada

A. M. Karydas · W. W. Seeley · L. T. Grinberg · B. L. Miller
Department of Neurology, University of California, 500 Parnassus Ave, San Francisco, CA 94143, USA

E. Finger · A. Kertesz
The University of Western Ontario, 1151 Richmond St, London, ON N6A 3K7, Canada

E. H. Bigio · S. Weintraub · M. Mesulam
Cognitive Neurology and Alzheimer's Disease Center, Northwestern University Feinberg School of Medicine, 320 East Superior Street, Chicago, IL 60611, USA

K. J. Hatanpaa · C. L. White III
University of Texas Southwestern Medical Center, 5323 Harry Hines Blvd, Dallas, TX 75390, USA

M. J. Strong
Molecular Brain Research Group, Robarts Research Institute, 100 Perth Drive, London, ON N6A 5K8, Canada

T. G. Beach
Banner Sun Health Research Institute, 10515 W Santa Fe Dr, Sun City, AZ 85351, USA

observed when focusing on those patients with frontotemporal lobar degeneration and TAR DNA-binding protein 43 inclusions (FTLD-TDP; OR 0.26, $p < 0.001$). Our study identifies *TMEM106B* as the first genetic factor modifying disease presentation in *C9ORF72* expansion carriers. Homozygosity for the minor allele protects carriers from developing FTD, but not from developing MND; similar effects are seen in FTLD-TDP patients with yet unknown genetic causes. These new findings show that the protective effects of *TMEM106B* are not confined to carriers of *GRN* mutations and might be relevant for prognostic testing, and as a promising therapeutic target for the entire spectrum of FTLD-TDP.

Keywords *C9ORF72* · *TMEM106B* · Frontotemporal dementia · Motor neuron disease · Amyotrophic lateral sclerosis · Disease modifier

Introduction

Repeat expansions in chromosome 9 open reading frame 72 (*C9ORF72*) are the most common known genetic cause of frontotemporal dementia (FTD), amyotrophic lateral sclerosis (ALS), and a combination of both diseases [FTD with motor neuron disease (FTD/MND)] [7, 21]. Previously, we showed that the repeat length of these expansions varies between tissues, and that repeat lengths in the cerebellum are associated with survival after disease onset [26]. Moreover, we demonstrated that additional mutations in FTD-associated genes [progranulin (*GRN*) and microtubule-associated protein tau (*MAPT*)] might also contribute to the phenotypic variability that is detected in patients with *C9ORF72* repeat expansions [25].

To date, other genetic factors that could act as disease modifiers have not been reported in *C9ORF72* expansion carriers. For this reason, we investigated a promising candidate: transmembrane protein 106 B (*TMEM106B*). Single nucleotide polymorphisms (SNPs) in *TMEM106B* are associated with the risk of frontotemporal lobar degeneration with TAR DNA-binding protein 43 inclusions (FTLD-TDP) [28], which is the predominant pathology in many FTD patients [15, 18]. Interestingly, this association is most prominent in patients with *GRN* mutations; in these patients, two copies of the minor allele of *TMEM106B* SNPs (top SNP rs1990622) appear to reduce disease penetrance, protecting individuals from developing FTD [8, 28]. Variant rs3173615, which is in linkage disequilibrium (LD) with rs1990622, dictates a change from threonine to serine at position 185 (p.T185S) [5, 19, 29]. This variant has been shown to regulate *TMEM106B* protein levels and was proposed as the functional variant underlying the genetic association with FTLD-TDP [19].

In our present study, we assessed the frequency of *TMEM106B* SNPs rs3173615 and rs1990622 in a large cohort of *C9ORF72* expansion carriers ($n = 325$), in an additional cohort of patients with FTD or FTD/MND without *C9ORF72* expansions ($n = 586$), and in control subjects ($n = 1,302$). Our results reveal that variants in *TMEM106B* protect against developing FTD in carriers of *C9ORF72* repeat expansions, and also in patients with currently unknown causes of FTLD-TDP, emphasizing crucial similarities between *C9ORF72*-related FTLD-TDP, *GRN*-related FTLD-TDP, and other causes of FTLD-TDP.

Materials and methods

Subjects

For cohort 1 we included all 325 carriers of *C9ORF72* repeat expansions available to us. These subjects were obtained through North American/Canadian institutions: the Mayo Clinic ($n = 121$), Coriell Research Institute ($n = 71$), University of British Columbia, Canada ($n = 58$), University of California, San Francisco ($n = 38$), Robarts Research Institute ($n = 11$), Northwestern University Feinberg School of Medicine ($n = 9$), Drexel University College of Medicine ($n = 7$), University of Western Ontario, Canada ($n = 7$), and Banner Sun Health Research Institute ($n = 3$). Subjects were clinicopathologically diagnosed with FTD ($n = 86$), FTD/MND ($n = 78$) or MND ($n = 127$), with another diagnosis ($n = 7$; e.g., Alzheimer's disease, alcohol abuse, or behavioral impairment), or were asymptomatic at the time of last evaluation ($n = 27$; age at evaluation 43.6 ± 12.7). Our primary analysis focused on the 260 unrelated probands with FTD ($n = 69$), FTD/MND

Z. K. Wszolek · K. B. Boylan · N. R. Graff-Radford
Department of Neurology, Mayo Clinic, 4500 San Pablo Road,
Jacksonville, FL 32224, USA

C. Lippa
Department of Neurology, Drexel University College
of Medicine, 2900 W Queen Ln, Philadelphia, PA 19129, USA

R. Caselli
Department of Neurology, Mayo Clinic, 5777 E Mayo Blvd,
Phoenix, AZ 85054, USA

K. A. Josephs · J. E. Parisi · D. S. Knopman · R. C. Petersen ·
B. F. Boeve
Department of Neurology, Mayo Clinic, 1216 2nd St SW,
Rochester, MN 55902, USA

I. R. Mackenzie
Department of Pathology and Laboratory Medicine,
University of British Columbia, 2329 W Mall, Vancouver,
BC V6T 1Z4, Canada

($n = 71$) or MND ($n = 120$), and a group of neurologically normal controls of similar age and gender provided by the Mayo Clinic ($n = 376$). In secondary analyses, we also examined the sensitivity of our results by including the 65 remaining expansion carriers who were family members or had received another diagnosis, and by including additional controls from our major sites [$n = 410$; Coriell Research Institute ($n = 264$), University of British Columbia, Canada ($n = 17$), University of California, San Francisco ($n = 116$), and Robarts Research Institute ($n = 13$)]. All *C9ORF72* expansion carriers and control subjects were screened for *TMEM106B* SNPs rs3173615 and rs1990622.

Cohort 2 was previously screened for *TMEM106B* SNP rs1990622 [8, 28]; however, these subjects were investigated before the identification of repeat expansions in *C9ORF72*. Patients in this cohort were selected according to the following criteria: a clinical diagnosis of behavioral variant FTD, semantic dementia or progressive nonfluent aphasia, or a pathologic diagnosis of FTLN-TDP. Based on our current findings, we reanalyzed our genotype data in this cohort after exclusion of patients with *C9ORF72* repeat expansions and *GRN* mutations, and also genotyped all patients and controls for rs3173615. Furthermore, we extended this cohort to include all additional pathologically confirmed FTLN-TDP and FTLN/MND patients without *C9ORF72* repeat expansions or *GRN* mutations ($n = 31$), who were recruited at our Mayo Clinic Brain

Bank since our original study in 2010. In total, cohort 2 comprised 586 patients who were ascertained at the Mayo Clinic ($n = 380$), University of California, San Francisco ($n = 117$), University of Western Ontario, Canada ($n = 31$), Drexel University College of Medicine ($n = 21$), Northwestern University Feinberg School of Medicine ($n = 21$), University of British Columbia, Canada ($n = 10$), and University of Texas Southwestern Medical Center ($n = 6$). All control subjects in cohort 2 ($n = 765$) were obtained from the Mayo Clinic (not included in cohort 1, $n = 516$; included in cohort 1, $n = 133$) and University of California, San Francisco ($n = 116$) [8]. In addition, we performed a joint analysis including the controls from cohort 1 (total number of controls, $n = 1,302$).

Characteristics of cases and controls included in both cohorts are shown in Table 1. All subjects agreed to be in the study, and biological samples were obtained after informed consent with ethical committee approval from the respective institutions.

Genotyping

TMEM106B SNPs rs3173615 and rs1990622 were genotyped in cohort 1 on a Sequenom MassArray iPLEX platform (San Diego, CA) and analyzed with Typer 4.0 software. In cohort 2, rs3173615 and rs1990622 were genotyped using Taqman SNP genotyping assays

Table 1 Characteristics of study cohorts

Group	<i>N</i>	Female gender (%)	Age	Age at onset	FTD only (%)	Pathological diagnosis (%)
Cohort 1—controls and <i>C9ORF72</i> repeat expansion carriers (FTD, FTD/MND, and MND, or other diagnosis)						
Controls	376	173 (46.0)	61.2 ± 10.2 (35–90)	N/A	N/A	N/A
All repeat expansion carriers	325	148 (45.5)	59.2 ± 9.9 (35–90)	56.8 ± 9.2 (34–83)	86 (26.5)	118 (36.3)
FTD, FTD/MND, and MND probands	260	114 (43.8)	59.5 ± 10.0 (35–90)	56.9 ± 9.0 (34–83)	69 (26.5)	107 (41.2)
FTD probands	69	28 (40.6)	63.1 ± 12.3 (35–90)	58.1 ± 9.6 (34–79)	69 (100.0)	40 (58.0)
FTD/MND probands	71	25 (35.2)	60.6 ± 8.5 (39–80)	56.2 ± 9.0 (34–74)	0 (0.0)	51 (71.8)
MND probands	120	61 (50.8)	56.9 ± 8.6 (37–83)	56.5 ± 8.7 (37–83)	0 (0.0)	16 (13.3)
Cohort 2—controls and FTD or FTD/MND patients without <i>C9ORF72</i> repeat expansions or <i>GRN</i> mutations						
Controls	765	363 (47.4)	67.1 ± 10.0 (20–95)	N/A	N/A	N/A
All FTD or FTD/MND patients	586	273 (46.6)	66.8 ± 10.4 (26–99)	62.4 ± 9.8 (26–95)	531 (90.6)	141 (24.1)
All FTD patients	531	251 (47.3)	67.1 ± 10.1 (26–99)	62.7 ± 9.5 (26–86)	531 (100.0)	101 (19.0)
Pathologically diagnosed	101	48 (52.5)	74.0 ± 11.2 (49–99)	64.5 ± 11.1 (39–86)	101 (100.0)	101 (100.0)
Clinically diagnosed	430	203 (47.2)	65.5 ± 9.1 (26–90)	62.4 ± 9.1 (26–85)	430 (100.0)	0 (0.0)

Continuous variables are summarized with the sample mean ± standard deviation (range). For cohort 1, the age provided is age at blood draw in controls, age at onset in clinically diagnosed patients, and age at death in pathologically diagnosed patients. For cohort 2, the age provided is age at blood draw in controls, age at diagnosis in clinically diagnosed patients, and age at death in pathologically diagnosed patients. In cohort 1, information was unavailable regarding age ($n = 42$) and age at onset ($n = 57$). In cohort 2, information was unavailable regarding age at onset ($n = 95$)

(C_27465458_10, C__11171598_10; Invitrogen, Carlsbad, CA) on the 7900HT Fast Real Time PCR system. Genotype calls were made using SDS v2.2 software (Applied Biosystems, Foster City, CA). In control subjects, there was no evidence of a departure from Hardy–Weinberg equilibrium for rs3173615 or rs1990622 in both cohorts (all $p > 0.010$). The two SNPs were in almost complete LD ($r^2 \geq 0.97$), and given this strong LD, for simplicity we focused our analysis on functional variant rs3173615.

Immunohistochemistry

Immunohistochemistry was performed in a blinded fashion on a preliminary group of nine patients who were diagnosed with FTLD-TDP and did not show signs of MND. We included all three patients from the Mayo Clinic Brain Bank for whom fixed tissue was available and who were homozygous for the minor protective *TMEM106B* allele (GG genotype for rs3173615; *C9ORF72* repeat expansion, $n = 1$; without *C9ORF72* repeat expansion, $n = 2$). In addition, we also randomly selected six FTLD-TDP patients without MND who were homozygous for the major risk *TMEM106B* allele (CC genotype for rs3173615; *C9ORF72* repeat expansion, $n = 3$; without *C9ORF72* repeat expansion, $n = 3$), to allow comparisons of TDP-43 burden between patients homozygous for the minor or major allele.

We stained 5- μ m-thick sections from the frontal cortex for TDP-43 (pS409/410, 1:5,000, mouse monoclonal, Cosmobio Co., Tokyo, Japan), and repeat-associated non-ATG translation peptides (C9RANT, Rb5823, 1:5,000, Mayo Clinic) [1]; the latter has recently been shown to be distinctive for *C9ORF72* expansions [1, 17]. Sections were cut from formalin-fixed paraffin-embedded blocks, deparaffinized in xylene, rehydrated in a graded series of ethanol, and washed in distilled water. To process stains, we used DAKO Autostainer Plus (DAKO, Carpinteria, CA) and DAKO EnVision+ System–horseradish peroxidase (diaminobenzidine). Immunostains were subsequently counterstained with Lerner’s hematoxylin, dehydrated, and coverslipped. Stained slides were viewed using an Olympus BX40 microscope (Olympus Corporation of the Americas, Center Valley, PA). TDP-43 pathologic subtype was assigned in accordance with the harmonized criteria (Type A–C) [14]. Neuronal lesion counts were performed in a blinded fashion on layer 2 of the midfrontal gyrus. Counts were performed in six 10 \times microscopic fields randomly selected over the gyrus and averaged for a composite pathologic burden score.

Statistical analysis

Separately for both cohorts, comparisons of *TMEM106B* SNPs with controls were made using logistic regression

models; odds ratios (ORs) and 95 % confidence intervals (CIs) were estimated. Given our previous studies related to *TMEM106B* in carriers of *GRN* mutations [8, 19], we hypothesized that *TMEM106B* SNPs would affect disease penetrance in *C9ORF72* expansion carriers under a recessive genotypic model (presence vs. absence of two copies of the minor allele). For this reason, we utilized this model in our primary analysis, though in secondary analyses we also examined other genotypic models (additive, dominant, and co-dominant).

In cohort 1, models were adjusted for gender, and the primary analysis involved unrelated probands with FTD, FTD/MND, MND, and controls. Comparisons with controls were made for all patients, and also separately for FTD, FTD/MND, and MND subgroups. Sensitivity of results to the inclusion of related family members and individuals with other diagnoses, and to the inclusion of additional controls were considered in secondary analyses. For cohort 2, models were adjusted for gender and age (age at blood draw in controls, age at diagnosis in clinically diagnosed patients, age at death in pathologically diagnosed patients). We compared controls to all patients and additionally made comparisons separately for the FTD subgroup and according to the method of diagnosis (clinical or pathological).

To evaluate associations of *TMEM106B* SNPs with age at onset, we also used linear regression models adjusted for gender and disease subgroup for cohort 1 (using only FTD, FTD/MND, and MND probands); our primary analysis involved a recessive genotypic model, while other genotypic models (additive, dominant, and co-dominant) were examined in secondary analyses.

To adjust for multiple testing in our main disease-association analysis using recessive genotypic models, we utilized a Holm step-down adjustment [11] separately for both cohorts. After this adjustment, $p \leq 0.025$ (cohort 1) and $p \leq 0.0125$ (cohort 2) were considered statistically significant. All statistical analyses were performed with R Statistical Software (version 2.14.0; R Foundation for Statistical Computing, Vienna, Austria).

Results

Cohort 1

We first compared the frequency of carriers homozygous for the minor rs3173615 *TMEM106B* allele [corresponding to homozygous p.185S carriers (GG genotype)] between our FTD, FTD/MND, or MND probands with *C9ORF72* expansions and our control subjects (Table 1). Homozygosity for the minor allele was detected in 11.9 % of the 260 cases as compared to 19.1 % of the 376 controls (OR

Table 2 Associations of *TMEM106B* rs3173615 with disease

Group	N	<i>TMEM106B</i> rs3173615 genotype information				Comparison with controls under a recessive model	
		MAF	CC (%)	CG (%)	GG (%)	OR (95 % CI)	p value
Cohort 1—controls and <i>C9ORF72</i> repeat expansion carrier probands (FTD, FTD/MND, MND)							
Controls	376	43.2 %	123 (32.7)	181 (48.1)	72 (19.1)	1.00 (reference)	N/A
FTD, FTD/MND, and MND patients	260	38.8 %	89 (34.2)	140 (53.8)	31 (11.9)	0.57 (0.36–0.90)	0.014
FTD patients	69	35.5 %	25 (36.2)	39 (56.5)	5 (7.2)	0.33 (0.13–0.85)	0.009
FTD/MND patients	71	42.3 %	17 (23.9)	48 (67.6)	6 (8.5)	0.38 (0.16–0.92)	0.017
MND patients	120	38.8 %	47 (39.2)	53 (44.2)	20 (16.7)	0.85 (0.49–1.46)	0.55
Cohort 2—controls and FTD or FTD/MND patients without <i>C9ORF72</i> repeat expansions or <i>GRN</i> mutations							
Controls	765	41.0 %	280 (36.6)	342 (44.7)	143 (18.7)	1.00 (reference)	N/A
FTD and FTD/MND patients	586	39.3 %	213 (36.3)	285 (48.6)	88 (15.0)	0.77 (0.58–1.03)	0.079
FTD patients	531	39.8 %	187 (35.2)	265 (49.9)	79 (14.9)	0.76 (0.56–1.03)	0.071
Pathologically diagnosed	101	30.6 %	45 (44.6)	50 (49.5)	6 (5.9)	0.26 (0.11–0.61)	<0.001
Clinically diagnosed	430	42.0 %	142 (33.0)	215 (50.0)	73 (17.0)	0.91 (0.67–1.25)	0.56

ORs, 95 % CIs, and *p* values result from logistic regression models where rs3173615 was considered under a recessive model. For cohort 1, models were adjusted for gender. For cohort 2, models were adjusted for age (age at blood draw in controls, age at diagnosis in clinically diagnosed patients, and age at death in pathologically diagnosed patients) and gender. After applying a Holm step-down adjustment for multiple testing, $p \leq 0.025$ are considered as statistically significant in cohort 1, and $p \leq 0.0125$ are considered as statistically significant in cohort 2

FTD frontotemporal dementia, MND motor neuron disease, OR odds ratio, CI confidence interval, MAF minor allele frequency

0.57, $p = 0.014$; Table 2). When including all 65 family members and individuals with other diagnoses who harbored *C9ORF72* expansions, the frequency of carriers homozygous for the minor allele remained lower in individuals with *C9ORF72* expansions than in controls (11.4 vs. 19.1 %).

Subsequently, we determined whether the lower frequency of homozygous carriers was present in all disease subgroups (Table 2). Interestingly, the lowest frequency was encountered in FTD patients (7.2 %, OR 0.33, $p = 0.009$) followed by FTD/MND patients (8.5 %, OR 0.38, $p = 0.017$), whereas no significant difference was detected in MND patients (16.7 %, OR 0.85, $p = 0.55$). These findings were consistent when including family members (FTD 7.0 %, FTD/MND 9.0 %, and MND 16.7 %).

A similar trend was seen for all aforementioned differences with controls, when including 410 extra controls from other sites to form a combined control group of 786 individuals (Online Resource Table 1), and when additionally adjusting the logistic regression models for age in the subset of individuals with that information available (data not shown). We also considered other genotypic models (additive, dominant, and co-dominant), but the differences stayed most significant under the recessive genotypic model (Online Resource Table 2).

Furthermore, we investigated whether homozygosity for the minor allele might influence age at onset. In the 240 FTD, FTD/MND or MND probands with age at onset

Table 3 Associations of *TMEM106B* rs3173615 with age at onset for probands in cohort 1

Model	Association between rs3173615 and age at onset	
	Regression coefficient (95 % CI)	p value
Recessive (GG vs. CC or CG)	−1.26 (−4.78, 2.25)	0.48
Additive (effect of each additional G allele)	−0.24 (−2.04, 1.55)	0.79
Dominant (CG or GG vs. CC)	0.16 (−2.30, 2.63)	0.90
Co-dominant		
CG vs. CC	0.45 (−2.12, 3.02)	0.73
GG vs. CC	−0.99 (−4.84, 2.85)	0.61

Regression coefficients, 95 % CIs, and *p* values result from linear regression models adjusted for age and disease group. Regression coefficients are interpreted as the change in mean age at onset corresponding to the presence of the GG genotype for rs3173615 (recessive model), each additional G allele for rs3173615 (additive model), and the CG or GG genotype for rs3173615 (dominant model). For the co-dominant model, regression coefficients are interpreted as the change in mean age at onset in comparison to the CC genotype for rs3173615

information available, the mean age at onset was 1.26 years lower in patients homozygous for the minor allele in comparison to other patients, when utilizing a recessive genotypic model (95 % CI 4.78 years lower to 2.25 years higher; Table 3); it is important to realize, however, that this did

not approach significance ($p = 0.48$). The lack of association between rs3173615 and age at onset was also apparent when examining other genotypic models (additive, dominant, and co-dominant; Table 3). All association analyses regarding disease and age at onset were comparable for rs1990622 (data not shown).

Cohort 2

Before the identification of repeat expansions in *C9ORF72*, we assessed a large cohort of FTD patients, FTD/MND patients, and control subjects for *TMEM106B* SNPs [8, 28]. For our present study, we excluded all carriers of *C9ORF72* expansions and *GRN* mutations from this cohort and reanalyzed our data. In addition, we extended our cohort with 31 newly ascertained FTLT-DTP patients. In this cohort of 586 patients and 765 controls, we showed that the frequency of carriers homozygous for the minor rs3173615 *TMEM106B* allele (GG genotype) did not differ significantly between groups, when using a recessive genotypic model (15.0 vs. 18.7 %, OR 0.77, $p = 0.079$; Table 2). Because our findings in cohort 1 were most pronounced in FTD patients, we subsequently focused our analysis on patients with FTD (without signs of MND); however, in this group we also did not detect a significant difference between patients and controls (14.9 vs. 18.7 %, OR 0.76, $p = 0.071$). While this group included both clinically diagnosed patients ($n = 430$) and pathologically diagnosed patients ($n = 101$), we next investigated these subgroups separately. Although no significant differences were observed when comparing clinically diagnosed patients and controls (17.0 vs. 18.7 %, OR 0.91, $p = 0.56$; reported diagnosis of behavioral variant FTD 14.3 %; remaining FTD patients 18.1 %), the frequency of carriers homozygous for the minor allele was significantly lower in patients who received a diagnosis of FTLT-DTP than in controls (5.9 vs. 18.7 %, OR 0.26, $p < 0.001$), even though patients with *C9ORF72* expansions and *GRN* mutations had already been excluded.

Comparable results were obtained after inclusion of all additional controls from cohort 1 to form a combined control group of 1,302 subjects (Online Resource Table 1). Differences between patients and controls were also examined under other genotypic models (additive, dominant, and co-dominant); however, differences were not as strong as those observed under a recessive genotypic model (Online Resource Table 2). All results were similar when evaluating associations with *TMEM106B* rs1990622 (data not shown).

Immunohistochemistry

All nine patients for whom immunohistochemistry was performed displayed TDP-43 pathology (Table 4; Fig. 1).

Table 4 TDP-43 burden of patients with TDP-43 Type A pathology

Group	<i>C9ORF72</i> expansion	Genotype	Age at death	Gender	TDP-43 count
Cohort 1					
a	Yes	GG	90	F	7.7
b	Yes	CC	83	F	10.5
c	Yes	CC	74	M	11.0
d	Yes	CC	71	M	20.5
Cohort 2					
e	No	GG	90	F	4.5
f	No	CC	84	M	6.0
g	No	CC	86	M	16.7
h	No	CC	77	M	24.2

Subjects are homozygous carriers of the minor *TMEM106B* allele (GG), or of the major *TMEM106B* allele (CC) in rs3173615. TDP-43 counts were performed in six 10× microscopic fields randomly selected over the gyrus and averaged for a composite pathologic burden score. Identifiers correspond to Fig. 1

Only one patient had pathologic lesions consistent with TDP-43 Type C (predominant long dystrophic neurites; patient i) and, therefore, this patient was excluded from further analyses. The remaining eight patients showed pathologic lesions consistent with TDP-43 Type A (neuronal cytoplasmic and intranuclear inclusions and small dystrophic neurites; patients a–h). When performing a very preliminary comparison in TDP-43 Type A burden within each of the cohorts, the lowest pathologic burden was consistently present in homozygous carriers of the minor, protective, *TMEM106B* allele. Of note, all patients with *C9ORF72* repeat expansions demonstrated C9RANT pathology, characterized by neuronal cytoplasmic inclusions, as opposed to patients without these expansions.

Discussion

Our results identify rs3173615 (corresponding to *TMEM106B* p.T185S) as an important genetic modifying factor in *C9ORF72* expansion carriers. We focused our analysis on a recessive genotypic model, which revealed that individuals who carry a *C9ORF72* repeat expansion and harbor two copies of the minor *TMEM106B* allele (GG genotype) are less likely to develop disease symptoms (OR 0.57), in particular FTD (OR 0.33). Although these findings suggest that the protective effects of *TMEM106B* variants are not limited to previously reported individuals with *GRN* mutations, these protective effects seem more pronounced in carriers of *GRN* mutations, who demonstrate a frequency of homozygous carriers as low as 2 % [8, 19, 30], corresponding to an OR of approximately 0.12 [8].

of S185 [19]. Taken together, these findings suggest that lysosomal pathways may play a critical role in the pathogenesis of FTD; for instance, they may affect the degradation and subsequent aggregation of TDP-43, and thereby influence the likelihood of developing FTD.

Because we did detect homozygous carriers of the minor *TMEM106B* allele in a small percentage of FTLD-TDP patients, we postulate that another as yet unknown factor (or combination of factors) has such a strong effect in these patients that the protective effects of the minor allele are lessened. Importantly, our highly preliminary immunohistochemical findings may support the hypothesis that *TMEM106B* variants affect TDP-43: patients with the protective allele who did develop FTLD-TDP seemed to show less TDP-43 burden than patients with the risk allele. Since homozygous carriers of protective alleles are rare in FTD patients, however, we were only able to investigate a very limited number of samples and, consequently, additional cases should be assessed to confirm these suggestive findings.

Based on our present results and the aforementioned studies, we speculate that a cascade of lysosomal dysfunction might contribute to *C9ORF72*-related FTD and possibly to other types of FTD. In fact, it has already been suggested that FTD-associated genes valosin-containing protein (*VCP*) and charged multivesicular body protein 2b (*CHMP2B*), which account for a small subgroup of FTD patients, are involved in accumulation of autophagosomes, affect endolysosomal sorting, regulate the maturation and size of early endosomes, impair endosome–lysosome fusion, and delay the degradation process of cargo proteins [10, 13, 20, 22, 24], and thus lysosomal dysfunction could represent a shared mechanism that underlies many types of FTD.

Our study has several limitations. Despite the relatively large sample size, for example, the possibility of a type II error (i.e., false-negative association) should be considered. Moreover, due to the low frequency of two copies of the minor *TMEM106B* allele, especially in FTD patients, detailed analysis of clinical and pathological characteristics was obviously hampered. This could have also affected differences in frequencies detected when utilizing a recessive genotypic model and may have contributed to the lack of association with age at onset. Furthermore, although we excluded individuals with known familial relationships from our primary analysis, and *C9ORF72* and *TMEM106B* are located on different chromosomes, we cannot exclude the possibility that some subjects may be distantly related and share loci. Nevertheless, our study was able to thoroughly assess *TMEM106B* variants in a unique cohort of both clinically and pathologically diagnosed *C9ORF72* expansion carriers, in an additional large cohort of FTD patients and controls, and in a disease spectrum ranging from FTD to MND.

To summarize, our findings reveal a new link between *C9ORF72*-related FTLD-TDP, *GRN*-related FTLD-TDP, and other causes of FTLD-TDP: a decreased frequency of carriers homozygous for the minor G allele of *TMEM106B* rs3173615, suggesting a protective effect of this particular genotype. In *C9ORF72* mutation carriers, *TMEM106B* protects specifically against the development of FTD, but not MND, providing the first genetic modifier factor involved in disease presentation in this important genetic subgroup. We hypothesize that *TMEM106B* variants might protect *C9ORF72* expansion carriers from developing FTD, potentially by affecting lysosomal pathways. These findings provide novel opportunities for the development of prognostic tests, and for new treatment strategies aiming at *C9ORF72* expansion carriers and the larger subgroup of patients within the disease spectrum of FTD with TDP-43 pathology.

Acknowledgments This project was supported by NIH grants R01 NS080882, R01 NS065782, R01 AG026251, P01 AG017586, P50 NS072187, P50 AG016574, P30 AG013854, P30 AG012300, P30 AG019610, U01 AG006786, the ALS Therapy Alliance, and the Consortium for Frontotemporal Dementia Research. Data collection at the University of British Columbia is supported by CIHR grant #179009. Dr. Van Blitterswijk is supported by the Milton Safenowitz Post-Doctoral Fellowship for ALS research from the ALS Association.

Conflict of interest Mrs. DeJesus-Hernandez and Dr. Rademakers hold a patent on methods to screen for the hexanucleotide repeat expansion in the *C9ORF72* gene; the other authors declare that they have no conflict of interest.

References

- Ash PE, Bieniek KF, Gendron TF, Caulfield T, Lin WL, DeJesus-Hernandez M, van Blitterswijk MM, Jansen-West K, Paul JW 3rd, Rademakers R, Boylan KB, Dickson DW, Petrucelli L (2013) Unconventional translation of C9ORF72 GGGGCC expansion generates insoluble polypeptides specific to c9FTD/ALS. *Neuron* 77(4):639–646. doi:10.1016/j.neuron.2013.02.004
- Baker M, Mackenzie IR, Pickering-Brown SM, Gass J, Rademakers R, Lindholm C, Snowden J, Adamson J, Sadovnick AD, Rollinson S, Cannon A, Dwosh E, Neary D, Melquist S, Richardson A, Dickson D, Berger Z, Eriksen J, Robinson T, Zehr C, Dickey CA, Crook R, McGowan E, Mann D, Boeve B, Feldman H, Hutton M (2006) Mutations in progranulin cause tau-negative frontotemporal dementia linked to chromosome 17. *Nature* 442(7105):916–919. doi:10.1038/nature05016
- Brady OA, Zheng Y, Murphy K, Huang M, Hu F (2013) The frontotemporal lobar degeneration risk factor, TMEM106B, regulates lysosomal morphology and function. *Hum Mol Genet* 22(4):685–695. doi:10.1093/hmg/ddt475
- Chen-Plotkin AS, Unger TL, Gallagher MD, Bill E, Kwong LK, Volpicelli-Daley L, Busch JI, Akle S, Grossman M, Van Deerlin V, Trojanowski JQ, Lee VM (2012) TMEM106B, the risk gene for frontotemporal dementia, is regulated by the microRNA-132/212 cluster and affects progranulin pathways. *J Neurosci* 32(33):11213–11227. doi:10.1523/JNEUROSCI.0521-12.2012

5. Cruchaga C, Graff C, Chiang HH, Wang J, Hinrichs AL, Spiegel N, Bertelsen S, Mayo K, Norton JB, Morris JC, Goate A (2011) Association of TMEM106B gene polymorphism with age at onset in granulin mutation carriers and plasma granulin protein levels. *Arch Neurol* 68(5):581–586. doi:10.1001/archneurol.2010.350
6. Cruts M, Gijssels I, van der Zee J, Engelborghs S, Wils H, Pirici D, Rademakers R, Vandenberghe R, Dermaut B, Martin JJ, van Duijn C, Peeters K, Sciot R, Santens P, De Pooter T, Mattheijssens M, Van den Broeck M, Cuijt I, Vennekens K, De Deyn PP, Kumar-Singh S, Van Broeckhoven C (2006) Null mutations in progranulin cause ubiquitin-positive frontotemporal dementia linked to chromosome 17q21. *Nature* 442(7105):920–924. doi:10.1038/nature05017
7. DeJesus-Hernandez M, Mackenzie IR, Boeve BF, Boxer AL, Baker M, Rutherford NJ, Nicholson AM, Finch NA, Flynn H, Adamson J, Kouri N, Wojtas A, Sengdy P, Hsiung GY, Karydas A, Seeley WW, Josephs KA, Coppola G, Geschwind DH, Wszolek ZK, Feldman H, Knopman DS, Petersen RC, Miller BL, Dickson DW, Boylan KB, Graff-Radford NR, Rademakers R (2011) Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* 72(2):245–256. doi:10.1016/j.neuron.2011.09.011
8. Finch N, Carrasquillo MM, Baker M, Rutherford NJ, Coppola G, DeJesus-Hernandez M, Crook R, Hunter T, Ghidoni R, Benussi L, Crook J, Finger E, Hantapan KJ, Karydas AM, Sengdy P, Gonzalez J, Seeley WW, Johnson N, Beach TG, Mesulam M, Forloni G, Kertesz A, Knopman DS, Uitti R, White CL 3rd, Caselli R, Lipka C, Bigio EH, Wszolek ZK, Binetti G, Mackenzie IR, Miller BL, Boeve BF, Younkin SG, Dickson DW, Petersen RC, Graff-Radford NR, Geschwind DH, Rademakers R (2011) TMEM106B regulates progranulin levels and the penetrance of FTD in GRN mutation carriers. *Neurology* 76(5):467–474. doi:10.1212/WNL.0b013e31820a0e3b
9. Gass J, Cannon A, Mackenzie IR, Boeve B, Baker M, Adamson J, Crook R, Melquist S, Kuntz K, Petersen R, Josephs K, Pickering-Brown SM, Graff-Radford N, Uitti R, Dickson D, Wszolek Z, Gonzalez J, Beach TG, Bigio E, Johnson N, Weintraub S, Mesulam M, White CL 3rd, Woodruff B, Caselli R, Hsiung GY, Feldman H, Knopman D, Hutton M, Rademakers R (2006) Mutations in progranulin are a major cause of ubiquitin-positive frontotemporal lobar degeneration. *Hum Mol Genet* 15(20):2988–3001. doi:10.1093/hmg/ddl241
10. Han JH, Ryu HH, Jun MH, Jang DJ, Lee JA (2012) The functional analysis of the CHMP2B missense mutation associated with neurodegenerative diseases in the endo-lysosomal pathway. *Biochem Biophys Res Commun* 421(3):544–549. doi:10.1016/j.bbrc.2012.04.041
11. Holm S (1979) A simple sequentially rejective multiple test procedure. *Scand J Stat* 6:65–70
12. Lang CM, Fellerer K, Schwenk BM, Kuhn PH, Kremmer E, Edbauer D, Capell A, Haass C (2012) Membrane orientation and subcellular localization of transmembrane protein 106B (TMEM106B), a major risk factor for frontotemporal lobar degeneration. *J Biol Chem* 287(23):19355–19365. doi:10.1074/jbc.M112.365098
13. Lee JA, Beigneux A, Ahmad ST, Young SG, Gao FB (2007) ESCRT-III dysfunction causes autophagosome accumulation and neurodegeneration. *Curr Biol* 17(18):1561–1567. doi:10.1016/j.cub.2007.07.029
14. Mackenzie IR, Neumann M, Baborie A, Sampathu DM, Du Plessis D, Jaros E, Perry RH, Trojanowski JQ, Mann DM, Lee VM (2011) A harmonized classification system for FTDL-TDP pathology. *Acta Neuropathol* 122(1):111–113. doi:10.1007/s00401-011-0845-8
15. Mackenzie IR, Neumann M, Bigio EH, Cairns NJ, Alafuzoff I, Kril J, Kovacs GG, Ghetti B, Halliday G, Holm IE, Ince PG, Kamphorst W, Revesz T, Rozemuller AJ, Kumar-Singh S, Akiyama H, Baborie A, Spina S, Dickson DW, Trojanowski JQ, Mann DM (2009) Nomenclature for neuropathologic subtypes of frontotemporal lobar degeneration: consensus recommendations. *Acta Neuropathol* 117(1):15–18. doi:10.1007/s00401-008-0460-5
16. Majounie E, Renton AE, Mok K, Dopper EG, Waite A, Rollinson S, Chio A, Restagno G, Nicolaou N, Simon-Sanchez J, van Swieten JC, Abramzon Y, Johnson JO, Sendtner M, Pamphlett R, Orrell RW, Mead S, Sidle KC, Houlden H, Rohrer JD, Morrison KE, Pall H, Talbot K, Ansorge O, Hernandez DG, Arepalli S, Sabatelli M, Mora G, Corbo M, Giannini F, Calvo A, Englund E, Borghero G, Floris GL, Remes AM, Laaksovirta H, McCluskey L, Trojanowski JQ, Van Deerlin VM, Schellenberg GD, Nalls MA, Drory VE, Lu CS, Yeh TH, Ishiura H, Takahashi Y, Tsuji S, Le Ber I, Brice A, Drepper C, Williams N, Kirby J, Shaw P, Hardy J, Tienari PJ, Heutink P, Morris HR, Pickering-Brown S, Traynor BJ (2012) Frequency of the C9orf72 hexanucleotide repeat expansion in patients with amyotrophic lateral sclerosis and frontotemporal dementia: a cross-sectional study. *Lancet Neurol* 11(4):323–330. doi:10.1016/S1474-4422(12)70043-1
17. Mori K, Weng SM, Arzberger T, May S, Rentzsch K, Kremmer E, Schmid B, Kretschmar HA, Cruts M, Van Broeckhoven C, Haass C, Edbauer D (2013) The C9orf72 GGGGCC repeat is translated into aggregating dipeptide-repeat proteins in FTDL/ALS. *Science* 339(6125):1335–1338. doi:10.1126/science.1232927
18. Neumann M, Sampathu DM, Kwong LK, Truax AC, Micseanyi MC, Chou TT, Bruce J, Schuck T, Grossman M, Clark CM, McCluskey LF, Miller BL, Masliah E, Mackenzie IR, Feldman H, Feiden W, Kretschmar HA, Trojanowski JQ, Lee VM (2006) Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Science* 314(5796):130–133. doi:10.1126/science.1134108
19. Nicholson AM, Finch NA, Wojtas A, Baker MC, Perkerson RB 3rd, Castanedes-Casey M, Rousseau L, Benussi L, Binetti G, Ghidoni R, Hsiung GY, Mackenzie IR, Finger E, Boeve BF, Ertekin-Taner N, Graff-Radford NR, Dickson DW, Rademakers R (2013) TMEM106B p. T185S regulates TMEM106B protein levels: implications for frontotemporal dementia. *J Neurochem* 126(6):781–791. doi:10.1111/jnc.12329
20. Ramanathan HN, Ye Y (2012) The p97 ATPase associates with EEA1 to regulate the size of early endosomes. *Cell Res* 22(2):346–359. doi:10.1038/cr.2011.80
21. Renton AE, Majounie E, Waite A, Simon-Sanchez J, Rollinson S, Gibbs JR, Schymick JC, Laaksovirta H, van Swieten JC, Myllykangas L, Kalimo H, Paetau A, Abramzon Y, Rimes AM, Kaganovich A, Scholz SW, Duckworth J, Ding J, Harmer DW, Hernandez DG, Johnson JO, Mok K, Ryten M, Trabzuni D, Guerreiro RJ, Orrell RW, Neal J, Murray A, Pearson J, Jansen IE, Sondervan D, Seelaar H, Blake D, Young K, Halliwell N, Callister JB, Toulson G, Richardson G, Gerhard A, Snowden J, Mann D, Neary D, Nalls MA, Peuralinna T, Jansson L, Isoviita VM, Kaivorinne AL, Holtta-Vuori M, Ikonen E, Sulkava R, Benatar M, Wu J, Chio A, Restagno G, Borghero G, Sabatelli M, Heckerman D, Rogaeva E, Zinman L, Rothstein JD, Sendtner M, Drepper C, Eichler EE, Alkan C, Abdullaev Z, Pack SD, Dutra A, Pak E, Hardy J, Singleton A, Williams NM, Heutink P, Pickering-Brown S, Morris HR, Tienari PJ, Traynor BJ (2011) A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 72(2):257–268. doi:10.1016/j.neuron.2011.09.010
22. Ritz D, Vuk M, Kirchner P, Bug M, Schutz S, Hayer A, Bremer S, Lusk C, Baloh RH, Lee H, Glatter T, Gstaiger M, Aebbersold R, Wehl CC, Meyer H (2011) Endolysosomal sorting of ubiquitylated caveolin-1 is regulated by VCP and UBXD1 and impaired by VCP disease mutations. *Nat Cell Biol* 13(9):1116–1123. doi:10.1038/ncb2301

23. Rollinson S, Mead S, Snowden J, Richardson A, Rohrer J, Halliwell N, Usher S, Neary D, Mann D, Hardy J, Pickering-Brown S (2011) Frontotemporal lobar degeneration genome wide association study replication confirms a risk locus shared with amyotrophic lateral sclerosis. *Neurobiol Aging* 32(4):758.e751–758. doi:[10.1016/j.neurobiolaging.2010.12.005](https://doi.org/10.1016/j.neurobiolaging.2010.12.005)
24. Urwin H, Authier A, Nielsen JE, Metcalf D, Powell C, Froud K, Malcolm DS, Holm I, Johannsen P, Brown J, Fisher EM, van der Zee J, Bruylant M, Van Broeckhoven C, Collinge J, Brandner S, Futter C, Isaacs AM (2010) Disruption of endocytic trafficking in frontotemporal dementia with CHMP2B mutations. *Hum Mol Genet* 19(11):2228–2238. doi:[10.1093/hmg/ddq100](https://doi.org/10.1093/hmg/ddq100)
25. van Blitterswijk M, Baker MC, DeJesus-Hernandez M, Ghidoni R, Benussi L, Finger E, Hsiung GY, Kelley BJ, Murray ME, Rutherford NJ, Brown PE, Ravenscroft T, Mullen B, Ash PE, Bieniek KF, Hatanpaa KJ, Karydas A, McCarty Wood E, Coppola G, Bigio EH, Lippa C, Strong MJ, Beach TG, Knopman DS, Huey ED, Mesulam M, Bird T, White CL 3rd, Kertesz A, Geschwind DH, Van Deerlin VM, Petersen RC, Binetti G, Miller BL, Petrucelli L, Wszolek ZK, Boylan KB, Graff-Radford NR, Mackenzie IR, Boeve BF, Dickson DW, Rademakers R (2013) C9ORF72 repeat expansions in cases with previously identified pathogenic mutations. *Neurology* 81(15):1332–1341. doi:[10.1212/WNL.0b013e3182a8250c](https://doi.org/10.1212/WNL.0b013e3182a8250c)
26. van Blitterswijk M, DeJesus-Hernandez M, Niemantsverdriet E, Murray ME, Heckman MG, Diehl NN, Brown PH, Baker MC, Finch NA, Bauer PO, Serrano G, Beach TG, Josephs KA, Knopman DS, Petersen RC, Boeve BF, Graff-Radford NR, Boylan KB, Petrucelli L, Dickson DW, Rademakers R (2013) Association between repeat sizes and clinical and pathological characteristics in carriers of C9ORF72 repeat expansions (Xpansize-72): a cross-sectional cohort study. *Lancet Neurol* 12(10):978–988. doi:[10.1016/S1474-4422\(13\)70210-2](https://doi.org/10.1016/S1474-4422(13)70210-2)
27. van Blitterswijk M, DeJesus-Hernandez M, Rademakers R (2012) How do C9ORF72 repeat expansions cause amyotrophic lateral sclerosis and frontotemporal dementia: can we learn from other noncoding repeat expansion disorders? *Curr Opin Neurol* 25(6):689–700. doi:[10.1097/WCO.0b013e32835a3efb](https://doi.org/10.1097/WCO.0b013e32835a3efb)
28. Van Deerlin VM, Sleiman PM, Martinez-Lage M, Chen-Plotkin A, Wang LS, Graff-Radford NR, Dickson DW, Rademakers R, Boeve BF, Grossman M, Arnold SE, Mann DM, Pickering-Brown SM, Seelaar H, Heutink P, van Swieten JC, Murrell JR, Ghetti B, Spina S, Grafman J, Hodges J, Spillantini MG, Gilman S, Lieberman AP, Kaye JA, Woltjer RL, Bigio EH, Mesulam M, Al-Sarraj S, Troakes C, Rosenberg RN, White CL 3rd, Ferrer I, Llado A, Neumann M, Kretschmar HA, Hulette CM, Welsh-Bohmer KA, Miller BL, Alzualde A, Lopez de Munain A, McKee AC, Gearring M, Levey AI, Lah JJ, Hardy J, Rohrer JD, Lashley T, Mackenzie IR, Feldman HH, Hamilton RL, Dekosky ST, van der Zee J, Kumar-Singh S, Van Broeckhoven C, Mayeux R, Vonsattel JP, Troncoso JC, Kril JJ, Kwok JB, Halliday GM, Bird TD, Ince PG, Shaw PJ, Cairns NJ, Morris JC, McLean CA, DeCarli C, Ellis WG, Freeman SH, Frosch MP, Growdon JH, Perl DP, Sano M, Bennett DA, Schneider JA, Beach TG, Reiman EM, Woodruff BK, Cummings J, Vinters HV, Miller CA, Chui HC, Alafuzoff I, Hartikainen P, Seilhean D, Galasko D, Masliah E, Cotman CW, Tunon MT, Martinez MC, Munoz DG, Carroll SL, Marson D, Riederer PF, Bogdanovic N, Schellenberg GD, Hakonarson H, Trojanowski JQ, Lee VM (2010) Common variants at 7p21 are associated with frontotemporal lobar degeneration with TDP-43 inclusions. *Nat Genet* 42(3):234–239. doi:[10.1038/ng.536](https://doi.org/10.1038/ng.536)
29. van der Zee J, Van Langenhove T, Kleinberger G, Slegers K, Engelborghs S, Vandenberghe R, Santens P, Van den Broeck M, Joris G, Brys J, Mattheijssens M, Peeters K, Cras P, De Deyn PP, Cruts M, Van Broeckhoven C (2011) TMEM106B is associated with frontotemporal lobar degeneration in a clinically diagnosed patient cohort. *Brain* 134(Pt 3):808–815. doi:[10.1093/brain/awr007](https://doi.org/10.1093/brain/awr007)
30. Vass R, Ashbridge E, Geser F, Hu WT, Grossman M, Clay-Falcone D, Elman L, McCluskey L, Lee VM, Van Deerlin VM, Trojanowski JQ, Chen-Plotkin AS (2011) Risk genotypes at TMEM106B are associated with cognitive impairment in amyotrophic lateral sclerosis. *Acta Neuropathol* 121(3):373–380. doi:[10.1007/s00401-010-0782-y](https://doi.org/10.1007/s00401-010-0782-y)