



Plasma oxidation status and antioxidant capacity in psoriatic children

Tiziana Bacchetti¹ · Oriana Simonetti² · Francesca Ricotti² · Annamaria Offidani² · Gianna Ferretti³

Received: 13 April 2019 / Revised: 16 July 2019 / Accepted: 30 August 2019 / Published online: 17 September 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Psoriasis, a chronic inflammatory skin disease, is associated with oxidative stress of serum lipoproteins. In psoriatic children we evaluated the activity and levels of myeloperoxidase, the activity of paraoxonase-1 (PON1) and biochemical markers of lipid peroxidation, to investigate whether an imbalance between oxidant–antioxidants occurs very early in psoriasis. A total of 52 patients affected by psoriasis and 48 sex–age–matched healthy controls were enrolled. Serum MPO levels were measured using ELISA method. MPO and PON1 activities (paraoxonase, arylesterase and lactonase) were evaluated by spectroscopic methods. Our results demonstrated a significant increase of MPO levels and activity in psoriatic subjects. PON1 activities were found to be significantly decreased. A positive correlation has been established between the MPO/PON1 ratio and levels of lipid peroxides in all psoriatic patients. These results suggest that an imbalance between MPO and PON1 can reflect in higher oxidative stress in serum lipoproteins.

Keywords Myeloperoxidase · Paraoxonase-1 · High-density lipoproteins · Pediatric psoriasis · Lipid peroxidation

Introduction

Psoriasis (PS) is a chronic inflammatory skin disease characterized by increased keratinocyte proliferation and alterations in dermal and epidermal T-cells, monocytes–macrophages and neutrophils [1]. Increased cardiovascular and metabolic co-morbidity [2, 3] related to alterations of plasma lipoprotein levels and composition has been described [4–8]. A relationship between psoriasis and oxidative stress has also been demonstrated with higher levels of biochemical markers of plasma lipid and protein oxidation [4, 9–11], including higher levels of lipid peroxidation markers in very

low density and low density lipoproteins in adult patients with a history of disease that lasts for years [12]. High titers of autoantibodies against oxidized low density lipoproteins (ox-LDL) confirm that *in vivo* oxidation of LDL occurs in plasma of psoriatic patients [13]. Furthermore, an accumulation of ox-LDL has been detected in the upper epidermis of the involved skin from psoriatic patients by direct immune-fluorescent method [14]. Compositional and functional alterations of high-density lipoproteins (HDL) have also been studied in psoriasis and recently reviewed [15–18]. A decrease of cholesterol efflux from macrophages, higher levels of lipid hydroperoxides and a decrease of the anti-inflammatory and antioxidant properties have been reported in HDL from adult psoriatic patients [16, 18]. In detail, the activity of paraoxonase 1 (PON1), a multitasking enzyme associated to HDL [19], is significantly lower in adult PS patients and a relationship with disease activity has been observed [11, 20].

The study of the molecular mechanisms that could trigger oxidative damage to lipoproteins *in vivo* in psoriasis has not been previously investigated. A role of the enzyme myeloperoxidase (MPO) in oxidative damage of plasma lipoproteins has been recently reported in atherosclerosis and inflammatory diseases [21]. Myeloperoxidase (MPO), secreted by activated monocytes, is a pro-oxidative and pro-inflammatory hemeprotein [22], which catalyzes the conversion of

Tiziana Bacchetti and Oriana Simonetti contributed equally to this work.

✉ Oriana Simonetti
o.simonetti@univpm.it

¹ Department of Life and Environmental Sciences-Biochemistry, Università Politecnica delle Marche, 60100 Ancona, Italy

² Department of Clinical and Molecular Sciences-Dermatology, Università Politecnica delle Marche, 60100 Ancona, Italy

³ Department of Clinical Experimental Science and Odontostomatology-Biochemistry, Università Politecnica delle Marche, 60100 Ancona, Italy

chloride and hydrogen peroxide to hypochlorite [22]. MPO generates other reactive oxygen species such as hydroxyl radical and singlet oxygen and cross-links proteins [21]. ROS generated by activated monocytes oxidatively modify LDL and HDL in vitro [22]. Previous studies have suggested that the ratio between serum levels of myeloperoxidase and paraoxonase 1 activity could be a potential indicator of dysfunctional high-density lipoprotein [23]. We studied MPO levels and activity, PON1 activity and biochemical markers of lipid peroxidation in pediatric patients to investigate whether an unbalance between oxidant–antioxidants occurs very early in psoriasis.

Materials and methods

Subjects

52 psoriatic children (29/23 F/M mean age 9.8 ± 3.7) affected by psoriasis were included through consecutive recruitment from new referrals and follow-up visits at the Clinic of Dermatology of the Department of Clinical and Molecular Sciences of the Polytechnic University of Marche.

Inclusion criteria for pediatric patients were: a confident clinical diagnosis of psoriasis, age between 0 and 14 years and onset of disease within the past 12 months. Only scalp involvement or diaper rash was not regarded as sufficient for diagnosis. Diaper rash psoriasis was not included because of the difficulty in separating psoriatic diaper rash from other diaper rashes. None among the patients showed any clinical sign, or reported a history of psoriatic arthritis. Subjects with diabetes, clinical evidence of cardiovascular diseases, or receiving lipid-lowering drugs or antioxidant supplements were excluded from the study to avoid possible interferences on PON1 activity and plasma lipids. All selected subjects had avoided UVBnb, UVBbb, UVA or PUVA treatments and direct sun exposure for almost 4 weeks. Disease severity was quantified by the Psoriasis Area and Severity Index (PASI) (Table 1).

48 pediatric subjects (25/23 F/M mean age, 10.3 ± 2.7 years), age and sex-matched, without skin or systemic inflammatory disease (e.g., with nevi, molluscum contagiosum, warts) and without a family history of psoriasis or psoriatic arthritis were also recruited.

Both controls and PS patients have comparable body mass index (BMI) (21.8 for control vs 22.5 for patients). All the procedures were in accordance with the Helsinki Declaration of 1975, as revised in 2000. The study was approved by “Ethics Committee, Ospedali Riuniti di Ancona”. Written informed consent for participation in the study was obtained from each patient’s parent with the approval of the attending physician.

Table 1 Clinical characteristics and plasma parameters in controls and subjects affected by psoriasis

	Healthy subjects	Psoriatic patients
Age (years)	9.8 ± 3.7	10.3 ± 2.7 years
Fasting glucose (mg/dL)	81.1 ± 7.7	73.4 ± 15.8
TG (mg/dL)	74.3 ± 18.4	75.2 ± 31.0
TC (mg/dL)	165.9 ± 26.7	158.2 ± 25.5
HDL-C (mg/dL)	51.6 ± 7.3	49.21 ± 6.5
LDL-C (mg/dL)	89.9 ± 20.9	90.1 ± 24.1
BMI	21.8 ± 3.5	22.7 ± 3.1
PASI	–	9.8 ± 4.3

TG triglycerides, TC total cholesterol, LDL-C LDL-cholesterol, HDL-C HDL-cholesterol

Sample collection

Blood samples of controls and PS patients were collected at 8 a.m., after overnight fasting. Venous blood (about 10 ml) was placed in two tubes. An aliquot was incubated with heparin and plasma was separated by low speed centrifugation (3000 rpm) at 4 °C for 20 min. Another aliquot was used for serum separation. A glass centrifuge tube containing blood (about 5 ml) without anticoagulant was left at room temperature for 30 min to allow formation of the clot, which was immediately removed. The tubes were centrifuged in a tabletop centrifuge for 10 min at a rotation speed of 2000 rpm. Both plasma and serum samples were divided in aliquots and stored immediately at – 80 °C.

Plasma lipid profile

Fasting levels of total cholesterol (TC), HDL-cholesterol (HDL-C), LDL-cholesterol (LDL-C) and triglycerides (TG) were measured in plasma of controls and PS patients using commercially available kits (Roche Diagnostics, Switzerland).

Serum lipid peroxidation

The levels of lipid hydroperoxides were determined in serum of controls and psoriatic patients using FOX2 assay as previously described [11]. The levels of lipid hydroperoxides were quantified using a stock solution of t-butyl hydroperoxide. The results are shown as μmol of lipid hydroperoxides for L of serum.

Serum total antioxidant capacity

Serum total antioxidant capacity was assessed using the oxygen radical absorbance capacity (ORAC) assay [24].

The oxygen radical absorbance capacity of serum employs the oxidative loss of intrinsic fluorescence of fluorescein induced by the free radical initiator 2,2'-azobis(2-amidinopropane) hydrochloride. Fluorescein fluorescence decay shows a lag or retardation in the presence of antioxidants, related to the antioxidant capacity of the sample. Trolox was used as a reference antioxidant for calculating the ORAC values. Results were expressed as μmol Trolox equivalents/L ($\mu\text{mol TE/L}$).

Serum PON1 activities

PON1 activities were evaluated in serum of controls and psoriatic patients using three substrates [25] paraoxon for paraoxonase activity, phenylacetate for arylesterase activity and dihydrocoumarin for lactonase. All assays were performed in a 96-well plate, in a total reaction volume of 200 μL .

Paraoxonase activity 10 μL of serum (non-diluted samples) were used. The basal assay mixture included 5 mmol/L Tris-HCl, pH 7.4 containing 1 mmol/L CaCl_2 and 1.0 mmol/L paraoxon. Paraoxon hydrolysis was spectrophotometrically monitored for 8 min (every 15 s) at 412 nm. Non-enzymatic hydrolysis of paraoxon was subtracted from the total rate of hydrolysis. One unit of PON1 paraoxonase activity was equivalent to 1 nmol of paraoxon hydrolyzed/min/mL.

Arylesterase activity

Serum samples were diluted 1:10 with 1 mmol/L CaCl_2 in 50 mmol/L Tris-HCl, pH 8.0 and then, 5 μL was taken for a total reaction volume of 200 μL . After addition of the substrate phenyl acetate (1 mmol/L), the hydrolysis was monitored at 270 nm for 3 min (every 15 s). One unit of arylesterase activity was equivalent to 1 μmol of phenyl acetate hydrolyzed/min/mL.

Lactonase activity

Serum samples were diluted 1:10 with 1 mmol/L CaCl_2 in 50 mmol/L Tris-HCl, pH 8.0 and 3 μL was then taken for the assay. After addition of the substrate dihydrocoumarin (DHC) (1.0 mmol/L), the hydrolysis was monitored at 270 nm for 10 min (every 15 s). Non-enzymatic hydrolysis of DHC was subtracted from the total rate of hydrolysis. One unit of lactonase activity was equivalent to 1 μmol of DHC hydrolyzed/min/mL.

All assays were performed in a 96-well plate, in a total reaction volume of 200 μL .

Myeloperoxidase levels and activity

Serum MPO levels were measured using a solid-phase two-site MPO ELISA Kit from Mercodia (Uppsala, Sweden) according to instructions provided by the manufacturer (26).

MPO activity was evaluated by a colorimetric Activity Assay Kit (Sigma-Aldrich).

Statistical analysis

The results are reported as mean \pm standard error (SE). For the comparison of normally distributed variables between groups, Student's *t* test was used. Paraoxonase activity showed a non-gaussian distribution; therefore we used a non-parametric test (Wilcoxon rank sum test). Pearson correlation coefficients and their significance levels were calculated for linear regression analysis. Differences were considered statistically significant at $p < 0.05$ (Microcal Origin 5.0, OriginLab, Northampton, MA).

Results

As shown in Table 1, plasma levels of total cholesterol, HDL-C, LDL-C were not significantly modified in pediatric patients compared to controls.

Lipid peroxides and total antioxidant capacity

Higher levels of lipid hydroperoxides (Table 2) and lower values of total antioxidant capacity were observed in serum of patients compared to sex-age-matched healthy control children. The differences were statistically significant.

Activity of PON1 and levels of MPO

As reported in Table 2, lower activities of PON1 (paraoxonase, arylesterase and lactonase) were observed in serum of patients compared to control subjects ($p < 0.05$). In the same patients, a significant increase of serum levels and activity of myeloperoxidase (MPO) was found in PS children ($p < 0.05$, Table 2). The comparison of the ratio between serum MPO level /PON1 paraoxonase activity (MPO/PON1 ratio) showed higher levels in patients compared to controls ($p < 0.05$, Table 2). In all groups of subjects, levels of serum PON1 paraoxonase activity were negatively correlated with serum MPO concentration (Table 3). Furthermore, a significant positive correlation was established between serum levels of lipid hydroperoxides and MPO/PON1 ratio (Table 3).

Table 2 PON1 activities (paraoxonase, arylesterase, lactonase), MPO activity and levels in controls and psoriatic patients * $p < 0.05$ vs healthy subjects

	Healthy subjects	Psoriatic patients
Lipid hydroperoxides ($\mu\text{mol/L}$)	2.56 ± 0.7	$5.2 \pm 1.4^*$
Total antioxidant capacity (mmol TE /L)	13.2 ± 1.9	$8.78 \pm 4.1^*$
PON-1—paraoxonase (U/mL)	211.1 ± 13.9	$94.7 \pm 9.6^*$
PON-1—arylesterase (U/mL)	76.3 ± 2.6	$51.2 \pm 3.0^*$
PON1—lactonase (U/mL)	27.2 ± 1.4	$18.3 \pm 1.5^*$
MPO levels (ng/mL)	101.8 ± 12.3	$149.6 \pm 14.8^*$
MPO activity	9 ± 3	$14 \pm 2^*$
MPO /PON1 (paraoxonase) ratio	0.7 ± 0.09	$1.85 \pm 0.31^*$

Table 3 Correlations between PON1 paraoxonase activity and MPO levels and between MPO/ PON1 ratio and lipid hydroperoxide levels in serum of control subjects and of patients affected by psoriasis

	Healthy subjects	Psoriatic patients
PON1 activity vs MPO levels	$r = -0.64, p < 0.004$	$r = -0.57, p < 0.01$
MPO/ PON1 ratio vs lipid hydroperoxides	$r = 0.46, p < 0.03$	$r = 0.69, p < 0.006$

Discussion

Higher levels of biomarkers of lipid peroxidation, lower total antioxidant capacity and lower PON1 activities in serum of psoriatic children have been demonstrated. The lower activity of PON1 is in good agreement with other studies in adult subjects [11, 15, 20]. Our results, in absence of significant changes of plasma lipids, suggest that oxidative stress and an unbalance between oxidant–antioxidants occur very early in psoriatic children.

Our study, for the first time, also demonstrates a significant increase in serum MPO levels and activity in psoriatic children and PON1 activity was negatively correlated with MPO levels. A higher ratio MPO/PON1 was observed in serum of pediatric patients compared to controls. Moreover, a positive correlation has been established between the MPO/PON1 ratio and levels of lipid peroxides in all psoriatic patients. These results suggest that an unbalance between MPO and PON1 can reflect in higher oxidative stress in serum lipoproteins. All psoriatic children included in our study had a mild form of the disease. No correlation has been established between PASI scores and biochemical indexes of lipid peroxidation or PON1 activities in psoriatic children. In adult patients, a negative correlation between PON1 activity and PASI score has been reported in subjects with PASI values ranging from moderate to severe disease by other authors [26–28].

However, other studies did not find significant relationship between PON1 activities and severity of the disease

[29]. These results are likely related to disease duration ranging from few months to several years.

Some hypotheses can be formulated to explain the potential molecular mechanisms involved in the alterations of PON1 activities in psoriatic children. We suggest that the higher MPO levels and activity could contribute to alterations of HDL and paraoxonase. This hypothesis is supported by previous studies which have demonstrated that among molecular mechanism able to trigger lipid peroxidation of lipoproteins, an emerging role is reserved to the enzyme MPO [21–23]. High MPO levels and/or activity have recently been demonstrated in a psoriatic animal model [30]. Moreover, in adult human subjects psoriatic lesions are characterized by a focal dense infiltration of neutrophils, T lymphocytes and macrophages that migrate from the vascular to the dermal compartment and through all layers of the epidermis up to the stratum corneum [31]. Furthermore, Dylek et al. [32] have recently demonstrated that neutrophils in psoriasis lesions are actively producing MPO [32]. In vitro studies confirm that MPO and reactive oxygen species generated by neutrophils trigger lipid peroxidation of LDL and HDL [22]. In particular, MPO mediates modifications of apoAI with formation of chlorotyrosine and nitrotyrosine from tyrosine residues [33–35]. Other mechanisms could be involved. A considerable number of inflammatory cytokines have been shown to be elevated in lesional psoriasis skin in adult subjects [36]. Previous studies have shown that inflammation in both humans and rabbits leads to the loss of PON1 activity and a decreased ability to protect LDL against oxidation [19]. TNF and IL-1 lead to a decrease in PON1 mRNA levels in liver and a decrease in serum PON1 activity [37].

In conclusion, our data demonstrate that psoriasis in pediatric patients, is associated with higher levels of MPO, oxidative damage of plasma lipids and lipoproteins, a decrease in antioxidant defenses and a significant decrease of PON1. The higher ratio MPO/ PON1 suggests dysfunctional high-density lipoprotein in psoriatic pediatric patients. HDL has a number of important functions (antioxidant, antithrombotic) that could reduce the risk of CVD and which may be adversely impacted by inflammation




Fig. 1 Schematic representation of potential molecular mechanisms involved in lipid peroxidation, lipoprotein dysfunction and alterations in inflamed skin in psoriasis. (1) The higher production of superoxide anion and MPO from activated PMN contributes to lipid peroxidation of LDL and HDL. (2) MPO catalyzes the conversion of chloride and hydrogen peroxide to hypochlorite, a strong reactive oxygen species

(ROS). (3) Oxidized LDL (ox-LDL) and oxidized HDL (ox-HDL) show compositional and functional alterations. In particular, ox-LDL exert pro-inflammatory properties and activate migration and degranulation of PMN. Ox-HDL have lower PON1 activity and exhibit lower antioxidant and anti-inflammatory activity

[38]. As aforementioned, PON1 is a multitasking protein localized at the HDL surface and is associated with protection against oxidative stress-related diseases [19, 39], in fact it exerts a protective role against lipid peroxidation of biological membranes, HDL and LDL [19, 39, 40]. We suggest that whenever the activity of the antioxidant enzyme PON1 is lowered, it will be unable to prevent oxidation of membranes and LDL and could be involved in the higher oxidative stress in psoriasis.

Figure 1 summarizes our hypothesis of the potential role of MPO and PON1 in inflammation and lipid peroxidation in psoriasis. The relationship between psoriasis, oxidative stress, inflammation and alteration of lipoprotein functions is supported by previous studies, which have demonstrated that drug treatments decrease inflammation, lipid peroxidation and recover HDL functions in psoriatic adult patients [17, 20]. It as to be stressed that modifications of HDL composition, PON1 activity and MPO levels realize early in pediatric patients, even in absence of modification of plasma lipids.

Although further studies are necessary to evaluate whether the lower PON1 activities are primary events or a consequence of abnormal metabolism of inflamed skin in pediatric patients, we suggest that an intensive multidisciplinary lifestyle intervention on inflammatory biomarkers could positively affect risk factors for cardiovascular disease in psoriatic children.

Funding None.

Compliance with ethical standards

Conflict of interest None.

References

1. JM Crow 2012 Psoriasis uncovered Nature 492 7429 S50 S51

2. M Augustin G Glaeske MA Radtke E Christophers K Reich I Schäfer 2010 Epidemiology and comorbidity of psoriasis in children *Br J Dermatol* 162 3 633 636
3. IM Miller C Ellervik S Yazdanyar GB Jemec 2013 Meta-analysis of psoriasis, cardiovascular disease, and associated risk factors *J Am Acad Dermatol* 69 6 1014 1024
4. G Ferretti R Alleva M Taus O Simonetti B Cinti A Offidani G Bossi G Curatola 1994 Abnormalities of plasma lipoprotein composition and fluidity in psoriasis *Acta Dermato Venereol* 74 3 171 175
5. C Koebnick M-H Black N Smith JK Der-Sarkissian AH Porter SJ Jacobsen 2011 The association of psoriasis and elevated blood lipids in overweight and obese children *J Pediatrics* 159 4 577 583
6. O Simonetti G Ferretti A Salvi A Offidani G Bossi 1992 Plasma lipid changes in psoriatic children *Dermatology* 185 2 96 100
7. A Pietrzak A Michalak-Stoma G Chodorowska JC Szeptietowski 2010 Lipid disturbances in psoriasis: an update *MediatInflamm* 2010 535 612
8. B Vanizor Kural A Orem G Cimsit YE Yandi M Calapoglu 2003 Evaluation of the atherogenic tendency of lipids and lipoprotein content and their relationships with oxidant-antioxidant system in patients with psoriasis *Clin Chim Acta* 328 1–2 71 82
9. M Matoshvili A Katsitadze T Sanikidze D Tophuria A Richetta S D'Epiro 2014 Alterations of redox-status during psoriasis *Georgian Med News* 232–233 60 64
10. C Yazici K Kose S Utas E Tanrikulu N Taslidere 2016 A novel approach in psoriasis: first usage of known protein oxidation markers to prove oxidative stress *Arch Dermatol Res* 308 3 207 212
11. G Ferretti T Bacchetti A Campanati O Simonetti G Liberati A Offidani 2012 Correlation between lipoprotein(a) and lipid peroxidation in psoriasis: role of the enzyme paraoxonase-1 *Br J Dermatol* 166 1 204 207
12. AM Offidani G Ferretti M Taus O Simonetti N Dousset P Valdiguie G Curatola G Bossi 1994 Lipoprotein peroxidation in adult psoriatic patients *Acta Dermato Venereol* 186 38 40
13. A Orem G Cimsit O Deger C Orem B Vanizor 1999 The significance of autoantibodies against oxidatively modified low-density lipoprotein (LDL) in patients with psoriasis *Clin Chim Acta* 284 1 81 88
14. NS Tekin O TekinI F Barut EY Sipahi 2007 Accumulation of oxidized low-density lipoprotein in psoriatic skin and changes of plasma lipid levels in psoriatic patients *Mediators Inflamm* 2007 78454
15. MJ Paiva-Lopes J Delgado Alves 2017 Psoriasis-associated vascular disease: the role of HDL *J Biomed Sci* 24 1 73
16. M Holzer P Wolf S Curcic R Birner-Gruenberger W Weger M Inzinger D El-Gamal C Wadsack A Heinemann C Marsche 2012 Psoriasis alters HDL composition and cholesterol efflux capacity *JLipid Res* 53 8 1618 1624
17. M Holzer P Wolf M Inzinger M Trieb S Curcic L Pasterk W Weger MA Heinemann G Marsche 2014 Anti-psoriatic therapy recovers high-density lipoprotein composition and function *J Invest Dermatol* 134 3 635 642
18. L He S Qin L Dang G Song S Yao N Yang 1841 Li Y (2014) Psoriasis decreases the anti-oxidation and anti-inflammation properties of high-density lipoprotein *Biochim Biophys Acta* 12 1709 1715
19. M Mackness B Mackness 2015 Human paraoxonase-1 (PON1): Gene structure and expression, promiscuous activities and multiple physiological roles *Gene* 567 1 12 21
20. T Bacchetti A Campanati G Ferretti O Simonetti G Liberati A Offidani 2013 Oxidative stress and psoriasis: the effect of anti-tumour necrosis factor- inhibitor treatment *Br J Dermatol* 168 5 984 989
21. JW Heinecke 1999 Mechanisms of oxidative damage by myeloperoxidase in atherosclerosis and other inflammatory disorders *J Lab Clin Med* 133 4 321 325
22. RL Zhang ML Brennan ZZ Shen JC MacPherson D Schmitt CE Molenda SL Hazen 2002 Myeloperoxidase functions as a major enzymatic catalyst for initiation of lipid peroxidation at sites of inflammation *J Biol Chem* 277 48 46116 46122
23. Y Haraguchi R Toh M Hasokawa H Nakajima T Honjo K Otsui K Mori M Miyamoto-Sasaki M Shinohara K Nishimura T Ishida K Hirata 2014 Serum myeloperoxidase/paraoxonase 1 ratio as potential indicator of dysfunctional high-density lipoprotein and risk stratification in coronary artery disease *Atherosclerosis* 234 2 288 294
24. BX Ou M Hampsch-Woodill RL Prior 2009 Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe *J Agric Food Chem* 49 10 4619 4626
25. W Rock M Rosenblat R Miller-Lotan AP Levy M Elias M Aviram 2008 Consumption of wonderful variety pomegranate juice and extract by diabetic patients increases paraoxonase 1 association with high-density lipoprotein and stimulates its catalytic activities *J Agric Food Chem* 56 18 8704 8713
26. G Ferretti T Bacchetti A Campanati O Simonetti G Liberati A Offidani 2012 Correlation between lipoprotein(a) and lipid peroxidation in psoriasis: role of the enzyme paraoxonase-1 *Br J Dermatol* 166 204 207
27. L He S Qin L Dang G Song S Yao N Yang Y Li 2014 Psoriasis decreases the anti-oxidation and anti-inflammation properties of high-density lipoprotein *Biochim Biophys Acta* 1841 12 1709 1715
28. E Attwa E Swelam 2011 Relationship between smoking-induced oxidative stress and the clinical severity of psoriasis *J Eur Acad Dermatol Venereol* 25 7 782 787
29. SD Pektas G Akoglu A Metin S Neselioglu O Erel 2013 Evaluation of systemic oxidant/antioxidant status and paraoxonase 1 enzyme activities in psoriatic patients treated by narrow band ultraviolet B phototherapy *Redox Rep* 18 5 200 204
30. JO Baek D Byamba WH Wu TG Kim MG Lee 2012 Assessment of an imiquimod-induced psoriatic mouse model in relation to oxidative stress *Arch Dermatol Res* 304 9 699 706
31. A Pietrzak M Koziol-Montewka B Lecewicz-Toruń D Krasowska 2000 Is there any correlation between the total number of neutrophils in plasma and concentration of interleukin-8 in psoriatic patients? *Med Sci Monit* 6 867 870
32. N Dilek AR Dilek Y Taskin T Erkinuresin O Yalcin Y Saral 2006 Contribution of myeloperoxidase and inducible nitric oxide synthase to pathogenesis of psoriasis *Postepy Dermatologii i Alergologii* 33 6 435 439
33. L Zheng B Nukuna ML Brennan M Sun M Goormastic M Settle D Schmitt X Fu L Thomson PL Fox H Ischiropoulos JD Smith M Kinter SL Hazen 2004 Apolipoprotein A-I is a selective target for myeloperoxidase-catalyzed oxidation and functional impairment in subjects with cardiovascular disease *J Clin Invest* 114 4 529 541
34. DQ Peng Z Wu G Brubaker L Zheng M Settle E Gross M Kinter SL Hazen JD Smith 2005 Tyrosine modification is not required for myeloperoxidase-induced loss of apolipoprotein A-I functional activities *J Biol Chem* 280 40 33775 33784
35. Y Huang ZP Wu M Riwanto SQ Gao BS Levison X Gu X Fu MA Wagner C Besler G Gerstenecker R Zhang XM Li AJ Donato Di V Gogonea WH Tang JD Smith EF Plow PL Fox DM Shih AJ Lulis EA Fisher JA DiDonato U Landmesser SL Hazen 2013 Myeloperoxidase, paraoxonase-1, and HDL form a functional ternary complex *J Clin Invest* 123 9 3815 3828
36. J Baliwag DH Barnes A Johnston 2015 Cytokines in psoriasis *Cytokine* 73 2 342 350

37. KR Feingold RA Memon AH Moser C Grunfeld 1998 Paraoxonase activity in the serum and hepatic mRNA levels decrease during the acute phase response *Atherosclerosis* 139 307 315
38. KR Feingold C Grunfeld 2016 Effect of inflammation on HDL structure and function *Curr Opin Lipidol* 27 5 521 530
39. JE Chambers 2008 PON1 multitasks to protect health *Proc Nat Acad Sci USA* 105 35 12639 12640
40. G Ferretti T Bacchetti S Masciangelo V Bicchiega 2010 HDL-paraoxonase and membrane lipid peroxidation: a comparison between healthy and obese subjects *Obesity* 18 6 1079 1084

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.