



Standardization for oncologic head and neck surgery

Ohad Ronen¹ · K. Thomas Robbins² · Remco de Bree³ · Orlando Guntinas-Lichius⁴ · Dana M. Hartl⁵ · Akihiro Homma⁶ · Avi Khafif⁷ · Luiz P. Kowalski^{8,9} · Fernando López^{10,11,12} · Antti A. Mäkitie¹³ · Wai Tong Ng¹⁴ · Alessandra Rinaldo¹⁵ · Juan P. Rodrigo^{10,11,12} · Alvaro Sanabria^{16,17} · Alfio Ferlito¹⁸

Received: 15 March 2021 / Accepted: 3 May 2021 / Published online: 12 May 2021
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2021

Abstract

The inherent variability in performing specific surgical procedures for head and neck cancer remains a barrier for accurately assessing treatment outcomes, particularly in clinical trials. While non-surgical modalities for cancer therapeutics have evolved to become far more uniform, there remains the challenge to standardize surgery. The purpose of this review is to identify the barriers in achieving uniformity and to highlight efforts by surgical groups to standardize selected operations and nomenclature. While further improvements in standardization will remain a challenge, we must encourage surgical groups to focus on strategies that provide such a level.

Keywords Standardization · Quality assurance · Head and neck cancer · Surgical oncology

Abbreviations

RT Radiotherapy
CT Chemotherapy

MDT Multidisciplinary team
HNSCC Head and neck squamous cell carcinoma

This article was written by members of the International Head and Neck Scientific Group (www.IHNSG.com).

✉ Ohad Ronen
ohadr@gmc.gov.il

¹ Department of Otolaryngology–Head and Neck Surgery, Galilee Medical Center, Affiliated with Azrieli Faculty of Medicine, Bar-Ilan University, Safed, Israel

² Department of Otolaryngology Head and Neck Surgery, Southern Illinois University Medical School, Springfield, IL, USA

³ Department of Head and Neck Surgical Oncology, University Medical Center Utrecht, Utrecht, The Netherlands

⁴ Department of Otorhinolaryngology, Institute of Phoniatriy/Pedaudiology, Jena University Hospital, Jena, Germany

⁵ Head and Neck Oncology Service, Gustave Roussy, Villejuif, France

⁶ Department of Otolaryngology–Head and Neck Surgery, Faculty of Medicine and Graduate School of Medicine, Hokkaido University, Sapporo, Hokkaido, Japan

⁷ Head and Neck Surgery and Oncology Unit, A.R.M. Center for Advanced Otolaryngology Head and Neck Surgery, Assuta Medical Center, Tel Aviv, Israel

⁸ Department of Otorhinolaryngology–Head and Neck Surgery, A.C. Camargo Cancer Center, São Paulo, Brazil

⁹ Department of Head and Neck Surgery, University of São Paulo Medical School, São Paulo, Brazil

¹⁰ Department of Otolaryngology, Hospital Universitario Central de Asturias-ISP, Oviedo, Spain

¹¹ University of Oviedo-IUOPA, Oviedo, Spain

¹² Head and Neck Cancer Unit, CIBERONC, Madrid, Spain

¹³ Department of Otorhinolaryngology–Head and Neck Surgery, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

¹⁴ Department of Clinical Oncology, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong, China

¹⁵ University of Udine School of Medicine, Udine, Italy

¹⁶ Department of Surgery, School of Medicine, Universidad de Antioquia/Hospital Universitario San Vicente Fundación, Medellín, Colombia

¹⁷ CEXCA Centro de Excelencia en Enfermedades de Cabeza Y Cuello, Medellín, Colombia

¹⁸ Coordinator of the International Head and Neck Scientific Group, Padua, Italy

Introduction

Comparisons of treatment outcomes for oncologic surgery are challenging because of the lack of uniformity among specific procedures. In contrast, modern radiotherapy (RT) techniques are more standardized with less variation among centers [1–6]. Both RT and chemotherapy (CT) have a long history in the development of guidelines for uniformity. In contrast, the sequence of steps and extent of surgical procedures vary greatly in head neck surgery, as do the expertise and preferences of individual surgeons. This variability presents a barrier in clinical trials research and in measuring the therapeutic efficacy of selected surgical interventions.

Distinct from achieving standardization of treatment modalities, quality assurance measurements involve assessment of adherence to standardized guidelines. Such initiatives have been implemented to achieve safe and affordable health care, reduced surgical complications, improved hygiene standards, improved patient satisfaction, and lowered health care costs [7–12]. Measuring patient satisfaction, producing cost-analysis comparisons, and reducing complications are all important parts of improving quality; according to the American Society for Quality Control, quality assurance or quality control includes all the planned and systematic activities implemented within the quality system that can be demonstrated to provide confidence that a product or service will fulfill requirements for quality [13]. However, in the absence of standards for surgical technique, the validity of such measures may be unreliable. The purpose of this manuscript is to analyze the challenges that compromise standardization of surgical procedures, highlight some successful projects, and to offer strategies for further progress toward improving standardization. The challenges with suggested solutions are summarized in Table 1.

Challenges for achieving standardization

Surgical training

Operative techniques are primarily learned over time, through hands-on experience with the master modeling behavior and supervising the apprentice. As an analogy, one cannot learn how to drive a car from a book. Courses provided by surgical societies such as the European Head and Neck Society and the American Head and Neck Society (EHNS and AHNS) aim to organize training sessions based on surgical approaches that are generally accepted and evidence-based. Surgeons are exposed to hands-on learning during residency and fellowship training. While techniques among experienced surgeons have common threads, it is not unusual to find variations among different institutions and countries. Moreover, in the absence of formal guidelines, surgical practices vary greatly, complicating efforts to compare outcomes or to perform quality assurance studies.

Host factors

There are various unavoidable obstacles that potentially may interfere with surgical outcomes such as abnormal body habitus, patient comorbidity, and even advanced stage of disease. Dysmorphia may present challenges with surgical exposure of target organs and structures and may interfere with postoperative wound healing. While obesity is detrimental to the general health of patients, a recent meta-analysis found that patients with higher body mass indices had increased overall survival and decreased disease-related mortality and recurrence rate [14]. Such considerations are less prominent when delivering radiotherapy or chemotherapy in a standardized fashion [14]. On the other hand, low skeletal muscle mass, often referred to as sarcopenia, is associated with an increased risk of wound complications and decreased survival [15]. The negative impact of various

Table 1 Recommended actions for achieving standardization

Issue	Suggested solution
Surgical training	Courses and training sessions conducted by surgical societies Broader exposure to multiple surgeons with varying philosophies and skills Use of simulation and technical skills laboratories
Host factors	Inclusion of patients associated factors in clinical trials
Experience	Inclusion of minimal case load and measurable surgery-related measures in clinical trials
Multidisciplinary treatment planning	Promotion of impartial, data-driven, patient-centered recommendations and discussions
Emerging technologies	Formal teaching and mentoring during learning curve period
Heterogeneous surgical procedures	Develop standardized terminology Develop consensus among surgeons from multiple institutions
Quality assurance	Application of surrogate markers and use of measurable outcomes

co-morbidities has been included in the American College of Surgeons National Surgical Quality Improvement Program (ACS NSQIP). The availability of its universal surgical risk calculator allows an online risk calculation derived from a real-world database [16].

Experience

One major factor that correlates with surgical expertise is the number of patients managed and the experience in treating rare or complex cases. This varies greatly across training programs. There is general agreement that treatment of rare cancers such as HNC should be concentrated in high volume, specialized and performed in multidisciplinary centers [17]. A recent paper showed that high volume academic centers had significantly better oncological results than smaller institutions. When compared with high-accruing centers, patients at low-accruing centers had worse overall survival (5 years: 51.0% v 69.1%; $P=0.002$). Treatment at low-accruing centers was associated with an increased death risk of 91% (hazard ratio [HR], 1.91; 95% CI, 1.37–2.65) [18]. In a retrospective study, authors from medical centers in North Carolina managed to show that HNSCC patients treated at academic hospitals, community cancer centers, and hospitals that were in the top third by case volume had more favorable outcomes [19]. From a surgical point of view, the influence of surgeon and/or hospital case volume on head and neck surgery outcomes [20] is lacking. An example of validating surgical quality in the setting of a prospective multicenter randomized trial was recently published by Ferris et al. [21] In this study, a credentials committee composed of ten experienced head and neck surgeons assessed the eligibility of applicant surgeons based on specific minimal case-loads, the maximal number of positive and close margins, the rate of surgery-related bleeding, and the nodal yield from lymphadenectomy. Surgeons were re-evaluated during the study and could be placed on hold based on predetermined criteria.

Multidisciplinary treatment planning

The process of multidisciplinary treatment planning is considered to be an important aspect of cancer treatment that may influence the outcome. Furthermore, it could be argued that such treatment planning may help to compensate for variations in surgical technique as well as for quality assurance [22]. Given the complexity of treating HNSCC patients and the involvement of many disciplines, MDT improves communication among many healthcare professionals who are involved in decision-making processes by bringing together the essential expertise to deliver high-quality cancer care [23, 24]. MDT offers the opportunity for optimal oncological outcomes with less adverse effects of treatment using coordinated professional efforts. Indeed, a balance between

cure and complications remains a central goal, requiring optimization of the therapeutic effect with prudent and individualized application of the various treatment modalities in appropriately selected patients [25]. However, reviews that examined the effect that multidisciplinary teams are mixed with some showing no improvement on survival while others did [26–30]. One possible explanation for this contradiction is the inherent issue of patient compliance for recommended treatment plans [31]. Another possible reason is the personality dynamics within the MDT. The team should be patient-centered and members should express recommendations that are impartial and data-driven.

Emerging technologies

The development of novel surgical techniques that employ advanced technology such a robotic manipulation present added pitfalls for standardization. Most senior surgeons did not receive formal training in the field of minimal access surgery which often involve either lasers or robots. In such situations, the technical expertise of the surgeon may have an influence on outcomes. For example, in trans-oral laser surgery, the experience of the operator has been shown to be inversely correlated with tumor recurrence [32]. Accompanying the challenges of minimal access surgery is the issue of piecemeal resections. Although cutting through tumor is less frequently considered a taboo, [33] its negative effect on clinical outcomes compared to ‘en bloc’ resection remains to be proven. Relevant to piecemeal resection, the advent of minimal access surgical techniques often precludes the removal of tumors as a mono bloc specimen. However, the evidence indicates that mono bloc procedures are not always essential and that tumors can be fragmented as long as radical removal of the tumor is accomplished at the end of the procedure [34, 35].

Progress toward standardization

Several efforts by surgical groups have engaged in projects intended to overcome the heterogeneity associated with surgical procedures. Some of these have focused on terminology and classifications while others have actually analyzed specific techniques to reduce variability and thereby improve accuracy for measuring outcomes in clinical trials.

One example of standardizing terminology was a project on neck dissection types enacted by the American Academy of Otolaryngology – Head and Neck Surgery [36–38]. The exact definitions of the various neck levels and sub-levels were defined in the clinical, [39] radiologic [4] and surgical settings [40]. While this was an important step toward standardization, surgeons still used their own discretion as to the extent of the surgery as well as the sequence of steps during surgery. A second example of efforts to improve terminology

is the work done by the European Laryngological Society [41, 42] in developing a classification for endoscopic cordectomies and supraglottic laryngectomies. The proposal included eight different types according to the surgical approach used and the degree of resection. The intent of this proposal was to reach a better consensus among clinicians and uniformity in reports of the extent and depth of resection in cordectomy procedures. Development of a common language in the head and neck surgical community could allow relevant comparisons of results of surgery within the literature and facilitate standardization of practice [41]. A 2007 revision of this classification, which added endoscopic cordectomy type VI, [43] is currently used in many centers. In 2017, The ELS Working Committee for Nomenclature proposed a clarification of classification, emphasizing the type of laser used and the route this laser is transmitted [44]. The ELS also proposed a systematic nomenclature for endoscopic and open partial horizontal laryngectomies [44, 45].

Categorization of transoral lateral oropharyngectomies is a third example where standardized nomenclature has been recommended [46]. The increase in incidence of human papilloma virus (HPV) related oropharyngeal squamous cell carcinoma was followed by a rise in surgical transoral excision of the lateral oropharynx. Similar to Remacle et al. [43] the authors managed to construct a comprehensive yet simple classification system of the depth of lateral pharyngeal resection and the direction of resection for disease extension.

With regard to standardizing operative techniques, the Japan Neck Dissection Study Group launched a collaborative national effort to reach a consensus on a uniform neck dissection procedure [47]. This was prompted by the lack of consensus by the authors in comparing the specific aspects of the various non-radical neck dissections. To unify the understanding of the procedure, a formal protocol was initiated. During the initial observation phase, the group found the technique for neck dissection to be highly variable among the participating institutions. In essence, surgeons used the same terminology to describe slightly different procedures. The group managed to compare and standardize the procedure mainly by requiring surgeons to observe their colleagues perform the technique in different institutions. A construct of 79 items was developed, thus allowing the procedures to be analysed based on general and detailed anatomic data. Through this approach, surgeons could observe subtle differences compared to their own practice and adjust their own techniques accordingly. After a second observation stage, improvements in standardization were documented.

Similarly, the effort to standardize sentinel node biopsy for patients with clinically node-negative oral cavity squamous cell carcinoma represents another example of unifying surgical guidelines based on available evidence and expert opinion. A consensus statement on this issue was developed during the eighth international symposium for sentinel node

biopsy in head and neck cancer held in London in 2018 [48]. In this international project, a wide variety of topics ranging from patient selection, surgical technique, to management was debated and agreed upon. Potential areas of development were highlighted for future prospective studies.

Strategies to expand standardization

While there remain a number of obstacles to achieving standardization in oncologic surgery, further improvements are possible. The continued application of indirect measures and surrogate markers, as demonstrated by Ferris et al. [21] represents one approach to assess uniformity and the quality of surgery. Also, there remain opportunities to improve the terminology for selected operations such as glossectomy, maxillectomy and mandibulectomy along with its array of variations [49–51]. For example, currently there are many suggested nomenclatures for mandibulectomy, but none is recognized as the standard [51–57]. Such variations make it difficult to compare reports of different outcomes, particularly with regard to surgical complications that affect not only the quality of life, but essential functions in the head and neck region [58]. Similar to the consensus reached for classifying neck dissection, agreements could be reached for other operative procedures that have numerous variations.

In terms of standardizing the specific components of a surgical procedure, projects like the Japan Neck Dissection Study Group [47] and the European Laryngologic Society represent significant contributions. Such examples lead one

Table 2 Potential surrogate markers and measurable outcomes for quality analysis in head and neck surgery

Quality surrogate markers and measurable outcomes
<i>I Pre-operative</i>
Multidisciplinary team meetings
High-volume surgeons
High-volume hospitals
<i>II Intra-operative</i>
Negative surgical margins
Nerve monitoring
Nodal yield
<i>III Immediate post-operative</i>
Early complications
Functional impairment
Mortality
Days of admission
<i>IV Late post-operative</i>
Late complications
Locoregional failure rate
Survival outcomes
Quality of life measurements

to surmise that such coordinated exercises can be extended to other operations. The question becomes what other procedures should be addressed and by whom. Definitely, whenever a specific surgical intervention becomes an influential component of treatment outcomes in a prospective clinical trial, there is an important need to standardize it. Under such circumstances, this can be achieved through the cooperation of the participating surgeons to reach agreements on technique, use of surrogate markers for quality assurance, and acceptable levels for complications and adverse surgical outcomes (Table 2).

Beyond the existing strategies applied to enhance uniformity, recent technologies and approaches offer novel opportunities for future progress. For example, simulation laboratories have become an integral component for teaching surgical techniques [59]. While the emphasis to date has been on training surgical residents, simulation technology could be redirected to improve standardization for selected procedures [60]. Also, simulation provides uniform and objective metrics for use in the assessment of technical skills [61]. One could imagine this approach becoming an important step in developing and activating a treatment protocol in which a surgical intervention is being measured within a multi-institutional setting. An additional strategy that has now moved beyond the realm of novelty is the application of virtual conferencing. In addition to increasing the opportunities to communicate both verbally and visually among participating surgeons, the virtual sharing of sound and imagery to outline selected operative techniques has become a reality. Such advances provide a sense of optimism for advancing a level of standardization within the field.

Author contributions OR: conceived the idea of the paper, OR and KTR: performed literature research, and drafted the manuscript; all authors: provided scientific revision of the manuscript.

References

- Chang ATY, Tan LT, Duke S, Ng W-T (2017) Challenges for quality assurance of target volume delineation in clinical trials. *Front Oncol* 7:221. <https://doi.org/10.3389/fonc.2017.00221>
- Hong TS, Tomé WA, Harari PM (2012) Heterogeneity in head and neck IMRT target design and clinical practice. *Radiother Oncol J Eur Soc Ther Radiol Oncol* 103:92–98. <https://doi.org/10.1016/j.radonc.2012.02.010>
- Rasch C, Steenbakkers R, van Herk M (2005) Target definition in prostate, head, and neck. *Semin Radiat Oncol* 15:136–145. <https://doi.org/10.1016/j.semradonc.2005.01.005>
- Brouwer CL, Steenbakkers RJHM, Bourhis J et al (2015) CT-based delineation of organs at risk in the head and neck region: DAHANCA, EORTC, GORTEC, HKNPCSG, NCIC CTG, NCRI, NRG Oncology and TROG consensus guidelines. *Radiother Oncol J Eur Soc Ther Radiol Oncol* 117:83–90. <https://doi.org/10.1016/j.radonc.2015.07.041>
- Grégoire V, Grau C, Lapeyre M, Maingon P (2018) Target volume selection and delineation (T and N) for primary radiation treatment of oral cavity, oropharyngeal, hypopharyngeal and laryngeal squamous cell carcinoma. *Oral Oncol* 87:131–137. <https://doi.org/10.1016/j.oraloncology.2018.10.034>
- Grégoire V, Ang K, Budach W et al (2014) Delineation of the neck node levels for head and neck tumors: a 2013 update. DAHANCA, EORTC, HKNPCSG, NCIC CTG, NCRI, RTOG, TROG consensus guidelines. *Radiother Oncol J Eur Soc Ther Radiol Oncol* 110:172–181. <https://doi.org/10.1016/j.radonc.2013.10.010>
- Ang D, McKenney M, Norwood S et al (2015) Benchmarking statewide trauma mortality using agency for healthcare research and quality's patient safety indicators. *J Surg Res* 198:34–40. <https://doi.org/10.1016/j.jss.2015.05.053>
- Cima RR, Lackore KA, Nehring SA et al (2011) How best to measure surgical quality? Comparison of the agency for healthcare research and quality patient safety indicators (AHRQ-PSI) and the American college of surgeons national surgical quality improvement program (ACS-NSQIP) postoperative adverse e. *Surgery* 150:943–949. <https://doi.org/10.1016/j.surg.2011.06.020>
- Sedman A, Harris JM 2nd, Schulz K et al (2005) Relevance of the agency for healthcare research and quality patient safety indicators for children's hospitals. *Pediatrics* 115:135–145. <https://doi.org/10.1542/peds.2004-1083>
- Simon C, Dietz A, Leemans CR (2019) Quality assurance in head and neck cancer surgery: where are we, and where are we going? *Curr Opin Otolaryngol Head Neck Surg* 27:151–156. <https://doi.org/10.1097/MOO.0000000000000519>
- Weber RS (2007) Improving the quality of head and neck cancer care. *Arch Otolaryngol Head Neck Surg* 133:1188–1192
- Yuan F, Chung KC (2016) Defining quality in health care and measuring quality in surgery. *Plast Reconstr Surg* 137:1635–1644. <https://doi.org/10.1097/PRS.0000000000002028>
- American Society for Quality. <https://asq.org/>. Accessed 14 Mar 2021
- den Hollander D, Kampman E, van Herpen CML (2015) Pre-treatment body mass index and head and neck cancer outcome: A review of the literature. *Crit Rev Oncol Hematol* 96:328–338. <https://doi.org/10.1016/j.critrevonc.2015.06.002>
- Economopoulou P, de Bree R, Kotsantis I, Psyrris A (2019) Diagnostic tumor markers in head and neck squamous cell carcinoma (HNSCC) in the clinical setting. *Front Oncol* 9:827. <https://doi.org/10.3389/fonc.2019.00827>
- Vosler PS, Orsini M, Enepekides DJ, Higgins KM (2018) Predicting complications of major head and neck oncological surgery: an evaluation of the ACS NSQIP surgical risk calculator. *J Otolaryngol Head Neck Surg* 47:21. <https://doi.org/10.1186/s40463-018-0269-8>
- Orlandi E, Alfieri S, Simon C et al (2019) Treatment challenges in and outside a network setting: Head and neck cancers. *Eur J Surg Oncol J Eur Soc Surg Oncol Br Assoc Surg Oncol* 45:40–45. <https://doi.org/10.1016/j.ejso.2018.02.007>
- Wuthrick EJ, Zhang Q, Machtay M et al (2015) Institutional clinical trial accrual volume and survival of patients with head and neck cancer. *J Clin Oncol Off J Am Soc Clin Oncol* 33:156–164. <https://doi.org/10.1200/JCO.2014.56.5218>
- Farquhar DR, Masood MM, Lenze NR et al (2020) Academic affiliation and surgical volume predict survival in head and neck cancer patients receiving surgery. *Laryngoscope*. <https://doi.org/10.1002/lary.28744>
- Alfieri S, Orlandi E, Bossi P (2017) the case volume issue in head and neck oncology. *Curr Treat Options Oncol* 18:65. <https://doi.org/10.1007/s11864-017-0507-8>

21. Ferris RL, Flamand Y, Holsinger FC et al (2020) A novel surgeon credentialing and quality assurance process using transoral surgery for oropharyngeal cancer in ECOG-ACRIN Cancer Research Group Trial E3311. *Oral Oncol* 110:104797. <https://doi.org/10.1016/j.oraloncology.2020.104797>
22. Takes RP, Halmos GB, Ridge JA et al (2020) Value and quality of care in head and neck oncology. *Curr Oncol Rep* 22:92. <https://doi.org/10.1007/s11912-020-00952-5>
23. Jalil R, Ahmed M, Green JSA, Sevdalis N (2013) Factors that can make an impact on decision-making and decision implementation in cancer multidisciplinary teams: an interview study of the provider perspective. *Int J Surg* 11:389–394. <https://doi.org/10.1016/j.ijso.2013.02.026>
24. Morton RP (2012) Toward comprehensive multidisciplinary care for head and neck cancer patients: quality of life versus survival. *Otolaryngol neck Surg Off J Am Acad Otolaryngol Neck Surg* 147:404–406. <https://doi.org/10.1177/0194599812450856>
25. Bossi P, Alfieri S (2016) The benefit of a multidisciplinary approach to the patient treated with (Chemo) radiation for head and neck cancer. *Curr Treat Options Oncol* 17:53. <https://doi.org/10.1007/s11864-016-0431-3>
26. Croke JM, El-Sayed S (2012) Multidisciplinary management of cancer patients: chasing a shadow or real value? An overview of the literature. *Curr Oncol* 19:e232–e238. <https://doi.org/10.3747/co.19.944>
27. Pillay B, Wootten AC, Crowe H et al (2016) The impact of multidisciplinary team meetings on patient assessment, management and outcomes in oncology settings: A systematic review of the literature. *Cancer Treat Rev* 42:56–72. <https://doi.org/10.1016/j.ctrv.2015.11.007>
28. Wang Y-H, Kung P-T, Tsai W-C et al (2012) Effects of multidisciplinary care on the survival of patients with oral cavity cancer in Taiwan. *Oral Oncol* 48:803–810. <https://doi.org/10.1016/j.oraloncology.2012.03.023>
29. Friedland PL, Bozic B, Dewar J et al (2011) Impact of multidisciplinary team management in head and neck cancer patients. *Br J Cancer* 104:1246–1248. <https://doi.org/10.1038/bjc.2011.92>
30. Prades J, Remue E, van Hoof E, Borrás JM (2015) Is it worth reorganising cancer services on the basis of multidisciplinary teams (MDTs)? A systematic review of the objectives and organisation of MDTs and their impact on patient outcomes. *Health Policy* 119:464–474. <https://doi.org/10.1016/j.healthpol.2014.09.006>
31. Rangabashyam MS, Lee SY, Tan SY et al (2020) Adherence of head and neck squamous cell carcinoma patients to tumor board recommendations. *Cancer Med* 9:5124–5133. <https://doi.org/10.1002/cam4.3097>
32. Vilaseca I, Nogués-Sabaté A, Avilés-Jurado FX et al (2019) Factors of local recurrence and organ preservation with transoral laser microsurgery in laryngeal carcinomas; CHAID decision-tree analysis. *Head Neck* 41:756–764. <https://doi.org/10.1002/hed.25422>
33. Robbins KT, Bradford CR, Rodrigo JP et al (2016) Removing the taboo on the surgical violation (Cut-Through) of Cancer. *JAMA Otolaryngol - Head Neck Surg* 142:1010–1013. <https://doi.org/10.1001/jamaoto.2016.1826>
34. Ambrosch P (2007) The role of laser microsurgery in the treatment of laryngeal cancer. *Curr Opin Otolaryngol Head Neck Surg* 15:82–88. <https://doi.org/10.1097/MOO.0b013e3280147336>
35. Snyderman CH, Carrau RL, Kassam AB et al (2008) Endoscopic skull base surgery: principles of endonasal oncological surgery. *J Surg Oncol* 97:658–664. <https://doi.org/10.1002/jso.21020>
36. Robbins KT, Medina JE, Wolfe GT et al (1991) Standardizing neck dissection terminology. Official report of the academy's committee for head and neck surgery and oncology. *Arch Otolaryngol Head Neck Surg* 117:601–605
37. Robbins KT, Clayman G, Levine PA et al (2002) Neck dissection classification update: revisions proposed by the American head and neck society and the American academy of otolaryngology-head and neck surgery. *Arch Otolaryngol Head Neck Surg* 128:751–758
38. Robbins KT, Shaha AR, Medina JE et al (2008) Consensus statement on the classification and terminology of neck dissection. *Arch Otolaryngol Head Neck Surg* 134:536–538. <https://doi.org/10.1001/archotol.134.5.536>
39. Shah JP, Strong E, Spiro RH, Vikram B (1981) Surgical grand rounds. Neck dissection: current status and future possibilities. *Clin Bull* 11:25–33
40. Ronen O, Samant S, Robbins KT (2021) Cummings Otolaryngology - Head and Neck Surgery. In: Flint PW, Haughey BH, Lund VJ et al (eds) Cummings Otolaryngology - Head and Neck Surgery, 7th edn. Elsevier Inc., Philadelphia, pp 1805–1830
41. Remacle M, Eckel HE, Antonelli A et al (2000) Endoscopic cordectomy. A proposal for a classification by the Working Committee, European Laryngological Society. *Eur Arch Oto-Rhino-Laryngol Off J Eur Fed Oto-Rhino-Laryngol Soc Affil with Ger Soc Oto-Rhino-Laryngol Head Neck Surg* 257:227–231. <https://doi.org/10.1007/s004050050228>
42. Remacle M, Hantzakos A, Eckel H et al (2009) Endoscopic supraglottic laryngectomy: a proposal for a classification by the working committee on nomenclature, European Laryngological Society. *Eur Arch Oto-Rhino-Laryngol Off J Eur Fed Oto-Rhino-Laryngol Soc Affil with Ger Soc Oto-Rhino-Laryngol - Head Neck Surg* 266:993–998. <https://doi.org/10.1007/s00405-008-0901-8>
43. Remacle M, Van Haverbeke C, Eckel H et al (2007) Proposal for revision of the European Laryngological Society classification of endoscopic cordectomies. *Eur Arch Oto-Rhino-laryngol Off J Eur Fed Oto-Rhino-Laryngol Soc Affil with Ger Soc Oto-Rhino-Laryngol - Head Neck Surg* 264:499–504. <https://doi.org/10.1007/s00405-007-0279-z>
44. Remacle M, Arens C, Eldin MB et al (2017) Laser-assisted surgery of the upper aero-digestive tract: a clarification of nomenclature. A consensus statement of the European Laryngological Society. *Eur Arch Oto-Rhino-laryngol Off J Eur Fed Oto-Rhino-Laryngol Soc Affil with Ger Soc Oto-Rhino-Laryngol - Head Neck Surg* 274:3723–3727. <https://doi.org/10.1007/s00405-017-4708-3>
45. Succo G, Peretti G, Piazza C et al (2014) Open partial horizontal laryngectomies: a proposal for classification by the working committee on nomenclature of the European Laryngological Society. *Eur Arch Oto-Rhino-laryngol Off J Eur Fed Oto-Rhino-Laryngol Soc Affil with Ger Soc Oto-Rhino-Laryngol - Head Neck Surg* 271:2489–2496. <https://doi.org/10.1007/s00405-014-3024-4>
46. De Virgilio A, Kim S-H, Magnuson JS et al (2019) Anatomical-based classification for transoral lateral oropharyngectomy. *Oral Oncol* 99:104450. <https://doi.org/10.1016/j.oraloncology.2019.104450>
47. Saikawa M, Kishimoto S (2010) Standardizing the extent of resection in nonradical neck dissections: The final report of the Japan neck dissection study group prospective study. *Int J Clin Oncol* 15:13–22. <https://doi.org/10.1007/s10147-009-0016-2>
48. Schilling C, Stoeckli SJ, Vigili MG et al (2019) Surgical consensus guidelines on sentinel node biopsy (SNB) in patients with oral cancer. *Head Neck* 41:2655–2664. <https://doi.org/10.1002/hed.25739>
49. Ansarin M, Bruschini R, Navach V et al (2019) Classification of glossectomies: Proposal for tongue cancer resections. *Head Neck* 41:821–827. <https://doi.org/10.1002/hed.25466>
50. Brown JS, Rogers SN, McNally DN, Boyle M (2000) A modified classification for the maxillectomy defect. *Head Neck* 22:17–26. [https://doi.org/10.1002/\(sici\)1097-0347\(200001\)22:1%3c17::aid-hed4%3e3.0.co;2-2](https://doi.org/10.1002/(sici)1097-0347(200001)22:1%3c17::aid-hed4%3e3.0.co;2-2)

51. Brown JS, Barry C, Ho M, Shaw R (2016) A new classification for mandibular defects after oncological resection. *Lancet Oncol* 17:e23–30. [https://doi.org/10.1016/S1470-2045\(15\)00310-1](https://doi.org/10.1016/S1470-2045(15)00310-1)
52. Baumann DP, Yu P, Hanasono MM, Skoracki RJ (2011) Free flap reconstruction of osteoradionecrosis of the mandible: a 10-year review and defect classification. *Head Neck* 33:800–807. <https://doi.org/10.1002/hed.21537>
53. David DJ, Tan E, Katsaros J, Sheen R (1988) Mandibular reconstruction with vascularized iliac crest: a 10-year experience. *Plast Reconstr Surg* 82:792–803. <https://doi.org/10.1097/00006534-198811000-00011>
54. Iizuka T, Häfliger J, Seto I et al (2005) Oral rehabilitation after mandibular reconstruction using an osteocutaneous fibula free flap with endosseous implants. Factors affecting the functional outcome in patients with oral cancer. *Clin Oral Implants Res* 16:69–79. <https://doi.org/10.1111/j.1600-0501.2004.01076.x>
55. Jewer DD, Boyd JB, Manktelow RT et al (1989) Orofacial and mandibular reconstruction with the iliac crest free flap: a review of 60 cases and a new method of classification. *Plast Reconstr Surg* 84:391–395
56. Pavlov BL (1974) Classification of mandibular defects. *Stomatologiia (Mosk)* 53:43–46
57. Urken ML, Weinberg H, Vickery C et al (1991) Oromandibular reconstruction using microvascular composite free flaps. Report of 71 cases and a new classification scheme for bony, soft-tissue, and neurologic defects. *Arch Otolaryngol Head Neck Surg* 117:733–744. <https://doi.org/10.1001/archotol.1991.01870190045010>
58. Semenov YR, Starmer HM, Gourin CG (2012) The effect of pneumonia on short-term outcomes and cost of care after head and neck cancer surgery. *Laryngoscope* 122:1994–2004. <https://doi.org/10.1002/lary.23446>
59. Sarker SK, Patel B (2007) Simulation and surgical training. *Int J Clin Pract* 61:2120–2125. <https://doi.org/10.1111/j.1742-1241.2007.01435.x>
60. Zevin B, Levy JS, Satava RM, Grantcharov TP (2012) A consensus-based framework for design, validation, and implementation of simulation-based training curricula in surgery. *J Am Coll Surg* 215:580–586.e3. <https://doi.org/10.1016/j.jamcollsurg.2012.05.035>
61. Wiet GJ, Stredney D, Wan D (2011) Training and simulation in otolaryngology. *Otolaryngol Clin North Am* 44:1333–1350. <https://doi.org/10.1016/j.otc.2011.08.009>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.