



Inter- α -inhibitor Ameliorates Endothelial Inflammation in Sepsis

Vandy P. Stober¹ · Yow-Pin Lim² · Steven Opal² · Lisheng Zhuo³ · Koji Kimata³ · Stavros Garantziotis¹

Received: 19 February 2019 / Accepted: 12 April 2019 / Published online: 26 April 2019

© This is a U.S. Government work and not under copyright protection in the US; foreign copyright protection may apply 2019

Abstract

Purpose Vascular endothelial cells demonstrate severe injury in sepsis, and a reduction in endothelial inflammation would be beneficial. Inter- α -Inhibitor (I α I) is a family of abundant plasma proteins with anti-inflammatory properties and has been investigated in human and animal sepsis with encouraging results. We hypothesized that I α I may protect endothelia from sepsis-related inflammation.

Methods I α I-deficient or sufficient mice were treated with endotoxin or underwent complement-induced lung injury. VCAM-1 and ICAM-1 expression was measured in blood and lung as marker of endothelial activation. Human endothelia were exposed to activated complement C5a with or without I α I. Blood from human sepsis patients was examined for VCAM-1 and ICAM-1 and levels were correlated with blood levels of I α I.

Results I α I-deficient mice showed increased endothelial activation in endotoxin/sepsis- and complement-induced lung injury models. In vitro, levels of endothelial pro-inflammatory cytokines and cell growth factors induced by activated complement C5a were significantly decreased in the presence of I α I. This effect was associated with decreased ERK and NF κ B activation. I α I levels were inversely associated with VCAM-1 and ICAM-1 levels in a human sepsis cohort.

Conclusions I α I ameliorates endothelial inflammation and may be beneficial as a treatment of sepsis.

Keywords Sepsis · Complement · Inter-alpha-inhibitor · Endothelial injury · Lipopolysaccharide

Introduction

Sepsis, defined as the systemic inflammatory response to infection, remains a significant health problem worldwide. In the United States, septic shock causes >200,000 deaths annually at a cost of >\$15 billion [1]. Sepsis treatment remains largely supportive, and mortality is high at 30–50% [2], mostly from multi-organ failure (MOF) [3].

Sepsis causes endothelial injury, which induces inflammatory mediators, tissue factors, and coagulation pathways, leading to tissue hypoxia and MOF. Several mechanisms lead to endothelial injury in sepsis, for example, recent evidence suggests that oxidative stress and inflammation lead to damage in the endothelial glycocalyx, a gel-like layer

covering the luminal surface of endothelial cells and shedding of glycocalyx components like heparin sulfate into the circulation [4]. This in turn leads to a disruption in the hemostatic function of endothelia and their ability to function effectively as a barrier [5]. The ultimate result is disruption of the microcirculation, leading to organ dysfunction which is a major hallmark of severe sepsis. Several upstream mediators promote endothelial injury, like bacterial endotoxins, reactive oxygen species, and inflammatory cytokines; however, activation of the complement cascade is a major contributor and resides at the apex of the pathophysiological processes leading to endothelial injury and organ failure [6].

Complement is a family of serum proteins acting via a catalytic cascade to amplify inflammation and induce opsonization and chemotaxis. Activated complement component 5 (C5a) is a central mediator of the complement cascade, and promotes a strong inflammatory response [7]. Blockade of C5a generation improves survival and decreases MOF in sepsis models [8].

Complement activation is central in endothelial injury [9]. C5a directly activates endothelial cells causing upregulation of cellular adhesion molecules E-selectin, vascular cell

✉ Stavros Garantziotis
garantziotis@niehs.nih.gov

¹ National Institute of Environmental Health Sciences, 111 TW Alexander Dr, Research Triangle Park, NC 27709, USA

² Brown University, East Providence, RI, USA

³ Research Complex for Medicine Frontiers, Aichi Medical University, Nagakute, Aichi, Japan

adhesion molecule-1 (VCAM-1), and intercellular adhesion molecule 1 (ICAM-1) [10]. These molecules assist inflammatory cell binding to activated endothelia and infiltration into the inflamed tissues [11–13].

Inter-alpha-inhibitor (I α I) is a family of plasma heavy- and light-chain anti-inflammatory proteins which have been investigated in sepsis with encouraging results (6). Two major forms of the protein family are found in circulation: Inter- α -inhibitor and Pre- α -inhibitor. Both proteins are collectively referred to as I α I in this report. I α I administration improves survival in sepsis models, e.g., after administration of lipopolysaccharide (LPS) [14–16]. Importantly, I α I administration in sepsis improves coagulation parameters which are markers of endothelial dysfunction [16]. We have shown that I α I inhibits complement activation and ameliorates lung injury in sepsis [17]. We now investigated the I α I effect on LPS- and C5a-induced endothelial activation. We show that I α I-deficient mice have increased markers of endothelial activation (ICAM-1, VCAM-1) in two sepsis models. We further show *in vitro* that I α I decreases ERK and NF κ B signaling and the downstream expression of inflammatory cytokines after C5a exposure. Finally, we demonstrate an inverse correlation of I α I serum levels with VCAM-1 and ICAM-1 levels in human sepsis patients. In aggregate, our results suggest that I α I protects endothelial cells from inflammation by activated complement, thus being beneficial in sepsis.

Materials and Methods

Experimental Animals

I α I-deficient mice have been described previously [17]. Mice were backcrossed > 10 generations into the C57BL/6J mouse strain before used. Experimental animals were provided water and chow *ad libitum* and were kept in an enriched environment with a 12-h day-night cycle. All protocols were approved by the National Institute of Environmental Health Sciences Animal Care Committee.

Endothelial Cell Exposures

HUVEC cells were plated at 5000 cells/cm² in 96-well dishes. Cells were incubated in a humidified incubator at 37 °C with 5% CO₂ until they are 80% confluent in EGM (Lonza, Anaheim, CA) with 2% FBS. 24 h prior to exposure, cells were serum starved with basal EGM and incubated with either C5a (50 nM), C5a (50 nM), and I α I (50 nM), or media only. Supernatant was assessed using Bioplex (Bio-rad, Hercules, CA).

Western Blot Analysis

Lysates were collected after washing the cell layer twice with 1 ml ice-cold TBS in RIPA (Sigma-Aldrich, St Louis, MO) with HALT proteinase inhibitors (Thermo Scientific, Waltham, MA). Protein levels were quantified by BCA (Thermo Scientific). Western blot membranes were blocked with 4% Milk/TBS overnight and incubated for 4 h in TBS/0.5% BSA. Primary antibodies were ERK1/2 p44/42, Phospho-p44/42 ERK1/2 (Thr202/Tyr204), and NF κ B p100/p52 (1:1000, Cell Signaling, Danvers, MA). Goat anti-Rabbit HRP secondary antibody was incubated at 1:10,000 and developed using Supersignal chemiluminescent substrate (Thermo-Fisher Scientific, Waltham, MA) and developed using BIOMax film (Kodak, Rochester, NY). Densitometry analysis was with ImageJ (National Institutes of Health, Bethesda, MD).

In Vivo Sepsis Model

LPS: Mice aged 6–8 weeks received *i.p.* LPS at 10 mg/kg (low dose), or 40 mg/kg (high dose) body weight and were sacrificed 24 h later by CO₂.

C5a: Anti-BSA rabbit-fractionated antiserum (Sigma-Aldrich) was reconstituted in sterile-filtered ddH₂O, diluted in LPS-free PBS to a concentration of 250 μ g in 50 μ l, and instilled retropharyngeally in isoflurane-anesthetized mice. Mice were then given 1 mg of LPS-free BSA (Equitech-Bio, Kerrville, TX) in 400 μ l of LPS-free PBS into the retro-orbital sinus and sacrificed 4 h later by CO₂.

For all mice, plasma was collected and lung tissue was snap frozen and stored at -80°C until used.

Quantitative RT-PCR

RNA was isolated from mouse lungs via the Trizol method. Expression of ICAM-1, VCAM-1, TNF- α , and E-selectin was examined (primers are listed in Table 1). Results were normalized to the ribosomal protein 18S.

Immunohistochemistry

Formalin-fixed mouse lungs were sectioned at 5 μ m and stained with rabbit anti-I α I (Dako, Carpinteria, CA), or biotinylated HA binding protein (Seikagaku America, East Falmouth, MA),

Human Clinical Sepsis Study

Plasma samples were from a previous prospective multi-center phase III clinical trial of 266 patients with severe

Table 1 Primer sequences for adhesion molecules and cytokines used in this study

Gene	Forward	Reverse
ICAM-1	GAG AGT GGA CCC AAC TGG AA	GCC ACA GTT CTC AAA GCA CA
VCAM-1	CAA TGG GGT GGT AAG GAA	GTC ACA GCG CAC AGG TAA GA
TNF α	TGT TGC CTC CTC TTT TGC TT	TGG TCA CCA AAT CAG CGT TA
E-selectin	AGT CTA GCG CCT GGA TGA AA	CCA GCG AGG AGA ACA AAA AC
18s	CGG CTA CCA CAT CCA AGG AA	GCT GGA ATT ACC GCG GCT

sepsis [18]. Specimens used for this study were exempt from IRB approval according to 45 CFR Part 46.1 01 (b) (4) because (a) this study used existing specimens collected for other research purposes or non-research activities; (b) subject identity and personal health information was de-identified and no clinical data were prospectively collected. Plasma samples were analyzed at admission (day 0) and day 5. Because some plasma samples had been used up, only 90 day-0 samples were analyzed for VCAM-1, 112 day-0 samples for ICAM-1, and 36 day-5 samples were analyzed for either. Investigators were blinded to plasma α I level results. DuoSet ELISA kits (R&D Systems, Minneapolis, MN) were used for sVCAM-1 and sICAM-1 measurements. Absorbance was measured at 490 nm/540 nm using Biotek Powerwave XS. Plasma α I levels were measured by investigators who were unaware of sVCAM-1/sICAM-1 results. α I levels were determined with a quantitative competitive ELISA using a single monoclonal antibody (MAb 69.26). Measurements were performed in triplicate.

Statistical Analysis and Regression Model

Analysis was performed using the GraphPad Prism 6 software (GraphPad Software, La Jolla, CA). ANOVA with Bonferroni post hoc test and Student's *t* test was used and $P < 0.05$ was considered significant.

Results

α I Deficiency is Associated with Increased Markers of Endothelial Injury in Sepsis Models

E-selectin, VCAM-1, and ICAM-1 are well-described markers of endothelial activation in sepsis, promote leukocyte adhesion on activated endothelia, and have been associated with morbidity, MOF, and prognosis [19, 20]. We determined the effect of α I deficiency on the expression of endothelial activation in two sepsis and organ injury models. Gene expression in lung tissue of α I-deficient mice showed significant increase in VCAM-1, ICAM-1, and E-selectin (Fig. 1a) when compared to α I-sufficient mice. Soluble VCAM-1 (sVCAM-1) is shed from activated endothelia, and in general is associated with severity of




Fig. 1 α I-deficient mice dosed with LPS had increased expression of vascular adhesion molecules compared to α I-sufficient mice. **a** Gene expression in lung tissue of VCAM-1, ICAM-1, and E-selectin was severalfold higher in α I-deficient mice. **b** Increased levels of sVCAM-1 in LPS-exposed α I-deficient mice for high- and low-LPS dose

endothelial inflammation in septic patients [20]. Because measurement of VCAM-1 tissue expression is not easily feasible in septic patients, many investigators use sVCAM-1 as a surrogate [19]. Since we investigated the association of sVCAM-1 with serum α I in sepsis patients (see below), we tested whether sVCAM-1 levels parallels tissue VCAM-1 expression in our mouse model. We found a significant increase in serum sVCAM-1 in α I-deficient mice compared to sufficient mice, 24 h post intraperitoneal exposure to 10 mg/Kg or 40 mg/Kg LPS (Fig. 1b), confirming that both tissue expression and serum levels are increased in the absence of α I. We also found significant upregulation of VCAM-1 and ICAM-1 in lung tissue of α I-deficient mice compared to sufficient mice (Fig. 2a) in a second sepsis model, immune complex lung injury [17]. We then assessed serum cytokines and chemokines linked to endothelial inflammation in the context of sepsis [21–27]. We found significantly elevated levels of VEGF,




Fig. 2 IαI-deficient mice in the C5a-induced lung injury model had increased endothelial injury and inflammation compared to IαI-sufficient mice. **a** Gene expression in lung tissue of VCAM-1 and ICAM-1 was 2- to 3-fold higher in IαI-deficient mice. **b** Endothelial growth factors and relevant cytokine levels in lung lavage fluid of exposed mice were higher in IαI-deficient mice compared to IαI-sufficient mice

PDGF, CXCL-9, M-CSF, IL-18, and GM-CSF in IαI-deficient mice relative to sufficient mice (Fig. 2b).

IαI Ameliorates Endothelial Inflammation Induced by C5a In Vitro

We then investigated the effect of IαI on endothelial inflammation induced by C5a, using an in vitro model of human umbilical vascular endothelial cells (HUVEC). We observed a strong upregulation of inflammatory cytokines after exposure of HUVEC to C5a. IαI significantly ameliorated C5a effects on IL-6, FGF, GM-CSF, MCP-1, PDGF, and IFN-γ. For some cytokines, e.g., IL-9, IαI did not have a protective

effect on C5a-induced upregulation (Fig. 3a, b). Taken together our results suggest that, in addition to inhibiting complement activation [17], IαI also inhibits C5a-induced inflammation in endothelial cells.

IαI Ameliorates NfκB And ERK Activation After Endothelial Exposure To C5a In Vitro

To further elucidate the mechanism of IαI protection against C5a-induced endothelial inflammation, we investigated the effect of IαI on signaling pathways which are known to be activated by C5a in endothelia, such as ERK [28] and NFκB [10], in the HUVEC model. We found that IαI significantly reduced ERK and NFκB activation after C5a exposure (Fig. 3c, d).

Plasma IαI Levels are Inversely Associated with sVCAM-1 and sICAM-1 Levels in Human Sepsis Patients

It has been previously shown that IαI plasma levels on admission are directly associated with survival and inversely correlated with organ failure scores and plasma IL-6 levels in a cohort of sepsis patients [18]. We utilized the same cohort to investigate whether there was an association between plasma IαI levels and plasma VCAM-1 and ICAM-1 levels, as markers of endothelial inflammation. We found a modest but significant inverse correlation between IαI levels and plasma VCAM-1 and ICAM-1 in sepsis patients on admission (day 0), but not at day 5 after admission (Fig. 4). This finding agreed with the experimental results in mouse sepsis models and in vitro endothelial injury models.

Discussion

It is increasingly recognized that sepsis therapy should not be limited to antimicrobial treatment. Instead, great health benefits may be realized by correcting the dysregulated immune response which underlies septic shock pathobiology [29]. Endothelial inflammation is a hallmark of sepsis [30–32] and leads to shock and MOF [33, 34]. Several cytokines produced by activated endothelia have effects on the endothelia themselves and immune cells that interact with them: GM-CSF promotes inflammatory cell adhesion to endothelia [35]; PDGF promotes endothelial integrity [36] and is associated with survival in sepsis [27]; IL-6 has a role in angiogenesis [37]; FGF-2 promotes endothelial migration through activation of ERK1/2 [38]; and MCP-1 expression by endothelia promotes the accumulation of monocytes in the injury site [39]. Endothelial activation induces expression of VCAM-1 and ICAM-1, which are involved in sepsis pathogenesis and are markers of end-organ




Fig. 3 In vitro exposure of HUVEC to C5a demonstrated amelioration of the inflammatory response by IαI. **a** IαI reduced levels of IL-6, FGF-2 and GM-CSF that were increased by C5a. **b** IαI increased levels of MCP-1, PDGF, and IFN-γ that were reduced by

C5a exposure. **c** IαI decreased activation of ERK and **d** IαI decreased activation of NFκB when dosed together with C5a. PBS=phosphate buffered saline vehicle

failure, morbidity, and mortality in severe sepsis [20, 40]. Thus, VCAM-1 and ICAM-1 are representative markers of endothelial injury and activation in sepsis [19, 20]. We have now demonstrated that IαI suppresses VCAM-1 and ICAM-1 in two distinct models of sepsis lung injury (LPS and C5a). IαI also normalized in vitro the expression of relevant cytokines and endothelial factors. In addition, IαI is inversely associated with soluble VCAM-1 and ICAM-1 levels in patients admitted with sepsis. IαI is abundant in the circulation (normal levels of 0.2–0.6 mg/ml) and can be found in direct contact with vascular endothelia (Fig. 5). Thus, IαI is an excellent candidate modifier protein for endothelial inflammation and dysfunction in sepsis.

VCAM-1 and ICAM-1 upregulation in sepsis is mediated by inflammatory stimuli present in the serum of septic animals [41], including cytokines such as TNFα, IFN-γ, and IL-1β, glycosaminoglycans like short fragments of hyaluronan [42] and bacterial components such as LPS [43], which lead to the activation of NFκB in endothelial cells [42]. IαI is a potent anti-inflammatory protein and ameliorates sepsis-induced inflammation through several pathways. The IαI light-chain inhibits LPS-induced increase of cytoplasmic free Ca⁺⁺ in immune and structural cells [44], and protects epithelia from injury after IL-1β and TNF-α exposure [45] through a Ca⁺⁺-inhibitory effect. Ca⁺⁺ inhibition was seen in several cell types [44, 46, 47]. IαI also inhibits the

Fig. 4 Inverse association between serum α I and sVCAM a or sICAM b levels at day 0 (admission, left panels) but not at day 5 (right panels)





Fig. 5 Mouse lung staining for hyaluronan (panel a, green), α I (panel b, red), and DAPI (panel c, blue) after i.p. endotoxin exposure shows deposition of hyaluronan and α I along the luminal endothelial surface of a venule (arrows). Panel d (transmitted light) is provided

for structural reference. Panels e and f show merged images without and with transmitted light image, respectively. Note immune cells rolling along the endothelium which partially stain for α I as well (arrowheads). Magnification: $\times 250$

activation of neutrophils after LPS exposure, by reducing ERK1/2 and p38 [48]. Furthermore, I α I has been shown to bind extracellular histones and neutralize histone-induced adverse effects in sepsis [49]. Finally, we recently described that I α I complexing with hyaluronan in septic animals leading to its rapid clearance and ameliorated sepsis mortality [50, 51]. This suggests that I α I is able to reduce the inflammatory cascade in injured endothelia through many pathways, ultimately leading to reduced VCAM-1 and ICAM-1 upregulation and secretion.

We investigated I α I in endothelial inflammation *in vitro* by using C5a exposure. We believe that this is a representative model since C5a is central to endothelial injury and MOF in sepsis [52–54]; furthermore, complement activation occurs throughout the course of systemic inflammation associated with sepsis [53]. A major pathway of C5a inflammation in endothelia is via ERK [28] and NF κ B [10] activation. These pathways may be connected, since ERK leads to NF κ B activation [55]. We demonstrate that I α I ameliorates both ERK and NF κ B activation in endothelia. The I α I light-chain bikunin inhibits ERK activation in LPS-treated monocytes *in vitro* and LPS-dosed rats *in vivo* [56]; the effect may be at least partly due to inhibition of Ca⁺⁺ influx [57], since bikunin inhibits LPS-induced Ca⁺⁺ influx in neutrophils [48] and HUVEC [47]. Ca⁺⁺ influx occurs after C5a binding to its receptor [58] and leads to the C5a-induced ERK activation [59]; thus, Ca⁺⁺ influx inhibition is a plausible mechanism of I α I effects. The hypothesis that I α I effects target ERK is supported by the observation that C5a-induced IL-9 upregulation, which is not modulated by ERK [60], was not changed by I α I. I α I heavy chains may also contribute to its effect. Endothelial cells express several integrins [61] which I α I heavy chains can bind to [62]. Furthermore, I α I heavy chains are necessary for binding to hyaluronan, a crucial component of the endothelial glycocalyx [63]. I α I heavy chains may provide the necessary attachment to endothelial cells (see Fig. 5).

We established translational relevance of our findings by replicating the inverse association of I α I with VCAM-1 and ICAM-1 that we found in mice in a cohort of sepsis patients which in a previous work had demonstrated an inverse association of I α I plasma levels with mortality, MOF score, and serum IL-6 levels [18]. We found that I α I and VCAM-1/ICAM-1 levels are inversely associated in sepsis patients on admission (day 0), but not at day 5 after admission. On admission, patients have received very few treatment interventions. After 5 days, patients have received several potentially confounding interventions and possible complications, which may explain the absence of association. Thus, it may be that the relative absence of confounders allowed us to unmask the association between I α I and endothelial injury during early sepsis. Alternatively, I α I effects may be superseded by other factors that influence endothelial activation in

later sepsis stages. Admission levels of I α I and soluble adhesion factors have been independently found to predict morbidity and survival in sepsis [18, 20] and our work suggests that there may be a causative connection between the two.

In conclusion, we show that I α I protects endothelia from injury caused by sepsis and the activated complement factor C5a. This find provides additional mechanistic insights into the effects of I α I in sepsis and supports the use of I α I as a treatment in sepsis.

Acknowledgements The authors thank Drs. Michael B. Fessler and John House for constructive criticism, and Ms. Ligon Perrow for excellent animal husbandry. This work was supported, in part, by the Division of Intramural Research, National Institute of Environmental Health Sciences, NIH. Yow-Pin Lim gratefully acknowledges support from NIAID grant 2R42AI062095.

Authors Contribution Conception and design: SG. Experiments, analysis, and interpretation: VPS, Y-PL, SO, KK, and LZ. Drafting of manuscript: VPS and SG. Review of manuscript: VPS, Y-PL, SO, KK, LZ, and SG.

Compliance with Ethical Standards

Conflict of interest Y-PL is the President and Chief Scientific Officer at ProThera Biologics, LLC, a company which explores the use of I α I in several systemic inflammatory diseases, including sepsis.

References

1. Angus DC et al (2001) Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. *Crit Care Med* 29(7):1303–1310
2. Kaukonen KM et al (2014) Mortality related to severe sepsis and septic shock among critically ill patients in Australia and New Zealand, 2000–2012. *Jama* 311(13):1308–1316
3. Ferreira FL et al (2001) Serial evaluation of the SOFA score to predict outcome in critically ill patients. *Jama* 286(14):1754–1758
4. Schmidt EP et al (2012) The pulmonary endothelial glycocalyx regulates neutrophil adhesion and lung injury during experimental sepsis. *Nat Med* 18(8):1217–1223
5. Ince C et al (2016) The endothelium in sepsis. *Shock* 45(3):259–270
6. Bosmann M, Ward PA (2012) Role of C3, C5 and anaphylatoxin receptors in acute lung injury and in sepsis. *Adv Exp Med Biol* 946:147–159
7. Czermak BJ et al (1999) Protective effects of C5a blockade in sepsis. *Nat Med* 5(7):788–792
8. Huber-Lang M et al (2001) Role of C5a in multiorgan failure during sepsis. *J Immunol* 166(2):1193–1199
9. Goldenberg NM et al (2011) Broken barriers: a new take on sepsis pathogenesis. *Sci Transl Med* 3(88):88ps25.
10. Albrecht EA et al (2004) C5a-induced gene expression in human umbilical vein endothelial cells. *Am J Pathol* 164(3):849–859
11. Kyan-Aung U, Haskard DO, Lee TH (1991) Vascular cell adhesion molecule-1 and eosinophil adhesion to cultured human umbilical vein endothelial cells *in vitro*. *Am J Respir Cell Mol Biol* 5(5):445–450
12. Kyan-Aung U et al (1991) Endothelial leukocyte adhesion molecule-1 and intercellular adhesion molecule-1 mediate the

- adhesion of eosinophils to endothelial cells in vitro and are expressed by endothelium in allergic cutaneous inflammation in vivo. *J Immunol* 146(2):521–528
13. Bochner BS et al (1991) Adhesion of human basophils, eosinophils, and neutrophils to interleukin 1-activated human vascular endothelial cells: contributions of endothelial cell adhesion molecules. *J Exp Med* 173(6):1553–1557
 14. Singh K et al (2010) Inter-alpha inhibitor protein administration improves survival from neonatal sepsis in mice. *Pediatr Res* 68(3):242–247
 15. Yang S et al (2002) Administration of human inter-alpha-inhibitors maintains hemodynamic stability and improves survival during sepsis. *Crit Care Med* 30(3):617–622
 16. Jourdain M et al (1997) Effects of inter-alpha-inhibitor in experimental endotoxemic shock and disseminated intravascular coagulation. *Am J Respir Crit Care Med* 156(6):1825–1833
 17. Garantziotis S et al (2007) Inter-alpha-trypsin inhibitor attenuates complement activation and complement-induced lung injury. *J Immunol* 179(6):4187–4192
 18. Opal SM et al (2007) Longitudinal studies of inter-alpha inhibitor proteins in severely septic patients: a potential clinical marker and mediator of severe sepsis. *Crit Care Med* 35(2):387–392
 19. Skibsted S et al (2013) Biomarkers of endothelial cell activation in early sepsis. *Shock* 39(5):427–432
 20. Zonneveld R et al (2014) Soluble adhesion molecules as markers for sepsis and the potential pathophysiological discrepancy in neonates, children and adults. *Crit Care* 18(1):204
 21. Jesmin S et al (2012) Time-dependent alterations of VEGF and its signaling molecules in acute lung injury in a rat model of sepsis. *Inflammation* 35(2):484–500
 22. Amatschek S et al (2011) CXCL9 induces chemotaxis, chemorepulsion and endothelial barrier disruption through CXCR2-mediated activation of melanoma cells. *Br J Cancer* 104(3):469–479
 23. Zsebo KM et al (1988) Vascular endothelial cells and granulopoiesis: interleukin-1 stimulates release of G-CSF and GM-CSF. *Blood* 71(1):99–103
 24. Fuste B et al (2004) Granulocyte colony-stimulating factor increases expression of adhesion receptors on endothelial cells through activation of p38 MAPK. *Haematologica* 89(5):578–585
 25. Burg J et al (2002) GM-CSF expression by human lung microvascular endothelial cells: in vitro and in vivo findings. *Am J Physiol Lung Cell Mol Physiol* 283(2):L460–L467
 26. Chandrasekar B et al (2004) Activation of intrinsic and extrinsic proapoptotic signaling pathways in interleukin-18-mediated human cardiac endothelial cell death. *J Biol Chem* 279(19):20221–20233
 27. Brueckmann M et al (2007) Prognostic value of platelet-derived growth factor in patients with severe sepsis. *Growth Factors* 25(1):15–24
 28. Monsinjon T et al (2003) Regulation by complement C3a and C5a anaphylatoxins of cytokine production in human umbilical vein endothelial cells. *Faseb J* 17(9):1003–1014
 29. Levy MM et al (2014) Surviving sepsis campaign: association between performance metrics and outcomes in a 7.5-year study. *Crit Care Med* 40(11):1623–1633.
 30. Vallet B (2003) Bench-to-bedside review: endothelial cell dysfunction in severe sepsis: a role in organ dysfunction? *Crit Care* 7(2):130–138
 31. Kubli S et al (2003) Endothelium-dependent vasodilation in the skin microcirculation of patients with septic shock. *Shock* 19(3):274–280
 32. Davis JS et al (2009) Sepsis-associated microvascular dysfunction measured by peripheral arterial tonometry: an observational study. *Crit Care* 13(5):R155
 33. Coletta C et al (2014) Endothelial dysfunction is a potential contributor to multiple organ failure and mortality in aged mice subjected to septic shock: preclinical studies in a murine model of cecal ligation and puncture. *Crit Care* 18(5):511
 34. Duffy MJ et al (2011) Impaired endothelium-dependent vasodilation is a novel predictor of mortality in intensive care. *Crit Care Med* 39(4):629–635
 35. Smith WB, Gamble JR, Vadas MA (1994) The role of granulocyte-macrophage and granulocyte colony-stimulating factors in neutrophil transendothelial migration: comparison with interleukin-8. *Exp Hematol* 22(3):329–334
 36. Lindahl P et al (1997) Pericyte loss and microaneurysm formation in PDGF-B-deficient mice. *Science* 277(5323):242–245
 37. Motro B et al (1990) Pattern of interleukin 6 gene expression in vivo suggests a role for this cytokine in angiogenesis. *Proc Natl Acad Sci USA* 87(8):3092–3096
 38. Pintucci G et al (2002) Lack of ERK activation and cell migration in FGF-2-deficient endothelial cells. *Faseb J* 16(6):598–600
 39. Colotta F et al (1994) Expression of monocyte chemotactic protein-1 by monocytes and endothelial cells exposed to thrombin. *Am J Pathol* 144(5):975–985
 40. Amalakuhan B et al (2016) Endothelial adhesion molecules and multiple organ failure in patients with severe sepsis. *Cytokine* 88:267–273
 41. Laudes IJ et al (2004) Disturbed homeostasis of lung intercellular adhesion molecule-1 and vascular cell adhesion molecule-1 during sepsis. *Am J Pathol* 164(4):1435–1445
 42. Oertli B et al (1998) Mechanisms of hyaluronan-induced up-regulation of ICAM-1 and VCAM-1 expression by murine kidney tubular epithelial cells: hyaluronan triggers cell adhesion molecule expression through a mechanism involving activation of nuclear factor-kappa B and activating protein-1. *J Immunol* 161(7):3431–3437
 43. Wuthrich RP (1992) Intercellular adhesion molecules and vascular cell adhesion molecule-1 and the kidney. *J Am Soc Nephrol* 3(6):1201–1211
 44. Kanayama N et al (1997) The role of chondroitin sulfate chains of urinary trypsin inhibitor in inhibition of LPS-induced increase of cytosolic free Ca²⁺ in HL60 cells and HUVEC cells. *Biochem Biophys Res Commun* 238(2):560–564
 45. El Maradny E et al (1994) Urinary trypsin inhibitor has a protective effect on the amnion. *Gynecol Obstet Invest* 38(3):169–172
 46. Kanayama N et al (1995) Urinary trypsin inhibitor prevents uterine muscle contraction by inhibition of Ca⁺⁺ influx. *Am J Obstet Gynecol* 173(1):192–199
 47. Kanayama N et al (1998) Urinary trypsin inhibitor suppresses vascular smooth muscle contraction by inhibition of Ca²⁺ influx. *Biochim Biophys Acta* 1381(2):139–146
 48. Kanayama S et al (2007) Bikunin suppresses expression of pro-inflammatory cytokines induced by lipopolysaccharide in neutrophils. *J Endotoxin Res* 13(6):369–376
 49. Chaaban H et al (2015) Inter-alpha inhibitor protein and its associated glycosaminoglycans protect against histone-induced injury. *Blood* 125(14):2286–2296
 50. Ni K et al (2019) Intravascular heavy chain-modification of hyaluronan during endotoxic shock. *Biochem Biophys Rep* 17:114–121
 51. Ni K et al (2018) Rapid clearance of heavy chain-modified hyaluronan during resolving acute lung injury. *Respir Res* 19(1):107
 52. Unnewehr H et al (2013) Changes and regulation of the C5a receptor on neutrophils during septic shock in humans. *J Immunol* 190(8):4215–4225
 53. Flierl MA, Schreiber H, Huber-Lang MS (2006) The role of complement, C5a and its receptors in sepsis and multiorgan dysfunction syndrome. *J Invest Surg* 19(4):255–265

54. Laudes IJ et al (2002) Expression and function of C5a receptor in mouse microvascular endothelial cells. *J Immunol* 169(10):5962–5970
55. Chen BC, Lin WW (2001) PKC- and ERK-dependent activation of I kappa B kinase by lipopolysaccharide in macrophages: enhancement by P2Y receptor-mediated CaMK activation. *Br J Pharmacol* 134(5):1055–1065
56. Molor-Erdene P et al (2005) Urinary trypsin inhibitor reduces LPS-induced hypotension by suppressing tumor necrosis factor-alpha production through inhibition of Egr-1 expression. *Am J Physiol Heart Circ Physiol* 288(3):H1265–1271
57. Kobayashi H et al (2003) A Kunitz-type protease inhibitor, bikunin, inhibits ovarian cancer cell invasion by blocking the calcium-dependent transforming growth factor-beta 1 signaling cascade. *J Biol Chem* 278(10):7790–7799
58. Monk PN, Partridge LJ (1993) Characterization of a complement-fragment-C5a-stimulated calcium-influx mechanism in U937 monocytic cells. *Biochem J* 295(Pt 3):679–684
59. Hawlisch H et al (2005) C5a negatively regulates toll-like receptor 4-induced immune responses. *Immunity* 22(4):415–426
60. Perumal NB, Kaplan MH (2011) Regulating Il9 transcription in T helper cells. *Trends Immunol* 32(4):146–150
61. Bader BL et al (1998) Extensive vasculogenesis, angiogenesis, and organogenesis precede lethality in mice lacking all alpha v integrins. *Cell* 95(4):507–519
62. Adair JE et al (2009) Inter-alpha-trypsin inhibitor promotes bronchial epithelial repair after injury through vitronectin binding. *J Biol Chem* 284(25):16922–16930
63. Lennon FE, Singleton PA (2011) Hyaluronan regulation of vascular integrity. *Am J Cardiovasc Dis* 1(3):200–213

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.