



Impact of prior interventional treatment on the complications after Frey procedure for chronic pancreatitis

A. Merdrignac¹ · D. Bergeat¹ · F. Robin¹ · E. Gagnard¹ · K. Turner¹ · M. Rayar^{1,2} · B. Meunier¹ · K. Boudjema^{1,2} · Laurent Sulpice^{1,2}

Received: 8 May 2019 / Accepted: 3 October 2019 / Published online: 25 October 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Purpose Chronic pancreatitis is an inflammatory disease responsible for pain partially explained by pancreatic duct dilatation. Early surgery has become the treatment of choice for hypertrophic pancreatic head with main pancreatic duct dilatation. Frey procedure (FP), combining both surgical resection and decompression, is one of the standard surgical procedures. However, a “step-up approach” with endoscopic or limited surgical procedures is still frequently proposed before referring to expert pancreatic centres. The aim of the study was to evaluate the impact of a prior treatment on post-operative complications of FP.

Methods All 61 consecutive patients who underwent FP between 2006 and 2017 were included. Perioperative data and outcomes were analyzed and compared according to the presence of a prior treatment.

Results Twenty-four patients did not receive any prior treatment and thirty-seven patients had a prior endoscopic or limited surgical treatment. Preoperative data and outcomes were similar between the 2 groups. The rate of biliary derivation during FP was significantly higher in the group without prior endoscopic procedure. A prior treatment was not a risk factor for major morbidity (Clavien grade \geq III).

Conclusions A first attempt of endoscopic or limited surgical procedures before FP may not influence post-operative complications. Even if not recommended, a “step-up approach” proposing a first less invasive treatment could still be proposed to the patients who want to delay a morbid surgical procedure.

Keywords Chronic pancreatitis · Pancreatic surgery · Frey procedure

Introduction

Chronic pancreatitis (CP) is characterized by pancreatic parenchymal lesions progressing from inflammation to fibrosis and calcifications. Pain is the main resulting symptom and its management is a therapeutic challenge [1]. Although the origin of pain is not clearly known and

certainly multifactorial, an enlargement of main pancreatic duct due to obstruction by strictures or stones is generally observed. The aim of both endoscopic and surgical procedures is to decrease pressure in the main pancreatic duct. Endoscopic treatment involves sphincterotomy, dilatation of strictures by temporary stent, removal of stones, and extracorporeal shock wave lithotripsy. Complications of endoscopic treatment are post-endoscopic retrograde cholangiopancreatography (ERCP) pancreatitis (3.5%), infection (1.4%), haemorrhage (1.3%), and perforation (0.6%) [2]. Post-ERCP complications lead to a death rate of less than 1% mostly due to severe pancreatitis.

Surgical techniques consist in drainage of main pancreatic duct, resection, or association of both. In cases of hypertrophic pancreatic head, surgical treatment mostly consists in hybrid procedures associating main pancreatic duct decompression and duodenum-preserving pancreatic head resection such as the Frey procedure (FP). Several previously published studies have demonstrated that FP

The study has been presented at the 13th French congress of digestive and hepato-bilio-pancreatic surgery.

✉ Laurent Sulpice
laurent.sulpice@chu-rennes.fr

¹ CHU Rennes, Service de Chirurgie Hépatobiliaire et Digestive, INSERM 1241, NuMeCan (Nutrition Metabolism and Cancer), Université de Rennes 1, 2 rue Henri Le Guilloux, F-35000 Rennes, France

² INSERM 1414, CIC, Université de Rennes 1, 2 rue Henri Le Guilloux, F-35000 Rennes, France

offers an effective pain relief and is a safe surgical technique in the management of CP [3]. Frey procedure is associated with 25% rate of post-operative morbidity and less than 1% rate of mortality [3, 4]. The most frequent specific complication after FP is post-operative bleeding (10–20%) and pancreatic fistula (8–10%) [3, 5].

For patients with obstructive CP and dilated pancreatic duct, surgery is superior to endoscopy in terms of pain relief [6, 7]. In a Cochrane review, morbidity and mortality seem not to differ between the two intervention modalities, but the power of the trials was not sufficient to detect the small differences expected in this outcome [6]. Surgical intervention at an early stage of CP is a promising approach in terms of pain relief and pancreatic function. Early surgery has become the strategy of choice for expert pancreatic centres [8–10]. A randomized controlled trial has been conducted to compare early surgery vs endoscopy but results have not been published yet (ESCAPE trial ISRCTN45877994) [10]. The development of central pain (sensitization) is a concern when surgical intervention is delayed [11]. A delayed surgical intervention will not be beneficial in terms of pain relief. Negi et al. conclude that patients should be referred for surgery before opiates are needed to relieve pain [12]. These results are confirmed by Ahmed Ali et al. who recommend surgery within 3 years of symptomatic CP [13]. In patients with an enlarged pancreatic head, a recent international survey of pancreatologists has shown that 58% preferred initial surgery (pylorus preserving pancreaticoduodenectomy) versus 42% initial endoscopy [14].

However, currently, some patients are still referred to expert pancreatic centres after a first “step-up approach.” The impact of a prior endoscopic treatment on post-operative complications is not studied and known. Moreover, some patients do not present initial cephalic pancreatic complications of chronic pancreatitis and require first distal, decompressive, or urgent pancreatic surgical procedures. The aim of the study was to evaluate the impact of a prior endoscopic or surgical treatment on the post-operative complications of Frey procedure for painful CP with hypertrophic pancreatic head.

Patients and methods

Patients

A database of all consecutive patients who underwent the FP in a tertiary centre has been created and maintained prospectively since 2006 after institutional approval and informed consent. Data were collected regarding demographics, preoperative procedures (endoscopic, surgical, radiologic), surgical variables, length of hospital stay, post-operative morbidity, and follow-up.

All patients operated between 2006 and 2017 were included in the analyses. No patient was excluded of the analysis. Diagnosis of CP was based on the clinical (clinical history, physical examination) and imaging findings (computed tomographic scan, magnetic resonance cholangiopancreatography, endoscopic ultrasound). Indication for surgery was systematically determined during a multidisciplinary meeting including surgeons, gastroenterologists, and radiologists. Patients were divided into two groups according to the presence of a prior endoscopic procedure (attempts of pancreatic and/or biliary catheterism) or pancreatic surgery for CP.

Surgical procedure

FP was performed by one senior pancreatic surgeon (LS) according to the technique first described in 1987 by Frey [15]. The need for biliary diversion was determined preoperatively or perioperatively when common bile duct (CBD) stricture persisted as observed by perioperative cholangiography. The type of biliary diversion chosen in case of CBD stricture depended on the accessibility of the CBD in the resection cavity and was left to the discretion of the surgeon.

The technique of CBD reinsertion combined a 1-cm longitudinal cut of the distal CBD in the pancreatic head resection cavity and an anastomosis to pancreatic parenchyma using 5-0 polydioxanone interrupted sutures. A Roux-en-Y choledochojejunostomy was performed if bilioenteric anastomosis was chosen. The same Roux-en-Y jejunal loop was used for the bilioenteric anastomosis and the pancreaticojejunal anastomosis. The distal extremity of the jejunal loop was first anastomosed with the pancreas.

Definitions

The post-operative morbidity was defined as any complication that occurred within 90 days after surgery. The complications were categorized according to the Dindo and Clavien classification [16]. Major complications were categorized as grades III to V in the Dindo and Clavien classification. Post-operative pancreatic fistula was classified into three groups according to the criteria given by the International Study Group of Pancreatic Fistula [17]. Post-operative delayed gastric emptying was classified according to the criteria given by the International Study Group of Pancreatic Surgery (ISGPS) but only grades B and C were used for this study as our centre policy was to maintain the nasogastric tube until at least post-operative day 5 [18]. Post-pancreatectomy haemorrhage including intra- and extra-luminal bleeding was classified according to the ISGPS definition [19]. Biliary fistula was defined as the appearance of bile in peritoneal drains. Post-operative pain was defined as controlled if no analgesic




Fig. 1 Distribution of all patients according to prior treatment for chronic pancreatitis

was necessary or only level 1 analgesic according to the World Health Organization pain ladder (neither opioid nor weak opioid).

Table 1 Preoperative data of all patients divided in two groups according to the presence of a prior endoscopic or surgical procedure for chronic pancreatitis (CP)

	Total n = 61	Prior surgical and/or endoscopic procedure n = 37	None prior procedure n = 24	p
Age (years, mean ± SD)	48.36 ± 10.02	48.56 ± 9.79	48.04 ± 10.57	0.95
Gender				1
Male	56	34	22	
Female	5	3	2	
Etiology of CP				1
Alcoholic	56	34	22	
Idiopathic	2	1	1	
Hereditary	2	1	1	
Post-radiation	1	1	0	
Duration of symptoms of CP (years, mean ± SD)	6.45 ± 7.28	6.85 ± 7.33	5.83 ± 7.31	0.46
Preoperative diabetes mellitus	16	7	9	0.14
BMI (mean ± SD)	17.34 ± 7.31	16.19 ± 9.31	19.11 ± 7.23	0.16
ASA				1
I	2	1	1	
II	50	30	20	
III	9	6	3	

SD, standard derivation; CP, chronic pancreatitis; BMI, body mass index; ASA, American Society of Anesthesiologists

Statistical analysis

Qualitative variables are expressed as numbers with percentages and were compared using the Fisher exact test. Quantitative variables are expressed as mean values ± the standard deviation (SD) and were compared using Mann–Whitney *U* test. *p* values less than 0.05 were considered statistically significant. Analyses were performed with R statistical software (available: <http://www.r-project.org/>).

Results

Sixty-one patients were operated for a FP between 2006 and 2017. Patients were divided in two groups: 37 patients with a prior endoscopic or pancreatic surgical procedure (60.7%) and 24 patients without prior interventional treatment for CP (39.3%). Nine patients underwent a prior pancreatic surgery: 3 cystogastric anastomosis, 3 distal pancreatectomies, 2 pancreaticojejunostomies, and 1 pancreatic necrosectomy. Twenty-eight patients have had one or more prior endoscopic procedure: 5 pancreatic stenting, 2 biliary stenting, 3 cystogastric drainages, and 21 attempts for endoscopic pancreatic drainage. The distribution of patients is presented in Fig. 1. Preoperative data were comparable between the two groups (Table 1). The majority were male with CP due to alcohol abuse.

Post-operative data are presented in Table 2. Operation time and hospitalization duration were similar. The rate of biliary derivation was significantly higher in the group without prior endoscopic procedure. The rate of major

Table 2 Post-operative data of all patients divided in two groups according to the presence of a prior endoscopic or surgical procedure for chronic pancreatitis

	Total <i>n</i> = 61	Prior surgical and/or endoscopic procedure <i>n</i> = 37	No prior procedure <i>n</i> = 24	<i>p</i>
Operation time (min, mean ± SD)	271 ± 55	264 ± 54	282 ± 55	0.22
Transfusion during operation	3	2	1	1
Biliary diversion combined to Frey procedure	24	8	16	0.005
Hospitalization duration (days, mean ± SD)	13.3 ± 14.1	12.29 ± 7.91	14.84 ± 20.44	0.44
Clavien				1
≥ III	12	7	5	
I	32	18	14	
II	17	12	5	
IIIa	1	1	0	
IIIb	5	5	0	
IVa	5	1	4	
IVb	1	0	1	
V	0	0	0	
Pancreatic fistula	6	3	3	0.67
Biliary fistula	2	1	1	1
Delayed gastric emptying	11	9	2	0.17
Haemorrhage	14	9	5	1
Readmission	10	5	5	0.49

The number of patients presenting with pancreatic fistula, delayed gastric emptying, or haemorrhage is equal to the sum of all grades to permit statistical analyses. *SD*, standard derivation

complication (Clavien ≥ III) was not different including pancreatic fistula and haemorrhage.

Mean follow-up was 24.6 ± 21.4 and 33 ± 30.3 months in the group with and without prior endoscopic or surgical procedure respectively (*p* = 0.3). Data on pain control at 2 years were available for 43 patients: 24 and 19 patients in the group with and without prior procedure respectively. Pain control was obtained for 95.8% and 84.2% patients (*p* = 0.3) in the group with and without prior procedure respectively. Data on diabetes mellitus at 2 years were available for 46 patients: 30 and 16 patients in the group with and without prior procedure respectively. Diabetes mellitus was present in 46.6% and 50% (*p* = 1) in the group with and without prior procedure respectively.

Table 3 Univariate analysis for identification of risk factors for complications with Clavien grade ≥ III

	Total <i>n</i> = 61	Clavien grade < III <i>n</i> = 49	Clavien grade ≥ III <i>n</i> = 12	<i>p</i>
Gender				0.5726
Male	56	44	12	
Female	5	5	0	
Age (years, mean ± SD)	48.36 ± 10.02	48.06 ± 10.17	49.58 ± 9.71	0.6048
BMI (mean ± SD)	20.99 ± 2.91	20.78 ± 2.9	21.84 ± 2.92	0.3183
Duration of symptoms (years, mean ± SD)	5.93 ± 6.27	6.17 ± 6.61	4.96 ± 4.72	0.5775
Preoperative diabetes mellitus	16	14	2	0.4899
ASA				0.5947
I	2	2	0	
II	50	41	9	
III	9	6	3	
Prior endoscopic or surgical procedure	37	30	7	1
Biliary derivation	24	20	4	0.73
Enucleation diameter (cm, mean ± SD)	4.12 ± 1.15	4.18 ± 1.21	3.87 ± 0.83	0.4988
Enucleation volume (cm ³ , mean ± SD)	26.48 ± 18.98	27.25 ± 19.99	23.44 ± 14.54	0.7393

SD, standard derivation

Univariate analysis for identification of risk factors for complications with Clavien grade ≥ III is presented in Table 3. Multivariate analysis was not realized because no risk factor was identified in univariate analysis.

Discussion

The treatment of painful CP with hypertrophic pancreatic head could involve both endoscopic and surgical procedures. However, current recommendations indicate that the timing of surgical intervention is an important factor in clinical outcomes.

The “step-up approach” consists with a less invasive endoscopic treatment as first-line therapy followed by surgical procedure in case of failure. Prolonged and repeated pancreatic stenting has been identified as a risk factor for failure after subsequent rescue surgery [13]. Moreover, the Dutch Pancreatitis Study Group has argued against the “step-up approach” on the basis that opioids, the mainstay of conservative treatment, do not alter disease progression and that reliance on them delays intervention [20].

Nevertheless, a first less invasive approach using endoscopic procedure is still frequently proposed to patients before referring to expert pancreatic centres or if patients refused surgery. The present study shows similar rate of post-operative complications after FP between the patients who had a prior endoscopic or surgical treatment and those who were free of prior interventional treatment. One can hypothesize that specific complications of endoscopic procedures could influence surgical outcomes. Post-ERCP pancreatitis leads to local inflammation that could increase morbidity of pancreatic surgery. Chronic pancreatitis may be a protector factor for post-ERCP pancreatitis [21]. Conversely, difficult or failed cannulation is a risk factor for post-ERCP pancreatitis. The majority of patients with a prior endoscopic procedure in this study underwent a failure of pancreatic drainage. The similar rates of surgical morbidity in the 2 groups of this study are in accordance with the protecting effect of CP on post-ERCP pancreatitis. Moreover, a prior endoscopic or surgical treatment is not a risk factor for complications with Clavien grade \geq III in the present cohort.

In the group of patients with a prior endoscopic or surgical procedure, the need for a biliary diversion during FP was significantly lower. Among these patients, two had a biliary stenting that could have avoided the need for a biliary derivation during surgery and explain the statistical difference between the 2 groups.

The main limits of the present study are the small number of patients and the long study period in a monocentric cohort. However, the majority of published cohorts do not exceed fifty patients. Pancreatic surgery is more frequently indicated for pancreatic neoplasms than for CP in Western Europe. This fact explains the difficulty to identify risk factors for complications after FP. Follow-up of patients with CP is often difficult and leads to biases due to missing data.

Conclusions

The management of painful CP with enlarged pancreatic head involves both endoscopic and surgical procedures after failure of medical treatment. Several studies have shown benefit for early surgery on pain relief. The present study shows that a first attempt of endoscopic procedure before surgery may not influence post-operative complications. A preoperative biliary

stenting could avoid the need for a surgical biliary derivation combined to the FP. A “step-up approach” could not be recommended but, in cases of patients referred after prior interventional treatment, patients could undergo Frey procedure without excess risk of post-operative complications.

Author’s contributions Study conception: AM and LS. Acquisition of data: AM, FR, EG, and KT. Analysis and interpretation of data: AM, DB, and MR. Drafting of manuscript: AM and LS. Critical revision of manuscript: AM, BM, KB, and LS.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Research involving human participants All procedures performed in the study involving human participants were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent Informed consent was obtained from all individual participants included in the study.

References

1. Braganza JM, Lee SH, McCloy RF, McMahon MJ (2011) Chronic pancreatitis. *Lancet* 377:1184–1197. [https://doi.org/10.1016/S0140-6736\(10\)61852-1](https://doi.org/10.1016/S0140-6736(10)61852-1)
2. Andriulli A, Loperfido S, Napolitano G, Niro G, Valvano MR, Spirito F, Pilotto A, Forlano R (2007) Incidence rates of post-ERCP complications: a systematic survey of prospective studies. *Am J Gastroenterol* 102:1781–1788. <https://doi.org/10.1111/j.1572-0241.2007.01279.x>
3. Zhou Y, Shi B, Wu L, Wu X, Li Y (2015) Frey procedure for chronic pancreatitis: evidence-based assessment of short- and long-term results in comparison to pancreatoduodenectomy and Beger procedure: a meta-analysis. *Pancreatol* 15:372–379. <https://doi.org/10.1016/j.pan.2015.05.466>
4. Suzumura K, Hatano E, Okada T, Asano Y, Uyama N, Nakamura I, Hai S, Fujimoto J (2018) Short- and long-term outcomes of the Frey procedure for chronic pancreatitis: a single-center experience and summary of outcomes in Japan. *Surg Today* 48:58–65. <https://doi.org/10.1007/s00595-017-1548-0>
5. Ahmed Ali U, Pahlplatz JM, Nealon WH et al (2015) Endoscopic or surgical intervention for painful obstructive chronic pancreatitis. In: *Cochrane Database of Systematic Reviews*. John Wiley & Sons, Ltd
6. Strobel O, Büchler MW, Werner J (2009) Surgical therapy of chronic pancreatitis: indications, techniques and results. *Int J Surg* 7:305–312. <https://doi.org/10.1016/j.ijsu.2009.05.011>
7. Yang CJ, Bliss LA, Schapira EF, Freedman SD, Ng SC, Windsor JA, Tseng JF (2014) Systematic review of early surgery for chronic pancreatitis: impact on pain, pancreatic function, and re-intervention. *J Gastrointest Surg* 18:1–7. <https://doi.org/10.1007/s11605-014-2571-8>
8. Yang CJ, Bliss LA, Freedman SD, Sheth S, Vollmer CM, Ng SC, Callery MP, Tseng JF (2015) Surgery for chronic pancreatitis: the role of early surgery in pain management. *Pancreas* 44:819–823. <https://doi.org/10.1097/MPA.0000000000000333>

9. Ke N, Jia D, Huang W et al (2018) Earlier surgery improves outcomes from painful chronic pancreatitis. *Medicine (Baltimore)* 97. <https://doi.org/10.1097/MD.00000000000010651>
10. Ahmed Ali U, Issa Y, Bruno MJ, van Goor H, van Santvoort H, Busch OR, Dejong CH, Nieuwenhuijs VB, van Eijck C, van Dullemen H, Fockens P, Siersema PD, Gouma DJ, van Hooft J, Keulemans Y, Poley JW, Timmer R, Besselink MG, Vleggaar FP, Wilder-Smith OH, Gooszen HG, Dijkgraaf MG, Boermeester MA, Dutch Pancreatitis Study Group (2013) Early surgery versus optimal current step-up practice for chronic pancreatitis (ESCAPE): design and rationale of a randomized trial. *BMC Gastroenterol* 13:49. <https://doi.org/10.1186/1471-230X-13-49>
11. Anderson MA, Akshintala V, Albers KM, Amann ST, Belfer I, Brand R, Chari S, Cote G, Davis BM, Frulloni L, Gelrud A, Guda N, Humar A, Liddle RA, Slivka A, Gupta RS, Szigethy E, Talluri J, Wassef W, Wilcox CM, Windsor J, Yadav D, Whitcomb DC (2016) Mechanism, assessment and management of pain in chronic pancreatitis: recommendations of a multidisciplinary study group. *Pancreatology* 16:83–94. <https://doi.org/10.1016/j.pan.2015.10.015>
12. Negi S, Singh A, Chaudhary A (2010) Pain relief after Frey's procedure for chronic pancreatitis. *Br J Surg* 97:1087–1095. <https://doi.org/10.1002/bjs.7042>
13. Ali UA, Nieuwenhuijs VB, van Eijck CH et al (2012) Clinical outcome in relation to timing of surgery in chronic pancreatitis: a nomogram to predict pain relief. *Arch Surg* 147:925–932. <https://doi.org/10.1001/archsurg.2012.1094>
14. Issa Y, van Santvoort HC, Fockens P, Besselink MG, Bollen TL, Bruno MJ, Boermeester MA, Collaborators (2017) Diagnosis and treatment in chronic pancreatitis: an international survey and case vignette study. *HPB* 19:978–985. <https://doi.org/10.1016/j.hpb.2017.07.006>
15. Frey CF, Smith GJ (1987) Description and rationale of a new operation for chronic pancreatitis. *Pancreas* 2:701–707
16. Dindo D, Demartines N, Clavien P-A (2004) Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 240:205–213
17. Bassi C, Dervenis C, Butturini G, Fingerhut A, Yeo C, Izbicki J, Neoptolemos J, Sarr M, Traverso W, Buchler M, International Study Group on Pancreatic Fistula Definition (2005) Postoperative pancreatic fistula: an international study group (ISGPF) definition. *Surgery* 138:8–13. <https://doi.org/10.1016/j.surg.2005.05.001>
18. Wente MN, Bassi C, Dervenis C, Fingerhut A, Gouma DJ, Izbicki JR, Neoptolemos JP, Padbury RT, Sarr MG, Traverso LW, Yeo CJ, Büchler MW (2007) Delayed gastric emptying (DGE) after pancreatic surgery: a suggested definition by the International Study Group of Pancreatic Surgery (ISGPS). *Surgery* 142:761–768. <https://doi.org/10.1016/j.surg.2007.05.005>
19. Wente MN, Veit JA, Bassi C, Dervenis C, Fingerhut A, Gouma DJ, Izbicki JR, Neoptolemos JP, Padbury RT, Sarr MG, Yeo CJ, Büchler MW (2007) Postpancreatectomy hemorrhage (PPH): an International Study Group of Pancreatic Surgery (ISGPS) definition. *Surgery* 142:20–25. <https://doi.org/10.1016/j.surg.2007.02.001>
20. Issa Y, van Santvoort HC, van Goor H, Cahen DL, Bruno MJ, Boermeester MA (2013) Surgical and endoscopic treatment of pain in chronic pancreatitis: a multidisciplinary update. *Dig Surg* 30:35–50. <https://doi.org/10.1159/000350153>
21. Freeman ML (2007) Pancreatic stents for prevention of post-endoscopic retrograde cholangiopancreatography pancreatitis. *Clin Gastroenterol Hepatol* 5:1354–1365. <https://doi.org/10.1016/j.cgh.2007.09.007>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.