



An oilseed rape WRKY-type transcription factor regulates ROS accumulation and leaf senescence in *Nicotiana benthamiana* and *Arabidopsis* through modulating transcription of *RbohD* and *RbohF*

Liu Yang¹ · Chaofei Ye¹ · Yuting Zhao¹ · Xiaolin Cheng¹ · Yiqiao Wang¹ · Yuan-Qing Jiang¹ · Bo Yang¹

Received: 23 September 2017 / Accepted: 1 March 2018 / Published online: 6 March 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Main conclusion Overexpression of *BnaWGR1* causes ROS accumulation and promotes leaf senescence. *BnaWGR1* binds to promoters of *RbohD* and *RbohF* and regulates their expression.

Manipulation of leaf senescence process affects agricultural traits of crop plants, including biomass, seed yield and stress resistance. Since delayed leaf senescence usually enhances tolerance to multiple stresses, we analyzed the function of specific MAPK–WRKY cascades in abiotic and biotic stress tolerance as well as leaf senescence in oilseed rape (*Brassica napus* L.), one of the important oil crops. In the present study, we showed that expression of one WRKY gene from oilseed rape, *BnaWGR1*, induced an accumulation of reactive oxygen species (ROS), cell death and precocious leaf senescence both in *Nicotiana benthamiana* and transgenic *Arabidopsis* (*Arabidopsis thaliana*). *BnaWGR1* regulates the transcription of two genes encoding key enzymes implicated in production of ROS, that is, respiratory burst oxidase homolog (*Rboh*) *D* and *RbohF*. A dual-luciferase reporter assay confirmed the transcriptional regulation of *RbohD* and *RbohF* by *BnaWGR1*. In vitro electrophoresis mobility shift assay (EMSA) showed that *BnaWGR1* could bind to W-box *cis*-elements within promoters of *RbohD* and *RbohF*. Moreover, *RbohD* and *RbohF* were significantly upregulated in transgenic *Arabidopsis* overexpressing *BnaWGR1*. In summary, these results suggest that *BnaWGR1* could positively regulate leaf senescence through regulating the expression of *RbohD* and *RbohF* genes.

Keywords *Brassica napus* · Leaf senescence · Rboh · ROS · Transcriptional regulation · WRKY

Abbreviations

CaMV Cauliflower mosaic virus
EMSA Electrophoretic mobility shift assay
GUS β -Glucuronidase
NBT Nitroblue tetrazolium

ROS Reactive oxygen species
SAG Senescence-associated gene

Introduction

Leaf senescence is a fine-tuned active degenerative process through degradation of macromolecules including chlorophylls, proteins and so on, which recycles nutrients released from the old organs to growing young leaves and developing fruits and seeds (Lim and Nam 2005). Leaf senescence is controlled by many factors including age, nutrition status, abiotic and biotic stress conditions, and plant hormones, which interplay in synergistic or antagonistic ways together with calcium signals (Lim et al. 2007). Reactive oxygen species (ROS), including singlet oxygen, superoxide anion, hydrogen peroxide (H₂O₂) and hydroxyl radical, are both

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00425-018-2868-z>) contains supplementary material, which is available to authorized users.

✉ Yuan-Qing Jiang
jiangyq@nwfufu.edu.cn

✉ Bo Yang
yangwl@nwfufu.edu.cn

¹ State Key Laboratory of Crop Stress Biology for Arid Areas, College of Life Sciences, Northwest A&F University, Yangling 712100, Shaanxi, China

signaling and toxic molecules, depending on the concentration (Mittler et al. 2004). At lower concentrations, ROS are utilized as signaling molecules to regulate development and various physiological responses. At higher concentrations, ROS can cause oxidative damage to proteins, DNA, and lipids (Mittler et al. 2004). Plants have evolved complex arrays of non-enzymatic and enzymatic detoxification mechanisms to detoxify excessive ROS (Apel and Hirt 2004). It is believed that ROS production arising from the disassembly of the chloroplast and the degradation of chlorophyll induces leaf senescence (Rogers and Munne-Bosch 2016). Among the ROS, H_2O_2 is most studied as it has a relatively longer half-life, facilitating research. It has been shown in *Arabidopsis* that the hydrogen peroxide level starts to increase at the time points when plants start to bolt and also senescence-associated genes (*SAGs*) are enhanced by the elevated level of ROS (Ye et al. 2000; Navabpour et al. 2003). It has been reported that there is a causal link between WRKY-mediated leaf senescence and ROS in *Arabidopsis* (Besseau et al. 2012). However, it remains elusive how the transcriptional activity of a WRKY protein results in modulating cellular ROS levels.

The respiratory burst oxidase homologues (*Rboh*s) are the most studied ROS-producing enzymes, serving as main molecular ‘hubs’ in ROS-mediated signaling including cell growth, plant development such as root hair, abiotic stress and immune responses (Marino et al. 2012). In *Arabidopsis thaliana*, there are 10 *Rboh* genes, which are *RbohA* through *RbohJ* (Marino et al. 2012). Of them, *AtRbohD* and *-F* are two prominent members and it has been shown that *AtRbohD* and *-F* are required in ABA-mediated ROS production and stomatal closure (Kwak et al. 2003), and in ROS burst of pathogen/microbe-associated molecular patterns (PAMP/MAMPs)-triggered immunity (PTI) and effector-triggered immunity (ETI) (Torres et al. 2002). Reports have demonstrated that the activity of *Rboh*s can be modulated at post-translational level through phosphorylation of the N-terminal cytosolic domain (Baxter et al. 2014). It has also been shown that *Rboh* genes could be regulated by specific NAC and WRKY transcription factors at the transcriptional level (Lee et al. 2012; Adachi et al. 2015). For instance, phospho-mimicking forms of tobacco WRKY7, -8, -9 and -11 belonging to Group I bind to the W-box element in the *RbohB* promoter to positively regulate its expression and the activation of *RbohB* by the MPK-WRKY pathway is involved in R3a/AVR3a-mediated ETI immunity and INF1-triggered immunity (Adachi et al. 2015). However, whether there is any other non-Group I WRKY protein that regulates *Rboh* transcription remains unknown.

WRKY transcription factors form a large family in either model plant or crops, with 72 members in *Arabidopsis* (Eulgem et al. 2000). WRKY transcription factors are classified into three major groups: WRKY proteins with two

WRKY domains and C2H2 zinc motifs belong to Group I, whereas those with one WRKY domain and C2H2 zinc motifs belong to Group II. WRKY proteins having only one WRKY domain but having C2HC zinc-finger motifs are categorized into Group III. Group II WRKY proteins are further divided into IIa, IIb, IIc, IId, and IIe based on the primary amino acid sequences (Eulgem et al. 2000). WRKYs function in mediating disease resistance and expression of defense-related genes in systemic acquired resistance (SAR) (Zheng et al. 2006; Knoth et al. 2007; Bhattarai et al. 2010; Chen et al. 2010; Birkenbihl et al. 2012), abiotic stress (Bakshi and Oelmüller 2014), and so on. WRKY transcription factors usually recognize and bind to a W-box element, which has a consensus TTGAC(C/T) sequence (Ciolkowski et al. 2008). After binding to the W-box element contained in the promoter region of a downstream target gene, a WRKY transcription factor may activate or repress the transcription of the target gene or both. In *Arabidopsis*, six WRKY proteins have been shown to positively or negatively regulate leaf senescence, including WRKY6 (Robatzek and Somssich 2001), -22 (Zhou et al. 2011), -53 (Miao et al. 2004), -54 (Besseau et al. 2012), -57 (Jiang et al. 2014) and -70 (Besseau et al. 2012). For instance, *AtWRKY6* positively regulated leaf senescence by specifically activating the senescence-induced receptor-like kinase (*SIRK*) gene (Robatzek and Somssich 2002). *AtWRKY53* played a regulatory role in early events of leaf senescence (Hinderhofer and Zentgraf 2001). *AtWRKY53* could bind to the promoters of *CAT2* and *-3* to transactivate their expression, which encode two ROS-scavenging enzymes (Miao et al. 2004). Usually, it is believed that the upregulation of ROS-scavenging enzymes is largely to prevent premature senescence when senescence initiates (Zimmermann et al. 2006). With the proceeding of the senescence, the H_2O_2 level increases to destroy the cellular integrity leading to the remobilization of the nutrient out of the senescent leaf. It was speculated that the first H_2O_2 peak might increase the level of *WRKY53* (Zimmermann et al. 2006). However, whether *AtWRKY53* could directly mediate ROS accumulation during the process of leaf senescence remains unknown.

Oilseed rape (*Brassica napus* L.) is one of the most important oil crops in the world. The regulatory mechanisms and key transcription factors controlling leaf senescence in oilseed rape remain unclear. Dissecting the senescence network in oilseed rape will not only provide approaches to control premature leaf senescence, but also offer alternative strategies of enhancing stress tolerance (Rivero et al. 2007). Previously, we performed a systematic identification, cloning and analysis of WRKY transcription factor genes and their upstream kinase genes in oilseed rape (Yang et al. 2009; Liang et al. 2013). In this study, we report the functional characterization of one of the WRKY-type transcription factor genes *BnaWGR1* (*WRKY generating ROS 1*) from

oilseed rape. In addition, we identified that its expression induced ROS accumulation in tobacco and leaf senescence in *Arabidopsis*, through modulating the expression of *RbohD* and *-F*.

Materials and methods

Plant materials and growth conditions

Seeds of oilseed rape (*Brassica napus* L. spring type, double haploid) were originally supplied by Dr. Michael Deyholos (UBCO, Canada) and those of *Arabidopsis thaliana* and *Nicotiana benthamiana* were from Lehle Seeds (Round Rock, TX, USA). Plants were grown in a growth chamber with 22/20 °C day/night temperatures under a light period of 14 h per day and a light intensity of 120 $\mu\text{mol m}^{-2} \text{s}^{-1}$. *Arabidopsis* ecotype Col-0 was used as the wild type (WT). Seeds were surface sterilized and plated on half-strength Murashige and Skoog (MS, Caisson Labs, Smithfield, UT, USA) medium (2.15 g L⁻¹ basal salts, 1% sucrose, pH 5.7 and 8 g L⁻¹ Phytoblend) and imbibed for 2 d at 4 °C before germinated in a growth chamber. Seven-day-old seedlings were transferred into a commercial soil mix (Pindstrup, Kongerslev, Denmark).

Promoter cloning and GUS staining

A DNA fragment containing a promoter region of 1.5 kb upstream of the initiation codon was cloned from oilseed rape genomic DNA using primers listed in Table S1. Primers were designed through mining the public bacterial artificial clone (BAC) sequences stored in the NCBI database. After sequencing, the promoter was cloned into the binary vector pCAMBIA1391Z. The resulting construct was introduced into the *Agrobacterium tumefaciens* strain GV3101 and transformed into wild-type *Arabidopsis* Col-0 (Clough and Bent 1998). To screen transgenic homozygous lines, T1 seeds were selected on hygromycin (Roche, Mannheim, Germany)-containing 1/2 × MS medium for 7 days before they were transferred to soil to get T2 seeds individually. Then, individual T2 seeds were plated on hygromycin-containing 1/2 × MS medium to evaluate homozygosity. Only homozygous T2 lines resistant to hygromycin were subjected to histochemical staining of GUS as described previously (Jiang and Deyholos 2009).

β-Glucuronidase (GUS) activity assay

Measurement of GUS activity was performed according to Kim et al. (2006). Briefly, tissue was ground in liquid nitrogen and mixed in 200 μL extraction buffer (50 mM sodium phosphate buffer, 10 mM EDTA, 0.1% SDS, 0.1% Triton

X-100 and 10 mM β-mercaptoethanol). 15 μg total protein was transferred to 980 μL of 1 mM 4-methylumbelliferyl-β-D-glucuronide (4-MUG) dissolved in the extraction buffer. The reaction was stopped by 0.2 M sodium carbonate buffer (pH11.3) at 5, 10, 20, 40 min, respectively, for fluorescence measurement on a multiple-well plate reader (Spectra Max M2, Molecular Devices, CA, USA) with an excitation wavelength of 365 nm and an emission at 455 nm. The resultant product, 4-methylumbelliferone (4-MU), was quantified against a standard curve.

Subcellular localization assay

To construct the *BnaWGR1* fused with *GFP* expression vectors, the full-length coding sequence of *BnaWGR1* was subcloned into a modified pYJGFP vector with C-terminal fusion with *GFP* and into a p35SNGFP with N-terminal fusion with *GFP*, respectively, using the primers listed in Supplementary Table S1. The NLS-mCherry expression cassette was constructed by fusing nuclear localization signal (NLS) from pGADT7 vector (Clontech, Mountain View, CA, USA) upstream of mCherry gene in binary vector pYJmCherry using the primers listed in Table S1. These two constructs were individually introduced into *A. tumefaciens* strain GV3101 and infiltrated into the abaxial surface of 21-day-old *N. benthamiana* leaves using 1-mL needleless syringes, together with P19, a gene silencing repressor (Liang et al. 2013). After infiltration, *N. benthamiana* was grown in a growth chamber for 24 h before 50 μM MG132 (Selleck, Houston, TX, USA) was infiltrated. GFP signals were observed 24 h later with a FV1000MPE confocal microscopy (Olympus, Shinjuku-ku, Tokyo, Japan).

Construction of plasmids and generation of transgenic plants

The coding region of *BnaWGR1* was cloned to pYWLHA vector through *Nco* I and *Sal* I sites under the control of cauliflower mosaic virus (CaMV) 35S promoter. After mobilization into *A. tumefaciens* strain GV3101 through the freeze–thaw method, pYWLHA-*BnaWGR1* was transformed into WT Col-0 through a floral dip protocol (Clough and Bent 1998). Transgenic lines were selected on 1/2 × MS medium containing 25 mg L⁻¹ hygromycin for two generations to harvest homozygous T₃ seeds. High-expression lines were screened through qRT-PCR.

Quantitative RT-PCR (qRT-PCR) assay

Leaf tissues of at least three biological replicates were harvested at the specified time points for RNA extraction using Plant RNA kit (Omega bio-tek, Salt Lake City, UT, USA) followed by removal of genomic DNA using DNA-free kit

(Ambion, Vilnius, Lithuania). The first-strand cDNA was synthesized from 2.5 µg of RNA using oligo(dT)₁₈ and RNase H-MMLV (TaKaRa, Kusatsu, Shiga, Japan). Relative qRT-PCR was performed as described earlier using tenfold-diluted cDNA and a SYBR Green I kit (CW BIO, Beijing, China) on a CFX96 real-time PCR system (Bio-Rad, Hercules, CA, USA). Each primer pair was examined for amplification specificity and efficiency. The fold changes, normalized to two internal reference genes, were calculated using the formula proposed by Pfaffl (2001). For *N. benthamiana*, *NbL23* (ribosomal large subunit 23) and *NbPP2A* (type 2A serine/threonine protein phosphatase) were used as reference genes (Liu et al. 2012). As for Arabidopsis, *UBIQUITIN CONJUGATING ENZYME 21* (*AtUBC21*) and *POLYUBIQUITIN 10* (*AtUBQ10*) were used as reference genes (Czechowski et al. 2005).

For absolute qRT-PCR, each gene was cloned into the pJET1.2 vector. Serial tenfold dilutions of plasmids were performed to generate standard curve of each gene by plotting log ratio of the copy number of template DNA against the Ct value (Bustin 2000). The regression equation was generated for each gene to calculate the corresponding template copies using the formula of N (copy numbers) = quantity of plasmid / (base numbers of plasmid × 660 Da) × N_A (6.02×10^{23} copies/mol). Primers are listed in Supplementary Table S1.

Transient expression and physiological assays

Plasmids of pYWLHA-*BnaWGR1* contained in GV3101 were used for transient expression in true leaves of 30-day-old *N. benthamiana* plants. Agrobacteria harboring P19, a silencing suppressor, and *BnaWGR1* plasmids, respectively, were mixed at a ratio of 1:1 before infiltration. Leaf discs were immersed in deionized water for 30 min at 25 °C. Electric conductivity was measured using a DDS-307 conductivity meter (Leici, Shanghai, China) (Oh and Martin 2011).

3,3'-Diaminobenzidine (DAB, MP Biomedicals, Solon, OH, USA) staining was performed according to Daudi et al. (2012). For nitroblue tetrazolium (NBT, Amresco, Solon, OH, USA) staining, leaves were cut from plants and placed in 9-cm Petri dishes containing 10 mM NaN₃ (Sigma-Aldrich, St. Louis, MO, USA) in 10 mM potassium phosphate buffer (pH 7.8) before vacuumed for 30 min. Then, leaves were transferred into 0.1% NBT solution and 10 mM potassium phosphate buffer (pH 7.8) and vacuumed for 30 min again. After incubation and shaking for 2 h at room temperature, leaves were decolorized in AGE solution (acetic acid:glycerol:ethanol, 1:1:3, by vol.) and boiled for 30 min before photographed (Jambunathan 2010). The content of H₂O₂ was quantified as described previously (Li et al. 2015). Measurement of chlorophyll was performed as described based on the method reported previously (Niu

et al. 2016). In brief, 0.1 g tissue was immersed in 5 mL absolute ethanol overnight followed by measuring the absorbance at 665, 649 and 470 nm on a spectrophotometer (Thermo Scientific, Waltham, MA, USA). Malondialdehyde (MDA) was extracted from 0.1 g leaf tissue by 0.1% trichloroacetic acid (TCA, Alfa Aesar, Tianjing, China) followed by centrifugation. One milliliter of supernatant was then transferred and mixed with 2 mL of 20% TCA and 2 mL of 0.5% TBA before boiling. After cooling down, the absorbance was measured at 532 and 600 nm, respectively (Jambunathan 2010).

Terminal deoxynucleotidyl transferase-mediated dUTP nick end labeling (TUNEL, Promega, Madison, WI, USA) assay was performed as described (Li et al. 2015).

Dual-luciferase reporter assay

Promoter regions plus 5'untranslated region (UTR) were amplified from genomic DNA using the primers listed in Table S1. After confirmed through sequencing, they were cloned upstream of firefly luciferase (LUC) gene in the pGreenII0800-LUC vector as reporter plasmids (Hellens et al. 2005). An expression cassette harboring an endogenous control Renilla luciferase (REN) gene driven by 35S promoter was also located in the T-DNA region of pGreenII0800-LUC vector. pYWLHA-*BnaWGR1* and pYJHA-*GFP* were used as effector plasmid and control plasmid, respectively. The dual-LUC reporter assay was performed on 30-day-old *N. benthamiana* leaves using a dual-LUC kit (Promega) as described previously (Niu et al. 2016).

Senescence assay

Seeds of different genotypes were harvested from plants grown under the same conditions and at the same time. Seven-day-old seedlings of different genotypes were transferred from 1/2 × MS medium to soil and grown as described previously. For age-dependent leaf senescence comparison, age-matched true leaves from 38- and 43-day-old plants were used for analyzing the percentage of each group, including green, green/yellow, fully yellow, and brown/dry. The 5th, 6th and 7th true leaves of 45-day-old plants were used for chlorophyll content measurement. A percentage of yellow leaves was analyzed from 6th to 10th true leaves when the plants were 59 days old.

Prokaryotic expression, purification and electrophoretic mobility shift assay (EMSA)

BnaWGR1 was subcloned to pGEX-4T-1 vector (Amersham, Fairfield, CT, USA) through the restriction sites of *BamH* I and *Xho* I for prokaryotic expression. *Glutathione S-transferase* (*GST*)-fused *BnaWGR1* was expressed and

purified from *E. coli*. Briefly, pGEX-4T-1-*BnaWGR1* was transformed into BL21 (DE3) Codonplus strain (Stratagene, Santa Clara, CA, USA) and fusion protein was induced by isopropyl thio- β -D-galactoside (IPTG) at 25 °C for 6 h in 500 mL of 2 \times YT medium. Cell pellets were harvested for lysing and purification was performed using GST-bind resin (Novagen, Madison, WI, USA). GST proteins expressed from the empty vector pGEX-4T-1 were also purified and used as a negative control.

Different DNA fragments containing W-box elements were PCR amplified using primers listed in Table S1. Purified fragments were labeled with biotin using the biotin 3' end DNA labeling kit (Pierce, Waltham, MA, USA). To perform the DNA–protein binding reaction, 2000 ng purified *BnaWGR1*-GST was incubated with 20 fmol 3'-biotin-labeled DNA and 50 ng Poly (dI-dC) in a total volume of 20 μ L for 20 min at room temperature. Binding reactions were stopped by adding loading buffer and resolved on a 4–6% native polyacrylamide gel run in 0.5 \times Tris–borate–EDTA (TBE) buffer. The DNA–protein complex was electrophoretically transferred to Hybond N⁺ nylon membrane (Amersham) followed by cross-linking at 120 mJ cm⁻² for 45–60 s. To detect biotin-labeled DNA, the membrane was stabilized with streptavidin–horseradish peroxidase conjugate in a blocking buffer with 1–300 dilution after blocking. After equilibrating with substrate equilibration buffer, the membrane was subjected to mixture of luminol/enhancer solution and stable peroxide solution in a 1–1 ratio according to the manufacturer's manual of LightShift Chemiluminescent EMSA Kit (Pierce). The membrane was then exposed on a ChemiDoc XRS+ system (Bio-Rad).

Statistical analysis

Statistical analysis was performed using SPSS Statistics 17.0, and differences were analyzed with one-way ANOVA followed by Duncan's multiple comparison test. Student's *t* test was performed in Microsoft Excel 2003. Graphs were edited in Photoshop CS (Adobe) or plotted by Sigmaplot 12.3 (Systat Software, Inc., San Jose, CA, USA).

Results

Expression pattern of *BnaWGR1* and subcellular localization

Previously, we cloned over 53 cDNA sequences of *WRKY* genes from oilseed rape and investigated their response towards different abiotic and biotic stresses (Yang et al. 2009; Liang et al. 2013). A further microarray profiling of transcriptomic change during leaf senescence in oilseed rape indicated that quite a few *WRKY* genes were highly

expressed in senescent leaves, one of which is *BnaWGR1*. The translated sequence of *BnaWGR1* contains one DNA-binding WRKY domain, which harbors a C₂HC-type zinc-finger motif at the C terminus. Thus, *BnaWGR1* belongs to Group III in the WRKY family. When compared to the 72 Arabidopsis WRKYs and about 100 rice WRKYs, *BnaWGR1* shows the highest homology to At4g23810 (*AtWRKY53*) and Os08g29660 (*OsWRKY69*), with an identity of 71.6 and 28.2%, respectively, at the amino acid levels.


Next, we examined the subcellular localization of *BnaWGR1* by both N- and C-terminal fusion with GFP using *Agrobacterium*-mediated infiltration in leaves of *N. benthamiana*. We observed water-soaking symptoms on leaves infiltrated with *BnaWGR1*-GFP and GFP-*BnaWGR1* plasmids, suggesting ROS accumulation and hypersensitive response-like cell death. We failed to detect any green signal in *N. benthamiana* leaves via agroinfiltration at the first try. We used a well-known 26S proteasome inhibitor, MG132, and co-infiltrated it into leaf cells 1.5 days after initial agroinfiltration and observed the GFP signal 12 h later. As a result, we successfully observed green signals in nuclei (Fig. 1a), which is consistent with the role of a transcription factor. Our results suggested that there was a rapid turnover of expressed *BnaWGR1* in leaf cells, which was likely attributed to proteasomal degradation. It is known that protein levels of some specific TFs are tightly controlled in plant cells and a recent study on a tomato NAC TF gene also indicates this (Huang et al. 2013). In contrast, the GFP alone was localized to both cytoplasm and nuclei (Fig. 1a).

To examine the tissue specificity of *BnaWGR1* expression, the promoter region together with 5'UTR was cloned and fused with β -glucuronidase (*GUS*) reporter gene. We examined the *GUS* expression driven by promoter of *BnaWGR1* in two representative transgenic T₂ lines. In 7-day-old seedlings, moderate *GUS* staining in vascular system of the cotyledons and in hypocotyls was observed (Fig. 1b). In 6-week-old mature plants, strong *GUS* expression in the sepals and rosette leaves was detected. In young siliques, moderate to strong *GUS* expression was detected in the two transgenic lines (Fig. 1b). At the age of 36 and 43 days, *GUS* activity of 9th–11th leaves was significantly higher than *GUS* activity of 9th–11th leaves at the age of 30 days (Fig. S1).

Expression of *BnaWGR1* induces ROS accumulation and cell death

To further study the function of *BnaWGR1*, we cloned it into a binary vector, in which *BnaWGR1* was under the control of CaMV 35S promoter. The transient expression of *BnaWGR1* in leaves of *N. benthamiana* began to induce H₂O₂ accumulation and cell death from 3 days post-infiltration (dpi)

Fig. 1 Assays of subcellular localization and promoter-*GUS* transgenic lines of *BnaWGR1*. **a** Subcellular localization assay of GFP and *BnaWGR1*-GFP in leaf cells of *N. benthamiana*. A plasmid harboring *NLS-mCherry* was co-transformed as a nuclear marker. GFP, bright and merge represent the GFP, bright and merged fields, respectively. Bars = 50 μ m. **b** Histochemical staining of *ProBnaWGR1-GUS* transgenic plants. Two independent transgenic lines (17-2 and 19-5) were examined. The left to right panels indicate 7-day-old seedling, flowers, rosette leaves and siliques of 6-week-old mature plants




compared to the *GFP* expression control (Fig. 2a). As time went on, the symptom of cell death became more severe.

We also performed NBT staining to monitor the change in the content of superoxide anion (O_2^-), which is highly mobile, instable and could be converted to H_2O_2 . Leaves expressing *BnaWGR1* showed obvious accumulation of O_2^- , while *GFP* expression did not have such an effect (Fig. 2b). The content of H_2O_2 in the *GFP*- and *BnaWGR1*-expressing

leaves was significantly higher in the *BnaWGR1*-expressing tissues compared to *GFP* control beginning from 3 dpi (Fig. 2c).

As there was an active genomic DNA degradation during cell death, we employed TUNEL assay to test this. The green signals of fluorescein isothiocyanate (FITC) were observed in nuclei of the *BnaWGR1*-expressing leaf tissue, while there was no signal of FITC signal in the control of

Fig. 2 Expression of *BnaWGR1* induces accumulation of ROS and cell death. **a** Symptom of leaves of *N. benthamiana* after expression of *BnaWGR1* and *GFP* at 2–9 days post-infiltration (dpi). DAB staining shows H₂O₂ accumulation. **b** NBT staining of elicited O₂⁻ in leaves expressing *BnaWGR1* and *GFP* at 3–7 h post-infiltration (hpi). **c** Quantitative assay of H₂O₂ contents at three time points. **d** TUNEL assay of nuclear fragmentation in leaf tissues. DAPI staining was used to show nuclei. Scale bars = 50 μm. **e** Quantitative assay of malondialdehyde (MDA) contents in *BnaWGR1*- and *GFP*-expressing leaf tissues. **f** Measurement of electrolyte leakages in leaf tissues expressing *BnaWGR1* and *GFP*. Data in **c**, **e** and **f** were averages of three biological replicates with error bars representing standard errors. Asterisks indicate significant difference between *BnaWGR1*- and *GFP*-expressing tissues (*P* < 0.05)



GUS-expressing tissue (Fig. 2d), indicative of fragmentation of genomic DNA induced by *BnaWGR1* expression. Moreover, malondialdehyde (MDA), from lipid peroxidation due to the attacked cell membrane by intracellular ROS, increased significantly at 5, 7 and 9 dpi in leaf tissues expressing *BnaWGR1* (Fig. 2e). In addition, ion leakage increased significantly in *BnaWGR1*-expressing tissues compared to *GFP* at the three time points examined (Fig. 2f), which further supported that *BnaWGR1* positively modulated ROS accumulation and cell death.

WGR1 regulates developmental leaf senescence

To further characterize the function of *BnaWGR1*, we first performed absolute quantitative RT-PCR to investigate the

expression of *BnaWGR1* in leaves of four different developmental stages of oilseed rape plants, which were represented by young leaves (YL), mature leaves (ML), early senescent (ES) and late senescent (LS) leaves (Fig. 3a). The four stages of oilseed rape leaves were also validated by monitoring transcript levels of three marker genes, which are *BnaSAG12*, *BnaSEN1* and *-4*. The standard curve of each gene was generated by log ratio of the copy number of template DNA plotted against the Ct values (Fig. S2). As we expected, *BnaSAG12* was specifically induced at LS, and *BnaSEN1* and *-4* were induced at ML, ES and LS (Fig. 3b). *BnaWGR1* was significantly induced at ES and reduced to normal level at LS (Fig. 3b).

Furthermore, we expressed *BnaWGR1* under the control of *CaMV35S* promoter in wild-type *Arabidopsis*. Among




Fig. 3 Expression analysis of *BnaWGR1* during leaf senescence in oilseed rape. **a** The fourth true leaves of oilseed rape at four different developmental stages. *YL* young leaves of 21-day-old seedlings, *ML* fully expanded mature leaves of 28-day-old seedlings, *ES* early senescent leaves of 35-day-old seedlings, *LS* late senescent leaves of 44-day-old seedlings, with >50% leaf area yellowing. **b** Abs-

olute qRT-PCR analysis of transcript levels of *BnaSAG12*, *BnaSEN1*, *BnaSEN4* and *BnaWGR1* in wild-type leaves at different developmental stages. Data represent means of three biological replicates \pm standard errors (SE). Different letters indicate significant difference ($P < 0.05$)

over 30 transgenic T_3 lines, two *BnaWGR1*-overexpressing lines, which are OX#24 and OX#26, were identified by qRT-PCR (Fig. S3). A categorization of four groups of leaves according to the leaf colors including green, green/yellow, fully yellow, and brown/dry was used. The percentage of each category was recorded from 16 individually grown plants for statistical analysis. Using this categorization, *BnaWGR1*-OX#24 and #26 exhibited accelerated leaf senescence compared to 38- and 43-day-old WT plants (Fig. 4a and b). The percentages of green/yellow leaves of OX#24 and OX#26 were significantly higher than that of WT (Fig. 4a), whereas the percentages of green leaves of OX#24 and OX#26 were lower than that of WT (Fig. 4a, b). At 45 and 49 days post-stratification, earlier leaf senescence occurred in the two overexpression lines compared to WT as evidenced by rosette leaves (Fig. 4c). The content of chlorophyll between overexpression lines and WT was also compared in the age-matched rosette leaves. In 5th and 6th rosette leaves, the two overexpression plants had a lower

chlorophyll content than WT at the age of 45 days (Fig. 4d). Similarly, the chlorophyll content in the 7th rosette leaves of *BnaWGR1*-overexpression plants was also lower than that of WT (Fig. 4e). A simple comparison of phenotypes of the 6th through 10th rosette leaves of 59-day-old soil-grown plants also supported the conclusion (Fig. 4f). We repeated the assay independently and the results were very similar, that is, OX#24 and OX#26 showed early leaf senescence and lower chlorophyll content, compared to WT (Fig. S4). These data indicate that *BnaWGR1* positively regulates leaf senescence.

BnaWGR1 induces the transcription of ROS and cell death-related genes

To better understand how *BnaWGR1* modulated ROS accumulation and cell death, we determined to screen the putative downstream components of the *BnaWGR1* signaling pathway through examining the transcript levels of genes coding for *Rboh* and *metacaspase (MC)* both in tobacco




Fig. 4 Overexpression of *BnaWGR1* accelerates age-dependent leaf senescence. **a, b** The percentages of green, green/yellow, yellow and brown/dry leaves of 16 independent plants of 38-day-old (**a**) and 43-day-old (**b**) genotypes were calculated and compared. **c** Photographs of a 45- and 49-day-old WT and two overexpression lines. Rosette leaves excised from age-matched plants were arranged in rows from bottom to top. **d, e** Quantitative assay of total chlorophyll in the 5th and 6th true leaves (**d**) and 7th true leaves (**e**) of three

45-day-old genotypes from three independent plants. **f** Percentages of yellow leaves from the 6th to 10th true leaves of three 59-day-old genotypes. 16 independent plants for each genotype were calculated. In **a, b** and **d–f**, different letters represent significant differences from ANOVA test. *FW* fresh weight. Values were averages ± SE (*n* = 3), and significant differences are indicated by different letters based on one-way ANOVA test (*P* < 0.05)

and transgenic *Arabidopsis* plants. qRT-PCR assay of leaf tissues of tobacco plants transiently expressing *BnaWGR1* and *GFP* showed that *RbohA* and *MC5* were significantly induced in *BnaWGR1*-expressing tissues at 2 dpi and 3 dpi compared to 0 dpi (Fig. 5a). *RbohB* was also induced at 2 dpi in *BnaWGR1*-expressing tissues (Fig. 5a). However, *MC8* was not significantly changed at all (Fig. 5a). Together, these data confirmed that *BnaWGR1* mediated ROS accumulation and cell death, possibly through affecting the expression of genes directly implicated in homeostasis of ROS and cell death.

Next, the expression levels of *RbohD* and *-F* were examined in the rosette leaves of 43-day-old transgenic *Arabidopsis* plants expressing *BnaWGR1*. qRT-PCR assay

demonstrated a significant increase of *AtRbohD* and *AtRbohF* in *BnaWGR1*-overexpressing lines compared to WT, while expression of *AtMC5* and *AtMC8* showed a significant increase only in one of the two overexpression lines (Fig. 5b, c, Fig. S5). In parallel, the expression level of a senescence marker gene *SAG12* was also significantly higher in transgenic plants compared to WT (Fig. S5).

We further utilized a dual-luciferase (LUC) system to investigate whether *BnaWGR1* directly activates the transcription of *RbohD* and *RbohF*, which are orthologs of *RbohB* and *RbohA* from tobacco, respectively. A 2.1 and 2.8 kb of promoter and 5'UTR fragments of *AtRbohD* and *AtRbohF*, respectively, were cloned into the reporter vector, in which a CaMV 35S promoter-controlled *Renilla*




Fig. 5 Screening of putative target genes of *BnaWGR1* through qRT-PCR and dual-LUC assays. **a** Identification of transcriptional changes of ROS- and cell death-related genes through qRT-PCR at 0, 2 and 3 dpi. Each value represents the mean \pm SE of four biological replicates. Asterisks denote significant differences (compared to the level at 0 dpi) by Student *t* test ($P < 0.05$). **b, c** Expression analysis of *AtRbohD* and *AtRbohF* in the two independent *BnaWGR1*-expression transgenic lines at the indicated different leaf ages of 43-day-old plants by qRT-PCR. Data are means of four independent biological replicates \pm SE. Significant differences are indicated by different letters based on one-way ANOVA test ($P < 0.05$). **d** Schematic diagrams of reporter and effector plasmids used in the dual-luciferase (LUC)

assay. Promoters of *RbohD* and *RbohF* driving firefly LUC reporter gene were used as reporters while the CaMV 35S promoter-controlled *Renilla luciferase* (*REN*) was used for internal normalization. The effector construct consists of either *BnaWGR1* or *GFP* driven by the CaMV 35S promoter. Black rectangles depict terminator sequences. **e, f** The dual-LUC assay results of transcriptional regulation of *AtRbohD* (**e**) and *AtRbohF* (**f**) mediated by *BnaWGR1* in leaves of *N. benthamiana*. The relative LUC activity was calculated as the ratio of LUC/REN. Data are means of three independent biological replicates \pm SE. Asterisks denote significant differences by Student's *t* test ($P < 0.05$)

luciferase (*REN*) was used as the internal normalization gene (Fig. 5d). The effector was *BnaWGR1* driven by CaMV 35S while that of *GFP* was used as the control (Fig. 5d). The different combinations of effector and reporter constructs were co-expressed in a transient system. The relative LUC activity represented by LUC/REN ratio was monitored at three time points. It showed that transient expression of *BnaWGR1* significantly induced the transcription of *LUC* driven by the promoters of *AtRbohD* and *-F* (Fig. 5e, f). The data suggest that *BnaWGR1* could

possibly directly regulate the transcription of *AtRbohD* and *AtRbohF*.

BnaWGR1 directly binds to promoters of *RbohD* and *RbohF*

To test if *BnaWGR1* could specifically bind to the W-box element, EMSA was performed. A recombinant *BnaWGR1* fused with GST tag (*BnaWGR1*-GST) was purified from a prokaryotic system and the tandem repeat W-box probe was

labeled with biotin (Fig. 6a). A thick band of protein–DNA complex with reduced mobility was observed when recombinant BnaWGR1-GST was incubated with 4xW-box probes, but not with a mutated 4xW-box, in which TTGACC was mutated to TTGAAC (Fig. 6b). Furthermore, a competition experiment using unlabeled 4xW-box probes confirmed the binding specificity of BnaWGR1 to W-box, but not to the mutated W-box sequence (Fig. 6b).

An analysis of the promoters of *AtRbohD* and *AtRbohF* indicates that there are five and two canonical W-box elements, respectively (Fig. S6). Hence, we speculated that BnaWGR1 might directly bind to some of these W-boxes during the transcriptional regulation of *AtRbohD* and *F*. Hence, we designed five probes, namely P1 through P5 for the *AtRbohD* (Fig. 7a). It is noted that two W-box elements are clustered in the P1 fragment. The results showed that shifted bands (retarded binding complex) for some probes were clearly detected (Fig. 7b). Moreover, weaker retarded bands for P1, P2 and P5 fragments were observed when unlabeled competitive probes were added (Fig. 7c). The results indicate that BnaWGR1 can recognize and bind to the W-box *cis*-elements in *AtRbohD* promoter.




Fig. 6 BnaWGR1 binds to W-box elements. **a** Schematic diagram of quadruple W-box and the mutated W-box (mW-box) sequences. **b** EMSA assay of binding of recombinant BnaWGR1-GST to canonical W-box sequence, but not to mW-box sequence. The BnaWGR1 fused to GST tag and GST protein (negative control) were used to detect the binding. Biotin-labeled probes were incubated with recombinant BnaWGR1-GST or GST proteins. 10- and 20-fold molar excess of unlabelled W-box or mW-box fragments were used as competitive probes

Similarly, the binding of BnaWGR1 to promoter of *AtRbohF* was tested through EMSA. Three fragments (P1 through P3) were amplified and labeled with biotin (Fig. 7d). The results showed that recombinant BnaWGR1-GST bound to P1 and P3, but not to P2 (Fig. 7e). In parallel, BnaWGR1-GST also bound to the quadruple W-box sequence (Fig. 7f). A further competition experiment showed that unlabelled probes successfully diminished the binding of BnaWGR1 to P1 and P3 fragments, which suggests the specificity of binding (Fig. 7f).


Moreover, promoter regions of *BnaRbohD* and *BnaRbohF* from oilseed rape were cloned and multiple W-box elements were also identified in the sequences (Fig. 8a, b, Fig. S5). Three and two fragments harboring W-box elements were prepared and subjected to binding assay with recombinant BnaWGR1-GST proteins. BnaWGR1 bound to all the fragments, although with different affinities (Fig. 8c). We further performed a dual-luciferase (LUC) assay to examine the transactivation of *BnaRbohD* and *F* by BnaWGR1. A 1.2 kb of promoter and 5'UTR fragments of *BnaRbohD* and *-F*, respectively, were cloned into the reporter vector (Fig. 8d). The relative LUC activity revealed that transient expression of *BnaWGR1* significantly induced the transcription of *LUC* driven by the promoters of *BnaRbohD* and *-F* at the three time points, compared to the control (Fig. 8e, f). Taken together, these data reveal that BnaWGR1 regulates the expression of *RbohD* and *RbohF* genes of Arabidopsis through directly binding to their promoters and also enhances the promoter activities of *RbohD* and *RbohF* orthologs in oilseed rape.

Discussion

Leaf senescence is a developmental programmed cell death process providing plants with phenotypic plasticity to help them to adapt to adverse environmental conditions (Lim et al. 2007). Leaf senescence also determines crop plant yields and, therefore, a better understanding of the essential genes regulating this process is important. In this study, we characterized the function and molecular mechanism of one of a WRKY-type transcription factor gene in regulating leaf senescence of the oil crop, oilseed rape.

Expression of *BnaWGR1* promotes leaf senescence

Previously, a transcriptomic profiling of early senescent leaves of oilseed rape identified *BnaWGR1* as a WRKY-type gene highly induced in early senescent leaves (unpublished data). Further, a transient expression of *BnaWGR1* in leaves of *N. benthamiana* indicated significant accumulation of ROS and hypersensitive response-like cell death (Fig. 2). A transcriptional profiling of *BnaWGR1* expression






Fig. 8 BnaWGR1 directly binds to the promoter region of *BnaRbohD* and *BnaRbohF*. **a**, **b** The schematic diagrams of the promoter regions of *BnaRbohD* (**a**) and *BnaRbohF* (**b**). Black rectangles indicate W-box and W-box-like sequences. The different fragments used in EMSA were indicated. **c** EMSA analysis of binding of recombinant BnaWGR1–GST to different fragments of promoters. The canonical tandem repeat W-box probe was used as a positive control. **d** Schematic diagrams of reporter and effector plasmids used in the dual-luciferase (LUC) assay. Promoters of *BnaRbohD* and *BnaRbohF* driving firefly luciferase (LUC) reporter gene were used as reporters

while the CaMV 35S promoter-controlled *Renilla luciferase* (*REN*) was used for internal normalization. The effector construct consists of either BnaWGR1 or GFP driven by the CaMV 35S promoter. Black rectangles depict terminator sequences. **e**, **f** The dual-LUC assay of transcriptional regulation of *BnaRbohD* (**e**) and *BnaRbohF* (**f**) mediated by BnaWGR1 in leaves of *N. benthamiana*. The relative LUC activity was calculated as the ratio of LUC/REN. Data are means of three independent biological replicates ± SE. Asterisks denote significant differences by Student’s *t* test (**P* < 0.05; ***P* < 0.01; ****P* < 0.001)

at the four different stages of leaf development revealed high expression levels only in the early senescent leaves (Fig. 3), suggesting a potential role of BnaWGR1 in process of leaf senescence. This observation was complemented with the GUS activity assay when senescence is beginning (Fig. S1). A stronger GUS activity of 9th–11th leaves of the transgenic lines harboring the Pro*BnaWGR1*-GUS construct was

observed at the age of 36 and 43 days (Fig. S1). In Arabidopsis, some TF genes are preferentially expressed in early senescencing leaves, while others in late senescencing leaves (Gepstein et al. 2003; Balazadeh et al. 2008). We, therefore, propose that BnaWGR1 may play an important role when leaf senescence procedure proceeds. Furthermore, one of the transgenic Arabidopsis expressing *BnaWGR1* showed

a significantly precocious leaf senescence compared to WT and the other transgenic line also showed the trend of earlier senescence than WT (Fig. 4, Fig. S4). This result indicates that *BnaWGR1* can positively regulate leaf senescence. It should be noted that the expression level of *BnaWGR1* might be post-translationally regulated or tightly controlled as a signal of GFP fused *BnaWGR1* was only detected when the 26S proteasome inhibitor MG132 was used (Fig. 1a).

BnaWGR1 positively regulates the expression of *RbohD* and *RbohF*

For a regulator mastering the early events of senescence, it is interesting to investigate the downstream targets. For example, three abscisic acid (ABA)-responsive element (ABRE)-binding transcription factors, ABF2, ABF3, and ABF4 act as positive regulators of chlorophyll catabolism genes to trigger ABA-dependent chlorophyll degradation and leaf senescence (Gao et al. 2016). AtNAP promotes chlorophyll degradation and leaf senescence via enhancing ABA synthesis mediated by abscisic aldehyde oxidase3 (AAO3) (Yang et al. 2014). OsNAP, an ortholog of AtNAP, also acts as a positive regulator of onset of senescence by modulating ABA biosynthesis and directly targeting *SAGs* in rice (Liang et al. 2014). As for AtWRKY53, it could bind to the promoters of a few transcription factor genes including *WRKY6*, *-62*, senescence-associated and defense-related genes (Miao et al. 2004). Surprisingly, AtWRKY53 can bind to promoters and transactivate the expression of *CAT2* and *-3*, which encode two ROS-scavenging enzymes decreasing the content of H₂O₂ (Miao et al. 2004). We observed that the expression of *BnaWGR1* induced significant ROS accumulation (Fig. 2). It implies that *BnaWGR1* may disturb ROS homeostasis by either repressing genes encoding ROS-scavenging enzymes or activating genes encoding ROS-producing enzymes. Using a combination of methods, we identified two possible new direct targets of *BnaWGR1*, which are ROS-producing genes encoding *RbohD* and *-F* to be activated by *BnaWGR1* through dual-LUC and EMSA assays.

ROS are highly reactive molecules being able to damage cellular components but are also proposed as signaling elements at lower concentrations (Apel and Hirt 2004). Plant homologs of mammalian NADPH oxidase (NOX), also known as Rbohs, act as a group of plasma membrane-localized enzymes producing H₂O₂ that mediate the switch of plants from growth to defense or adapt to environmental changes (Marino et al. 2012). Previously, it has been demonstrated that NbRbohA and NbRbohB, orthologs of AtRbohF and AtRbohD, respectively, are the two main members responsible for ROS production in *N. benthamiana* (Yoshioka et al. 2003). AtRbohD and *-F* are two famous enzymes mediating ABA promotion of ROS production, ABA-induced cytosolic Ca²⁺ increase, ABA-activated

plasma membrane Ca²⁺-permeable channels in guard cells, and ABA-induced stomatal closing, as well as plant immunity (Torres et al. 2002; Kadota et al. 2014; Adachi et al. 2015). The diverse functions of Rbohs suggest they may serve as important molecular “hubs” in ROS signaling in plants (Marino et al. 2012). However, the role of *RbohD* and *RbohF* in leaf senescence has not been reported yet. Here, we provide supporting evidence that *BnaWGR1* may regulate the expression of *RbohD* and *-F* to accelerate leaf senescence through increasing ROS levels.

In conclusion, this study reveals that the oilseed rape *WGR1* transcription factor acts as a possible positive regulator of leaf senescence potentially through promoting the expression of two central ROS-producing enzyme genes, *RbohD* and *-F*, by directing binding to their promoters to advance ROS production. This is a finding not reported for any WRKY transcription factor gene in plants. It could be possible that other sources of ROS within cells may be present and hence make a global analysis of ROS levels and interactions more challenging. Collectively, the current work uncovers the potential role of an oilseed rape *BnaWGR1* in leaf senescence and will be helpful for a better understanding and manipulation of leaf senescence in oilseed rape.

Author contribution statement BY and YQJ conceived and designed research. LY, CY and YZ conducted experiments. XC and YW contributed materials. BY and YQJ analyzed data. BY and YQJ wrote the manuscript. All authors read and approved the manuscript.

Acknowledgements This work was supported by the National Natural Science Foundation of China [No. 31301648 to BY and 31270293 to Y-QJ]. Natural Science Foundation of Shaanxi Province [no. 2014JQ3102 to BY]. The Fundamental Research Funds for the Central Universities of China [no. 2109021137, nos. 2452015413, 2452016009, to BY]. We thank Profs. Jian-Ye Chen (South China Agricultural University) and Benke Kuai (Fudan University) for technical assistance.

Compliance with ethical standards

Conflict of interest The authors declare that they have no competing interests.

References

- Adachi H, Nakano T, Miyagawa N, Ishihama N, Yoshioka M, Katou Y, Yaeno T, Shirasu K, Yoshioka H (2015) WRKY transcription factors phosphorylated by MAPK regulate a plant immune NADPH oxidase in *Nicotiana benthamiana*. *Plant Cell* 27:2645–2663
- Apel K, Hirt H (2004) Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu Rev Plant Biol* 55:373–399
- Bakshi M, Oelmüller R (2014) WRKY transcription factors: Jack of many trades in plants. *Plant Signal Behav* 9:e27700

- Balazadeh S, Riano-Pachon DM, Mueller-Roeber B (2008) Transcription factors regulating leaf senescence in *Arabidopsis thaliana*. *Plant Biol (Stuttg)* 10(Suppl 1):63–75
- Baxter A, Mittler R, Suzuki N (2014) ROS as key players in plant stress signalling. *J Exp Bot* 65:1229–1240
- Besseau S, Li J, Palva ET (2012) WRKY54 and WRKY70 co-operate as negative regulators of leaf senescence in *Arabidopsis thaliana*. *J Exp Bot* 63:2667–2679
- Bhattacharai KK, Atamian HS, Kaloshian I, Eulgem T (2010) WRKY72-type transcription factors contribute to basal immunity in tomato and *Arabidopsis* as well as gene-for-gene resistance mediated by the tomato *R* gene *Mi-1*. *Plant J* 63:229–240
- Birkenbihl RP, Diezel C, Somssich IE (2012) *Arabidopsis* WRKY33 is a key transcriptional regulator of hormonal and metabolic responses toward *Botrytis cinerea* infection. *Plant Physiol* 159:266–285
- Bustin SA (2000) Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J Mol Endocrinol* 25:169–193
- Chen L, Zhang L, Yu D (2010) Wounding-induced WRKY8 is involved in basal defense in *Arabidopsis*. *Mol Plant Microbe Interact* 23:558–565
- Ciolkowski I, Wanke D, Birkenbihl RP, Somssich IE (2008) Studies on DNA-binding selectivity of WRKY transcription factors lend structural clues into WRKY-domain function. *Plant Mol Biol* 68:81–92
- Clough SJ, Bent AF (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* 16:735–743
- Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible WR (2005) Genome-wide identification and testing of superior reference genes for transcript normalization in *Arabidopsis*. *Plant Physiol* 139:5–17
- Daudi A, Cheng Z, O'Brien JA, Mammarella N, Khan S, Ausubel FM, Bolwell GP (2012) The apoplastic oxidative burst peroxidase in *Arabidopsis* is a major component of pattern-triggered immunity. *Plant Cell* 24:275–287
- Eulgem T, Rushton PJ, Robatzek S, Somssich IE (2000) The WRKY superfamily of plant transcription factors. *Trends Plant Sci* 5:199–206
- Gao S, Gao J, Zhu X, Song Y, Li Z, Ren G, Zhou X, Kuai B (2016) ABF2, ABF3, and ABF4 promote ABA-mediated chlorophyll degradation and leaf senescence by transcriptional activation of chlorophyll catabolic genes and senescence-associated genes in *Arabidopsis*. *Mol Plant* 9:1272–1285
- Gepstein S, Sabehi G, Carp MJ, Hajouj T, Neshor MF, Yariv I, Dor C, Bassani M (2003) Large-scale identification of leaf senescence-associated genes. *Plant J* 36:629–642
- Hellens RP, Allan AC, Friel EN, Bolitho K, Grafton K, Templeton MD, Karunairatnam S, Gleave AP, Laing WA (2005) Transient expression vectors for functional genomics, quantification of promoter activity and RNA silencing in plants. *Plant Methods* 1:13
- Hinderhofer K, Zentgraf U (2001) Identification of a transcription factor specifically expressed at the onset of leaf senescence. *Planta* 213:469–473
- Huang W, Miao M, Kud J, Niu X, Ouyang B, Zhang J, Ye Z, Kuhl JC, Liu Y, Xiao F (2013) SINAC1, a stress-related transcription factor, is fine-tuned on both the transcriptional and the post-translational level. *New Phytol* 197:1214–1224
- Jambunathan N (2010) Determination and detection of reactive oxygen species (ROS), lipid peroxidation, and electrolyte leakage in plants. *Methods Mol Biol* 639:292
- Jiang Y, Deyholos MK (2009) Functional characterization of *Arabidopsis* NaCl-inducible WRKY25 and WRKY33 transcription factors in abiotic stresses. *Plant Mol Biol* 69:91–105
- Jiang Y, Liang G, Yang S, Yu D (2014) *Arabidopsis* WRKY57 functions as a node of convergence for jasmonic acid- and auxin-mediated signaling in jasmonic acid-induced leaf senescence. *Plant Cell* 26:230–245
- Kadota Y, Sklenar J, Derbyshire P, Stransfeld L, Asai S, Ntoukakis V, Jones JD, Shirasu K, Menke F, Jones A, Zipfel C (2014) Direct regulation of the NADPH oxidase RBOHD by the PRR-associated kinase BIK1 during plant immunity. *Mol Cell* 54:43–55
- Kim K, Franceschi V, Davin L, Lewis N (2006) β -Glucuronidase as reporter gene. Advantages and limitations. In: Salinas J, Sanchez-Serrano JJ (eds) *Arabidopsis* protocols. Humana Press Inc, Totowa, pp 263–273
- Knoth C, Ringler J, Dangl JL, Eulgem T (2007) *Arabidopsis* WRKY70 is required for full RPP4-mediated disease resistance and basal defense against *Hyaloperonospora parasitica*. *Mol Plant Microbe Interact* 20:120–128
- Kwak JM, Mori IC, Pei ZM, Leonhardt N, Torres MA, Dangl JL, Bloom RE, Bodde S, Jones JD, Schroeder JI (2003) NADPH oxidase *AtrbohD* and *AtrbohF* genes function in ROS-dependent ABA signaling in *Arabidopsis*. *EMBO J* 22:2623–2633
- Lee S, Seo PJ, Lee HJ, Park CM (2012) A NAC transcription factor NTL4 promotes reactive oxygen species production during drought-induced leaf senescence in *Arabidopsis*. *Plant J* 70:831–844
- Li L, Ye C, Zhao R, Li X, Liu WZ, Wu F, Yan J, Jiang YQ, Yang B (2015) Mitogen-activated protein kinase kinase kinase (MAP-KKK) 4 from rapeseed (*Brassica napus* L.) is a novel member inducing ROS accumulation and cell death. *Biochem Biophys Res Commun* 467:792–797
- Liang W, Yang B, Yu B, Zhou Z, Li C, Sun Y, Zhang Y, Jia M, Wu F, Zhang H, Wang B, Deyholos M, Jiang Y-Q (2013) Identification and analysis of MKK and MPK gene families in canola (*Brassica napus* L.). *BMC Genom* 14:392
- Liang C, Wang Y, Zhu Y, Tang J, Hu B, Liu L, Ou S, Wu H, Sun X, Chu J, Chu C (2014) OsNAP connects abscisic acid and leaf senescence by fine-tuning abscisic acid biosynthesis and directly targeting senescence-associated genes in rice. *Proc Natl Acad Sci USA* 111:10013–10018
- Lim PO, Nam HG (2005) The molecular and genetic control of leaf senescence and longevity in *Arabidopsis*. *Curr Top Dev Biol* 67:49–83
- Lim PO, Kim HJ, Nam HG (2007) Leaf senescence. *Annu Rev Plant Biol* 58:115–136
- Liu D, Shi L, Han C, Yu J, Li D, Zhang Y (2012) Validation of reference genes for gene expression studies in virus-infected *Nicotiana benthamiana* using quantitative real-time PCR. *PLoS ONE* 7:e46451
- Marino D, Dunand C, Puppo A, Pauly N (2012) A burst of plant NADPH oxidases. *Trends Plant Sci* 17:9–15
- Miao Y, Laun T, Zimmermann P, Zentgraf U (2004) Targets of the WRKY53 transcription factor and its role during leaf senescence in *Arabidopsis*. *Plant Mol Biol* 55:853–867
- Mittler R, Vanderauwera S, Gollery M, Van Breusegem F (2004) Reactive oxygen gene network of plants. *Trends Plant Sci* 9:490–498
- Navabpour S, Morris K, Allen R, Harrison E, Mackerness SAH, Buchanan-Wollaston V (2003) Expression of senescence-enhanced genes in response to oxidative stress. *J Exp Bot* 54:2285–2292
- Niu F, Wang C, Yan J, Guo X, Wu F, Yang B, Deyholos MK, Jiang YQ (2016) Functional characterization of NAC55 transcription factor from oilseed rape (*Brassica napus* L.) as a novel transcriptional activator modulating reactive oxygen species accumulation and cell death. *Plant Mol Biol* 92:89–104
- Oh CS, Martin GB (2011) Tomato 14-3-3 protein TFT7 interacts with a MAP kinase kinase to regulate immunity-associated programmed

- cell death mediated by diverse disease resistance proteins. *J Biol Chem* 286:14129–14136
- Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 29:e45
- Rivero RM, Kojima M, Gepstein A, Sakakibara H, Mittler R, Gepstein S, Blumwald E (2007) Delayed leaf senescence induces extreme drought tolerance in a flowering plant. *Proc Natl Acad Sci USA* 104:19631–19636
- Robatzek S, Somssich IE (2001) A new member of the *Arabidopsis* WRKY transcription factor family, *AtWRKY6*, is associated with both senescence- and defence-related processes. *Plant J* 28:123–133
- Robatzek S, Somssich IE (2002) Targets of *AtWRKY6* regulation during plant senescence and pathogen defense. *Genes Dev* 16:1139–1149
- Rogers H, Munne-Bosch S (2016) Production and scavenging of reactive oxygen species and redox signaling during leaf and flower senescence: similar but different. *Plant Physiol* 171:1560–1568
- Torres MA, Dangl JL, Jones JD (2002) *Arabidopsis* gp91^{phox} homologues *AtrbohD* and *AtrbohF* are required for accumulation of reactive oxygen intermediates in the plant defense response. *Proc Natl Acad Sci USA* 99:517–522
- Yang B, Jiang Y, Rahman MH, Deyholos MK, Kav NN (2009) Identification and expression analysis of *WRKY* transcription factor genes in canola (*Brassica napus* L.) in response to fungal pathogens and hormone treatments. *BMC Plant Biol* 9:68
- Yang J, Worley E, Udvardi M (2014) A NAP-AAO3 regulatory module promotes chlorophyll degradation via ABA biosynthesis in *Arabidopsis* leaves. *Plant Cell* 26:4862–4874
- Ye Z, Rodriguez R, Tran A, Hoang H, de los Santos D, Brown S, Vellanoth RL (2000) The developmental transition to flowering represses ascorbate peroxidase activity and induces enzymatic lipid peroxidation in leaf tissue in *Arabidopsis thaliana*. *Plant Sci* 158:115–127
- Yoshioka H, Numata N, Nakajima K, Katou S, Kawakita K, Rowland O, Jones JD, Doke N (2003) *Nicotiana benthamiana* gp91^{phox} homologs *NbrbohA* and *NbrbohB* participate in H₂O₂ accumulation and resistance to *Phytophthora infestans*. *Plant Cell* 15:706–718
- Zheng Z, Qamar SA, Chen Z, Mengiste T (2006) *Arabidopsis* WRKY33 transcription factor is required for resistance to necrotrophic fungal pathogens. *Plant J* 48:592–605
- Zhou X, Jiang Y, Yu D (2011) WRKY22 transcription factor mediates dark-induced leaf senescence in *Arabidopsis*. *Mol Cells* 31:303–313
- Zimmermann P, Heinlein C, Orendi G, Zentgraf U (2006) Senescence-specific regulation of catalases in *Arabidopsis thaliana* (L.) Heynh. *Plant Cell Environ* 29:1049–1060