



Nitro-fatty acids: electrophilic signaling molecules in plant physiology

Luciano Di Fino¹ · Andrés Arruebarrena Di Palma¹ · Enzo A. Perk¹ · Carlos García-Mata¹ · Francisco J. Schopfer² · Ana M. Laxalt¹

Received: 24 June 2021 / Accepted: 30 October 2021 / Published online: 13 November 2021
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2021

Abstract

Main conclusion Nitro fatty acids (NO₂-FA) have relevant physiological roles as signaling molecules in biotic and abiotic stress, growth, and development, but the mechanism of action remains controversial. The two main mechanisms involving nitric oxide release and thiol modification are discussed.

Abstract Fatty acids (FAs) are major components of membranes and contribute to cellular energetic demands. Besides, FAs are precursors of signaling molecules, including oxylipins and other oxidized fatty acids derived from the activity of lipoxygenases. In addition, non-canonical modified fatty acids, such as nitro-fatty acids (NO₂-FAs), are formed in animals and plants. The synthesis NO₂-FAs involves a nitration reaction between unsaturated fatty acids and reactive nitrogen species (RNS). This review will focus on recent findings showing that, in plants, NO₂-FAs such as nitro-linolenic acid (NO₂-Ln) and nitro-oleic acid (NO₂-OA) have relevant physiological roles as signaling molecules in biotic and abiotic stress, growth, and development. Moreover, since there is controversy on mechanisms of action of NO₂-FAs as signaling molecules, we will provide evidence showing why this aspect needs further evaluation.

Keywords Nitro-fatty acids · Reactive nitrogen species · Nitric oxide

Introduction

Nitro-fatty acids (NO₂-FAs) are the product of the reaction between nitric oxide (NO)- and nitrite (NO₂⁻)-derived reactive nitrogen species (RNS) and unsaturated fatty acids. The formation of nitro-lipids was initially proposed based on the observation that NO inhibited lipid oxidation propagation reactions (Rubbo et al. 1994). The resulting species remained elusive, but the incorporation of a NO₂ group into linoleic acid was reported (O'Donnell et al. 1999). Nitrated linoleic acid was then reported in humans (Baker et al. 2004; Lima et al. 2002). NO₂-FAs were later detected in plants (Fazzari et al. 2014) and proposed as signal molecules

involved in oxidative stress responses, plant development, and defense responses (Mata-Perez et al. 2016a; Arruebarrena Di Palma et al. 2020; Vollár et al. 2020; Di Fino et al. 2020).

Mechanisms of NO₂-FA formation

The signaling events triggered by NO₂-FAs have been extensively studied, contrasting with the lower number of reports evaluating the formation and characterization of endogenous species. In animals, reactions in the gastric compartment promoted by dietary or salivary NO₂⁻ are the major contributor to systemic levels (Delmastro-Greenwood et al. 2015). A second mechanism includes NO-driven nitration reactions during inflammation (Bonacci et al. 2012; Vitturi et al. 2013; Villacorta et al. 2018). A third mechanism includes the reaction of NO₂⁻-derived species during ischemia/reperfusion events. Independent of the mechanism, the nitration of fatty acids occurs through an initial addition of the nitrogen dioxide radical ([•]NO₂) to a double bond with the formation of a β-nitroalkyl radical (Trostchansky and Rubbo 2008;

Communicated by Gerhard Leubner.

✉ Ana M. Laxalt
amlaxalt@mdp.edu.ar

¹ Instituto de Investigaciones Biológicas, CONICET-Universidad Nacional de Mar del Plata, Mar del Plata, Argentina

² Department of Pharmacology and Chemical Biology, University of Pittsburgh, Pittsburgh, PA, USA




Fig. 1 Role of reactive nitrogen species in fatty acid nitration. Nitration of fatty acids occurs via the addition of $\bullet\text{NO}_2$ (nitrogen dioxide radical) to a double bond. NO-dependent nitration (dark grey boxes). Under oxidative conditions with concomitant formation superoxide anion ($\text{O}_2^{\bullet-}$) and $\bullet\text{NO}$, these molecules react at diffusion-limited reaction rates to yield peroxynitrite (ONOO^-). Through protonation and then homolysis, ONOO^- produces $\bullet\text{NO}_2$. Under conditions with high local NO concentrations and the presence of oxygen, the auto-oxidation of NO leads to the formation of nitrogen dioxide. Heme peroxidase-catalyzed formation (light grey boxes). NO_2^- , the proximal product of $\bullet\text{NO}$ oxidation and nitrate (NO_3^-) reduction, can be oxidized to $\bullet\text{NO}_2$ in a one-electron oxidation reaction catalyzed by heme peroxidases. Acid-catalyzed formation (white boxes). The protonation of NO_2^- to HNO_2 under acidic conditions (pKa 3.4) leads to its dimerization to form dinitrogen tetroxide ($\text{H}_2\text{N}_2\text{O}_4$), which, upon dehydration, forms N_2O_3 . This species is unstable and undergoes homolysis to form $\bullet\text{NO}$ and $\bullet\text{NO}_2$

Schopfer et al. 2019). Figure 1 summarizes the chemical mechanisms of NO_2 -FA formation. Thus, the different biological scenarios in which NO_2 -FA are formed are defined by the mechanism involved in the formation of $\bullet\text{NO}_2$. This has important consequences, as the different mechanisms involve reaction milieus that differ on hydrophobicity, the rate and level of different nitrogen oxides formation, oxygen tension, and contributions of other oxidants and anti-oxidants. These factors will modulate the rates, yields, and products formed during the nitration reactions. The main nitration pathways relevant to biological conditions are depicted in different colours in Fig. 1:

- NO-dependent nitration: NO plays a central role in the formation of nitrated fatty acids. Under oxidative conditions with concomitant formation of superoxide anion ($\text{O}_2^{\bullet-}$) and $\bullet\text{NO}$, these molecules react at diffusion-limited reaction rates to yield peroxynitrite (ONOO^-). Upon protonation to peroxynitrous acid at physiological pH values, ONOO^- can undergo homolytic cleavage to form ($\bullet\text{NO}_2$ and $\bullet\text{OH}$). Under conditions with high

local $\bullet\text{NO}$ concentrations and the presence of oxygen, the auto-oxidation of NO leads to the formation of nitrogen dioxide.

- Heme peroxidase-catalyzed formation of $\bullet\text{NO}_2$: NO_2^- , the proximal product of $\bullet\text{NO}$ oxidation and nitrate (NO_3^-) reduction, can be oxidized to $\bullet\text{NO}_2$ in a one-electron oxidation reaction catalyzed by heme peroxidases (van der Vliet et al. 1997).
- Acid-catalyzed formation of $\bullet\text{NO}_2$: The protonation of NO_2^- to HNO_2 under acidic conditions (pKa 3.4) leads to its dimerization to form dinitrogen tetroxide ($\text{H}_2\text{N}_2\text{O}_4$), which, upon dehydration, forms N_2O_3 . This species is unstable and undergoes homolysis to form NO and $\bullet\text{NO}_2$ (Vitturi et al. 2015).

The nitration of fatty acids proceeds through the addition of the $\bullet\text{NO}_2$ radical to a double bond present in fatty acids. This is an extremely fast reaction, but results in an unstable product and the $\bullet\text{NO}_2$ is consequently eliminated with the reformation of the double bond (Pryor et al. 1981; Gallon and Pryor 1993). This futile reaction can lead to double bond isomerization, as demonstrated for arachidonic acid (Balazy and López-Fernández 2003). When the addition reaction occurs on a conjugated double bond, the intermediary product is stabilized by resonance, providing an opportunity for other reactions to occur. Through a series of complex reactions, this radical is finally oxidized to a nitroalkene (Fazzari et al. 2014). Consequently, nitration yields are largely increased ($> 10^5$ fold) in the presence of conjugated double bonds (Bonacci et al. 2012). In plants, similar scenarios can be foreseen; however, it is unclear whether any of the mechanisms mentioned above play a role or if these reactions take place in specific cellular or subcellular compartments.

NO_2 -FAs detection

Nitrated fatty acids were initially described in plants as cysteine adducts of nitro-oleic acid (NO_2 -OA) (50 pmol NO_2 -OA-Cys/g FW) and as conjugated nitro-linoleic acid (NO_2 -cLA) (9- and 12- NO_2 -cLA isomers), as described in olive fruit and extra virgin olive oil (EVOO), respectively (Fazzari et al. 2014). Later, in *Arabidopsis thaliana*, nitro-linolenic acid (NO_2 -Ln) was detected in seeds (11 pmol/g FW), 14-old days seedling (3.8 pmol/g FW) and in leaves of 45-day old plants (0.54 pmol/g FW) (Mata-Pérez et al. 2016a). The same group later reported the presence of NO_2 -Ln in rice (*Oryza sativa*) and Pea (*Pisum sativum*). However, some methodological details were missing in the manuscript (e.g., growth plant condition, stage of the plants, techniques for isolation and detection of NO_2 -FAs), making the interpretation of the results more difficult (Mata-Pérez

et al. 2017). Recently, another group reported the presence of NO₂-OA in *Brassica napus* seeds (1.69 nmol/g FW) and 7-day-old seedling plants (0.66 nmol/g FW) (Vollár et al. 2020). The published data makes it clear that the highest content of NO₂-FAs as free fatty acid (non-esterified) is found in seeds, likely due to the high fatty acid content found in seeds (Mata-Pérez et al. 2016a; Vollár et al. 2020). In mammals, the levels of esterified NO₂-FAs account for over 95% of total circulating levels, with accumulation in the adipose tissue (Fazzari et al. 2017). Thus, these reports provide support to the presence and regulation of NO₂-FAs levels in plants.

Despite these findings, the detection, characterization, and quantification of NO₂-FAs in plants have proven extremely challenging. Several factors have contributed to this fact and are analyzed below:

Formation of Michael addition products: NO₂-FAs form reversible covalent adducts with nucleophilic amino acids of proteins, particularly cysteine and histidine (Rudolph et al. 2009). The cellular cytoplasm is a highly reducing environment and, in animals, the cysteine content in proteins amounts to 10–50 mM and glutathione (GSH) levels are between 4 and 12 mM (Rudolph et al. 2009; Turell et al. 2013). In animal-derived cells, it has been calculated that approximately 99% of NO₂-cLA is adducted, leaving only a small proportion of the molecule free for detection (Turell et al. 2017). Moreover, the proportion of free fatty acids in plant cells is low compared to the total fraction of fatty acid esterified to membranes phospholipids and triglycerides. Thus, to account for the presence of NO₂-FA in membranes, investigators treated the lipid extracts with a lipase cocktail (Mata-Pérez et al. 2016a, 2018).

Matrix effects: plant samples are very rich in pigments of lipidic nature, such as chlorophylls and carotenoids, that can affect extraction yields and further reduce detection of NO₂-FAs by dampening ionization when measured by liquid chromatography with tandem mass spectrometry (LC-MS/MS) (Mata-Pérez et al. 2018). Although specific experiments to address this problem have not been conducted, the high content of these pigments might be partially responsible for the low concentration detected in pigmented tissues compared to non-pigmented tissues.

Currently, the number of identified nitrated lipid species in plants (NO₂-OA, NO₂-Ln and NO₂-cLA) is smaller than those reported in humans (Salvatore et al. 2020). NO₂-cLA is the most abundant NO₂-FA detected in human urine and plasma, followed closely by NO₂-cLn (Salvatore et al. 2020; Bonacci et al. 2012). In vitro assays revealed that conjugated LA (cLA) yielded 10⁵ greater nitration products than LA (Bonacci et al. 2012). The presence of NO₂-cLA in olive is intriguing given the scarcity of cLA when compared to other fatty acids (Fazzari et al. 2014). Apart from the cLA isomers, the isomers of α -linolenic acid (Ln) containing

conjugated dienes and trienes are the second most prevalent species in humans (Salvatore et al. 2020). Conjugated linolenic acid (cLn) isomers are present at high levels in certain plant species and are the primary natural source of conjugated trienes (Hennessy et al. 2016), providing an interesting and abundant pool of nitration targets. Fatty acids with conjugated triene systems present in plants are mainly represented by (i) eleostearic acid (9-*cis*,11-*trans*,13-*trans*-octadecatrienoic) found in a tung plant *Aleurites fordii* (Takagi and Itabashi 1981) and chinese bitter gourd *Momordica charantia* (Liu et al. 1997); (ii) Calendic acid (*trans*-8,*trans*-10,*cis*-12-octadecatrienoic acid) found throughout the genus *Calendula* (Qiu et al. 2001) and (iii) Punicic acid (*cis*-9,*trans*-11,*cis*-13-octadecatrienoic acid) found in pomegranate seeds (*Punica granatum*) (Takagi and Itabashi 1981) and in *Trichosanthes*, *Mormodica balsamina*, *Fevillea trilobata* oil and *Ecballium elaterium* (Hennessy et al. 2016). Thus, the presence of cLn in different plant species makes them attractive candidates for searching and identifying new nitro-fatty acids in plant species.

Mechanisms of NO₂-FA action

Nitro-lipids are electrophilic molecules that participate in reversible Michael addition with proteins involved in cell signaling, metabolism, and transcription (Schopfer et al. 2019). The main intracellular targets are soft nucleophiles such as cysteine present in proteins and/or GSH, but proteomic approaches also revealed the formation of histidine adducts (Schopfer et al. 2011). In animals and humans, the mechanism of action of NO₂-FAs involves nitro-alkylation, a post-translational modification that regulates stability, activity, and localization of the target proteins (Rubbo and Radi 2008; Rudolph and Freeman 2009; Geisler and Rudolph 2012). Nitro-alkylation is a reversible process with the ability to regulate physiological processes by modulating the activity of transcription factors and enzymatic reactions. In plants, the first evidence of nitro-alkylation was reported in olive fruit, where NO₂-OA was found adducted to cysteine residues (Fazzari et al. 2014). The significance of nitro-alkylation reactions in plants was later established by the targeting of cytosolic Ascorbate peroxidase (APX) in *Arabidopsis* (Aranda-Caño et al. 2019).

Nitro-alkylation resembles the mode of action of reactive electrophile species (RES) produced by plant cells under different stresses (Farmer and Davoine 2007; Farmer and Mueller 2013). Interestingly, transcriptional and physiological responses are shared between RES and NO₂-FA. First, the RES-oxilipins OPDA (12-oxophytodienoic acid) (Muench et al. 2016) and electrophilic isothiocyanates (Øverby et al. 2015; Ferber et al. 2020) induced the activation of heat shock factors and transcription of several heat shock (HS)

proteins as described for NO₂-Ln (Mata-Pérez et al. 2016a). In addition, inhibition of primary root growth and cellular cycle was reported for RES-oxylipins, electrophilic isothiocyanates (Almeras et al. 2003; Mueller et al. 2008; Pauwels et al. 2008; Åsberg et al. 2015; Urbancsok et al. 2017, 2018) and NO₂-OA treatments in Arabidopsis (Di Fino et al. 2020). Considering the similarities between these reactive electrophilic species, future studies should focus on evaluating possible common signaling and/or mechanisms of action on plant physiology and divergent activities between these reactive species. More research is necessary to expand our understanding of target proteins and shed light on the signaling pathway in which NO₂-FA participate.

Another mechanism of action reported for NO₂-FAs relates to their ability to act as NO donors. This mechanism was first described in biochemical systems, cell culture, ex vivo and in vitro, and later translated to plant physiology (Schopfer, et al. 2005; Lima et al. 2003; Gorczynski et al. 2007; Mata-Pérez et al. 2016b, c; Grippo et al. 2020; Sánchez-Calvo et al. 2013). Nonetheless, the mechanism(s) whereby NO₂-FAs release NO have not been definitively established and remain controversial. A possible mechanism may include a modified Nef reaction, where, under neutral aqueous conditions, NO₂-FAs form a nitronate anion, that upon a sequence of protonation and deprotonation steps forms a nitroso intermediate characterized by a weak C–N bond that decomposes to release NO (Schopfer, et al. 2005). Using electron paramagnetic resonance (EPR) spectroscopy, NO₂-cLA, NO₂-LA and NO₂-OA have been shown to decompose and generate NO in aqueous solutions. By comparing the EPR signal intensities of spin-adducts, NO₂-cLA generated a significantly greater yield of NO than NO₂-LA and NO₂-OA (Grippo et al. 2020). In vitro, NO₂-LA releases less than 0.1% of NO in an aqueous solution. The NO release was inhibited in the presence of non-ionic detergent micelles as hydrophobic environments stabilize NO₂-LA (Schopfer et al. 2005). Other studies demonstrated that both NO₂-OA isomers yields, under optimal conditions, less than 0.1% of NO, with a 46 times lower release compared to NO donor SNP when using a 50 times higher concentration of NO₂-OA (Gorczynski et al. 2007). In an aqueous solution, NO₂-Ln releases NO with a peak within the first hour but 6 times lower than the rate reported for the NO donor GSNO (Mata-Pérez et al. 2016b; Ederli et al. 2009). The deprotonation of the bisallylic carbon is the first step in releasing NO; consequently, NO₂-Ln and NO₂-LA display higher NO release rates than NO₂-OA. Nonetheless, the rates and yields observed in biochemical systems remain very low. These results clearly show that nitro-alkenes have an objectively lower ability to release NO than SNP or GSNO donors. More importantly, NO₂-FA NO release reactions compete against in vivo alkylation reactions. The high intracellular thiol concentration, the rapid formation of GSH and protein

adducts, and the increased stability by incorporation into phospholipids and glycerolipids further reduces their ability to release NO (Schopfer et al. 2019). However, subcellular compartmentalization and/or enzymatic reactions could promote NO release under specific environments and cellular contexts.

NO formation was detected in vivo in Arabidopsis leaves and roots treated with 1 mM NO₂-Ln (Sánchez-Calvo et al. 2013), and in primary root treated with 100 μM of NO₂-Ln (Mata-Pérez et al. 2016c). Curiously, those NO₂-Ln treatments triggered different physiological responses than those triggered by NO donors (further discussed in the next section). A dual response was reported in Brassica seeds treated with NO₂-OA: at 50 μM, NO₂-OA did not release NO; however, at 100 μM, NO₂-OA NO was formed, while 500 μM NO₂-OA did not produce NO (Vollár et al. 2020). In contrast, we failed to detect NO release from Arabidopsis roots or tomato culture cells treated with NO₂-OA (Arruebarrena Di Palma et al. 2020; Di Fino et al. 2020). In aggregate, while all nitro-alkenes are electrophilic and react with cellular nucleophiles, some of them might release NO under specific conditions. The elucidation of the mechanism of action for NO₂-FA under physiological conditions will require considering NO release and Michael addition reactions and incorporate new approaches to dissect the individual contributions of these pathways to the biological response.

Differentiation from NO-dependent activities

In higher plants, NO is involved in a wide range of biological processes, including germination, plant metabolism, senescence, cellular death, stomatal movement, photosynthesis, gravitropism, and primary root growth (Del Castello et al. 2020; Lamattina et al. 2003). Some molecular targets of NO₂-FAs are chemically similar to those described for NO signaling, including redox-sensitive cysteine residues. This signaling crossover complicates the analysis of the responses obtained with NO donors and NO₂-FAs and the evaluation of the mechanism responsible for the biological actions.

NO participates in seed germination regulation by interacting with the ABA signaling during seed dormancy (Liu and Zhang 2009). For instance, pharmacological approaches demonstrated that most known NO donors promote dormancy-releasing and subsequent germination, while NO scavengers favour dormancy (Bethke et al. 2007). It was reported that in *Brassica napus*, NO₂-OA releases NO, which in turn increases germination rate (Vollár et al. 2020). However, this effect was only observed when seeds were treated with 100 μM NO₂-OA, with higher or lower concentrations failing to increase NO levels and affect germination.

Further work will be necessary to elucidate the mechanisms of action and the signaling pathway involved in this process.

NO upregulates APX activity through S-nitrosation of the conserved cysteine 32 residue (Correa-Aragunde et al. 2013; Begara-Morales et al. 2014; Yang et al. 2015). Conversely, in vitro NO₂-Ln nitro-alkylate APX's histidine residues inhibiting its enzymatic activity (Aranda-Caño et al. 2019). However, although NO₂-Ln has been proposed to act through NO release (Mata-Pérez et al. 2016b), S-nitrosation of the APX cysteine residue upon in vitro NO₂-Ln treatments has not been reported (Aranda-Caño et al. 2019). The overlapping activities as an NO donor and alkylation agent highlight the complication of establishing a mechanism of action and remind us to be cautious when considering NO₂-Ln exclusively as a NO donor.

NO inhibits primary root growth in *Arabidopsis thaliana* (Fernández-Marcos et al. 2011). Seedlings treated with NO donors show an imbalance in auxin levels (decrease in auxin response reporter DR5:GUS activity) due to NO-dependent alteration of the auxin efflux protein PIN-FORMED1 localization in the cellular membrane of meristematic cells (Fernández-Marcos et al. 2011). *Arabidopsis* seedlings treated with NO₂-OA showed primary root length inhibition without affecting endogenous NO levels and auxin homeostasis (Di Fino et al. 2020). In contrast, NO₂-OA reduced root growth by inhibiting the cell cycle in the meristem zone (Di Fino et al. 2020). This provides further evidence that NO and NO₂-FAs can induce the same morphological responses through independent signaling pathways.

RNAseq analysis of *Arabidopsis* cell suspensions treated with NO₂-Ln showed that several upregulated genes are related to the heat shock response (Mata-Pérez et al. 2016a). This response was first reported in NO₂-FAs-treated endothelial cells (Kansanen et al. 2009). To study whether NO₂-Ln modulates gene expression via exclusive NO production, we compared RNAseq data from NO₂-Ln (Mata-Pérez et al. 2016a) vs. NO treated *Arabidopsis* leaves (Hussain et al. 2016) (Fig. 2). Among the most representative shared genes, heat shock protein (HSPs) expression was strongly up-regulated while RNA transcription factors were down-regulated (Fig. 2, a complete set of data is shown in Supplemental Table S1, and a complete list of GO terms for biological processes in Supplemental Table S2). Despite this similarity, it is worth mentioning that only 122 genes are shared between both treatments and that the expression of 194 genes regulated by NO₂-Ln was unaffected by the NO treatment under the analyzed conditions. Altogether, the evidence presented above points towards a NO₂-FA-specific mechanism of action in different plant physiological systems.




Fig. 2 Overlap of genes whose expression is affected by treatments with NO₂-Ln and NO. Venn Diagram showing the overlap of differential expressed genes (p -value < 0.05 and $\log_2 \geq 1$ or ≤ -1) between treatment with NO₂-Ln (blue circle) and treatment with NO (yellow circle) was performed with MapMan Software. On the overlapping area are the principal groups of genes shared between the treatments. Those groups of genes were extracted from the BIN analysis performed with MapMan Software. On the unshared area genes related to stress, development and transcription whose expression is only affected in one of the two treatments. A complete list can be found in Supplemental Table 1. RNA seq data was obtained from: *Arabidopsis* cells treated with from 100 μ M of NO₂-Ln for 1 h (Mata-Pérez et al. 2016a) and *Arabidopsis* leaves infiltrated with 1 mM CysNO for 6 h (Hussain et al. 2016)

NO₂-OA as regulators of ROS and Ca²⁺

A recent report showed that exogenous application of NO₂-OA induced ROS production on both tomato cell suspensions and tomato and *Arabidopsis* leaves (Fig. 3, Arruebarrena Di Palma et al. 2020). Pharmacological and reverse genetic experiments showed that NO₂-OA-induced ROS were generated specifically by NADPH oxidase RBOH isoform D. The NADPH oxidase RBOHD is responsible for the rapid and robust production of ROS upon pathogen-associated molecular patterns perception (Kadota et al. 2015). In line with the activation of RBOHD, we tested the expression of genes related to pathogen responses observing up-regulation of *SIPR1a*, *SIHSR203J* and *SIPAL* (Arruebarrena Di Palma et al. 2020). Multilayered regulations tightly control RBOHD activity. It is mainly activated via direct binding of Ca²⁺ to EF-hand motifs, by Ca²⁺-dependent and independent protein kinases, and by the signal lipid phosphatidic acid (PA) (Kadota et al. 2015; Zhang et al. 2009). No Nitric Oxide was detected under these experimental conditions, suggesting that NO₂-FA-derived NO was not a primary contributor to the observed biological responses. Moreover, as RBOHD was reported to be negatively regulated




Fig. 3 Schematic representation of the physiological responses induced by NO_2 -FAs in Arabidopsis. Exogenous application of nitrooleic acid (NO_2 -OA) induces both reactive oxygen species (ROS) and cytosolic calcium levels in Arabidopsis leaf discs (Arruebarrena Di Palma et al. 2020). In addition, NO_2 -OA induces stomatal closure (Arruebarrena Di Palma et al. 2020), seed germination (Vollár et al. 2020), and cell cycle regulation (Di Fino et al. 2020). Another NO_2 -FA, nitro-linolenic acid (NO_2 -Ln) induces a specific set of genes that codify for heat shock proteins (HSP) and heat shock factors (HSF) in Arabidopsis cell cultures (Mata-Peréz et al. 2016a). The green box shows the biological process involved. Arrow indicates induction

by NO via S-nitrosation of Cys890 (Yun et al. 2011), a NO role is unlikely as we observed enzymatic activation upon treatment with NO_2 -OA (Arruebarrena Di Palma et al. 2020). In addition, RBOHD and F were required for NO_2 -OA-dependent induction of stomatal closure (Fig. 3, Arruebarrena Di Palma et al. 2020). However, the mechanism involved in the NADPH oxidase activation by NO_2 -OA remains unclear, and it could be hypothesized that NO_2 -FA form adducts directly with RBOHD protein regulating its activity. Nevertheless, we found that calcium and protein kinase activation are required to induce maximal ROS production upon NO_2 -OA treatments (Arruebarrena Di Palma et al. 2020). Thus, it might be possible that NO_2 -FA directly regulate protein kinases and/or Ca^{2+} channels and, therefore, Ca^{2+} -dependent protein kinases.

Cytosolic-free calcium elevation is an early signal in the developmental process and response to abiotic and biotic stresses in plants. Membrane-localized Ca^{2+} permeable ion channels modulate the influx of Ca^{2+} , leading to elevation in cytosolic Ca^{2+} concentrations (Kong et al. 2020). Results obtained in our lab using leaf discs that express the cytosolic calcium sensor aequorin showed that $[\text{Ca}^{+2}]_{\text{cyt}}$ increases

within 60–90 min in a dose-dependent manner between 5 and 25 μM NO_2 -OA treatments (unpublished result). The specific target and the mechanistic regulation of NO_2 -FA on calcium intracellular concentrations have not been explored, but cyst residues in target proteins might be key molecular targets. Recently, the H_2O_2 -induced Ca^{2+} increase 1 (HPCA1, a leucine-rich-repeat receptor kinase) was shown to act as an extracellular H_2O_2 sensor (Wu et al. 2020). Interestingly, HPCA1 contains four Cys residues acting as conserved redox-active sites. The molecular mechanism of H_2O_2 sensing involves the oxidation of these cysteines that induce a conformational change of the receptor. Nitrolipid targeting and regulation of HPCA1 might hold the key to understanding the relationship between Ca^{2+} and NO_2 -FA.

Redox regulators as targets of NO_2 -FAs?

Activation or deactivation of signaling pathways and enzymatic activities mediated by NO_2 -FA involves the reaction with cysteines of various cellular targets (Schopfer et al. 2019).

A well-described signaling pathway regulated by NO_2 -FAs in mammals is Nfr2/Keap1 (nuclear factor (erythroid-derived 2)-like 2)/kelch-like ECH-associated protein 1). Keap1 functions as a negative regulator of Nrf2 by keeping Nrf2 in the cytoplasmic compartment, promoting ubiquitination and downstream degradation by the proteasome. Under metabolic and inflammatory stress conditions, the formation of NO_2 -FA leads to the nitro-alkylation of Keap1 and destabilization of Nfr2-Keap1 complex. This inhibits Nrf2 proteasomal degradation and causes nuclear translocation and accumulation of newly synthesized Nrf2 protein, inducing the gene transcription of antioxidant and detoxification enzymes that protect against oxidative stress, inflammation, and drug toxicity (Magesh et al. 2012; Suzuki et al. 2013; Joshi et al. 2012; Kansanen et al. 2011). The central redox regulator NPR1 (non expressor of pathogenesis-related genes 1) is the plant counterpart of the mammalian Nfr2-Keap1 pathway, with which they share a common redox sensing mechanism (González-Bosch 2018). Both are activated by modifications (oxidation or alkylations) of their thiol groups that promote their nuclear translocation and activation of defense genes. In both pathways, proteasomal degradation prevents their activation and transcription activity (González-Bosch 2018). Arabidopsis NPR1 has 17 cysteines, with 10 of them being highly conserved among different species (Mou et al. 2003). Upon pathogen infection or accumulation of salicylic acid (SA), changes in cellular redox potential lead to the reduction of cysteines through the activity of thioredoxins (TRX-h3 and TRX-h5), promoting NPR1 translocation to the nucleus (Withers and Dong 2016). Unlike Nrf2, NPR1 does not contain a DNA-binding domain

but acts as a transcriptional coactivator by interacting with several transcriptional factors leading to the expression of pathogenesis-related (PR) proteins (Withers and Dong 2016). Indeed, Farmer and Mueller (Farmer and Mueller 2013) proposed NPR1 as a possible target of RES. Considering the similarities between both signaling mechanisms, it will be interesting to focus future research on NPR1 as a possible target of NO₂-FA action in plants.

Another relevant cellular target reported in animals is glutathione (GSH) (Schopfer et al. 2019). Through enzymatic and non-enzymatic reactions, GSH plays an active role in plant cell redox control systems. Glutathione protects cells against oxidative damage induced by environmental challenges (Hernández et al. 2017). The concentration of GSH in Arabidopsis leaf cells is between 0.05 mM in vacuole to near 15 mM in mitochondria (Gill et al. 2013), making it a likely target of NO₂-FA in plants. Exogenous treatment of tomato cell suspension with NO₂-OA results in the formation of GS-NO₂-OA adducts, with a concomitant reduction in GSH cellular concentration (Arruebarrena Di Palma et al. 2018). Also, it was reported that the release of ¹⁵N-labeled NO from exogenously applied ¹⁵NO₂-Ln (1 mM) induces the generation of GS¹⁵NO (Mata-Pérez et al. 2020). The authors indicated that the formation of NO from NO₂-Ln probably led to the modulation of the intracellular GSH pool, although the exact mechanism of GSNO formation from NO₂-Ln and GSH remains to be elucidated. Moreover, GSH was reported to interplay with different plant hormones like JA, MeJA, SA, and ethylene (Hasanuzzaman et al. 2017). Although GSH has multiple roles in plant physiology, the pathways or mechanisms involved in transducing GSH-induced responses have only been partially addressed (Hasanuzzaman et al. 2017). Given the concentration and reactivity of GSH with NO₂-FA, it would be important to establish the role that adduct formation plays in modulating plant signaling and physiology.

Final considerations

NO₂-FA has emerged as new molecules regulating physiological processes in plants. Different NO₂-FA were identified and detected in a variety of species. Our group has expanded the knowledge about this molecule on plant physiology, studying both their role in development (Di Fino et al. 2020) as well as signaling (Arruebarrena Di Palma et al. 2020) (Fig. 3). Depending on the nitrolipid, the exogenous application may exert a role as an electrophile (NO₂-OA), an NO donor (NO₂-Ln), or both. The identification of 342 protein adducted to NO₂-FA (working as an electrophile) has been reported, but unfortunately, their identities have not

been revealed so far (Aranda-Caño et al. 2019). The study of nitrolipid function in non-model plants and the identification of the cellular targets of NO₂-FA will shed light on its mechanism of action and the plant physiological responses that they might regulate.

Author contribution statement AML and LDF conceived the review. LDF wrote the initial draft. AADP wrote about the physiological role of NO₂-FA. EAP performed the RNAseq analysis. CGM, AML and FJS contributed to writing and manuscript editing.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00425-021-03777-z>.

Acknowledgements This work was supported by the UNMdP, Consejo Nacional de Investigaciones Científicas y Técnicas, Agencia Nacional de Promoción Científica y Tecnológica, ANPCyT; PICT-Raíces 2013-0800 and PICT 2017-0601 (to AML), PICT 2016-3084 (to AADP) and NIH GM125944 and DK112854 (to FJS).

Data availability All data generated or analyzed during this study are already published. Data sharing does not apply to this article as no datasets were generated during the current study. The comparison of the two different already published RNAseq is, however, presented as a supplemental table and available to the readers.

Declarations

Conflict of interest FJS acknowledges financial interest in Creegh Inc. No other competing financial interests are noted.

References

- Akaike T, Nishida M, Fujii S (2013) Regulation of redox signalling by an electrophilic cyclic nucleotide. *J Biochem* 153:131–138. <https://doi.org/10.1093/jb/mvs145>
- Almeras E, Stolz S, Vollenweider S, Reymond P, Mene-Saffrane L, Farmer EE (2003) Reactive electrophile species activate defense gene expression in Arabidopsis. *Plant J* 34:205–216. <https://doi.org/10.1046/j.1365-3113x.2003.01718.x>
- Aranda-Caño L, Sánchez-Calvo B, Begara-Morales JC, Chaki M, Mata-Pérez C, Padilla MN, Valderrama R, Barroso JB (2019) Post-translational modification of proteins mediated by nitro-fatty acids in plants: nitroalkylation. *Plants (basel)* 8(4):82. <https://doi.org/10.3390/plants8040082>
- Arruebarrena Di Palma A, Di Fino LM, Salvatore SR, D'Ambrosio JM, Grozoff GEG, García-Mata C, Schopfer F, Laxalt AM (2018) Nitro-oleic acid induced reactive oxygen species formation and plant defense signaling in tomato cell suspensions. *BioRxiv* 2:e263
- Arruebarrena Di Palma A, Di Fino LM, Salvatore S, R, D'Ambrosio, J. M, García-Mata, C, Schopfer, F. J, Laxalt AM (2020) Nitro-oleic acid triggers ROS production via NADPH oxidase activation in plants: A pharmacological approach. *J. Plant Physiol.* 246-247:153128. <https://doi.org/10.1016/j.jplph.2020.153128>

- Åsberg SE, Bones AM, Øverby A (2015) Allyl isothiocyanate affects the cell cycle of *Arabidopsis thaliana*. *Front Plant Sci* 6:364. <https://doi.org/10.3389/fpls.2015.00364>
- Baker PR, Schopfer FJ, Sweeney S, Freeman BA (2004) Red cell membrane and plasma linoleic acid nitration products: synthesis, clinical identification, and quantitation. *Proc Natl Acad Sci* 101(32):11577–11582. <https://doi.org/10.1073/pnas.0402587101>
- Balazy M, López-Fernández J (2003) Isomerization and nitro-oxidation of arachidonic acid by NO₂. *Adv Exp Med Biol* 525:173–176. https://doi.org/10.1007/978-1-4419-9194-2_36
- Begara-Morales JC, Sánchez-Calvo B, Chaki M, Valderrama R, Mata-Pérez C, López-Jaramillo J, Padilla MN, Carreras A, Corpas FJ, Barroso JB (2014) Dual regulation of cytosolic ascorbate peroxidase (APX) by tyrosine nitration and S-nitrosylation. *J Exp Bot* 65(2):527–538. <https://doi.org/10.1093/jxb/ert396>
- Bethke PC, Libourel IG, Aoyama N, Chung YY, Still DW, Jones RL (2007) The *Arabidopsis* aleurone layer responds to nitric oxide, gibberellin, and abscisic acid and is sufficient and necessary for seed dormancy. *Plant Physiol* 143(3):1173–1188. <https://doi.org/10.1104/pp.106.093435>
- Bonacci G, Baker PR, Salvatore SR, Shores D, Khoo NK, Koenitzer JR, Vitturi DA, Woodcock SR, Golin-Bisello F, Cole MP, Watkins S, St Croix C, Batthyany CI, Freeman BA, Schopfer FJ (2012) Conjugated linoleic acid is a preferential substrate for fatty acid nitration. *J Biol Chem* 287(53):44071–44082. <https://doi.org/10.1074/jbc.M112.401356>
- Correa-Aragunde N, Foresi N, Delledonne M, Lamattina L (2013) Auxin induces redox regulation of ascorbate peroxidase I activity by S-nitrosylation/denitrosylation balance resulting in changes of root growth pattern in *Arabidopsis*. *J Exp Bot* 64(11):3339–3349. <https://doi.org/10.1093/jxb/ert172>
- Del Castello F, Nejamkin A, Foresi N, Lamattina L, Correa-Aragunde N (2020) Chimera of globin/nitric oxide synthase: toward improving nitric oxide homeostasis and nitrogen recycling and availability. *Front Plant Sci* 11:575651. <https://doi.org/10.3389/fpls.2020.575651>
- Delmastro-Greenwood M, Hughan KS, Vitturi DA, Salvatore SR, Grimes G, Potti G, Shiva S, Schopfer FJ, Gladwin MT, Freeman BA, Gelhaus Wendell S (2015) Nitrite and nitrate-dependent generation of anti-inflammatory fatty acid nitroalkenes. *Free Radical Biol Med* 89:333–341. <https://doi.org/10.1016/j.freeradbiomed.2015.07.149>
- Di Fino LM, cerrudo I, salvatore sr, schopfer fj, garcía-mata c, laxalt am (2020) exogenous nitro-oleic acid treatment inhibits primary root growth by reducing the mitosis in the meristem in *Arabidopsis thaliana*. *Front Plant Sci* 11:1059. <https://doi.org/10.3389/fpls.2020.01059>
- Ederli L, Reale L, Madeo L, Ferranti F, Gehring C, Fornaciari M, Romano B, Pasqualini S (2009x) NO release by nitric oxide donors in vitro and in planta. *Plant Physiol Biochem* 47(1):42–48. <https://doi.org/10.1016/j.plaphy.2008.09.008>
- Farmer EE, Davoine C (2007) Reactive electrophile species. *Curr Opin Plant Biol* 10(4):380–386. <https://doi.org/10.1016/j.pbi.2007.04.019>
- Farmer EE, Mueller MJ (2013) ROS-mediated lipid peroxidation and RES-activated signaling. *Ann Rev Plant Biol* 64(1):429–450. <https://doi.org/10.1146/annurev-arplant-050312-120132>
- Fazzari M, Trostchansky A, Schopfer FJ, Salvatore SR, Sánchez-Calvo B, Vitturi D, Valderrama R, Barroso JB, Radi R, Freeman BA, Rubbo H (2014) Olives and olive oil are sources of electrophilic fatty acid nitroalkenes. *PLoS ONE* 9(1):e84884. <https://doi.org/10.1371/journal.pone.0084884>
- Fazzari M, Khoo NK, Woodcock SR, Jorkasky DK, Li L, Schopfer FJ, Freeman BA (2017) Nitro-fatty acid pharmacokinetics in the adipose tissue compartment. *J Lipid Res* 58(2):375–385. <https://doi.org/10.1194/jlr.M072058>
- Ferber E, Gerhards J, Sauer M, Krischke M, Dittrich MT, Müller T, Berger S, Agnes F, Mueller MJ (2020) Chemical priming by isothiocyanates protects against intoxication by products of the mustard oil bomb. *Front Plant Sci* 11:887. <https://doi.org/10.3389/fpls.2020.00887>
- Fernández-Marcos M, Sanz L, Lewis DR, Muday GK, Lorenzo O (2011) Nitric oxide causes root apical meristem defects and growth inhibition while reducing PIN-FORMED 1 (PIN1)-dependent acropetal auxin transport. *Proc Natl Acad Sci USA* 108(45):18506–18511. <https://doi.org/10.1073/pnas.1108644108>
- Gallon AA, Pryor WA (1993) The identification of the allylic nitrite and nitro derivatives of methyl linoleate and methyl linolenate by negative chemical ionization mass spectroscopy. *Lipids* 28(2):125–133. <https://doi.org/10.1007/BF02535776>
- Geisler AC, Rudolph TK (2012) Nitroalkylation a redox sensitive signaling pathway. *Biochem Biophys Acta* 1820(6):777–784. <https://doi.org/10.1016/j.bbagen.2011.06.014>
- Gill SS, Anjum NA, Hasanuzzaman M, Gill R, Trivedi DK, Ahmad I, Pereira E, Tuteja N (2013) Glutathione and glutathione reductase: a boon in disguise for plant abiotic stress defense operations. *Plant Physiol Biochem* 70:204–212. <https://doi.org/10.1016/j.plaphy.2013.05.032>
- González-Bosch C (2018) Priming plant resistance by activation of redox-sensitive genes. *Free Radical Biol Med* 122:171–180. <https://doi.org/10.1016/j.freeradbiomed.2017.12.028>
- Gorczyński MJ, Huang J, Lee H, King SB (2007) Evaluation of nitroalkenes as nitric oxide donors. *Bioorg Med Chem Lett* 17(7):2013–2017. <https://doi.org/10.1016/j.bmcl.2007.01.016>
- Grippio V, Mojovic M, Pavicevic A, Kabelac M, Hubatka F, Turanek J, Zatloukalova M, Freeman BA, Vacek J (2020) Electrophilic characteristics and aqueous behavior of fatty acid nitroalkenes. *Redox Biol* 38:101756. <https://doi.org/10.1016/j.redox.2020.101756>
- Hasanuzzaman M, Nahar K, Anee TI, Fujita M (2017) Glutathione in plants: biosynthesis and physiological role in environmental stress tolerance. *Physiol Mol Biol Plants Int J Funct Plant Biol* 23(2):249–268. <https://doi.org/10.1007/s12298-017-0422-2>
- Hennessy AA, Ross PR, Fitzgerald GF, Stanton C (2016) Sources and bioactive properties of conjugated dietary fatty acids. *Lipids* 51(4):377–397. <https://doi.org/10.1007/s11745-016-4135-z>
- Hernández JA, Barba-Espín G, Diaz-Vivancos P (2017) Glutathione-mediated biotic stress tolerance in plants. In: Hossain M, Mostofa M, Diaz-Vivancos P, Burritt D, Fujita M, Tran LS (eds) *Glutathione in plant growth, development, and stress tolerance*. Springer, Cham
- Hussain A, Mun BG, Imran QM, Lee SU, Adamu TA, Shahid M, Kim KM, Yun BW (2016) Nitric Oxide Mediated Transcriptome Profiling Reveals Activation of Multiple Regulatory Pathways in *Arabidopsis thaliana*. *Front Plant Sci* 7:975
- Joshi G, Johnson JA (2012) The Nrf2-ARE pathway: a valuable therapeutic target for the treatment of neurodegenerative diseases. *Recent Pat CNS Drug Dis* 7(3):218–229. <https://doi.org/10.2174/157488912803252023>
- Kadota Y, Shirasu K, Zipfel C (2015) Regulation of the NADPH oxidase RBOHD during plant immunity. *Plant Cell Physiol* 56(8):1472–1480. <https://doi.org/10.1093/pcp/pcv063>
- Kansanen E, Jyrkkänen HK, Volger OL, Leinonen H, Kiveliä AM, Häkkinen SK, Woodcock SR, Schopfer FJ, Horrevoets AJ, Ylä-Herttuala S, Freeman BA, Levonen AL (2009) Nrf2-dependent and -independent responses to nitro-fatty acids in human endothelial cells: identification of heat shock response as the major pathway activated by nitro-oleic acid. *J Biol Chem* 284(48):33233–33241. <https://doi.org/10.1074/jbc.M109.064873>

- Kansanen E, Bonacci G, Schopfer FJ, Kuosmanen SM, Tong KI, Leinonen H, Woodcock SR, Yamamoto M, Carlberg C, YläHerttua S, Freeman BA, Levonen AL (2011) Electrophilic nitro-fatty acids activate NRF2 by a KEAP1 cysteine 151-independent mechanism. *J Biol Chem* 286(16):14019–14027. <https://doi.org/10.1074/jbc.M110.190710>
- Kong X, Xu L, Jamieson P (2020) Plant sense: the rise of calcium channels. *Trends Plant Sci* 25(9):838–841. <https://doi.org/10.1016/j.tplants.2020.06.002>
- Lamattina L, García-Mata C, Graziano M, Pagnussat G (2003) Nitric oxide: the versatility of an extensive signal molecule. *Annu Rev Plant Biol* 54:109–136. <https://doi.org/10.1146/annurev.arplant.54.031902.134752>
- Lima ES, Di Mascio P, Rubbo H, Abdalla DS (2002) Characterization of linoleic acid nitration in human blood plasma by mass spectrometry. *Biochemistry* 41(34):10717–10722. <https://doi.org/10.1021/bi025504j>
- Lima ES, Di Mascio P, Abdalla DS (2003) Cholesteryl nitrolinoleate, a nitrated lipid present in human blood plasma and lipoproteins. *J Lipid Res* 44(9):1660–1666. <https://doi.org/10.1194/jlr.M200467-JLR200>
- Liu Y, Zhang J (2009) Rapid accumulation of NO regulates ABA catabolism and seed dormancy during imbibition in Arabidopsis. *Plant Signal Behav* 4(9):905–907. <https://doi.org/10.4161/psb.4.9.9532>
- Liu L, Hammond EG, Nikolau BJ (1997) In vivo studies of the biosynthesis of [alpha]-eleostearic acid in the seed of momordica charantia L. *Plant Physiol* 113(4):1343–1349. <https://doi.org/10.1104/pp.113.4.1343>
- Magesh S, Chen Y, Hu L (2012) Small molecule modulators of Keap1-Nrf2-ARE pathway as potential preventive and therapeutic agents. *Med Res Rev* 32(4):687–726. <https://doi.org/10.1002/med.21257>
- Mata-Pérez C, Sánchez-Calvo B, Padilla MN, Begara-Morales JC, Luque F, Melguizo M, Jimenez-ruiz J, Fierro-Risco J, Penas-Sanjuan A, Valderrama R, Corpas FJ, Barroso JB (2016a) Nitro-fatty acids in plant signaling: nitro-linolenic acid induces the molecular chaperone network in Arabidopsis. *Plant Physiol* 170:686–701. <https://doi.org/10.1104/pp.15.01671>
- Mata-Pérez C, Sánchez-Calvo B, Begara-Morales JC, Carreras A, Padilla MN, Melguizo M, Valderrama R, Corpas FJ, Barroso JB (2016b) Nitro-linolenic acid is a nitric oxide donor. *Nitric Oxide Biol Chem* 57:57–63. <https://doi.org/10.1016/j.niox.2016.05.003>
- Mata-Pérez C, Sánchez-Calvo B, Begara-Morales JC, Padilla MN, Valderrama R, Corpas FJ, Barroso JB (2016c) Nitric oxide release from nitro-fatty acids in Arabidopsis roots. *Plant Signal Behav* 11(3):e1154255. <https://doi.org/10.1080/15592324.2016.1154255>
- Mata-Pérez C, Sánchez-Calvo B, Padilla MN, Begara-Morales JC, Valderrama R, Corpas FJ, Barroso JB (2017) Nitro-fatty acids in plant signaling: new key mediators of nitric oxide metabolism. *Redox Biol* 11:554–561. <https://doi.org/10.1016/j.redox.2017.01.002>
- Mata-Pérez C, Padilla MN, Sánchez-Calvo B, Begara-Morales JC, Valderrama R, Corpas FJ, Barroso JB (2018) Nitro-fatty acid detection in plants by high-pressure liquid chromatography coupled to triple quadrupole mass spectrometry. In: Mengel A, Lindermayr C (eds) Nitric oxide. Methods in molecular biology, vol 1747. Humana Press, New York
- Mata-Pérez C, Padilla MN, Sánchez-Calvo B, Begara-Morales JC, Valderrama R, Chaki M, Aranda-Caño L, Moreno-González D, Molina-Díaz A, Barroso JB (2020) Endogenous biosynthesis of s-nitrosoglutathione from nitro-fatty acids in plants. *Front Plant Sci* 11:962. <https://doi.org/10.3389/fpls.2020.00962>
- Mou Z, Fan W, Dong X (2003) Inducers of plant systemic acquired resistance regulate NPR1 function through redox changes. *Cell* 113(7):935–944. [https://doi.org/10.1016/s0092-8674\(03\)00429-x](https://doi.org/10.1016/s0092-8674(03)00429-x)
- Mueller S, Hilbert B, Dueckershoff K, Roitsch T, Krischke M, Mueller MJ, Berger S (2008) General detoxification and stress responses are mediated by oxidized lipids through TGA transcription factors in Arabidopsis. *Plant Cell* 20(3):768–785. <https://doi.org/10.1105/tpc.107.054809>
- Muench M, Hsin CH, Ferber E, Berger S, Mueller MJ (2016) Reactive electrophilic oxylipins trigger a heat stress-like response through HSF1 transcription factors. *J Exp Bot* 67(21):6139–6148. <https://doi.org/10.1093/jxb/erw376>
- O'Donnell VB, Eiserich JP, Chumley PH, Jablonsky MJ, Krishna NR, Kirk M, Barnes S, Darley-Usmar VM, Freeman BA (1999) Nitration of unsaturated fatty acids by nitric oxide-derived reactive nitrogen species peroxynitrite, nitrous acid, nitrogen dioxide, and nitronium ion. *Chem Res Toxicol* 12:83–92
- Øverby A, Bævre MS, Thangstad OP, Bones AM (2015) Disintegration of microtubules in Arabidopsis thaliana and bladder cancer cells by isothiocyanates. *Front Plant Sci* 6:6. <https://doi.org/10.3389/fpls.2015.00006>
- Pauwels L, Morreel K, De Witte E, Lammertyn F, Van Montagu M, Boerjan W, Inzé D, Goossens A (2008) Mapping methyl jasmonate-mediated transcriptional reprogramming of metabolism and cell cycle progression in cultured Arabidopsis cells. *Proc Natl Acad Sci* 105(4):1380–1385. <https://doi.org/10.1073/pnas.0711203105>
- Pryor WA, Lightsey JW (1981) Mechanisms of nitrogen dioxide reactions: initiation of lipid peroxidation and the production of nitrous Acid. *Science* 214(4519):435–437. <https://doi.org/10.1126/science.214.4519.435>
- Qiu X, Reed DW, Hong H, MacKenzie SL, Covello PS (2001) Identification and analysis of a gene from Calendula officinalis encoding a fatty acid conjugase. *Plant Physiol* 125(2):847–855. <https://doi.org/10.1104/pp.125.2.847>
- Rubbo H, Radi R (2008) Protein and lipid nitration: role in redox signaling and injury. *Biochim Biophys Acta* 1780:1318–1324. <https://doi.org/10.1016/j.bbagen.2008.03.007>
- Rubbo H, Radi R, Trujillo M, Telleri R, Kalyanaraman B, Barnes S, Kirk M, Freeman BA (1994) Nitric oxide regulation of superoxide and peroxynitrite-dependent lipid peroxidation. Formation of novel nitrogen-containing oxidized lipid derivatives. *J Biol Chem* 269:26066–26075
- Rudolph V, Freeman BA (2009) Cardiovascular consequences when nitric oxide and lipid signaling converge. *Circ Res* 105(6):511–522. <https://doi.org/10.1161/CIRCRESAHA.109.202077>
- Rudolph V, Schopfer FJ, Khoo NK, Rudolph TK, Cole MP, Woodcock SR, Bonacci G, Groeger AL, Golin-Bisello F, Chen CS, Baker PR, Freeman BA (2009) Nitro-fatty acid metabolome: saturation, desaturation, beta-oxidation, and protein adduction. *J Biol Chem* 284(3):1461–1473. <https://doi.org/10.1074/jbc.M802298200>
- Salvatore SR, Rowart P, Schopfer FJ (2020) Mass spectrometry-based study defines the human urine nitrolipidome. *Free Rad Biol Med*. <https://doi.org/10.1016/j.freeradbiomed.2020.10.305>
- Sánchez-Calvo B, Barroso JB, Corpas FJ (2013) Hypothesis: Nitro-fatty acids play a role in plant metabolism. *Plant Sci Int J Exp Plant Biol* 199–200:1–6. <https://doi.org/10.1016/j.plantsci.2012.10.006>
- Schopfer FJ, Khoo N (2019) Nitro-fatty acid logistics: formation, bio-distribution, signaling, and pharmacology. *Trends Endocrinol Metab* 30(8):505–519. <https://doi.org/10.1016/j.tem.2019.04.009>
- Schopfer FJ, Baker PR, Giles G, Chumley P, Batthyany C, Crawford J, Patel RP, Hogg N, Branchaud BP, Lancaster JR, Freeman BA (2005) Fatty acid transduction of nitric oxide signaling. Nitrolinoleic acid is a hydrophobically stabilized nitric oxide donor. *J Biol Chem* 280(19):19289–19297. <https://doi.org/10.1074/jbc.M414689200>
- Schopfer FJ, Cipollina C, Freeman BA (2011) Formation and signaling actions of electrophilic lipids. *Chem Rev* 111(10):5997–6021. <https://doi.org/10.1021/cr200131e>

- Suzuki T, Motohashi H, Yamamoto M (2013) Toward clinical application of the Keap1-Nrf2 pathway. *Trends Pharmacol Sci* 34(6):340–346. <https://doi.org/10.1016/j.tips.2013.04.005>
- Takagi T, Itabashi Y (1981) Occurrence of mixtures of geometrical isomers of conjugated octadecatrienoic acids in some seed oils: analysis by open-tubular gas liquid chromatography and high performance liquid chromatography. *Lipids* 16(7):546–551
- Trostchansky A, Rubbo H (2008) Nitrated fatty acids: mechanisms of formation, chemical characterization, and biological properties. *Free Radic Biol Med* 44:1887–1896. <https://doi.org/10.1016/j.freeradbiomed.2008.03.006>
- Turell L, Radi R, Alvarez B (2013) The thiol pool in human plasma: the central contribution of albumin to redox processes. *Free Radical Biol Med* 65:244–253. <https://doi.org/10.1016/j.freeradbiomed.2013.05.050>
- Turell L, Vitturi DA, Coitiño EL, Lebrato L, Möller MN, Sagasti C, Salvatore SR, Woodcock SR, Alvarez B, Schopfer FJ (2017) The chemical basis of thiol addition to nitro-conjugated linoleic acid, a protective cell-signaling lipid. *J Biol Chem* 292(4):1145–1159. <https://doi.org/10.1074/jbc.M116.756288>
- Urbancsok J, Bones AM, Kissen R (2017) Glucosinolate-derived isothiocyanates inhibit Arabidopsis growth and the potency depends on their side chain structure. *Int J Mol Sci* 18(11):2372. <https://doi.org/10.3390/ijms18112372>
- Urbancsok J, Bones AM, Kissen R (2018) Arabidopsis mutants impaired in glutathione biosynthesis exhibit higher sensitivity towards the glucosinolate hydrolysis product allyl-isothiocyanate. *Sci Rep* 8(1):1–13. <https://doi.org/10.1038/s41598-018-28099-1>
- van der Vliet A, Eiserich JP, Halliwell B, Cross CE (1997) Formation of reactive nitrogen species during peroxidase-catalyzed oxidation of nitrite. A potential additional mechanism of nitric oxide-dependent toxicity. *J Biol Chem* 272(12):7617–7625. <https://doi.org/10.1074/jbc.272.12.7617>
- Villacorta L, Minarrieta L, Salvatore SR, Khoo NK, Rom O, Gao Z, Berman RC, Jobbagy S, Li L, Woodcock SR, Chen YE, Freeman BA, Ferreira AM, Schopfer FJ, Vitturi D (2018) In situ generation, metabolism and immunomodulatory signaling actions of nitro-conjugated linoleic acid in a murine model of inflammation. *Redox Biol* 15:522–531. <https://doi.org/10.1016/j.redox.2018.01.005>
- Vitturi DA, Chen CS, Woodcock SR, Salvatore SR, Bonacci G, Koentzer JR, Stewart NA, Wakabayashi N, Kensler TW, Freeman BA, Schopfer FJ (2013) Modulation of nitro-fatty acid signaling: prostaglandin reductase-1 is a nitroalkene reductase. *J Biol Chem* 288(35):25626–25637. <https://doi.org/10.1074/jbc.M113.486282>
- Vitturi DA, Minarrieta L, Salvatore SR, Postlethwait EM, Fazzari M, Ferrer-Sueta G, Lancaster JR, Freeman BA, Schopfer FJ (2015) Convergence of biological nitration and nitrosation via symmetrical nitrous anhydride. *Nat Chem Biol* 11(7):504–510. <https://doi.org/10.1038/nchembio.1814>
- Vollár M, Feigl G, Oláh D, Horváth A, Molnár Á, Kúsz N, Ördög A, Csopor D, Kolbert Z (2020) Nitro-oleic acid in seeds and differently developed seedlings of *Brassica napus* L. *Plants* (basel, Switzerland) 9(3):406. <https://doi.org/10.3390/plants9030406>
- Withers J, Dong X (2016) Posttranslational modifications of NPR1: a single protein playing multiple roles in plant immunity and physiology. *PLoS Pathog* 12(8):e1005707. <https://doi.org/10.1371/journal.ppat.1005707>
- Wu F, Chi Y, Jiang Z et al (2020) Hydrogen peroxide sensor HPCA1 is an LRR receptor kinase in Arabidopsis. *Nature* 578(7796):577–581. <https://doi.org/10.1038/s41586-020-2032-3>
- Yang H, Mu J, Chen L, Feng J, Hu J, Li L, Zhou JM, Zuo J (2015) S-nitrosylation positively regulates ascorbate peroxidase activity during plant stress responses. *Plant Physiol* 167(4):1604–1615. <https://doi.org/10.1104/pp.114.255216>
- Yun BW, Feechan A, Yin M, Saidi NB, Le Bihan T, Yu M, Moore JW, Kang JG, Kwon E, Spoel SH, Pallas JA, Loake GJ (2011) S-nitrosylation of NADPH oxidase regulates cell death in plant immunity. *Nature* 478:264–268
- Zhang H, Fang Q, Zhang Z, Wang Y, Zheng X (2009) The role of respiratory burst oxidase homologues in elicitor-induced stomatal closure and hypersensitive response in *Nicotiana benthamiana*. *J Exp Bot* 60(11):3109–3122. <https://doi.org/10.1093/jxb/erp146>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.