

Independent risk factors for recurrence of apparent life-threatening events in infants

Riyo Ueda¹ · Osamu Nomura² · Takanobu Maekawa¹ · Hirokazu Sakai¹ · Satoshi Nakagawa^{2,3} · Akira Ishiguro^{1,3}

Received: 21 July 2016 / Revised: 28 December 2016 / Accepted: 9 January 2017 / Published online: 20 January 2017
© Springer-Verlag Berlin Heidelberg 2017

Abstract The risk factors for recurrent apparent life-threatening event (ALTE) are unclear although the risk of recurrent ALTE is an important consideration for the management of ALTE patients. This study aimed to identify the risk factors for recurrent ALTE. We conducted a secondary analysis of the data from a single center retrospective cohort study in Japan conducted from March 2002 to January 2012, which included children diagnosed with ALTE at a pediatric emergency department (ED) in Tokyo. Among 112 ALTE patients, 18 (16%) had recurrences and 94 (84%) did not. Symptoms of respiratory tract infection (RTI) were more frequent in the recurrent group than in the non-recurrent group (44 vs. 14% $p = 0.0055$), and the proportion of patients triaged as level 1 was larger in the recurrent group than in the non-recurrent group (31 vs. 7%, $p = 0.0312$). Pallor was observed more frequently in the recurrent group than in the non-recurrent group (100 vs. 76%, $p = 0.0216$). Multivariate analysis dem-

onstrated that the independent risk factors of recurrent ALTE were respiratory tract infection symptoms (OR, 5.02; 95% CI, 1.48–16.98).

Conclusion: ALTE patients who had RTI symptoms at the ED visit for first ALTE should be admitted for close observation of potential recurrences.

What is Known:

- Approximately 10% of ALTE patients experienced recurrence of ALTE episodes.
- The risk of recurrent ALTE is one of the major consideration for the management of ALTE patients at the ED because these patients have higher rates of serious underlying diseases which require interventions.

What is New:

- Respiratory infection symptoms at ED presentation can be independent risk factors for recurrent ALTE.
-

Communicated by Mario Bianchetti

Electronic supplementary material The online version of this article (doi:10.1007/s00431-017-2855-6) contains supplementary material, which is available to authorized users.

✉ Osamu Nomura
osamunomura@hotmail.com

Riyo Ueda
uedariyo@ncnp.go.jp

Takanobu Maekawa
makawa-t@ncchd.go.jp

Hirokazu Sakai
sakai@saitama-med.ac.jp

Satoshi Nakagawa
nakagawa-s@ncchd.go.jp

Akira Ishiguro
ishiguro-a@ncchd.go.jp

¹ Department of General Pediatrics and Interdisciplinary Medicine, National Center for Child Health and Development, Tokyo, Japan

² Department of Critical Care and Anesthesia, National Center for Child Health and Development, Tokyo, Japan

³ Department of Postgraduate Education and Training, National Center for Child Health and Development, Tokyo, Japan

Keywords Infantile apparent life-threatening event · Recurrence · Risk factors

Abbreviations

| | |
|------|----------------------------------|
| ALTE | Apparent life-threatening event |
| BRUE | Brief resolved unexplained event |
| ED | Emergency department |
| PICU | Pediatric intensive care unit |
| RTI | Respiratory tract infection |

Introduction

An apparent life-threatening event (ALTE) was defined as “an episode that is frightening to the observer and that is characterized by some combination of apnea (central or occasionally obstructive), color change (usually cyanotic or pallid but occasionally erythematous or plethoric), marked change in muscle tone (usually marked limpness), choking, or gagging” at a consensus development conference convened by the National Institutes of Health in 1986 [1]. In May 2016, the American Academy of Pediatrics (AAP) published a new practice guideline for ALTE [13]. The guideline recommended the replacement of the term ALTE with “brief resolved unexplained event (BRUE)” because most of the ALTE patients had favorable outcomes in spite of its “life-threatening” nature. However, there are differences between the core concepts of managing ALTE and BRUE, and the clinical definition of BURE has yet to be widely accepted [13].

In recent years, the etiology and pathophysiology of ALTE are becoming established. Evidence has shown that most of the patients with ALTE had good outcome [4]. Therefore, researchers proposed that “low risk” ALTE patients who visited the emergency department (ED) could be managed without hospitalization [8]. For clinicians, the risk of recurrent ALTE is one of the major consideration for the management of ALTE patients at the ED because these patients have higher rates of serious underlying diseases which require interventions [10]. Several studies have shown that 10% of ALTE patients experienced recurrence of ALTE episodes [4, 7, 12], and one study suggested that patients with recurrent ALTE should be admitted for close observation and further investigation for underlying disease [7].

To date, the risk factors for recurrent ALTE have not been fully investigated. The aim of this study was to identify these risk factors.

Subjects and methods

Design

This was a secondary analysis of data from a single center retrospective cohort study in Japan conducted from

March 2002 to January 2012 and published in Japanese [14]. We included children diagnosed with ALTE at our ED during the study period but excluded children older than 1 year. The definition of ALTE was based on the National Institutes of Health Consensus Development Conference Statement [1]. We defined recurrent ALTE here as an episode sharing the same clinical features as the first episode of ALTE experienced by the patient. In particular, in-hospital recurrent ALTE was also defined as a clinical state that needed resuscitation to stabilize compromised respiratory and/or circulation, convulsions, and bradycardia due to hypoxia during hospitalization.

Chart review method

The chart review was conducted using a structured method [6]. We trained the designated abstractor (UR) in the abstraction method before data abstraction, constructing the abstraction form, and defining the variables through discussions with the investigators (UR, ON, and TM). After data abstraction was begun, we periodically held meetings (UR, ON and TM) to examine the abstracted patient data randomly.

Study setting

The National Center for Child Health and Development is a tertiary care pediatric and perinatal hospital in Tokyo, Japan. Each year, about 35,000 children visit the emergency department of the hospital. The hospital has 320 beds, including 40 neonatal intensive care unit beds and 20 pediatric intensive care unit (PICU) beds.

Data collection

Patients characteristics

We investigated the age, gender, birth history, underlying diseases, and symptoms of patients observed by their caregiver at the time of ALTE. Those symptoms were categorized into “pallor,” “hypotonia,” “breathing difficulties,” “blue/cyanotic,” “abnormal eye movement,” “excessive mucus production,” “red, hypertonic/overdistention/crying,” and “vomiting and jerkin limbs,” according to Semmekrot et al. [11]. The presence of respiratory tract infection (RTI) symptoms such as cough and nasal discharge was evaluated by physicians at the ED. The triage level of the patients, PICU admission, and definitive diagnosis of the patients at discharge were also investigated. For the determination of the patients’ triage level, trained nurses triaged all the patients who visited the ED according to the Canadian Pediatric Triage and Acuity Scale (P-CTAS) [15]. In general, triage level 1 is assigned to patients who present with severely unstable vital signs and to those who require immediate interventions to be stabilized.

Recurrence of ALTE and prognosis

We reviewed the incidence of recurrent ALTE by reviewing patient charts for the 6 months after the first incident of ALTE. The prognosis of the patients was also studied. Patients were identified as alive without recurrent ALTE if they had follow-up outpatient visit in the 6 months after the first incident and if there was no record of recurrent ALTE. Patients who did not come to the hospital for a follow-up visit during that time were deemed as unknown outcome.

Eventually, we compared the characteristics of the patients who experienced recurrent ALTE with those who did not.

Analysis

Data were analyzed using JMP11 (SAS Institute Inc., Cary, NC). In univariate analysis, continuous variables were compared by the Mann-Whitney *U* test and categorical variables by the Fisher's exact test. The significance level was set at $p < 0.05$. Logistic regression was used to determine the independent risk factors for recurrent ALTE. We selected candidate variables for analysis based on a review of the literature. From among these, we then identified variables for logistic regression analysis by calculating Akaike's Information Criterion (AIC) by forward selection.

Results

Patient characteristics

A total of 112 patients diagnosed with ALTE were identified during the study period (Table 1). Fifty-five (49%) patients were male, and there were nine (8%) patients whose gestational age was less than 36 weeks. Sixteen (14%) patients had underlying disease and 62 (55%) patients were less than 2 months old at the time of ALTE. At the ED presentation,

21 (19%) patients had RTI symptoms. Eleven (10%) patients were triaged as level 1 because of extremely unstable vital signs due to respiratory failure (six patients), compromised cardiopulmonary state (three patients), disturbed consciousness (one patient), and hypothermia (one patient). Among 11 patients assigned to triage level 1, two patients had history of premature birth (gestational age < 36 weeks) and eight patients were younger than 2 months of age at presentation. Five patients were intubated due to respiratory failure but no patients needed cardiopulmonary resuscitation. RTI was definitively diagnosed in five patients, gastroesophageal reflux in two patients, epilepsy in one patient, choking in one patient, and hypothermia in one patient. The diagnosis was indeterminate in one patient. Of 102 (92%) hospitalized patients, 15 (13%) were admitted to the PICU. Among the 15 patients who were admitted to the PICU, two patients had history of premature birth and nine patients were younger than 2 months of age at presentation. Six patients were intubated due to respiratory failure but no patients needed cardiopulmonary resuscitation. RTI was definitively diagnosed in seven patients, idiopathic apnea in three patients, gastroesophageal reflux in one patient, and choking in one patient. The diagnosis was indeterminate in three patients.

Recurrence of ALTEs


Eighteen (16%) patients experienced recurrent ALTE (Fig. 1). Among them, 15 experienced recurrence during hospitalization and three did so after discharge following hospitalization. Among the 15 patients who had recurrent ALTE during hospitalization, we confirmed that ten patients were alive 6 months after the first ALTE incident. Four patients could not be followed up (unknown) and one patient died. The patient who expired was a previously healthy 1-month-old female. She was initially brought to the ED because her parents saw that she had apnea. On arrival, she presented repeated apnea during the examination in the ED and was intubated.

Table 1 Characteristics of patients with ALTE

| | Total (<i>n</i> = 112) | Recurrent (<i>n</i> = 18) | Non-recurrent (<i>n</i> = 94) | <i>P</i> |
|---------------------------------|----------------------------|-------------------------------|-----------------------------------|----------|
| PICU admission | 15/112 (13) | 11 (61) | 4 (4) | <0.0001 |
| RTI symptoms at ED presentation | 21/112 (19) | 8 (44) | 13 (14) | 0.0055 |
| Level 1 triage | 11/112 (10) | 5 (31) | 6 (7) | 0.0132 |
| Male | 55/112 (49) | 11 (61) | 44 (47) | 0.3105 |
| Underlying disease | 16/112 (14) | 1 (6) | 15 (16) | 0.4615 |
| Age <2 months at the incident | 62/112 (55) | 11 (61) | 51 (54) | 0.6173 |
| Gestational age <36 weeks | 9/111 (8) | 2 (11) | 7 (8) | 0.6392 |

Values are numbers (%)

ALTE apparent life-threatening event, PICU pediatric intensive care unit, RTI respiratory tract infection, ED emergency department

Fig. 1 Derivation of study cohort

Finally, she was admitted to the PICU and received intensive care for 2 weeks. The definitive diagnosis in her case was respiratory failure due to a viral respiratory tract infection. She was discharged without any symptoms after 1 month's hospitalization. However, a month thereafter, she was found in the morning apneic and pulseless by her parents who had been cosleeping with her and was brought to the ED in a state of cardiopulmonary arrest. She failed to respond to resuscitation and expired. Although autopsy was performed, the cause of death was not determined.

All three patients who had recurrent ALTEs after discharge were confirmed to be alive. Of the 94 patients without recurrence, 56 were confirmed to be alive 6 months after the first ALTE and 38 were unknown.

Gender, birth weight, gestational age, age, presence of underlying disease, and nutrition were similar between the patients with recurrent ALTE and the patients without. RTI symptoms were more frequent in the recurrent ALTE group (44 vs. 14%, $p = 0.0055$). The proportions of patients triaged as level 1 (31 vs. 7%, $p = 0.0132$) and of those who needed intensive care (61 vs. 4%, $p < 0.0001$) were larger in the recurrent ALTE group than those in the non-recurrent group (Table 1). As for the symptoms observed shortly after the episode, pallor was observed more frequently in the recurrent ALTE group than in the non-recurrent group (100 vs. 76%, $p = 0.0216$). The frequency of hypotonia was slightly higher in the recurrent ALTE group than in the non-recurrent group (61 vs. 39%, $p = 0.1191$), although the difference was not statistically significant (Table 2). There was no statistical difference in the prevalence of definitive diagnosis between the two groups (Table 3).

Multivariate analysis showed that RTI symptoms at ED presentation (odds ratio [OR], 5.02; 95% confidence interval [CI], 1.48–16.98) was an independent risk factor

significantly associated with recurrent ALTE. Prolonged unstable vital signs (level 1 triage) were also likely to be a risk factor; although, the analysis did not show statistical significance (odds ratio [OR], 4.32; 95% confidence interval [CI], 0.97–18.47) (Table 4).

Table 2 Symptoms of the patients shortly after ALTE

| | Recurrent ($n = 18$) | Non-recurrent ($n = 94$) | p |
|----------------------------|---------------------------|-------------------------------|--------|
| Pallor | 18 (100) | 71 (76) | 0.0216 |
| Hypotonia | 11 (61) | 37 (39) | 0.1191 |
| Excessive mucus production | 1 (6) | 20 (21) | 0.1870 |
| Red | 0 (0) | 14 (15) | 0.1204 |
| Hypertonia | 1 (6) | 8 (9) | 1.0000 |
| Vomiting | 1 (5) | 6 (6) | 1.0000 |
| Jerkin limbs | 1 (6) | 3 (3) | 0.5090 |

Values are numbers (%)

ALTE apparent life-threatening event

Table 3 Definitive diagnosis of the patients

| | Recurrent ($n = 18$) | Non-recurrent ($n = 94$) | p |
|-----------------------------|---------------------------|-------------------------------|--------|
| Breath-holding spell | 3 (17) | 4 (4) | 0.0811 |
| Respiratory tract infection | 5 (28) | 11 (11) | 0.1322 |
| Choking | 0 (0) | 12 (13) | 0.2089 |
| GER | 3 (17) | 27 (29) | 0.3900 |
| Apnea | 3 (17) | 11 (12) | 0.6959 |
| Epilepsy | 0 (0) | 5 (5) | 1.0000 |

Values are numbers (%)

GER gastroesophageal reflux

Table 4 Adjusted risk of recurrent ALTE

| Characteristics | Odds ratio | 95% CI | <i>p</i> |
|---------------------------------|------------|------------|----------|
| RTI symptoms at ED presentation | 5.02 | 1.48–16.98 | 0.0102 |
| Level 1 triage | 4.32 | 0.97–18.47 | 0.0546 |

ALTE apparent life-threatening event, RTI respiratory tract infection, ED emergency department

Discussion

To the best of our knowledge, this is the first study that investigated the risk factors for recurrent ALTE. We identified that RTI symptoms at ED presentation were an independent risk factor for recurrent ALTE.

The literature has highlighted that recurrent ALTE may indicate the underlying pathology of patients [2, 5, 9, 12]. A multicenter prospective study of ALTE developed a prediction model for the need of hospitalization and identified three variables [7]. They are “obvious need for admission,” “significant medical history,” and “the recurrence of ALTE within 24 hours”. In the study, “obvious need for admission” was defined as requiring supplemental oxygen, intubation, ventilation, intravenous antibiotics for serious bacterial infections, antiepileptic drugs for status epilepticus, or if the patient was tested positive for respiratory syncytial virus or pertussis. Another retrospective study reported that the presence of upper respiratory tract infection symptoms was one of the predictors of extreme events in infants hospitalized for ALTE [3]. The findings of these studies corroborate with ours which showed that patients who had RTI symptoms and/or prolonged unstable vital signs at ED presentation were likely to experience recurrent ALTE.

As previous studies have suggested, recurrent ALTE was a significant predictive factor for hospitalization [7]. However, those studies did not indicate which patients were likely to experience recurrence. Our findings may be helpful for physicians to predict the risks for recurrent ALTE.

There are several limitations in this study. First, this was a retrospective study conducted in a single pediatric hospital in Japan. Thus, the generalizability of our findings is unclear. Second, the abstractor was not blinded to the study objective and patient assignment in the data collection process, thereby possibly causing a bias in the analysis [6]. Third, the follow-up duration for the study cohort was inadequate. Notably, the prognosis for nearly 40% of the patients was unknown. If those patients had recurrent ALTE beyond the follow-up period of this study, the results might have been different. Nonetheless, our hospital is the only one in western Tokyo area that has a pediatric ED and we routinely instruct ALTE patients to return to our ED in case of recurrences. Although we failed to follow up on a certain proportion of patients, it was unlikely that it had a major influence on our results.

In conclusion, this study illustrates that RTI symptoms and/or prolonged unstable vital signs at the first ALTE can be independent risk factors for recurrent ALTE. Patients with ALTE who had those symptoms should be admitted for close observation of potential recurrences.

Acknowledgements We would like to thank Dr. Julian Tang of the Department of Education for Clinical Research, National Center for Child Health and Development, for proofreading and editing this manuscript.

Authors’ contributions RU: All facets of the study from study design to data collection and analysis and finally, completion of the manuscript. ON: Study concept development, design, data interpretation, and manuscript review. TM: data analysis and manuscript revision. HS and SN: Interpretation of the data and manuscript review. AI: Interpretation of the data and manuscript revision and editing. All authors read and approved the final manuscript.

Compliance with ethical standards

Funding This study was supported by a grant from The Ministry of Health, Labor and Welfare of Japan to Satoshi Nakagawa.

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval This study was approved by the Ethics Committee at the The National Center for Child Health and Development.

References

- (1987) National Institutes of Health Consensus Development Conference on Infantile Apnea and Home Monitoring, Sept 29 to Oct 1, 1986. *Pediatrics* 79:292–299
- Al Khushi N, Cote A (2011) Apparent life-threatening events: assessment, risks, reality. *Paediatr Respir Rev* 12:124–132
- Al-Kindy HA, Gélinas J-F, Hatzakis G, Côté A (2009) Risk factors for extreme events in infants hospitalized for apparent life-threatening events. *J Pediatr* 154:332–337 e332
- Claudius I, Keens T (2007) Do all infants with apparent life-threatening events need to be admitted? *Pediatrics* 119:679–683
- Fu LY, Moon RY (2012) Apparent life-threatening events: an update. *Pediatr Rev* 33:361–368 quiz 368–369
- Gilbert EH, Lowenstein SR, Koziol-McLain J, Barta DC, Steiner J (1996) Chart reviews in emergency medicine research: where are the methods? *Ann Emerg Med* 27:305–308
- Kaji AH, Claudius I, Santillanes G, Mittal MK, Hayes K, Lee J, Gausche-Hill M (2013) Apparent life-threatening event: multicenter prospective cohort study to develop a clinical decision rule for admission to the hospital. *Ann Emerg Med* 61:379–387 e374
- Mittal MK, Sun G, Baren JM (2012) A clinical decision rule to identify infants with apparent life-threatening event who can be safely discharged from the emergency department. *Pediatr Emerg Care* 28:599–605
- Poets CF, Poets A (2013) Management of apparent life-threatening events (ALTE). *Paediatr Child Health* 23:200–206
- Sarohia M, Platt S (2014) Apparent life-threatening events in children: practical evaluation and management. *Pediatr Emerg Med Pract* 11:1–14 quiz 15

11. Semmekrot BA, van Sleuwen BE, Engelberts AC, Joosten KF, Mulder JC, Liem KD, Rodrigues Pereira R, Bijlmer RP, L'Hoir MP (2010) Surveillance study of apparent life-threatening events (ALTE) in the Netherlands. *Eur J Pediatr* 169:229–236
12. Tieder JS, Altman RL, Bonkowsky JL, Brand DA, Claudius I, Cunningham DJ, DeWolfe C, Percelay JM, Pitetti RD, Smith MB (2013) Management of apparent life-threatening events in infants: a systematic review. *J Pediatr* 163(94–99):e91–e96
13. Tieder JS, Bonkowsky JL, Etzel RA, Franklin WH, Gremse DA, Herman B, Katz ES, Krilov LR, Merritt JL, Norlin C, Percelay J, Sapien RE, Shiffman RN, Smith MBH (2016) Brief resolved unexplained events (formerly apparent life-threatening events) and evaluation of lower-risk infants. *Pediatrics* 137:e20160590
14. Ueda R, Maekawa T, Nomura O, Ishiguro A, Sakai H, Nakagawa S (2014) Clinical characteristics of infants who experienced apparent life-threatening events. *J Jpn Pediatr Soc* 118:1213–1218
15. Warren DW, Jarvis A, LeBlanc L, Gravel J, Group CNW, Canadian Association of Emergency P, National Emergency Nurses A, Association des Medecins d'Urgence du Q, Canadian Paediatric S, Society of Rural Physicians of C (2008) Revisions to the Canadian Triage and Acuity Scale paediatric guidelines (PaedCTAS). *CJEM* 10:224–243