



# Discovery of shared genomic loci using the conditional false discovery rate approach

Olav B. Smeland<sup>1</sup> · Oleksandr Frei<sup>1</sup> · Alexey Shadrin<sup>1</sup> · Kevin O'Connell<sup>1</sup> · Chun-Chieh Fan<sup>2,3</sup> · Shahram Bahrami<sup>1</sup> · Dominic Holland<sup>3,5,6</sup> · Srdjan Djurovic<sup>7,8</sup> · Wesley K. Thompson<sup>4</sup> · Anders M. Dale<sup>2,3,5,6</sup> · Ole A. Andreassen<sup>1</sup>

Received: 20 November 2018 / Accepted: 8 August 2019 / Published online: 13 September 2019  
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

## Abstract

In recent years, genome-wide association study (GWAS) sample sizes have become larger, the statistical power has improved and thousands of trait-associated variants have been uncovered, offering new insights into the genetic etiology of complex human traits and disorders. However, a large fraction of the polygenic architecture underlying most complex phenotypes still remains undetected. We here review the conditional false discovery rate (condFDR) method, a model-free strategy for analysis of GWAS summary data, which has improved yield of existing GWAS and provided novel findings of genetic overlap between a wide range of complex human phenotypes, including psychiatric, cardiovascular, and neurological disorders, as well as psychological and cognitive traits. The condFDR method was inspired by Empirical Bayes approaches and leverages auxiliary genetic information to improve statistical power for discovery of single-nucleotide polymorphisms (SNPs). The cross-trait condFDR strategy analyses separate GWAS data, and leverages overlapping SNP associations, i.e., cross-trait enrichment, to increase discovery of trait-associated SNPs. The extension of the condFDR approach to conjunctive FDR (conjFDR) identifies shared genomic loci between two phenotypes. The conjFDR approach allows for detection of shared genomic associations irrespective of the genetic correlation between the phenotypes, often revealing a mixture of antagonistic and agonistic directional effects among the shared loci. This review provides a methodological comparison between condFDR and other relevant cross-trait analytical tools and demonstrates how condFDR analysis may provide novel insights into the genetic relationship between complex phenotypes.

---

The condFDR/conjFDR software is available on <https://github.com/precimed/pleiofdr> as a MATLAB package, under GPL v3 license.

---

✉ Olav B. Smeland  
o.b.smeland@medisin.uio.no

✉ Ole A. Andreassen  
o.a.andreassen@medisin.uio.no

<sup>1</sup> NORMENT Centre, Division of Mental Health and Addiction, Oslo University Hospital and Institute of Clinical Medicine, University of Oslo, Kirkeveien 166, 0424 Oslo, Norway

<sup>2</sup> Department of Cognitive Science, University of California San Diego, La Jolla, San Diego, CA 92093, USA

<sup>3</sup> Department of Radiology, University of California of San Diego, La Jolla, San Diego, CA 92093, USA

## Introduction

Most human traits and disorders have a complex etiology, which is influenced by multiple environmental and genetic factors. While some phenotypes follow simple patterns of Mendelian inheritance, large-scale genome-wide

<sup>4</sup> Department of Family Medicine and Public Health, University of California San Diego, La Jolla, San Diego, CA 92093, USA

<sup>5</sup> Department of Neuroscience, University of California San Diego, La Jolla, San Diego, CA 92093, USA

<sup>6</sup> Center for Multimodal Imaging and Genetics, University of California San Diego, La Jolla, San Diego, CA 92037, USA

<sup>7</sup> Department of Medical Genetics, Oslo University Hospital, Oslo, Norway

<sup>8</sup> NORMENT Centre, Department of Clinical Science, University of Bergen, Bergen, Norway

association studies (GWAS) conducted during the last decade have shown that most phenotypes have a complex polygenic architecture, in which genetic risk is accounted for by a large number of genetic variants, each with small effect (Visscher et al. 2017). Accumulating evidence from GWAS demonstrates that many genetic variants influence more than one phenotype, i.e., they exhibit allelic pleiotropy (Sivakumaran et al. 2011; Solovieff et al. 2013). Identification of shared genetic influences between human traits and disorders can be highly valuable to inform disease nosology, epidemiological associations, and diagnostic classification systems, improve treatment strategies, provide biological insights and uncover shared biological underpinnings (Sivakumaran et al. 2011; Solovieff et al. 2013; Visscher et al. 2017). For example, it is now evident that psychiatric disorders share a large proportion of their genetic architecture (Brainstorm et al. 2018; Cross-Disorder Group of the Psychiatric Genomics C et al. 2013), suggesting that their etiologies are not fully distinct and hence challenging existing diagnostic guidelines (Smoller et al. 2018).

GWAS typically consist of genome-wide scans of millions of common genetic variants (tag single-nucleotide polymorphisms [SNPs]), estimating the strength of their association with the phenotype of interest in massively univariate regression analyses. Given the large numbers of SNPs tested, a GWAS must correct for multiple testing and applies a genome-wide significance threshold of  $p < 5 \times 10^{-8}$  to avoid false positive findings. The consequence is that only a subset of all involved genetic variants is revealed (i.e., many false negative findings) with a large fraction of the polygenic architecture remaining to be uncovered. This phenomenon was previously labeled “the missing heritability” (Manolio et al. 2009). With increasing GWAS sample sizes, statistical power has improved and more genetic variants have been uncovered (Visscher et al. 2017). However, despite the assembly of very large GWAS samples, often involving hundreds of thousands of participants, most of the polygenic architecture underlying complex human phenotypes remain undetected (Holland et al. 2019). The number of participants needed for a GWAS to fully uncover all genetic variants influencing a given phenotype depends on the unique polygenic architecture underlying that phenotype, which is determined by the number of causal variants involved and the distribution of effect sizes (Holland et al. 2019). For example, it has been estimated that to uncover most of the genetic variants influencing the complex disorders schizophrenia and bipolar disorder, genotypes from more than one million individuals are required (Holland et al. 2019).

## Improved discovery of shared loci using conditional false discovery rate

Although the successive incremental increases in GWAS sample sizes have effectively improved the discovery of trait-associated loci, an alternative and more cost-efficient approach is to apply statistical tools that improve the yield of existing GWAS. The conditional false discovery rate (condFDR) is such an approach, which boosts GWAS discovery by leveraging auxiliary genetic information to re-adjust the GWAS test statistics in a primary phenotype (Andreassen et al. 2013b; Schork et al. 2016). The condFDR method is a model-free strategy for analysis of GWAS summary statistics inspired by the Empirical Bayes statistical framework, which is designed for situations with dense elements, such as the large number of small genetic effects seen in polygenic traits and disorders. Most commonly, the condFDR method has been applied for cross-trait analysis, by leveraging overlapping SNP associations (i.e., cross-trait enrichment) between separate GWAS to re-rank the test statistics in a primary phenotype conditional on the associations in a secondary phenotype (Andreassen et al. 2013b; Schork et al. 2016). Other auxiliary enrichment sources, such as genomic annotations (Schork et al. 2013), can also be leveraged using condFDR (Lo et al. 2017; Wang et al. 2016b). Since its introduction in 2013 (Andreassen et al. 2013a), the condFDR method has increased genetic discovery in a wide spectrum of complex human traits and disorders, including psychiatric, cardiovascular and neurological disorders, as well as metabolic, psychological and cognitive traits, among others (see Table 1 for a selection of cross-trait condFDR studies) (Andreassen et al. 2013a, c, 2014a, b, c; Broce et al. 2018, 2019; Desikan et al. 2015; Drange et al. 2019; Ferrari et al. 2017; Hu et al. 2018; Karch et al. 2018; Le Hellard et al. 2017; LeBlanc et al. 2015; Liu et al. 2013; Lv et al. 2017; McLaughlin et al. 2017; Mufford et al. 2019; Shadrin et al. 2018; Smeland et al. 2017a, b, 2018; 2019; van der Meer et al. 2018; Wang et al. 2016a; Winsvold et al. 2017; Witoelar et al. 2017; Yokoyama et al. 2016, 2017; Zuber et al. 2018).

The present review focuses on the cross-trait condFDR approach, which returns a condFDR value for each SNP, defined as the probability that a SNP is null in the first phenotype (i.e., that it has no association with the phenotype) given that the  $p$  values in the first and second phenotypes are as small as or smaller than the observed ones. The condFDR estimates are obtained for each nominal SNP  $p$  value in the primary phenotype after computing the stratified empirical cumulative distribution functions (cdfs) of the nominal  $p$  values (Sun et al. 2006; Yoo et al. 2009). The separate strata are determined by the

**Table 1** Selected cross-trait conditional false discovery rate studies


Primary phenotype	Secondary phenotype	Novel loci for primary phenotype	Citation
Schizophrenia	Cardiovascular-disease risk factors	14 at condFDR < 0.01	Andreassen et al. (2013a)
Primary sclerosing cholangitis	Autoimmune diseases	33 at condFDR < 0.001	Liu et al. (2013)
Bipolar disorder	Schizophrenia	2 at condFDR < 0.01	Andreassen et al. (2013c)
Schizophrenia	Multiple sclerosis	5 at condFDR < 0.01	Andreassen et al. (2014a)
Systolic blood pressure	Comorbid traits and diseases	42 at condFDR < 0.01	Andreassen et al. (2014b)
Alzheimer disease	C-reactive protein, plasma lipids	55 at condFDR < 0.05	Desikan et al. (2015)
Coronary artery disease	Cardiovascular-disease risk factors	67 at condFDR < 0.01	LeBlanc et al. (2015)
Alzheimer disease	Autoimmune diseases	Not available	Yokoyama et al. (2016)
Amyotrophic lateral sclerosis	Schizophrenia	5 at condFDR < 0.01	McLaughlin et al. (2017)
Schizophrenia	Educational attainment	23 at condFDR < 0.01	Le Hellard et al. (2017)
Sporadic frontotemporal dementia	Alzheimer disease, Parkinson disease	13 at condFDR < 0.05	Ferrari et al. (2017)
Schizophrenia	Cognitive traits	13 at conjFDR < 0.05	Smeland et al. (2017a)
Corticobasal degeneration	Progressive supranuclear palsy, frontotemporal dementia	3 at conjFDR < 0.05	Yokoyama et al. (2017)
Amyotrophic lateral sclerosis	Neurodegenerative disorders	22 at condFDR < 0.05	Karch et al. (2018)
Frontotemporal dementia	Autoimmune diseases	5 at conjFDR < 0.05	Broce et al. (2018)
Schizophrenia	Subcortical brain volumes	3 at conjFDR < 0.05	Smeland et al. (2018)
Attention-deficit/hyperactivity disorder	Educational attainment	4 at condFDR < 0.01, 1 at conjFDR < 0.05	Shadrin et al. (2018)
Alzheimer disease	Cardiovascular-disease risk factors	4 at conjFDR < 0.05	Broce et al. (2019)
Schizophrenia, bipolar disorder	Intelligence	20 schizophrenia loci and 4 bipolar disorder loci at conjFDR < 0.01	Smeland et al. (2019)

relative enrichment of SNP associations as a function of increased nominal SNP  $p$  values in a secondary phenotype. The standard FDR framework derives from a model that assumes that the distribution of test statistics in a GWAS can be formulated as a mixture of null and non-null effects, with true associations (non-null effects) having more extreme test statistics than false associations (null effects) on average. Given a statistical genetic relationship between two phenotypes, stratification of the test statistics in a primary phenotype based on the genetic associations with a secondary phenotype will result in a reduction in the FDR at a given nominal  $p$  value relative to the FDR computed from the unstratified distribution of the primary phenotype  $p$  values alone, and thus, re-rank the test statistics.

The first step in the condFDR procedure is to construct conditional quantile–quantile (Q–Q) plots, which extends the standard Q–Q plots commonly applied in GWAS. Standard Q–Q plots visualize the enrichment of statistical association relative to that expected under the global null hypothesis by plotting the nominal  $-\log_{10} p$  values of the single SNP association statistics versus their empirical distribution. Conditional Q–Q plots help visualize the cross-trait enrichment between two phenotypes and are constructed by creating subsets of SNPs based on the level of association with the secondary phenotype. Under the global null hypothesis, the nominal  $p$  values will form a straight line plotted as a function of their empirical distribution. Under

polygenic association, standard Q–Q plots will be deflected leftwards, while cross-trait enrichment can be seen as successive leftward deflections in conditional Q–Q plots as levels of SNP associations with the secondary phenotype increase. Figure 1a presents a conditional Q–Q plot demonstrating SNP enrichment for the psychiatric disorder bipolar disorder ( $n = 51,710$ ) (Stahl et al. 2019) as a function of the association with intelligence ( $n = 269,867$ ) (Savage et al. 2018), adapted from Smeland et al. (2019). A complementary way to assess for cross-trait enrichment is to construct fold-enrichment plots, which provide a more direct visualization of the polygenic enrichment (Fig. 1b). The fold enrichment is calculated as the ratio between the  $-\log_{10}(p)$  cumulative distribution for a given stratum and the cumulative distribution for all SNPs. Figure 1b shows that for SNPs with  $p$  values below 0.001 in intelligence, there was up to 60-fold enrichment of stronger SNP associations with bipolar disorder in comparison to all SNPs. The enrichment seen in conditional Q–Q plots and fold-enrichment plots reflects increased tail probabilities in the distribution of test statistics and an overabundance of low  $p$  values compared to that expected by chance, which can be directly interpreted in terms of a Bayesian interpretation of the true discovery rate ( $\text{TDR} = 1 - \text{FDR}$ ; see Box 1 for mathematical framework) (Efron 2010). This is illustrated in Fig. 1c.

To control for spurious (i.e., non-generalizable) enrichment due to population stratification or cryptic relatedness




**Fig. 1** Cross-trait enrichment between bipolar disorder (BD;  $n=51,710$ ) (Stahl et al. 2019) and intelligence ( $n=269,867$ ) (Savage et al. 2018), adapted from Smeland et al. (2019). **a** Conditional Q–Q plot displaying the nominal  $-\log_{10} p$  values of the single SNP association statistics versus their empirical distribution in BD below the standard GWAS threshold of  $p < 5 \times 10^{-8}$  as a function of significance of association with intelligence at the level of  $p \leq 0.1$ ,  $p \leq 0.01$ ,  $p \leq 0.001$ , respectively. The blue line indicates all SNPs. The dashed line indicates the null hypothesis. **b** Fold-enrichment plot of enrich-

ment versus nominal  $-\log_{10} p$  values in BD as a function of association with intelligence. **c** Conditional true discovery rate (TDR) plot illustrating the increase in TDR associated with increased enrichment in BD conditioned on intelligence. The test statistics were corrected for genomic inflation, SNPs were randomly pruned across 500 iterations using a linkage disequilibrium  $r^2$  threshold of 0.1, and the extended major histocompatibility complex region and chromosomal region 8p.23.1 were excluded (Smeland et al. 2019)

(Devlin and Roeder 1999), all test statistics are corrected using a genomic inflation control procedure leveraging intergenic SNPs, which are relatively depleted for true associations (Schork et al. 2013). Conditional Q–Q plots and the condFDR computation are conducted after random pruning to approximate independence, by selecting one random SNP per LD block (defined by an  $r^2 > 0.1$ ) averaged over at least 100 iterations (Andreassen et al. 2013b; Schork et al. 2016). Similar to previously described stratified FDR procedures (Sun et al. 2006; Yoo et al. 2009), the condFDR value is then determined for each SNP by constructing a

two-dimensional FDR look-up table where the FDR for SNP associations with the primary phenotype is computed conditionally on the nominal  $p$  values for SNP associations with the secondary phenotype (Box 1). Figure 2a presents the respective condFDR look-up table for bipolar disorder conditional on intelligence, corresponding to the cross-trait enrichment observed in Fig. 1.

The conjFDR (conjFDR) is an extension of the condFDR, which allows for discovery of SNPs significantly associated with two phenotypes simultaneously (Andreassen et al. 2013a; Schork et al. 2016). The conjFDR is determined



**Fig. 2** **a** Conditional false discovery rate (condFDR) 2D look-up table for SNP associations with bipolar disorder (BD) conditional on SNP associations with intelligence, corresponding to the cross-trait enrichment observed in Fig. 1. The FDR in BD SNPs are computed conditionally on the nominal intelligence  $p$  values. **b** condFDR 2D look-up

table for SNP associations with intelligence conditional on SNP associations with BD. **c** Corresponding conjunctive FDR (conjFDR) 2D look-up table for SNP associations shared between BD and intelligence. The color refers to the FDR values

after inverting the roles of the primary and secondary phenotypes and repeating the condFDR procedure. Based on previous conjunction tests for  $p$  value statistics (Nichols et al. 2005), the conjFDR is defined as the maximum of the two condFDR values, providing a conservative estimate of the FDR for an SNP association with both phenotypes jointly (Fig. 2c). Thus, in combination, the condFDR/conjFDR approaches both improve SNP discovery rates (condFDR) and enable detection of shared genomic loci (conjFDR), respectively. Since the condFDR/conjFDR estimates are based on nominal  $p$  values only, these methods are agnostic to the effect directions of the individual SNPs, and can detect overlapping SNP associations irrespective of the genome-wide genetic correlation between phenotypes. However, after detecting likely overlapping SNPs, the directional SNP effects in the loci can be determined post hoc by comparing the effect sizes ( $z$  scores or odds ratios) between the phenotypes.

The condFDR/conjFDR approaches have some limitations. Although all SNPs are randomly pruned using an LD  $r^2$  threshold of 0.1, complex correlations among the test statistics may bias the condFDR estimates (Schwartzman and Lin 2011). Hence, given strong SNP associations within long-range LD regions, such as the extended major histocompatibility complex (MHC) region, chromosomal region 8p.23.1, the microtubule-associated tau protein (*MAPT*) region or the *APOE* region (Price et al. 2008), these regions should be excluded to avoid artificially inflated genetic enrichment. The condFDR/conjFDR procedures are agnostic about the specific causal variants underlying the overlapping genomic associations, which could arise from both shared and separate causal variants, or “mediated pleiotropy”, where one phenotype is causative of the other (Solovieff et al. 2013). Given that the cross-trait enrichment both reflects the extent of polygenic overlap between the phenotypes and the power of the two GWAS analyzed, cross-trait enrichment will be harder to detect if one or both investigated GWAS are inadequately powered. Another important limitation of the condFDR method is that a large fraction of overlapping participants between the investigated GWAS may inflate the cross-trait enrichment, and shared participants should, therefore, be reduced to a minimum. An extension of condFDR, allowing shared controls, has been proposed (Liley and Wallace 2015).

## Comparison to other cross-trait analytical tools

A large number of tools for cross-trait analysis using GWAS data have been developed in recent years, which have been reviewed in detail elsewhere (Gratten and Visscher 2016; Hackinger and Zeggini 2017; Pasaniuc and Price 2017;

Schork et al. 2016). In short, the methods differentiate in terms of the data analyzed (summary statistics versus individual genotype data), the underlying mathematical framework and assumptions, whether they are bivariate or multivariate in nature, and whether they measure overlap at the genome-wide level or across individual SNPs or loci/regions. Here, we compare the condFDR/conjFDR approach to a selection of relevant cross-trait analytical tools.

The most common approaches for evaluating genetic overlap at the genome-wide level include tools such as polygenic risk scores (Purcell et al. 2009), mixed-model approaches (Cross-Disorder Group of the Psychiatric Genomics C et al. 2013; Lee et al. 2012) and LD score regression (Bulik-Sullivan et al. 2015a), which return a single estimate of shared genetic risk between phenotypes. Polygenic risk scores are per-individual risk profiles based on the sum of alleles associated with a phenotype weighted by their effect sizes (Purcell et al. 2009). The polygenic risk score approach uses summary statistics as training data and requires individual genotype data in an independent target sample to test how well the polygenic risk score explains phenotypic variation in the target phenotype. Another traditional measure that estimates the degree of pleiotropy is the genetic correlation, which is defined as the correlation between the genetic influences for a pair of traits, thus indicating the proportion of variance that the two traits share due to genetic causes. Mixed-model approaches (Lee et al. 2012), originally implemented in the Genome-wide Complex Trait Analysis software (GCTA), obtained unbiased estimates of the genetic correlation using individual genotype data, relaxing several limitations of traditional studies based on pedigree data. Estimates of genetic correlation can also be quantified from GWAS summary statistics, using cross-trait LD score regression (Bulik-Sullivan et al. 2015a) and its multivariate extension Genomic SEM (Grotzinger et al. 2019). LD score regression aims to distinguish confounding from polygenicity by regressing the association statistics of SNPs on their ‘LD scores’, which is a measure of the amount of genetic variation the SNP represents (Bulik-Sullivan et al. 2015b). Application of LD score regression to the bivariate framework estimates the co-variance in the SNP heritability between two phenotypes, allowing sample overlap (Bulik-Sullivan et al. 2015a). An alternative approach estimating local genetic correlations based on the fixed effects model is also available (Shi et al. 2017). The condFDR approach is fundamentally different to these tools by aiming for discovery of specific genomic loci. However, the condFDR approach similarly focuses on the polygenic fraction that did not reach genome-wide significance to uncover cross-trait enrichment. To fully disentangle the genetic relationship between complex phenotypes, it is necessary to complement measures of genetic overlap at the genome-wide level with cross-trait analytical tools allowing




detection of individual shared loci regardless of their directional effects. For instance, a recent condFDR study demonstrated substantial cross-trait enrichment between bipolar disorder (Stahl et al. 2019) and intelligence (Savage et al. 2018) (Fig. 1) and uncovered a balanced pattern of concordant and discordant directional effects among 79 shared loci identified at  $\text{conjFDR} < 0.05$  (Fig. 3) (Smeland et al. 2019). These findings extend and comply with prior genetic studies reporting no significant genome-wide genetic correlation between the phenotypes (Brainstorm et al. 2018; Davies et al. 2018; Hill et al. 2016; Lencz et al. 2014; Savage et al. 2018; Sniekers et al. 2017; Stahl et al. 2019).

There is a large class of cross-trait methods aiming to discover specific genomic loci unique or shared between phenotypes inspired by the meta-analysis technique (Willer et al. 2010) and its extensions dealing with sample overlap (Han et al. 2016; Lin and Sullivan 2009). For example, the COMBINE approach (Ellinghaus et al. 2012) consists of two separate runs of a same-effect and opposite-effect meta-analysis, both using the inverse variance-weighted procedure. In the opposite-effect meta-analysis, the minor and major alleles are flipped in the second dataset to capture bi-allelic SNPs with opposite-effect directions in the two phenotypes investigated. This method was later refined and extended to multiple heterogeneous traits using restricted and weighted subset search (ASSET) (Bhattacharjee et al. 2012), which exhaustively explore subsets of studies to achieve the best

possible trade-off between specificity and sample size. Its successor, compare-and-contrast meta-analysis (CCMA) (Baurecht et al. 2015), further improved the power to discover associations by combining the subset search approach with trans-ethnic meta-analysis (MANTRA) (Morris 2011). Several alternative approaches explore additional information, including individual-level genotypes (MultiPhen) (O'Reilly et al. 2012), phenotypic correlations (TATES) (van der Sluis et al. 2013) or estimated genetic correlations (MTAG) (Turley et al. 2018). A common feature of all techniques based on a meta-analysis framework is that the analysis is performed independently for each SNP, thus requiring a follow-up mechanism to control for multiple testing, such as Bonferroni correction, to avoid false positive findings. The condFDR analysis, on the other hand, directly works with the entire original set of  $p$  values from the two GWAS and intrinsically incorporates multiple testing via the FDR framework (Efron 2010).

Another class of methods aims at disentangling LD structure to reveal underlying causal genetic mechanisms. Mendelian randomization aims to distinguish true pleiotropy from mediated pleiotropy by investigating whether one phenotype is causative to the other (Hernan and Robins 2006; Lawlor et al. 2008; Smith and Ebrahim 2003; Zhu et al. 2018). Mendelian randomization assigns genetic variants, which are expected to be independent of confounding factors, as instrumental variables to



**Fig. 3** Common genetic variants jointly associated with bipolar disorder (BD;  $n=51,710$ ) and intelligence ( $n=269,867$ ) at conjunctional false discovery rate ( $\text{conjFDR} < 0.05$ ), adapted from Smeland et al. (2019). Manhattan plot showing the  $-\log_{10}$  transformed  $\text{conjFDR}$  values for each SNP on the y-axis and chromosomal positions

along the x-axis. The dotted horizontal line represents the threshold for significant shared associations ( $\text{conjFDR} < 0.05$ , i.e.,  $-\log_{10}(\text{conjFDR}) > 1.3$ ). Independent lead SNPs are encircled in black. For details, see Supplementary Table 9 in Smeland et al. (2019)

test for causality. Several available Bayesian approaches (Giambartolomei et al. 2014; Pickrell et al. 2016) explore whether two association signals in the same genomic region obtained from two different GWAS share a single causal variant or multiple causal variants. Frei and colleagues performed a similar analysis at the genome-wide level, estimating the proportion of phenotype-specific causal variants and shared variants between complex phenotypes using GWAS summary data, while controlling for shared participants (Frei et al. 2019). The analysis demonstrates how the shared polygenic component may constitute a large fraction of the genetic architecture of one phenotype, while constituting a smaller fraction of the architecture of a phenotype with larger polygenicity. While the condFDR/conjFDR approach is agnostic about the causal variants underlying the identified associations, it complements these methods by improving the discovery of genomic loci, which can be used to prioritize downstream analysis.

## Conclusion

Accumulating evidence has shown that genetic pleiotropy is pervasive among complex human traits and disorders, providing important insights into etiological relationships. Since its introduction in 2013, application of the condFDR/conjFDR approach has increased yield of existing GWAS and aided the discovery of overlapping genomic loci between polygenic phenotypes. Given that large fractions of the polygenic architecture underlying most complex phenotypes still remain undetected, the condFDR/conjFDR approach represents a cost-effective powerful strategy useful for improving GWAS discovery and help elucidating shared genetic etiologies.

**Acknowledgements** National Institutes of Health (NS057198; EB00790); National Institutes of Health NIDA/NCI: U24DA041123; the Research Council of Norway (229129; 213837; 248778; 223273; 249711); the South-East Norway Regional Health Authority (2017-112); KG Jebsen Stiftelsen (SKGJ-2011-36).

## Compliance with ethical standards

**Conflict of interest** OA.A. has received speaker's honorarium from Lundbeck and is a consultant for Healthlytix. C.C.F. is under employment of Multimodal Imaging Service, dba Healthlytix, in addition to his research appointment at the University of California, San Diego. A.M.D. is a founder of and holds equity interest in CorTechs Labs and serves on its scientific advisory board. He is also a member of the Scientific Advisory Board of Healthlytix and receives research funding from General Electric Healthcare (GEHC). The terms of these arrangements have been reviewed and approved by the University of California, San Diego in accordance with its conflict of interest policies. Remaining authors have no conflicts of interest to declare.

## Box 1: Conditional and conjunctive false discovery rate

The 'enrichment' seen in the conditional Q–Q plots can be directly interpreted in terms of a Bayesian interpretation of the true discovery rate (TDR = 1 – false discovery rate (FDR)) (Efron 2010). More specifically, for a given  $p$  value, under a simple two-group (null and non-null) model, Bayes rule gives the posterior probability of being null as:

$$\text{FDR}(p) = \pi_0 F_0(p) / F(p), \quad (1)$$

where  $\pi_0$  is the proportion of null SNPs,  $F_0$  is the cumulative distribution function (cdf) of the null SNPs, and  $F$  is the cdf of all SNPs, both null and non-null (Efron 2007). Here, we assume the SNP  $p$  values are a priori independent and identically distributed. Under the null hypothesis,  $F_0$  is the cdf of the uniform distribution on the unit interval  $[0, 1]$ , so that Eq. (1) reduces to:

$$\text{FDR}(p) = \pi_0 p / F(p). \quad (2)$$

$F$  can be estimated by the empirical cdf  $q = N_p / N$ , where  $N_p$  is the number of SNPs with  $p$  values less than or equal to  $p$ , and  $N$  is the total number of SNPs. Replacing  $F$  by  $q$  in Eq. (2), we get:

$$\text{Estimated FDR}(p) = \pi_0 p / q, \quad (3)$$

which is biased upwards as an estimate of the FDR (Efron and Tibshirani 2002). Replacing  $\pi_0$  in Eq. (3) with unity gives an estimated FDR that is further biased upward:

$$q^* = p / q. \quad (4)$$

If  $\pi_0$  is close to one, the increase in bias going from Eqs. (3–4) is minimal. The quantity  $1 - p/q$  is, therefore, biased downward, and hence a conservative estimate of the TDR. Referring to the Q–Q plots, we see that  $q^*$  is equivalent to the nominal  $p$  value divided by the empirical quantile, as defined earlier. We can thus read the FDR estimate directly off the Q–Q plot as:

$$-\log_{10}(q^*) = \log_{10}(q) - \log_{10}(p), \quad (5)$$

i.e., the horizontal shift of the curves in the Q–Q plots from the expected line  $x = y$ , with a larger shift corresponding to a smaller FDR. To estimate the conditional FDR of a given SNP, we repeat the above procedure for a subset of SNPs with  $p$  values in the secondary GWAS equal to or lower than that observed for the given SNP. Formally, this is given by:

$$\text{FDR}(p_1 | p_2) = \pi_0(p_2) p_1 / F(p_1 | p_2), \quad (6)$$

where  $p_1$  is the  $p$  value for the first phenotype,  $p_2$  is the  $p$  value for the second, and  $F(p_1 | p_2)$  is the conditional cdf and  $\pi_0(p_2)$  the conditional proportion of null SNPs for the first phenotype, given that  $p$  values for the second phenotype are  $p_2$  or smaller. The condFDR framework is closely related to the stratified FDR method developed by Sun et al.

(2006). Whereas they propose computing FDR separately conditional on membership in pre-defined discrete strata of  $p$  values, here, we condition the estimated FDR on a continuous random variable, the SNP  $p$  values with respect to a second phenotype.

To identify SNPs jointly associated with two phenotypes using conjunctive FDR, the conditional FDR procedure is repeated after inverting the roles of the primary and secondary phenotypes. Similar to previous conjunction tests for  $p$  value statistics (Nichols et al. 2005), the conjunctive FDR estimate is defined as the maximum of both conditional FDR values, which minimizes the effect of a single phenotype driving the common association signal. Formally, the conjunctive FDR is given by:

$$\begin{aligned} \text{FDR}_{\text{Phenotype1}\&\text{Phenotype2}}(p_1, p_2) &= \pi_0 F_0(p_1, p_2) / F(p_1, p_2) \\ &+ \pi_1 F_1(p_1, p_2) / F(p_1, p_2) \\ &+ \pi_2 F_2(p_1, p_2) / F(p_1, p_2), \end{aligned} \quad (7)$$

where  $\pi_0$  is the a priori proportion of SNPs null for both phenotypes simultaneously and  $F_0(p_1, p_2)$  is the joint null cdf,  $\pi_1$  is the a priori proportion of SNPs non-null for the first phenotype and null for the second with  $F_1(p_1, p_2)$  the joint cdf of these SNPs, and  $\pi_2$  is the a priori proportion of SNPs non-null for the second phenotype and null for the first, with joint cdf  $F_2(p_1, p_2)$ .  $F(p_1, p_2)$  is the joint overall mixture cdf for all phenotype 1 and 2 SNPs.

Conditional empirical cdfs provide a model-free method to obtain conservative estimates of Eq. (7). This can be seen as follows: estimate the conjunction FDR by:

$$\begin{aligned} \text{Estimated FDR}_{\text{Phenotype1}\&\text{Phenotype2}} &= \max \left\{ \text{Estimated FDR}_{\text{Phenotype1}|\text{Phenotype2}}, \right. \\ &\left. \text{Estimated FDR}_{\text{Phenotype2}|\text{Phenotype1}} \right\}, \end{aligned} \quad (8)$$

where  $\text{Estimated FDR}_{\text{Phenotype1}|\text{Phenotype2}}$  and  $\text{Estimated FDR}_{\text{Phenotype2}|\text{Phenotype1}}$  are conservative (upwardly biased) estimates of Eq. (6). Thus, Eq. (8) is a conservative estimate of  $\max \{p_1/F(p_1|p_2), p_2/F(p_2|p_1)\} = \max \{p_1 F_2(p_2)/F(p_1, p_2), p_2 F_1(p_1)/F(p_1, p_2)\}$ , with  $F_1(p_1)$  and  $F_2(p_2)$  the marginal non-null cdfs of SNPs for phenotypes 1 and 2, respectively. For enriched samples,  $p$  values will tend to be smaller than predicted from the uniform distribution, so that  $F_1(p_1) \geq p_1$  and  $F_2(p_2) \geq p_2$ . Then,  $\max \{p_1 F_2(p_2)/F(p_1, p_2), p_2 F_1(p_1)/F(p_1, p_2)\} \geq [\pi_0 + \pi_1 + \pi_2] \max \{p_1 F_2(p_2)/F(p_1, p_2), p_2 F_1(p_1)/F(p_1, p_2)\} \geq [\pi_0 p_1 p_2 + \pi_1 p_2 F_1(p_1) + \pi_2 p_1 F_2(p_2)]/F(p_1, p_2)$ .

Under the assumption that SNPs are independent if one or both are null, reasonable for disjoint samples, this last quantity is precisely the conjunctive FDR given in Eq. (7). Thus,

Eq. (8) is a conservative model-free estimate of the conjunctive FDR.

## References

- Andreassen OA et al (2013a) Improved detection of common variants associated with schizophrenia by leveraging pleiotropy with cardiovascular-disease risk factors. *Am J Hum Genet* 92:197–209. <https://doi.org/10.1016/j.ajhg.2013.01.001>
- Andreassen OA, Thompson WK, Dale AM (2013b) Boosting the power of schizophrenia genetics by leveraging new statistical tools. *Schizophr Bull*. <https://doi.org/10.1093/schbul/sbt168>
- Andreassen OA et al (2013c) Improved detection of common variants associated with schizophrenia and bipolar disorder using pleiotropy-informed conditional false discovery rate. *PLoS Genet* 9:e1003455. <https://doi.org/10.1371/journal.pgen.1003455>
- Andreassen OA et al (2014a) Genetic pleiotropy between multiple sclerosis and schizophrenia but not bipolar disorder: differential involvement of immune-related gene loci. *Mol Psychiatry* 20:207–214. <https://doi.org/10.1038/mp.2013.195>
- Andreassen OA et al (2014b) Identifying common genetic variants in blood pressure due to polygenic pleiotropy with associated phenotypes. *Hypertension* 63:819–826. <https://doi.org/10.1161/hypertensionaha.113.02077>
- Andreassen OA et al (2014c) Shared common variants in prostate cancer and blood lipids. *Int J Epidemiol* 43:1205–1214. <https://doi.org/10.1093/ije/dyu090>
- Baurecht H et al (2015) Genome-wide comparative analysis of atopic dermatitis and psoriasis gives insight into opposing genetic mechanisms. *Am J Hum Genet* 96:104–120. <https://doi.org/10.1016/j.ajhg.2014.12.004>
- Bhattacharjee S et al (2012) A subset-based approach improves power and interpretation for the combined analysis of genetic association studies of heterogeneous traits. *Am J Hum Genet* 90:821–835. <https://doi.org/10.1016/j.ajhg.2012.03.015>
- Brainstorm C et al (2018) Analysis of shared heritability in common disorders of the brain. *Science*. <https://doi.org/10.1126/science.aap8757>
- Broce I et al (2018) Immune-related genetic enrichment in frontotemporal dementia: an analysis of genome-wide association studies. *PLoS Med* 15:e1002487. <https://doi.org/10.1371/journal.pmed.1002487>
- Broce IJ et al (2019) Dissecting the genetic relationship between cardiovascular risk factors and Alzheimer's disease. *Acta Neuropathol* 137:209–226. <https://doi.org/10.1007/s00401-018-1928-6>
- Bulik-Sullivan B et al (2015a) An atlas of genetic correlations across human diseases and traits. *Nat Genet* 47:1236–1241. <https://doi.org/10.1038/ng.3406>
- Bulik-Sullivan BK et al (2015b) LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet* 47:291–295. <https://doi.org/10.1038/ng.3211>
- Cross-Disorder Group of the Psychiatric Genomics C et al (2013) Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. *Nat Genet* 45:984–994. <https://doi.org/10.1038/ng.2711>
- Davies G et al (2018) Study of 300,486 individuals identifies 148 independent genetic loci influencing general cognitive function. *Nat Commun* 9:2098. <https://doi.org/10.1038/s41467-018-04362-x>
- Desikan RS et al (2015) Polygenic overlap between C-reactive protein, plasma lipids, and alzheimer disease. *Circulation* 131:2061–2069. <https://doi.org/10.1161/CIRCULATIONAHA.115.015489>



- Devlin B, Roeder K (1999) Genomic control for association studies. *Biometrics* 55:997–1004
- Drange OK et al (2019) Genetic overlap between alzheimer's disease and bipolar disorder implicates the MARK2 and VAC14 genes. *Front Neurosci* 13:220. <https://doi.org/10.3389/fnins.2019.00220>
- Efron B (2007) Size, power and false discovery rates. *Ann Stat* 35:1351–1377
- Efron B (2010) Large-scale inference: empirical Bayes methods for estimation, testing, and prediction. Institute of mathematical statistics monographs, vol 1. Cambridge University Press, Cambridge
- Efron B, Tibshirani R (2002) Empirical bayes methods and false discovery rates for microarrays. *Genet Epidemiol* 23:70–86. <https://doi.org/10.1002/gepi.1124>
- Ellinghaus D et al (2012) Combined analysis of genome-wide association studies for Crohn disease and psoriasis identifies seven shared susceptibility loci. *Am J Hum Genet* 90:636–647. <https://doi.org/10.1016/j.ajhg.2012.02.020>
- Ferrari R et al (2017) Genetic architecture of sporadic frontotemporal dementia and overlap with Alzheimer's and Parkinson's diseases. *J Neurol Neurosurg Psychiatry* 88:152–164. <https://doi.org/10.1136/jnnp-2016-314411>
- Frei O et al (2019) Bivariate causal mixture model quantifies polygenic overlap between complex traits beyond genetic correlation. *Nat Commun* 10:2417. <https://doi.org/10.1038/s41467-019-10310-0>
- Giambartolomei C, Vukcevic D, Schadt EE, Franke L, Hingorani AD, Wallace C, Plagnol V (2014) Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet* 10:e1004383. <https://doi.org/10.1371/journal.pgen.1004383>
- Gratten J, Visscher PM (2016) Genetic pleiotropy in complex traits and diseases: implications for genomic medicine. *Genome Med* 8:78. <https://doi.org/10.1186/s13073-016-0332-x>
- Grotzinger AD et al (2019) Genomic structural equation modelling provides insights into the multivariate genetic architecture of complex traits. *Nat Hum Behav*. <https://doi.org/10.1038/s41562-019-0566-x>
- Hackinger S, Zeggini E (2017) Statistical methods to detect pleiotropy in human complex traits. *Open Biol*. <https://doi.org/10.1098/rsob.170125>
- Han B, Duong D, Sul JH, de Bakker PI, Eskin E, Raychaudhuri S (2016) A general framework for meta-analyzing dependent studies with overlapping subjects in association mapping. *Hum Mol Genet* 25:1857–1866. <https://doi.org/10.1093/hmg/ddw049>
- Hernan MA, Robins JM (2006) Instruments for causal inference: an epidemiologist's dream? *Epidemiology (Cambridge, Mass)* 17:360–372. <https://doi.org/10.1097/01.ede.0000222409.00878.37>
- Hill WD, Davies G, Group CCW, Liewald DC, McIntosh AM, Deary IJ (2016) Age-dependent pleiotropy between general cognitive function and major psychiatric disorders. *Biol Psychiat* 80:266–273. <https://doi.org/10.1016/j.biopsych.2015.08.033>
- Holland D et al (2019) Beyond SNP heritability: polygenicity and discoverability of phenotypes estimated with a univariate gaussian mixture model. *bioRxiv*. <https://doi.org/10.1101/133132>
- Hu Y et al (2018) Identification of novel potentially pleiotropic variants associated with osteoporosis and obesity using the cFDR method. *J Clin Endocrinol Metab* 103:125–138. <https://doi.org/10.1210/jc.2017-01531>
- Karch CM et al (2018) Selective genetic overlap between amyotrophic lateral sclerosis and diseases of the frontotemporal dementia spectrum. *JAMA Neurol* 75:860–875. <https://doi.org/10.1001/jaman.2018.0372>
- Lawlor DA, Harbord RM, Sterne JA, Timpson N, Davey Smith G (2008) Mendelian randomization: using genes as instruments for making causal inferences in epidemiology. *Stat Med* 27:1133–1163. <https://doi.org/10.1002/sim.3034>
- Le Hellard S et al (2017) Identification of gene loci that overlap between schizophrenia and educational attainment. *Schizophr Bull* 43:654–664. <https://doi.org/10.1093/schbul/sbw085>
- LeBlanc M et al (2015) Identifying novel gene variants in coronary artery disease and shared genes with several cardiovascular risk factors. *Circ Res*. <https://doi.org/10.1161/circresaha.115.306629>
- Lee SH, Yang J, Goddard ME, Visscher PM, Wray NR (2012) Estimation of pleiotropy between complex diseases using single-nucleotide polymorphism-derived genomic relationships and restricted maximum likelihood. *Bioinformatics* 28:2540–2542. <https://doi.org/10.1093/bioinformatics/bts474>
- Lencz T et al (2014) Molecular genetic evidence for overlap between general cognitive ability and risk for schizophrenia: a report from the cognitive genomics consortium (COGENT). *Mol Psychiatry* 19:168–174. <https://doi.org/10.1038/mp.2013.166>
- Liley J, Wallace C (2015) A pleiotropy-informed Bayesian false discovery rate adapted to a shared control design finds new disease associations from GWAS summary statistics. *PLoS Genet* 11:e1004926. <https://doi.org/10.1371/journal.pgen.1004926>
- Lin DY, Sullivan PF (2009) Meta-analysis of genome-wide association studies with overlapping subjects. *Am J Hum Genet* 85:862–872. <https://doi.org/10.1016/j.ajhg.2009.11.001>
- Liu JZ et al (2013) Dense genotyping of immune-related disease regions identifies nine new risk loci for primary sclerosing cholangitis. *Nat Genet* 45:670–675. <https://doi.org/10.1038/ng.2616>
- Lo MT et al (2017) Modeling prior information of common genetic variants improves gene discovery for neuroticism. *Hum Mol Genet* 26:4530–4539. <https://doi.org/10.1093/hmg/ddx340>
- Lv WQ et al (2017) Novel common variants associated with body mass index and coronary artery disease detected using a pleiotropic cFDR method. *J Mol Cell Cardiol* 112:1–7. <https://doi.org/10.1016/j.yjmcc.2017.08.011>
- Manolio TA et al (2009) Finding the missing heritability of complex diseases. *Nature* 461:747–753. <https://doi.org/10.1038/nature08494>
- McLaughlin RL et al (2017) Genetic correlation between amyotrophic lateral sclerosis and schizophrenia. *Nat Commun* 8:14774. <https://doi.org/10.1038/ncomms14774>
- Morris AP (2011) Transethnic meta-analysis of genomewide association studies. *Genet Epidemiol* 35:809–822. <https://doi.org/10.1002/gepi.20630>
- Mufford M et al (2019) Concordance of genetic variation that increases risk for tourette syndrome and that influences its underlying neurocircuitry. *Transl Psychiatry* 9:120. <https://doi.org/10.1038/s41398-019-0452-3>
- Nichols T, Brett M, Andersson J, Wager T, Poline JB (2005) Valid conjunction inference with the minimum statistic. *Neuroimage* 25:653–660. <https://doi.org/10.1016/j.neuroimage.2004.12.005>
- O'Reilly PF, Hoggart CJ, Pomyen Y, Calboli FCF, Elliott P, Jarvelin M-R, Coin LJM (2012) MultiPhen: joint model of multiple phenotypes can increase discovery in GWAS. *PLOS One* 7:e34861. <https://doi.org/10.1371/journal.pone.0034861>
- Pasaniuc B, Price AL (2017) Dissecting the genetics of complex traits using summary association statistics. *Nat Rev Genet* 18:117–127. <https://doi.org/10.1038/nrg.2016.142>
- Pickrell JK, Berisa T, Liu JZ, Séguérel L, Tung JY, Hinds DA (2016) Detection and interpretation of shared genetic influences on 42 human traits. *Nat Genet* 48:709–717. <https://doi.org/10.1038/ng.3570>
- Price AL et al (2008) Long-range LD can confound genome scans in admixed populations. *Am J Hum Genet* 83:132–135. <https://doi.org/10.1016/j.ajhg.2008.06.005>

- Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, Sullivan PF, Sklar P (2009) Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature* 460:748–752. <https://doi.org/10.1038/nature08185>
- Savage JE et al (2018) Genome-wide association meta-analysis in 269,867 individuals identifies new genetic and functional links to intelligence. *Nat Genet* 50:912–919. <https://doi.org/10.1038/s41588-018-0152-6>
- Schork AJ et al (2013) All SNPs are not created equal: genome-wide association studies reveal a consistent pattern of enrichment among functionally annotated SNPs. *PLoS Genet* 9:e1003449. <https://doi.org/10.1371/journal.pgen.1003449>
- Schork AJ, Wang Y, Thompson WK, Dale AM, Andreassen OA (2016) New statistical approaches exploit the polygenic architecture of schizophrenia—implications for the underlying neurobiology. *Curr Opin Neurobiol* 36:89–98. <https://doi.org/10.1016/j.conb.2015.10.008>
- Schwartzman A, Lin X (2011) The effect of correlation in false discovery rate estimation. *Biometrika* 98:199–214. <https://doi.org/10.1093/biomet/asq075>
- Shadrin AA et al (2018) Novel loci associated with attention-deficit/hyperactivity disorder are revealed by leveraging polygenic overlap with educational attainment. *J Am Acad Child Adolesc Psychiatry* 57:86–95. <https://doi.org/10.1016/j.jaac.2017.11.013>
- Shi H, Mancuso N, Spendlove S, Pasaniuc B (2017) Local genetic correlation gives insights into the shared genetic architecture of complex traits. *Am J Hum Genet* 101:737–751. <https://doi.org/10.1016/j.ajhg.2017.09.022>
- Sivakumaran S et al (2011) Abundant pleiotropy in human complex diseases and traits. *Am J Hum Genet* 89:607–618. <https://doi.org/10.1016/j.ajhg.2011.10.004>
- Smeland OB et al (2017a) Identification of genetic loci jointly influencing schizophrenia risk and the cognitive traits of verbal-numerical reasoning, reaction time, and general cognitive function. *JAMA Psychiatry* 74:1065–1075. <https://doi.org/10.1001/jamapsychiatry.2017.1986>
- Smeland OB et al (2017b) Identification of genetic loci shared between schizophrenia and the big five personality traits. *Sci Rep* 7:2222. <https://doi.org/10.1038/s41598-017-02346-3>
- Smeland OB et al (2018) Genetic overlap between schizophrenia and volumes of hippocampus, putamen, and intracranial volume indicates shared molecular genetic mechanisms. *Schizophr Bull* 44:854–864. <https://doi.org/10.1093/schbul/sbx148>
- Smeland OB et al (2019) Genome-wide analysis reveals extensive genetic overlap between schizophrenia, bipolar disorder, and intelligence. *Mol Psychiatry*. <https://doi.org/10.1038/s41380-018-0332-x>
- Smith GD, Ebrahim S (2003) ‘Mendelian randomization’: can genetic epidemiology contribute to understanding environmental determinants of disease? *Int J Epidemiol* 32:1–22
- Smoller JW, Andreassen OA, Edenberg HJ, Faraone SV, Glatt SJ, Kendler KS (2018) Psychiatric genetics and the structure of psychopathology. *Mol Psychiatry*. <https://doi.org/10.1038/s41380-017-0010-4>
- Sniekers S et al (2017) Genome-wide association meta-analysis of 78,308 individuals identifies new loci and genes influencing human intelligence. *Nat Genet* 49:1107–1112. <https://doi.org/10.1038/ng.3869>
- Solovieff N, Cotsapas C, Lee PH, Purcell SM, Smoller JW (2013) Pleiotropy in complex traits: challenges and strategies. *Nat Rev Genet* 14:483–495. <https://doi.org/10.1038/nrg3461>
- Stahl EA et al (2019) Genome-wide association study identifies 30 loci associated with bipolar disorder. *Nat Genet* 51:793–803. <https://doi.org/10.1038/s41588-019-0397-8>
- Sun L, Craiu RV, Paterson AD, Bull SB (2006) Stratified false discovery control for large-scale hypothesis testing with application to genome-wide association studies. *Genet Epidemiol* 30:519–530. <https://doi.org/10.1002/gepi.20164>
- Turley P et al (2018) Multi-trait analysis of genome-wide association summary statistics using MTAG. *Nat Genet* 50:229–237. <https://doi.org/10.1038/s41588-017-0009-4>
- van der Meer D et al (2018) Brain scans from 21,297 individuals reveal the genetic architecture of hippocampal subfield volumes. *Mol Psychiatry*. <https://doi.org/10.1038/s41380-018-0262-7>
- van der Sluis S, Posthuma D, Dolan CV (2013) TATES: efficient multivariate genotype-phenotype analysis for genome-wide association studies. *PLoS Genet* 9:e1003235. <https://doi.org/10.1371/journal.pgen.1003235>
- Visscher PM, Wray NR, Zhang Q, Sklar P, McCarthy MI, Brown MA, Yang J (2017) 10 years of GWAS discovery: biology, function, and translation. *Am J Hum Genet* 101:5–22. <https://doi.org/10.1016/j.ajhg.2017.06.005>
- Wang Y et al (2016a) Genetic overlap between multiple sclerosis and several cardiovascular disease risk factors. *Mult Scler* 22:1783–1793. <https://doi.org/10.1177/1352458516635873>
- Wang Y et al (2016b) Leveraging genomic annotations and pleiotropic enrichment for improved replication rates in schizophrenia GWAS. *PLoS Genet* 12:e1005803. <https://doi.org/10.1371/journal.pgen.1005803>
- Willer CJ, Li Y, Abecasis GR (2010) METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 26:2190–2191. <https://doi.org/10.1093/bioinformatics/btq340>
- Winsvold BS et al (2017) Shared genetic risk between migraine and coronary artery disease: A genome-wide analysis of common variants. *PLoS One* 12:e0185663. <https://doi.org/10.1371/journal.pone.0185663>
- Witoelar A et al (2017) Genome-wide pleiotropy between parkinson disease and autoimmune diseases. *JAMA Neurol* 74:780–792. <https://doi.org/10.1001/jamaneurol.2017.0469>
- Yokoyama JS et al (2016) Association between genetic traits for immune-mediated diseases and alzheimer disease. *JAMA neurology* 73:691–697. <https://doi.org/10.1001/jamaneurol.2016.0150>
- Yokoyama JS et al (2017) Shared genetic risk between corticobasal degeneration, progressive supranuclear palsy, and frontotemporal dementia. *Acta Neuropathol* 133:825–837. <https://doi.org/10.1007/s00401-017-1693-y>
- Yoo YJ, Pinnaduwa D, Waggott D, Bull SB, Sun L (2009) Genome-wide association analyses of North American rheumatoid arthritis consortium and Framingham heart study data utilizing genome-wide linkage results. *BMC Proc* 3(Suppl 7):S103
- Zhu Z et al (2018) Causal associations between risk factors and common diseases inferred from GWAS summary data. *Nat Commun* 9:224. <https://doi.org/10.1038/s41467-017-02317-2>
- Zuber V et al (2018) Identification of shared genetic variants between schizophrenia and lung cancer. *Sci Rep* 8:674. <https://doi.org/10.1038/s41598-017-16481-4>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.