

Renming Song · Zoran Vondraček

Potential theory of subordinate killed Brownian motion in a domain

Received: 4 April 2002 / Revised version: 1 July 2002 /
Published online: 19 December 2002 – © Springer-Verlag 2002

Abstract. Subordination of a killed Brownian motion in a bounded domain $D \subset \mathbb{R}^d$ via an $\alpha/2$ -stable subordinator gives a process Z_t whose infinitesimal generator is $-(-\Delta|_D)^{\alpha/2}$, the fractional power of the negative Dirichlet Laplacian. In this paper we study the properties of the process Z_t in a Lipschitz domain D by comparing the process with the rotationally invariant α -stable process killed upon exiting D . We show that these processes have comparable killing measures, prove the intrinsic ultracontractivity of the semigroup of Z_t , and, in the case when D is a bounded $C^{1,1}$ domain, obtain bounds on the Green function and the jumping kernel of Z_t .

1. Introduction

Let X_t be a d -dimensional Brownian motion in \mathbb{R}^d and let T_t be an $\alpha/2$ -stable subordinator starting at zero, $0 < \alpha < 2$. It is well known that $Y_t = X_{T_t}$ is a rotationally invariant α -stable process whose generator is $-(-\Delta)^{\alpha/2}$, the fractional power of the negative Laplacian. The potential theory corresponding to the process Y is the Riesz potential theory of order α .

Suppose that D is a domain in \mathbb{R}^d , that is, an open connected subset of \mathbb{R}^d . We can kill the process Y upon exiting D . The killed process Y^D has been extensively studied in the last five years and various deep properties have been obtained. For instance, when D is a bounded $C^{1,1}$ domain, sharp estimates on the Green function of Y^D were established in [5] and [15], while the intrinsic ultracontractivity of the semigroup corresponding to Y^D was proved in [4], [6] and [16].

R. Song: Department of Mathematics, University of Illinois, Urbana, IL 61801, USA. e-mail: rsong@math.uiuc.edu

Z. Vondraček: Department of Mathematics, University of Zagreb, Zagreb, Croatia. e-mail: vondra@math.hr

This work was completed while the authors were in the Research in Pairs program at the Mathematisches Forschungsinstitut Oberwolfach. We thank the Institute for the hospitality.

The research of the first author is supported in part by NSF Grant DMS-9803240.

The research of the second author is supported in part by MZT grant 037008 of the Republic of Croatia.

Mathematics Subject Classification (2000): Primary 60J45; Secondary 60J75, 31C25

Key words or phrases: Killed Brownian motions – Stable processes – Subordination – Fractional Laplacian

Let $\Delta|_D$ be the Dirichlet Laplacian in D . The fractional power $(-\Delta|_D)^{\alpha/2}$ of the negative Dirichlet Laplacian is a very useful object in analysis and partial differential equations, see, for instance, [22] and [18]. There is a Markov process Z corresponding to $(-\Delta|_D)^{\alpha/2}$ which can be obtained as follows: We first kill the Brownian motion X at τ_D , the first exit time of X from the domain D , and then we subordinate the killed Brownian motion using the $\alpha/2$ -stable subordinator T_t . Note that in comparison with Y^D the order of killing and subordination has been reversed. The difference between the processes Y^D and Z can be explained as follows: Look at a path of the Brownian motion in \mathbb{R}^d , and put a mark on it at all the times given by the subordinator T_t . In this way we observe a trajectory of the process Y . The corresponding trajectory of Z is given by all the marks on the Brownian path prior to τ_D . There is the first mark on the Brownian path following the exit time τ_D . If this mark happens to be in D , the process Y has not been killed yet, and the mark corresponds to a point on the trajectory of Y^D , but not to a point on the trajectory of Z . If, on the other hand, the first mark on the Brownian path following the exit time τ_D happens to be in D^c , then trajectories of Z and Y^D are equal.

Despite its importance, the process Z has not been studied much. In [12], a relation between the harmonic functions of Z and the classical harmonic functions in D was established. In [14] (see also [10]) the domain of the Dirichlet form of Z was identified when D is a bounded smooth domain and $\alpha \neq 1$.

In this paper we study the process Z and some of its potential-theoretic properties. One way to understand the process Z is to describe its killing and jumping measures. It turns out that, at least when D is Lipschitz, the killing measure is comparable with the killing measure of the process Y^D . This fact, shown in Sections 2 and 3 of the paper, follows from an analysis of the lifetimes of processes Z and Y^D . In order to do that, we have to give a precise description of both processes in terms of the underlying Brownian motion X_t and the subordinator T_t . The process Z , being a symmetric Markov process, has an associated Dirichlet form. By using the comparability of killing measures of Z and Y^D , we show that the corresponding Dirichlet forms are also comparable. This fact is then used in Section 4 to prove the intrinsic ultracontractivity of the semigroup corresponding to Z . As a consequence of this result we derive a lower bound on the Green function of Z in terms of the first eigenfunction of the Dirichlet Laplacian $\Delta|_D$. In the last section we derive upper bounds on the Green function of Z for $C^{1,1}$ domains. These bounds may not be sharp, but they show that the behaviors of the Green functions of Z and Y^D are very different. In the same vein we obtain bounds for the jump kernel of Z which confirm that the jump kernel vanishes near the boundary of D .

2. Subordinate killed Brownian motion

Let $X^1 = (\Omega^1, \mathcal{F}^1, \mathcal{F}_t^1, X_t^1, \theta_t^1, \mathbb{P}_x^1)$ be a d -dimensional Brownian motion in \mathbb{R}^d , and let $T^2 = (\Omega^2, \mathcal{G}^2, T_t^2, \mathbb{P}^2)$ be an $\alpha/2$ -stable subordinator starting at zero, $0 < \alpha < 2$. We will consider both processes on the product space $\Omega = \Omega^1 \times \Omega^2$. Thus we set $\mathcal{F} = \mathcal{F}^1 \times \mathcal{G}^2$, $\mathcal{F}_t = \mathcal{F}_t^1 \times \mathcal{G}^2$, and $\mathbb{P}_x = \mathbb{P}_x^1 \times \mathbb{P}^2$. Moreover, we define $X_t(\omega) = X_t^1(\omega^1)$, $T_t(\omega) = T_t^2(\omega^2)$, and $\theta_t(\omega) = \theta_t^1(\omega^1)$, where $\omega = (\omega^1, \omega^2) \in$

Ω . Then $X = (\Omega, \mathcal{F}, \mathcal{F}_t, X_t, \theta_t, \mathbb{P}_x)$ is a d -dimensional \mathcal{F}_t -Brownian motion, and $T = (\Omega, \mathcal{G}, T_t, \mathbb{P}_x)$ is an $\alpha/2$ -stable subordinator starting at zero, independent of X for every \mathbb{P}_x . From now on, all processes and random variables will be defined on Ω .

Let $A_t = \inf\{s > 0 : T_s \geq t\}$ be the inverse of T . Since (T_t) is strictly increasing, (A_t) is continuous. Further, $A_{T_t} = t$ and $T_{A_s-} \leq s \leq T_{A_s}$.

We define a process Y subordinate to X by $Y_t = X_{T_t}$. It is well known that Y is a rotationally invariant α -stable process in \mathbb{R}^d . If $\mu_t^{\alpha/2}$ is the distribution of T_t (i.e., $(\mu_t^{\alpha/2}, t \geq 0)$ is the one-sided $\alpha/2$ -stable convolution semigroup), and $(P_t, t \geq 0)$ the semigroup corresponding to the Brownian motion X , then for any nonnegative Borel function f on \mathbb{R}^d , $\mathbb{E}_x(f(Y_t)) = \mathbb{E}_x(f(X_{T_t})) = \mathbb{E}_x(\int_0^\infty f(X_s) \mu_t^{\alpha/2}(ds)) = \int_0^\infty P_s f(x) \mu_t^{\alpha/2}(ds)$.

Let $D \subset \mathbb{R}^d$ be a bounded domain, and let $\tau_D^Y = \inf\{t > 0 : Y_t \notin D\}$ be the exit time of Y from D . The process Y killed upon exiting D is defined by

$$Y_t^D = \begin{cases} Y_t, & t < \tau_D^Y \\ \partial, & t \geq \tau_D^Y \end{cases} = \begin{cases} X_{T_t}, & t < \tau_D^Y \\ \partial, & t \geq \tau_D^Y \end{cases}$$

where ∂ is an isolated point serving as a cemetery.

Let $\tau_D = \inf\{t > 0 : X_t \notin D\}$ be the exit time of X from D . The Brownian motion killed upon exiting D is defined as

$$X_t^D = \begin{cases} X_t, & t < \tau_D \\ \partial, & t \geq \tau_D \end{cases}$$

We define now the subordinate killed Brownian motion as the process obtained by subordinating X^D via the $\alpha/2$ -stable subordinator T_t . More precisely, let $Z_t = (X^D)_{T_t}, t \geq 0$. Then

$$Z_t = \begin{cases} X_{T_t}, & T_t < \tau_D \\ \partial, & T_t \geq \tau_D \end{cases} = \begin{cases} X_{T_t}, & t < A_{\tau_D} \\ \partial, & t \geq A_{\tau_D} \end{cases}$$

where the last equality follows from the fact $\{T_t < \tau_D\} = \{t < A_{\tau_D}\}$. Note that A_{τ_D} is the lifetime of the process Z . Moreover, it holds that $A_{\tau_D} \leq \tau_D^Y$. Indeed, if $s < A_{\tau_D}$, then $T_s < \tau_D$, implying that $Y_s = X_{T_s} \in D$. Hence, $s < \tau_D^Y$. Therefore, the lifetime of Z is less than or equal to the lifetime of Y^D .

On the next page is a very rough picture illustrating the differences between the processes Z and Y^D . In this picture the curve is a Brownian path, the points on the path marked by the little crosses, circles and squares represent a trajectory of Y , the points on the path marked by the little crosses and circles represent a trajectory of Y^D , and the points on the path marked by the little crosses represent a trajectory of Z .

We compare now the semigroups corresponding to Y^D and Z . For any nonnegative Borel function f on D , let

$$Q_t f(x) = \mathbb{E}_x[f(Y_t^D)] = \mathbb{E}_x[f(Y_t), t < \tau_D^Y] = \mathbb{E}_x[f(X_{T_t}), t < \tau_D^Y]$$

$$R_t f(x) = \mathbb{E}_x[f(Z_t)] = \mathbb{E}_x[f(X^D)_{T_t}] = \mathbb{E}_x[f(X_{T_t}), t < A_{\tau_D}]$$




Fig. 1. trajectories of Z and Y^D .

Since $A_{\tau_D} \leq \tau_D^Y$, it follows that $R_t f(x) \leq Q_t f(x)$ for all $t \geq 0$.

The following result will be needed in order to compare the killing functions of the processes Z and Y^D .

Proposition 2.1. *Suppose that there exists $C \in (0, 1)$ such that $\mathbb{P}_x(X_t \in D) \leq C$ for every $t > 0$ and every $x \in \partial D$. Then*

$$(1 - C)(1 - R_t 1(x)) \leq 1 - Q_t 1(x) \leq 1 - R_t 1(x) \tag{2.1}$$

for every $t > 0$ and every $x \in D$.

Proof. Let $\tau_D^1(\omega^1) = \inf\{t > 0, X_t^1(\omega^1) \notin D\}$, i.e., $\tau_D^1(\omega^1) = \tau_D(\omega)$. Then $\mathcal{F}_{\tau_D^1}^1 \times \mathcal{G}^2 \subset \mathcal{F}_{\tau_D}$. Indeed, for $A_1 \in \mathcal{F}_{\tau_D^1}^1$ and $A_2 \in \mathcal{G}^2$, $(A_1 \times A_2) \cap \{\tau_D \leq t\} = (A_1 \cap \{\tau_D^1 \leq t\}) \times A_2 \in \mathcal{F}_t^1 \times \mathcal{G} = \mathcal{F}_t$. Thus, $A_1 \times A_2 \in \mathcal{F}_{\tau_D}$. Since such sets generate $\mathcal{F}_{\tau_D^1}^1 \times \mathcal{G}^2$, the claim follows.

We want to show that $T_{A_{\tau_D}}$ is $\mathcal{F}_{\tau_D^1}^1 \times \mathcal{G}^2$ -measurable. Note first that τ_D and $T_t, t \geq 0$, are $\mathcal{F}_{\tau_D^1}^1 \times \mathcal{G}^2$ -measurable. Therefore, $\{T_t < \tau_D\} \in \mathcal{F}_{\tau_D^1}^1 \times \mathcal{G}^2$. Since $\{A_{\tau_D} > t\} = \{T_t < \tau_D\}$, it follows that $\{A_{\tau_D} > t\}$ is $\mathcal{F}_{\tau_D^1}^1 \times \mathcal{G}^2$ -measurable. Clearly, A_s is $\mathcal{F}_0 \times \mathcal{G}^2 \subset \mathcal{F}_{\tau_D^1}^1 \times \mathcal{G}^2$ -measurable. Therefore, $\{T_{A_{\tau_D}} \geq s\} = \{A_s \geq A_{\tau_D}\} \in \mathcal{F}_{\tau_D^1}^1 \times \mathcal{G}^2$.

For any nonnegative Borel function f on \mathbb{R}^d , let $N_t(x, f) = \mathbb{E}_x(f(X_t))$. Since τ_D and $T_{A_{\tau_D}}$ are $\mathcal{F}_{\tau_D^1}^1 \times \mathcal{G}^2$ -measurable, and $T_{A_{\tau_D}} = \tau_D + (T_{A_{\tau_D}} - \tau_D)$, by an extended version of the strong Markov property (see [2], pp. 43–44),

$$\mathbb{E}_x[1_D(X_{T_{A\tau_D}})|\mathcal{F}_{\tau_D}] = \mathbb{E}_x[1_D(X_{\tau_D+(T_{A\tau_D}-\tau_D)})|\mathcal{F}_{\tau_D}] = N_{T_{A\tau_D}-\tau_D}(X_{\tau_D}, 1_D) \text{ a.s.} \tag{2.2}$$

By using the assumption of the proposition, we get $N_t(y, 1_D) = \mathbb{P}_y(X_t \in D) \leq C$ for every $t > 0$ and every $y \in \partial D$. From (2.2) we obtain that $\mathbb{P}_x(X_{T_{A\tau_D}} \in D|\mathcal{F}_{\tau_D}) \leq C$ a.s. for every $x \in D$. Since $\mathcal{F}_{\tau_D}^1 \times \mathcal{G}^2 \subset \mathcal{F}_{\tau_D}$, it follows that $\mathbb{P}_x(X_{T_{A\tau_D}}|\mathcal{F}_{\tau_D}^1 \times \mathcal{G}^2) \leq C$ a.s. Further,

$$\begin{aligned} \mathbb{P}_x(A_{\tau_D} \leq t < \tau_D^Y) &\leq \mathbb{P}_x(A_{\tau_D} \leq t, A_{\tau_D} < \tau_D^Y) \\ &= \mathbb{P}_x(A_{\tau_D} \leq t, X_{T_{A\tau_D}} \in D) \\ &= \mathbb{P}_x[\mathbb{P}_x(A_{\tau_D} \leq t, X_{T_{A\tau_D}} \in D|\mathcal{F}_{\tau_D}^1 \times \mathcal{G}^2)] \\ &= \mathbb{E}_x[1_{(A_{\tau_D} \leq t)}\mathbb{P}_x(X_{T_{A\tau_D}} \in D|\mathcal{F}_{\tau_D}^1 \times \mathcal{G}^2)] \\ &\leq C\mathbb{P}_x(A_{\tau_D} \leq t). \end{aligned}$$

It follows that

$$\begin{aligned} \mathbb{P}_x(A_{\tau_D} \leq t) &= \mathbb{P}_x(\tau_D^Y \leq t) + \mathbb{P}_x(A_{\tau_D} \leq t < \tau_D^Y) \\ &\leq \mathbb{P}_x(\tau_D^Y \leq t) + C\mathbb{P}_x(A_{\tau_D} \leq t), \end{aligned}$$

hence

$$\begin{aligned} \mathbb{P}_x(A_{\tau_D} \leq t) &= \mathbb{P}_x(\tau_D^Y \leq t) + \mathbb{P}_x(A_{\tau_D} \leq t < \tau_D^Y) \\ &\leq \mathbb{P}_x(\tau_D^Y \leq t) + C\mathbb{P}_x(A_{\tau_D} \leq t). \end{aligned}$$

Since $\mathbb{P}_x(A_{\tau_D} \leq t) = 1 - R_t 1(x)$ and $\mathbb{P}_x(\tau_D^Y \leq t) = 1 - Q_t 1(x)$, (2.1) follows. □

A domain $D \subset \mathbb{R}^d$ is said to satisfy an exterior cone condition if there exist a cone K with vertex at the origin and a positive constant r_0 , such that for each point $x \in \partial D$, there exist a translation and a rotation taking the cone K into a cone K_x with the vertex at x such that

$$K_x \cap B(x, r_0) \subset D^c \cap B(x, r_0).$$

Here $B(x, r_0)$ denotes the ball of radius r_0 centered at x . We show now that the condition in Proposition 2.1 is true for a bounded domain $D \subset \mathbb{R}^d$ satisfying an exterior cone condition. Let $K_x(r_0) = K_x \cap B(x, r_0)$ and $K(r_0) = K \cap B(0, r_0)$. Then we have for each $x \in \partial D$,

$$\mathbb{P}_x(X_t \notin D) \geq \mathbb{P}_x(X_t \in K_x(r_0)) = \mathbb{P}_0(X_t \in K(r_0)).$$

By scaling,

$$\mathbb{P}_0(X_t \in K(r_0)) = \mathbb{P}_0(X_1 \in \frac{1}{\sqrt{t}}K(r_0)) \geq \mathbb{P}_0(X_1 \in K(r_0)) =: C_1 \in (0, 1)$$

for every $t \in (0, 1]$, where for any $\rho > 0$, $\rho K(r_0)$ is defined to be the set $\{\rho x : x \in K(r_0)\}$. The last two displays show that $\mathbb{P}_x(X_t \notin D) \geq C_1$, for every $t \in (0, 1]$ and every $x \in \partial D$. Since D is bounded, there exists $R > 0$ such that for every $x \in \partial D$, $D \subset B(x, R)$. Hence,

$$\mathbb{P}_x(X_t \notin D) \geq \mathbb{P}_0(|X_t| > R) \geq \mathbb{P}_0(|X_1| > R) =: C_2 \in (0, 1)$$

for every $t \geq 1$ and every $x \in \partial D$. Let $C = 1 - \min\{C_1, C_2\}$. Then $C \in (0, 1)$ and $\mathbb{P}_x(X_t \in D) \leq C$ for every $t > 0$ and every $x \in \partial D$.

It is well known (see [4], for instance) that the transition semigroup Q_t corresponding to the killed stable process has a density with respect to the Lebesgue measure. Let $q(t, x, y)$ be this density. Let $r(t, x, y)$ be the density of R_t and let $p^D(t, x, y)$ be the transition density of the killed Brownian motion X^D . The density $r(t, x, y)$ is given by the formula

$$r(t, x, y) = \int_0^\infty p^D(s, x, y) \mu_t^{\alpha/2}(ds), \tag{2.3}$$

where $(\mu_t^{\alpha/2}, t \geq 0)$ is the one-sided $\alpha/2$ -stable convolution semigroup. Let $G_D(x, y)$ and $G_D^Y(x, y)$ denote the Green functions of Z and Y^D respectively. The Green function of Z is given by

$$G_D(x, y) = \int_0^\infty r(t, x, y) dt = \frac{1}{\Gamma(\alpha/2)} \int_0^\infty p^D(t, x, y) t^{\alpha/2-1} dt. \tag{2.4}$$

Proposition 2.2. *Let D be a bounded domain in \mathbb{R}^d .*

- (i) *The transition density $r(t, x, y)$ of Z is jointly continuous in (x, y) for each fixed t . Further, $r(t, x, y) \leq q(t, x, y)$ for all $t > 0$ and all $(x, y) \in D \times D$.*
- (ii) *When $d \geq 2$ or $\alpha \leq 1 = d$, the Green function $G_D(x, y)$ is finite and continuous on $D \times D \setminus \{(x, x), x \in D\}$. When $\alpha > 1 = d$, the Green function $G_D(x, y)$ is finite and continuous on $D \times D$. Further, $G_D(x, y) \leq G_D^Y(x, y)$ on $D \times D$.*

Proof. (i) Note that $p^D(s, x, y) \leq (2\pi s)^{-d/2} \exp\{-|x - y|^2/2s\} \leq (2\pi s)^{-d/2}$ for all $x, y \in D$. It follows from the asymptotic behavior near zero of the density of $\mu_t^{\alpha/2}$ given in [20] that the integral $\int_0^\infty s^{-d/2} \mu_t^{\alpha/2}(ds)$ is finite. So the continuity of $r(t, \cdot, \cdot)$ follows from the dominated convergence theorem. Since $R_t f(x) \leq Q_t f(x)$ for every $x \in D$ and every nonnegative Borel function f , we get $r(t, x, y) \leq q(t, x, y)$ for all $y \in D \setminus N(x)$ with $N(x)$ having zero Lebesgue measure. By continuity, the inequality holds for all $x, y \in D$.

(ii) The fact that $G_D(x, y) \leq G_D^Y(x, y)$ follows immediately from $r(t, x, y) \leq q(t, x, y)$. We now prove the continuity of G_D by treating three cases separately. (a) The case when $d \geq 2$ or when $\alpha < 1 = d$. Let $x, y \in D$, $|x - y| > 2\eta > 0$. Let (x_n, y_n) be a sequence in $D \times D$ converging to (x, y) such that $|x_n - y_n| > \eta$. Note that,

$$\begin{aligned} p^D(t, x_n, y_n) t^{-1+\alpha/2} &\leq (2\pi t)^{-d/2} \exp\{-|x_n - y_n|^2/2t\} t^{-1+\alpha/2} \\ &\leq c_1 t^{-d/2+\alpha/2-1} \exp\{-\eta^2/2t\} \end{aligned}$$

which is integrable on $(0, \infty)$. The continuity now follows from the dominated convergence theorem. (b) The case when $\alpha = 1 = d$. Let $x, y \in D, |x - y| > 2\eta > 0$. Let (x_n, y_n) be a sequence in $D \times D$ converging to (x, y) such that $|x_n - y_n| > \eta$. Using the intrinsic ultracontractivity of the killed Brownian semigroup on a bounded interval and Theorem 4.2.5 of [8], we know that there exists a $T > 0$ such that for any $t \geq T$,

$$p^D(t, x, y) \leq \frac{3}{2} e^{-\lambda_0 t} \phi_0(x)\phi_0(y), \quad x, y \in D,$$

where $-\lambda_0 < 0$ and ϕ_0 are the first eigenvalue and eigenfunction of the Dirichlet Laplacian in D respectively. Thus in this case, the functions $p^D(t, x_n, y_n)t^{\alpha/2-1} = p^D(t, x_n, y_n)t^{-1/2}$ is dominated by the function

$$g(t) = \begin{cases} c_1 t^{-1} \exp\{-\eta^2/2t\}, & t \leq T \\ c_2 t^{-1} e^{-\lambda_0 t}, & t \geq T \end{cases}$$

which is integrable on $(0, \infty)$. Now we can repeat the argument in the first case to arrive at the claimed continuity. (c) The case when $\alpha > 1 = d$. In this case, the family of functions $\{p^D(t, \cdot, \cdot)t^{\alpha/2-1} : x, y \in D\}$ is dominated by the function

$$h(t) = \begin{cases} c_1 t^{-3/2+\alpha/2}, & t \leq T \\ c_2 t^{-3/2+\alpha/2} e^{-\lambda_0 t}, & t \geq T \end{cases}$$

which is integrable on $(0, \infty)$. The continuity now follows from the dominated convergence theorem. □

3. The Dirichlet form of the subordinate killed Brownian motion

Recall that Y is a rotationally invariant α -stable process in \mathbb{R}^d with $\alpha \in (0, 2)$. It is well known that the Dirichlet form $(\mathcal{E}^Y, \mathcal{F})$ associated with Y is given by

$$\mathcal{E}^Y(u, v) = \frac{1}{2} A(d, -\alpha) \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{d+\alpha}} dx dy$$

$$\mathcal{F} = \left\{ u \in L^2(\mathbb{R}^d) : \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(u(x) - u(y))^2}{|x - y|^{d+\alpha}} dx dy < \infty \right\},$$

where

$$A(d, -\alpha) = \frac{\alpha \Gamma(\frac{d+\alpha}{2})}{2^{1-\alpha} \pi^{d/2} \Gamma(1 - \frac{\alpha}{2})}.$$

It follows from Remark 4 in Section 2.5.1 of [21] that \mathcal{F} is the same as the space $W^{\alpha/2,2}(\mathbb{R}^d)$.

Recall that, for any $s \in \mathbb{R}$, the classical Bessel potential space $H^s(\mathbb{R}^d)$ is defined to be

$$H^s(\mathbb{R}^d) = \{u \in S'(\mathbb{R}^d) : \int_{\mathbb{R}^d} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi < \infty\},$$

where $S'(\mathbb{R}^d)$ stands for the space of tempered distributions on \mathbb{R}^d and \hat{u} stands for the Fourier transform of u . Using Fourier analysis, one can easily show (cf.

Example 1.4.1 of [11]) that the spaces $W^{\alpha/2,2}(\mathbb{R}^d)$ and $H^{\alpha/2}(\mathbb{R}^d)$ are the same. Hence we have $\mathcal{F} = W^{\alpha/2,2}(\mathbb{R}^d) = H^{\alpha/2}(\mathbb{R}^d)$.

In this section we assume that D is a bounded domain in \mathbb{R}^d . The Dirichlet space on $L^2(D, dx)$ of the killed rotationally invariant α -stable process Y^D is $(\mathcal{E}^Y, H_0^{\alpha/2}(D))$ (cf. Theorem 4.4.3 of [11]), where

$$H_0^{\alpha/2}(D) = \{f \in H^{\alpha/2}(\mathbb{R}^d) : f = 0 \text{ q.e. on } D^c\}.$$

Here q.e. is the abbreviation for quasi-everywhere with respect to the Riesz capacity determined by $(\mathcal{E}^Y, W^{\alpha/2,2}(\mathbb{R}^d))$ (cf. [11]). The space $H_0^{\alpha/2}(D)$ can also be characterized as the \mathcal{E}^Y -closure of $C_0^\infty(D)$, the space of smooth functions with compact support in D . For $u \in H_0^{\alpha/2}(D)$,

$$\begin{aligned} \mathcal{E}^Y(u, v) &= \int_D \int_D (u(x) - u(y))(v(x) - v(y))J^Y(x, y) dx dy \\ &\quad + \int_D u(x)v(x)\kappa^Y(x)dx, \end{aligned}$$

where

$$J^Y(x, y) = \frac{1}{2}A(d, -\alpha)|x - y|^{-(d+\alpha)} \tag{3.1}$$

$$\kappa^Y(x) = A(d, -\alpha) \int_{D^c} \frac{1}{|x - y|^{d+\alpha}} dy \tag{3.2}$$

are the densities of the jumping and killing measures of Y^D .

Recall that Z is the process obtained by subordinating the killed Brownian motion on D with the one-sided $\alpha/2$ -stable process. Z is a symmetric Markov process and so there is a Dirichlet form $(\mathcal{E}, D(\mathcal{E}))$ associated with Z . It follows further from Theorem 1.18.10 of [21] that the domain $D(\mathcal{E})$ of \mathcal{E} is the complex interpolation space $[L^2(D), H_0^1(D)]_{\alpha/2}$. It follows from Proposition 2.2 of [7] that, when D is a bounded Lipschitz domain, $[L^2(D), H_0^1(D)]_{\alpha/2} = H_0^{\alpha/2}(D)$. Recall that Hilbert spaces are identified if they coincide in the set theoretical sense and if they have equivalent norms. Therefore, there exists a constant C such that for any $u \in H_0^{\alpha/2}(D)$,

$$C^{-1}(\mathcal{E}^Y(u, u) + (u, u)) \leq \mathcal{E}(u, u) + (u, u) \leq C(\mathcal{E}^Y(u, u) + (u, u)).$$

One immediate consequence of the comparability above is that for a Borel subset A of D , A is polar for Z is equivalent to that A is polar for the killed rotationally invariant α -stable process Y^D , which in turn is equivalent to that A is polar for the rotationally invariant α -stable process Y .

Let P_t^D be the transition semigroup corresponding to the Brownian motion killed upon exiting D and recall that the corresponding transition density is denoted by $p^D(t, x, y)$. It follows from [3] and [17] (see also [13]) that the jumping

measure $J(dx, dy)$ and the killing measure $\kappa(dx)$ of the process Z have densities $J(x, y)$ and $\kappa(x)$ given by the following formulae respectively:

$$J(x, y) = \int_0^\infty p^D(t, x, y) \nu(dt) \tag{3.3}$$

$$\kappa(x) = \int_0^\infty (1 - P_t^D 1(x)) \nu(dt) \tag{3.4}$$

Here

$$\nu(dt) = \frac{\alpha/2}{\Gamma(1 - \alpha/2)} t^{-\alpha/2-1} dt$$

is the Lévy measure of the $\alpha/2$ -stable subordinator.

It is easy to see from (3.3) that $J(x, y) \leq J^Y(x, y)$ for every $x, y \in D$. Now we are going to compare $\kappa(x)$ with $\kappa^Y(x)$. To do that we are going to use the following simple result.

Lemma 3.1. *Let (X_t, \mathbb{P}_x) be a d -dimensional Brownian motion, and let τ_D be the exit time of X from D . Then*

$$\kappa(x) = \frac{1}{\Gamma(1 - \alpha/2)} \mathbb{E}_x(\tau_D^{-\alpha/2}) \tag{3.5}$$

for every $x \in \mathbb{R}^d$.

Proof. Let F denote the \mathbb{P}_x -distribution function of τ_D . Note that $1 - P_t^D 1(x) = \mathbb{P}_x(\tau_D \leq t) = F(t)$. By using (3.4)

$$\begin{aligned} \kappa(x) &= \frac{\alpha/2}{\Gamma(1 - \alpha/2)} \int_0^\infty F(t) t^{-\alpha/2-1} dt \\ &= \frac{\alpha/2}{\Gamma(1 - \alpha/2)} \int_0^\infty \int_s^\infty t^{-\alpha/2-1} dt dF(s) \\ &= \frac{1}{\Gamma(1 - \alpha/2)} \int_0^\infty s^{-\alpha/2} dF(s) = \frac{1}{\Gamma(1 - \alpha/2)} \mathbb{E}_x(\tau_D^{-\alpha/2}) \end{aligned}$$

□

It was proved in [12] that $x \rightarrow \mathbb{E}_x(\tau_D^{-\alpha/2})$ is continuous, hence κ is a continuous function. We will use this fact in the next result.

Proposition 3.2. *Suppose that there exists $C \in (0, 1)$ such that $\mathbb{P}_x(X_t \in D) \leq C$ for every $t > 0$ and every $x \in \partial D$. Then*

$$(1 - C)\kappa(x) \leq \kappa^Y(x) \leq \kappa(x), \text{ for every } x \in D. \tag{3.6}$$

Proof. By the proof of Lemma 4.5.2 in [11], there exists a sequence $t_n \downarrow 0$ such that

$$\begin{aligned} \lim_{t_n \rightarrow 0} \frac{1}{t_n} \int_D f(x)(1 - R_{t_n} 1(x)) dx &= \int_D f(x)\kappa(x) dx \\ \lim_{t_n \rightarrow 0} \frac{1}{t_n} \int_D f(x)(1 - Q_{t_n} 1(x)) dx &= \int_D f(x)\kappa^Y(x) dx \end{aligned}$$

for every $f \in C_0(D)$. By Proposition 2.1 this implies that

$$\int_D f(x)(1 - C)\kappa(x) dx \leq \int_D f(x)\kappa^Y(x) dx \leq \int_D f(x)\kappa(x) dx ,$$

for every nonnegative $f \in C_0(D)$. Since both κ and κ^Y are continuous, the last relation implies that

$$(1 - C)\kappa(x) \leq \kappa^Y(x) \leq \kappa(x) , \quad x \in D .$$

□

Remark 3.3. Let $\delta(x)$ be the distance between x and ∂D . When D is a bounded Lipschitz domain, it follows easily from (3.2) that there exists a positive constant C_1 such that

$$C_1^{-1}(\delta(x))^{-\alpha} \leq \kappa^Y(x) \leq C_1(\delta(x))^{-\alpha} .$$

By using this and Proposition 3.2 it follows that there exists a constant C_2 such that

$$C_2^{-1}(\delta(x))^{-\alpha} \leq \kappa(x) \leq C_2(\delta(x))^{-\alpha} .$$

4. Intrinsic ultracontractivity

In this section we assume that D is a bounded Lipschitz domain and Z is the subordinate killed Brownian motion on D . The generator of Z is $(-\Delta|_D)^{\alpha/2}$, where $\Delta|_D$ is the Dirichlet Laplacian in D . It is well known that if $\{-\lambda_k, k = 0, 1, \dots\}$ are the eigenvalues of $\Delta|_D$ written in decreasing order and each repeated according to its multiplicity, and if $\{\phi_k, k = 0, 1, \dots\}$ are the corresponding eigenfunctions, then $\{-\lambda_k^{\alpha/2}, k = 0, 1, \dots\}$ are the eigenvalues of $(-\Delta|_D)^{\alpha/2}$ written in decreasing order and each repeated according to its multiplicity, and $\{\phi_k, k = 0, 1, \dots\}$ are the corresponding eigenfunctions.

Similar to Theorem 4.1 of [4], we have the following result.

Theorem 4.1. *For any $\eta > 0$ and $f \in H_0^{\alpha/2}(D) \cap L^\infty(D, dx)$, we have*

$$\int_D f^2 \log |f| dx \leq \eta \mathcal{E}(f, f) + \beta(\eta) \|f\|_2^2 + \|f\|_2^2 \log \|f\|_2 ,$$

with

$$\beta(\eta) = -\frac{d}{2\alpha} \log \eta + c$$

for some constant $c > 0$.

Proof. It follows from Proposition 2.2 that $r(t, x, y) \leq q(t, x, y)$, hence there exists a $c > 0$ such that $r(t, x, y) \leq ct^{-d/\alpha}$. Now we can repeat the proof of Theorem 4.1 of [4] to arrive at the conclusion. □

The following lemma appears on p.71 of [14]. The key ingredient in the proof there is an inequality (inequality (4.1) of [14]) proved in [19]. We include an elementary proof based on the behavior of the killing function of Z .

Lemma 4.2. *There exists a constant $C_1 > 0$ such that*

$$C_1^{-1} \mathcal{E}^Y(u, u) \leq \mathcal{E}(u, u) \leq C_1 \mathcal{E}^Y(u, u), \quad u \in H_0^{\alpha/2}(D).$$

Proof. Recall that the killing measures of Z and Y^D have densities κ and κ^Y respectively, which are both of the order $\delta(x)^{-\alpha}$. This implies that there is a constant c_1 such that

$$\int_D u^2(x) dx \leq c_1 \int_D u^2(x) \kappa(x) dx \tag{4.1}$$

From the last section we know that there exists a constant $c_2 > 0$ such that

$$c_2^{-1} (\mathcal{E}^Y(u, u) + (u, u)) \leq \mathcal{E}(u, u) + (u, u) \leq c_2 (\mathcal{E}^Y(u, u) + (u, u)), \quad u \in H_0^{\alpha/2}(D).$$

Therefore the 1-norms are equivalent to the 0-norms for both forms. Thus there is a constant C_1 such that

$$C_1^{-1} \mathcal{E}^Y(u, u) \leq \mathcal{E}(u, u) \leq C_1 \mathcal{E}^Y(u, u), \quad u \in H_0^{\alpha/2}(D).$$

□

Recall that for any domain D in \mathbb{R}^d , the quasi-hyperbolic distance between any two points x_1 and x_2 in D is defined by

$$\rho_D(x_1, x_2) = \inf_{\gamma} \int_{\gamma} \frac{ds}{\delta(x)}$$

where the infimum is taken over all rectifiable curves γ joining x_1 to x_2 in D and $\delta(x)$ is the Euclidean distance between x and ∂D . Fix a point $x_0 \in D$ which we call the center of D and we may assume without loss of generality that $\delta(x_0) = 1$.

Lemma 4.3. *There is a constant $C_2 = C_2(D) > 0$ such that for any $\beta > 0$*

$$\int_D (\rho_D(x_0, x))^{\beta} u^2(x) dx \leq C_2 \mathcal{E}(u, u), \quad u \in H_0^{\alpha/2}(D)$$

Proof. It follows from Lemma 3.2 of [6] that there is a constant $c = c(D) > 0$ such that for any $\beta > 0$

$$\int_D (\rho_D(x_0, x))^{\beta} u^2(x) dx \leq c \mathcal{E}^Y(u, u), \quad u \in H_0^{\alpha/2}(D).$$

Now the result follows from Lemma 4.2. □

Repeating the argument of Theorem 3.3 of [6](see also [1]), we get

Theorem 4.4. *For any $\varepsilon > 0$ and any $\sigma > 0$ we have*

$$\int_D f^2 \log \frac{1}{\phi_0} dx \leq \varepsilon \mathcal{E}(f, f) + \beta(\varepsilon) \|f\|_2^2, \quad f \in H_0^{\alpha/2}(D)$$

with

$$\beta(\varepsilon) = C_3 \varepsilon^{-\sigma} + C_4$$

for some positive constants C_3 and C_4 .

Combining Theorems 4.1 and 4.4 we get

Theorem 4.5. *For any $\varepsilon > 0$ and any $\sigma > 0$ we have*

$$\int_D f^2 \log \frac{|f|}{\varphi_0} dx \leq \eta \mathcal{E}(f, f) + \beta(\eta) \|f\|_2^2 + \|f\|_2^2 \log \|f\|_2$$

for all $f \in H_0^{\alpha/2}(D) \cap L^\infty(D, dx)$, with

$$\beta(\varepsilon) = -\frac{d}{2\alpha} \log \varepsilon + C_5 \varepsilon^{-\sigma} + C_6$$

for some positive constants C_5 and C_6 .

Using this and Corollary 2.2.8 of [8] we immediately get

Theorem 4.6. *The semigroup corresponding to the subordinate killed Brownian motion Z is intrinsic ultracontractive.*

Here is an immediate corollary of the intrinsic ultracontractivity. Recall that the Green function of the process Z is given by the formula (2.4).

Corollary 4.7. *There exists a constant C_8 such that for all $x, y \in D$,*

$$\begin{aligned} G_D(x, y) &\geq C_8 \phi_0(x) \phi_0(y), \\ J(x, y) &\geq C_8 \phi_0(x) \phi_0(y). \end{aligned}$$

Proof. The first inequality follows immediately from the intrinsic ultracontractivity and Theorem 4.2.5 of [8]. Now we show the second inequality. Since the semigroup of the killed Brownian motion in D is intrinsic ultracontractive, Theorem 4.2.5 of [8] implies that there exists $T > 1$ such that for all $t \geq T$,

$$p^D(t, x, y) \geq \frac{1}{2} e^{-\lambda_0 t} \phi_0(x) \phi_0(y), \quad xy \in D.$$

Thus

$$\begin{aligned} J(x, y) &= c_1 \int_0^\infty p^D(t, x, y) t^{-\alpha/2-1} dt \\ &\geq \frac{c_1}{2} \int_T^\infty e^{-\lambda_0 t} \phi_0(x) \phi_0(y) dt \\ &= c_2 \phi_0(x) \phi_0(y). \end{aligned}$$

□

Note that these lower bounds of G_D and J are of no use when x, y are away from the boundary. The next result gives lower bound when x and y are away from the boundary and it does not need the Lipschitz assumption.

Proposition 4.8. *For any bounded domain D in \mathbb{R}^d , there exists a constant $C_9 = C_9(\alpha, d)$ such that if $x, y \in D$ satisfy $|x - y| \leq \max\{\delta(x)/2, \delta(y)/2\}$, then*

$$G_D(x, y) \geq C_9 |x - y|^{\alpha-d}, \tag{4.2}$$

$$J(x, y) \geq C_9 |x - y|^{-\alpha-d}. \tag{4.3}$$

Proof. We prove the first inequality. The second is proved in the same way. Let $x, y \in D$ such that $|x - y| \leq \max\{\delta(x)/2, \delta(y)/2\}$. By using (2.4) and the formula for the transition density of the killed Brownian motion X^D , we get that

$$G_D(x, y) = \frac{1}{\Gamma(\alpha/2)} \int_0^\infty p(s, x, y) s^{\alpha/2-1} ds - \frac{1}{\Gamma(\alpha/2)} \mathbb{E}_x \int_{\tau_D}^\infty p(s, X_{\tau_D}, y) s^{\alpha/2-1} ds, \tag{4.4}$$

where $p(s, x, y)$ denotes the transition density of the Brownian motion X . Since $|X_{\tau_D} - y| \geq \delta(y)$ for each $y \in D$, we obtain the estimate

$$\begin{aligned} & \frac{1}{\Gamma(\alpha/2)} \mathbb{E}_x \int_{\tau_D}^\infty p(s, X_{\tau_D}, y) s^{\alpha/2-1} ds \\ & \leq \frac{(2\pi)^{-d/2}}{\Gamma(\alpha/2)} \int_0^\infty s^{-d/2+\alpha/2-1} \exp\{-\delta(y)^2/2s\} ds \\ & \leq c_1 \delta(y)^{\alpha-d} \\ & \leq c_1 2^{\alpha-d} |x - y|^{\alpha-d}. \end{aligned}$$

The estimate (4.2) follows from (4.4) and the last display. □

5. Upper bounds on the Green function and the jumping kernel

For any bounded domain D in R^d , we have seen that

$$G_D(x, y) \leq G_D^Y(x, y), \quad J(x, y) \leq J^Y(x, y), \quad x, y \in D.$$

Recall that G_D^Y and J^Y are the Green function and jumping function of Y^D respectively. These estimates are not useful near the boundary of D . Now we are going to derive estimates that are useful near the boundary when D is a bounded $C^{1,1}$ domain.

Theorem 5.1. *Suppose that D is a bounded $C^{1,1}$ domain in R^d . Then there exists a constant C_1 such that for all $x, y \in D$,*

$$\begin{aligned} G_D(x, y) & \leq C_1 \frac{\phi_0(x)\phi_0(y)}{|x - y|^{d+2-\alpha}}, \\ J(x, y) & \leq C_1 \frac{\phi_0(x)\phi_0(y)}{|x - y|^{d+2+\alpha}}. \end{aligned}$$

Proof. The proof of these two inequalities are very similar. We only give the proof of the first. It is well known that when D is a bounded $C^{1,1}$ domain, there exists a constant c_1 such that

$$c_1^{-1} \delta(x) \leq \phi_0(x) \leq c_1 \delta(x), \quad x \in D.$$

Now we can repeat the proof of Theorem 4.6.9 of [8] to get that the density p^D of the killed Brownian motion on D satisfies the following estimate

$$p^D(t, x, y) \leq c_2 t^{-(d+2)/2} \phi_0(x) \phi_0(y) e^{-\frac{|x-y|^2}{6t}}, \quad t > 0, x, y \in D,$$

where c_2 is some constant independent of $t, x,$ and y . Now using (2.4) we get that

$$\begin{aligned} G_D(x, y) &\leq c_2 \phi_0(x) \phi_0(y) \int_0^\infty t^{-(d+2)/2} e^{-\frac{|x-y|^2}{6t}} t^{\alpha/2-1} dt \\ &\leq c_3 \frac{\phi_0(x) \phi_0(y)}{|x-y|^{d+2-\alpha}}. \end{aligned}$$

□

Remark 5.2. If we only assume that D is a bounded Lipschitz domain, then we can get a similar upper bound for G_D with $d + 2$ replaced by some number $\mu \geq d$, where μ depends on the Lipschitz characteristics of D .

Remark 5.3. The estimates in the theorem above can also be written as

$$\begin{aligned} G_D(x, y) &\leq C_2 \frac{\delta(x) \delta(y)}{|x-y|^{d+2-\alpha}}, \quad x, y \in D \\ J(x, y) &\leq C_2 \frac{\delta(x) \delta(y)}{|x-y|^{d+2+\alpha}}, \quad x, y \in D, \end{aligned}$$

for some positive constant C_2 .

Summarizing our estimates on the Green function and the jumping kernel, we have the following:

Theorem 5.4. *Suppose that D is a bounded $C^{1,1}$ domain in R^d . Then there exist positive constants C_3 and C_4 such that for all $x, y \in D$,*

$$\begin{aligned} C_3 \delta(x) \delta(y) \leq G_D(x, y) &\leq C_4 \min\left(\frac{1}{|x-y|^{d-\alpha}}, \frac{\delta(x) \delta(y)}{|x-y|^{d+2-\alpha}}\right), \\ C_3 \delta(x) \delta(y) \leq J(x, y) &\leq C_4 \min\left(\frac{1}{|x-y|^{d+\alpha}}, \frac{\delta(x) \delta(y)}{|x-y|^{d+2+\alpha}}\right) \end{aligned}$$

Comparing the estimates on the Green function of Z with the estimates on the Green function of Y^D obtained in [5] and [15], we see that their boundary behaviors are different.

Acknowledgements. We thank H. Šikić for several useful discussions and Z.-Q. Chen for his comments on the first version of this paper. We thank the referee for pointing out a mistake in the first version of this paper and also for his detailed and helpful comments.

References

- [1] Bañuelos, R.: Intrinsic ultracontractivity and eigenfunction estimates for Schrödinger operators. *J. Funct. Anal.* **100**, 181–206 (1991)
- [2] Blumenthal, R.M., Gettoor, R.K.: *Markov Processes and Potential Theory*. Academic Press, New York 1968
- [3] Bouleau, N.: Quelques résultats probabilistes sur la subordination au sens de Bochner, in *Seminar on potential theory*, Paris, No. 7, 54–81, *Lecture Notes in Math.*, 1061, Springer, Berlin, 1984
- [4] Chen, Z.-Q., Song, R.: Intrinsic ultracontractivity and conditional gauge for symmetric stable processes. *J. Funct. Anal.* **150**, 204–239 (1997)
- [5] Chen, Z.-Q., Song, R.: Estimates on Green functions and Poisson kernels for symmetric stable processes. *Math. Ann.* **312**, 465–501 (1998)
- [6] Chen, Z.-Q., Song, R.: Intrinsic ultracontractivity, conditional lifetimes and conditional gauge for symmetric stable processes on rough domains. *Ill. J. Math.* **44**, 138–160 (2000)
- [7] Chen, Z.-Q., Song, R.: Hardy inequality for censored stable processes, *Tohoku Math. J.* (to appear), 2002
- [8] Davies, E.B.: *Heat kernels and spectral theory*. Cambridge, Cambridge University Press, 1989
- [9] Davis, B.: Intrinsic ultracontractivity and the Dirichlet Laplacian. *J. Funct. Anal.* **100**, 163–180 (1991)
- [10] Farkas, W., Jacob, N.: Sobolev spaces on non-smooth domains and Dirichlet forms related to subordinate reflecting diffusions. *Math. Nachr.* **224**, 75–104 (2001)
- [11] Fukushima, M., Oshima, Y., Takeda, M.: *Dirichlet forms and symmetric Markov processes*. Walter De Gruyter, Berlin, 1994
- [12] Glover, J., Rao, M., Šikić, H., Song, R.: Γ -potentials, in *Classical and modern potential theory and applications* (Chateau de Bonas, 1993), 217–232, *Kluwer Acad. Publ.*, Dordrecht, 1994
- [13] Ikeda, N., Watanabe, S.: On some relations between the harmonic measure and the Lévy measure for a certain class of Markov processes. *J. Math. Kyoto Univ.* **2**, 79–95 (1962)
- [14] Jacob, N., Schilling, R.: Some Dirichlet spaces obtained by subordinate reflected diffusions. *Rev. Mat. Iberoamericana* **15**, 59–91 (1999)
- [15] Kulczycki, T.: Properties of Green function of symmetric stable processes. *Probab. Math. Statist.* **17**, 339–364 (1997)
- [16] Kulczycki, T.: Intrinsic ultracontractivity for symmetric stable processes. *Bull. Polish Acad. Sci. Math.* **46**, 325–334 (1998)
- [17] Miyake, A.: The subordination of Lévy system for Markov processes. *Proc. Japan Acad.* **45**, 601–604 (1969)
- [18] Pazy, A.: *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Springer-Verlag, New York, 1983
- [19] Schilling, R.L.: On the domain of the generator of a subordinate semigroup. In: J. Král et al. (Eds.). *Potential Theory— ICPT 94*. *Proceedings Intl. Conf. Potential Theory*, Kouty (CR), 1994. de Gruyter, 449–462 (1996)
- [20] Skorohod, A.V.: Asymptotic formulas for stable distribution laws. In: *Selected Transl. Math. Statist. Probab.*, Vol. 1, AMS, Providence, Rhode Island, 157–161 (1961)
- [21] Triebel, H.: *Interpolation Theory, Function Spaces, Differential Operators*, Second Edition. Johann Ambrosius Barth, 1995
- [22] Yosida, K.: *Functional Analysis*. 6th edition, *springer-Verlag*, Berlin 1980