



Laser-capture microdissection of murine lung for differential cellular RNA analysis

Jagadish Loganathan¹ · Roshni Pandey¹ · Nilesh Sudhakar Ambhore¹ · Pawel Borowicz² · Venkatachalem Sathish¹

Received: 23 October 2018 / Accepted: 15 January 2019 / Published online: 2 February 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

The lung tissue contains a heterogeneous milieu of bronchioles, epithelial, airway smooth muscle (ASM), alveolar, and immune cell types. Healthy bronchiole comprises epithelial cells surrounded by ASM cells and helps in normal respiration. In contrast, airway remodeling, or plasticity, increases surrounding of bronchial epithelium during inflammation, especially in asthmatic condition. Given the profound functional difference between ASM, epithelial, and other cell types in the lung, it is imperative to separate and isolate different cell types of lungs for genomics, proteomics, and molecular analysis, which will improve the diagnostic and therapeutic approach to treat cell-specific lung disorders. Laser capture microdissection (LCM) is the technique generally used for the isolation of specific cell populations under direct visual inspection, which plays a crucial role to evaluate cell-specific effect in clinical and preclinical setup. However, maintenance of tissue RNA quality and integrity in LCM studies are very challenging tasks. It is obvious to believe that the major factor affecting the RNA quality is tissue-fixation method. The prime focus of this study was to address the RNA quality factors within the lung tissue using the different solvent system to fix tissue sample to obtain high-quality RNA. Paraformaldehyde and Carnoy's solutions were used for fixing the lung tissue and compared RNA integrity in LCM captured lung tissue samples. To further confirm the quality of RNA, we measured cellular marker genes in collected lung tissue samples from control and mixed allergen (MA)-induced asthmatic mouse model using qRT-PCR technique. RNA integrity number showed a significantly better quality of RNA in lung tissue samples fixed with Carnoy's solution compared to paraformaldehyde solution. Isolated RNA from MA-induced asthmatic murine lung epithelium, smooth muscle, and granulomatous foci using LCM showed a significant increase in remodeling gene expression compared to control which confirm the quality and integrity of isolated RNA. Overall, the study concludes tissue fixation solvent can alter the quality of RNA in the lung and the outcome of the results.

Keywords Carnoy's solution · RNA integrity · Lung tissue fixation · Laser capture microdissection · qRT-PCR

Introduction

Histological section of tissue samples under the microscope shows different characterized cells with various genetic profiles. These cells also communicate with each other to conduct the physiological function in normal and diseased condition (Arthur 2016; Corces et al. 2016; Lander 1996; MacArthur

et al. 2014; Oellrich and Smedley 2014). However, the expression levels of genes and their function would significantly vary between different cells (Antanaviciute et al. 2015; Fatkin et al. 2014; Zaidi et al. 2013). Thus, identifying the specific gene expression in a particular cell type or tissue from a specific cell type under normal and diseased conditions would significantly expand our knowledge in understanding the role of gene expression in diagnostic and therapeutic approach (Dillon et al. 2001; Domazet et al. 2008; Emmert-Buck et al. 1996). Laser capture microdissection (LCM) is a method to obtain samples containing specific cell types or cells of interest from sectioned tissue subjected to next-generation sequencing (NGS) or microarray to understand the entire RNA content of a single cell for functional elucidations (Ardekani et al. 2008; Brioschi et al. 2014; Venter et al. 2001). LCM is a highly efficient and accepted technique to obtain specific

✉ Venkatachalem Sathish
s.venkatachalem@ndsu.edu

¹ Department of Pharmaceutical Sciences, School of Pharmacy, North Dakota State University, Sudro Hall, Room 203, Fargo, ND 58108-6050, USA

² Department of Animal Sciences, North Dakota State University, Fargo, ND, USA

cells, however, it is more vulnerable to RNase contamination (Kerk et al. 2003). Most microdissection-based mRNA expression studies have been performed on frozen tissue samples which require more attention to avoid damage to the tissue during transportation, handling, tissue sectioning, and during post-LCM RNA isolation steps. More or less, all these steps are responsible for poor quality of RNA, thereby, reducing the efficacy for proceeding further downstream applications (Amini et al. 2017; Takahashi et al. 2010).

Most of the pathophysiology laboratories in clinical setup apply formalin or paraformaldehyde solution for tissue fixation. Among other fixatives, Carnoy's solution is commonly used in cancer-related studies because of its easy handling and more accurate staging (Luz et al. 2008). However, very limited or no clinical studies have reported the use of Carnoy's solution in the fixation of lung tissue specimens to study downstream analysis. In order to overcome these factors, we tried to optimize the best system for tissue fixation using mouse lung samples.

The lung, a well-developed organ, is essential for terrestrial life. The lung tissue is composed of heterogeneous cells; comprises tissues such as epithelial, mesenchymal, airway smooth muscle (ASM), alveolar, and immune cells. Together, these cells are responsible for the normal functioning of respiration (Hines and Sun 2014). Chronic inflammation centrally affecting the lung airway, which impairs the functions of multiple cells, including ASM, which plays a key role in airway remodeling via altered proliferation and deposition of extracellular matrix (ECM) (Anderson 2010; Bousquet et al. 1996; O'Byrne and Inman 2003). Airway walls are composed of more than one cell type, which alone or in combination contributes to airway hyperresponsiveness (AHR), inflammation, and remodeling, a characteristic features of asthma. It gives immense therapeutic importance to identify particular cells involved in the pathophysiology of asthma. Here, LCM technique acts as a potent tool to isolate individual cell types to understand its role in lung pathophysiology by subsequent molecular analysis.

In this study, we described the various technical approaches and updated LCM methods that we have used to isolate different lung cell types and their mRNA expression involved in airway remodeling in the lung. Furthermore, we optimized the lung fixation method using Carnoy's solution to maintain RNA quality and integrity in cells collected through LCM. Measured airway cellular gene expression levels by qRT-PCR further confirm the quality and integrity of isolated RNA.

Materials and methods

Animals

Murine lung tissue samples were collected from our ongoing pulmonary in vivo study, which is approved by the Institutional Animal Care and Use Committee at North

Dakota State University and conducted in accordance with guidelines derived from the National Institutes of Health Guide for the Care and Use of Laboratory Animals. We randomly selected six C57BL/6J mice each from control and mixed allergen (MA, allergic asthma model)-treated groups. Animals were anesthetized using isoflurane (volatile anesthesia) for 10 s to calm and lower the movement of mice during intranasal PBS or MA administration. Control mice received PBS and MA-treated mice received alternate day intranasal (i.n.) administration of a mixture of equal amounts (10 µg) of ovalbumin and extracts from *Alternaria alternata*, *Aspergillus fumigatus*, and *Dermatophagoides farinae* (house dust mite) for 4 weeks (alternative day exposure) in PBS to induce asthma (Yarova et al. 2015). At the end of the study, animals were euthanized by using an excess dose of sodium pentobarbital (200 mg/kg). Both male and female mice were used in this study. Mice were always housed under constant temperature and 12 h light and dark cycles provided with food and water ad libitum.

Histopathology using hematoxylin and eosin staining

The left lung was ligated and the right lung was inflated with 0.8 ml of either paraformaldehyde or Carnoy's solution and fixed overnight. Paraffin-embedded tissues were cut into 6-µm sections using a tissue microtome (Leica, UK). The sections were stained with hematoxylin and eosin stain (H&E) for histological observations to identify epithelium, airway smooth muscle, and granulomatous foci. The stained tissues were then mounted onto a light microscope and examined at × 40 magnification.

Laser capture microdissection

The lung tissue was collected from mouse and fixed using two different methods: (a) 4% paraformaldehyde and (b) Carnoy's fixative solution (Pereira et al. 2015). Tissue blocks were embedded in paraffin wax and sectioned using a rotary microtome to obtain 10 µm tissue sections placed on RNase free slides. The paraffin slides were washed in xylene for 9 min and ethanol for 30 s and then the excess xylene was blotted from the slide. Laser capture microdissection (LCM) was performed using a Zeiss Axiomager Z1 PALM Microbeam laser capture microscope system (Zeiss, Thornwood, NY). The microdissection process was visualized with an AxioCam Icc camera coupled to a computer and controlled by a Palm-Robo software (Zeiss, Thornwood, NY). The slides were then visualized under bright-field microscopy with a × 10 objective. An area of 0.3 or 1.0 mm² was selected using the PALM-Robo software. Identified granulomatous foci and different airway cells were marked using an on-screen tool of PALM MicroBeam. Laser cut was performed along the marking followed by catapulting the tissue. To confirm specific cell

type, hematoxylin and eosin stained (H&E) section of same tissue was used as a reference. Catapulted cells were collected into on RNase free microtube containing 0.5 ml single cell PicoPure® RNA extraction buffer. Tissues were either stored temporarily at -80°C or directly used for RNA isolation. The capture success of microdissected tissue was confirmed by looking at the microtube cap under the microscope.

RNA extraction and DNase treatment

Total RNA extraction was performed by using the PicoPure® RNA Isolation Kit (Thermo Fisher Scientific, Waltham, MA, USA). At first step, 100 μl of extraction buffer were added in each microtube containing tissue. For granulomatous foci, 50 μl of extraction buffer was added to avoid excessive dilution of RNA due to the small amount of microdissected tissue. Genomic DNA was removed by on-column DNase digestion, treating bound RNA in the column for 20 min with 5 μl of DNase I diluted in 35 μl of DNase digestion buffer (RNase-Free DNase Set, Zymoresearch, CA). Around 13 μl of elution buffer was pipetted directly onto the membrane of the column to elute the RNA into an RNase free tube. Collected RNA was then stored immediately at -80°C .

Measurement of RNA yield and integrity

The yield and integrity of total isolated RNA from microdissected samples measured using a 2100 Bioanalyzer (Agilent Technologies). RNA concentration of (1 μl per sample) was measured with the 2100 Bioanalyzer, then obtained concentration was divided by the tissue area value to calculate the RNA yield. Additionally, the RNA Integrity Number (RIN) was calculated by the 2100 Bioanalyzer according to an algorithm from Agilent Technologies (Kerman et al. 2006; Schroeder et al. 2006). RIN values obtained for each sample on a scale of 1–10 as an indication of RNA quality.

qRT-PCR analysis

Isolated and quantified RNA was used to synthesize cDNA using OneScript cDNA Synthesis Kit (Applied Biological Materials Inc., Richmond, BC, Canada) for each sample. Bright Green Master mix qPCR kit (Applied Biological Materials Inc., Richmond, BC, Canada; Cat# MasterMix-S) protocol was followed using QuantStudio 3 qRT-PCR system as per the manufacturer's instructions. The qRT-PCR products were loaded into a 2% agarose gel with gel green nucleic acid stain (Biotium), separated by electrophoresis and visualized licor D-Digit gel scanner. Following primer sequences were used to qRT-PCR analysis; For TNF- α , forward 5'-CTCC TCCTCCTGACCTTCTAATAAT-3', reverse 5'-CTCC TCTGCCAAGTTCATATC C-3'; for α -smooth muscle actin (α -SMA) forward 5'-GAGGCACCACTGAACCCTAA-3',

reverse 5'-CATCTCCAGAGTCC AGCACA-3'; for E-cadherin forward 5'-CACCTGGAGAGAGGCCATGT-3', reverse 5'-TGGGAAACATG AGCAGCTCT-3'; for fibronectin forward 5'-CGAGGTGACAGAGACCACAA-3', reverse 5'-CTGGAGTCAAGCCAGACACA-3'; for Vimentin forward 5'-CTTGAACGGAAA GTGGAATCCT-3', reverse 5'-GTCAGGCTTGGAACGTCC-3'; for s16 forward 5'-GATGAAGTCCGAGCTGGTAAA-3', reverse 5'-GGAG TGACAGGCAACTATGAA-3'. The fold changes in mRNA expression were calculated by normalization of cycle threshold [C(t)] value of target genes to reference gene s16 using the $\Delta\Delta\text{Ct}$ method.

Statistical analysis

All groups consisted of minimum six mice. “*n*” values represent the number of samples from individual animals. All qRT-PCR experiments were repeated at least two times. Statistical analysis was performed using Sigma Plot (SYSTAT, San Jose, CA) software statistical package using individual one-way ANOVA with Bonferroni post hoc test. All data are expressed as mean \pm SEM. Statistical significance was tested at minimum of $p < 0.05$ level.

Results

Effect of fixation on tissue capture success and RNA yield and integrity after LCM

Six control (PBS only) and six mixed allergen (MA)-induced asthmatic mice were used for the current study. The lung tissue samples were collected after 28 days of PBS and MA challenge. Mouse tissue samples cut to the thickness of 10 μm and mounted onto glass slides (glass or PEN-membrane coated) to isolate airway smooth muscle cells, epithelial, and granulomatous foci. The workflow for this study is tissue preparation, LCM, and isolation of RNA followed by RNA analysis by qRT-PCR (Fig. 1). We used PALM MicroBeam system (Zeiss) Laser capture microdissection system to isolate different airway cells from normal and inflammation-induced lung tissue of mouse. The cells isolated was ensured microscopical observation before and after excision. Specific cell types are identified through H&E-stained reference slides that are prepared from the tissue blocks used for making LCM slides (Fig. 2).

Effect of tissue fixation solvent on RNA quality

Mouse lung tissue specimens fixed into Carnoy's or paraformaldehyde solution and LCM isolated different airway cell types used to measure RNA yield with integrity to compare the effectiveness of fixative solution. The obtained RNA yield

Fig. 1 Workflow for the laser-capture microdissection (LCM) of lung tissue, isolation, and analysis of RNA. Airway smooth muscle (ASM) cells, epithelium, and lymphatic granules from normal and inflammation-induced lung tissue collected by LCM and further processed for RNA isolation. (a) Selected region of interest before LCM, (b) after LCM, (c) collected cells in lysis buffer. H&E stained tissue section used as a reference to identify different cell types





Fig. 2 Representative images of lung tissue specimens collected after mixed allergen exposure to induce asthma. Hematoxylin and eosin (H&E) staining used as a reference to identify different cells types; laser capture microdissected murine lung tissue of (a) ASM, (b) epithelium, and (c) granulomatous foci (10 µm section). Tissue before laser dissection (d) ASM, (e) epithelium, and (f) granulomatous foci; tissue after laser dissection (g) ASM, (h) epithelium, and (i) granulomatous foci. Scale bar, 50 µM



and integrity were compared for both fixative solvent in granulomatous foci, ASM, and epithelium (Fig. 3). The RNA yield in Carnoy's solution showed prominently more in comparison to paraformaldehyde (Table 1), which indicates that the tissue fixation with Carnoy's provided better yield and more integrity.

qRT-PCR for gene expression analysis in different cell types of mouse lung tissue

Mouse lung tissue collected from control and MA-induced asthmatic mouse model and fixed with Carnoy's solution resulted in better quality RNA isolated from three different cell types including airway smooth muscle cells, epithelium, and granulomatous foci. To evaluate the effectiveness of isolated RNA, we evaluated changes in various cellular specific gene expression in normal and pathological conditions in different cell types using qRT-PCR using *s16* as a housekeeping gene. The primary focus of this current study was to optimize the LCM method to obtain good quality RNA to further study the downstream genetic changes in different airway cells during inflammation and normal condition. qRT-PCR data from RNA isolated from ASM cells revealed a significant increase ($P < 0.001$) in α -smooth muscle actin, fibronectin, and E-cadherin in MA-challenged mice compared to PBS-treated

mice (Fig. 4a). Furthermore, epithelial cells isolated from airway showed a significant increase in expression of E-cadherin, an important epithelium marker and significant low expression of α -SMA in MA-challenged mice compared to PBS, indicating the homogenous cells with less contamination (Fig. 4b). In an inflammatory condition, the granulomatous foci showed an increase in TNF- α gene expression. Concurrent with this, we observed that the expression of TNF- α was significantly ($p < 0.001$) high in granulomatous foci in MA-challenged mice than in PBS-treated mice (Fig. 4c).

Discussion

LCM is a powerful technique that permits rapid dissection and isolation of specific tissues, cell organelles, and their fragments, which allow accurate examination of cellular genomic profiles by using qRT-PCR, RNA sequencing, genomics, and proteomics analysis. To process the prospective clinical tissue specimens, the most commonly used method in LCM studies is frozen tissue section. The major difficulty in carrying out this method is the tedious steps that required lots of attention which gives the high possibility of losing the yield and integrity of RNA prior to fixation and stabilization. Also,

Fig. 3 RNA integrity of specific cells isolated from LCM-derived tissue measured using Agilent bioanalyzer. RNA integrity of (a) ASM, (a') epithelium, and (a'') granulomatous foci from paraformaldehyde-fixed lung tissue. RNA integrity of (b) ASM, (b') epithelium, and (b'') granulomatous foci from Carnoy's fixed lung tissue. The average RIN values for different cell types shown with graphical representation. The RIN value indicates RNA integrity number which is better in Carnoy's fixed tissue than paraformaldehyde-fixed tissue. Data represented as mean \pm SEM of at least 06 mice per group; *** $p < 0.001$, * $p < 0.01$ paraformaldehyde vs Carnoy's solution




Table 1 Measurement of microdissected tissue area, RNA concentration, and RNA integrity

Tissue	Paraformaldehyde solution			Carnoy's solution		
	ASM	Epi	GF	ASM	Epi	GF
Number of microdissected area (n)	10.21 ± 2.81	12.01 ± 3.1	4.6 ± 1.14	11.22 ± 2.7	13.18 ± 2.42	4.08 ± 1.24
Concentration (ng/μl)	0.42 ± 0.03	0.72 ± 0.11	1.04 ± 0.12	1.44 ± 0.13***	1.98 ± 0.24***	2.18 ± 0.46***

ASM airway smooth muscle, Epi epithelium, GF granulomatous foci. Data represented as mean ± SEM of at least six mice samples per group

*** $p < 0.001$ paraformaldehyde vs Carnoy's solution for respective group

processing postoperative surgical specimens is very difficult with this method as it requires a multi-step process (Gautam et al. 2016). Multiple retrospectively collected clinical sample-based studies require easy access, long-term storage, and keeping up sample integrity. Limited studies have suggested the use of Carnoy's solution for tissue fixation to overcome these limitations and provide the RNA stability and integrity (Cox et al. 2006).

Tissue fixation and processing method, together, play an important role in the protection and maintenance of tissue morphology at the cellular level that affects RNA quality. Investigators used different fixation systems depending on tissue type used to preserve the RNA integrity. Brown and Smith (2009) suggested using high-salt buffer solution to fix rat brain tissue samples, whereas Santoro et al. (Santoro et al. 2018) used RNALater solution (Applied Biosystem, Foster City, CA, USA) to preserve skin biopsies for 1 week at 4 °C to get stabilized RNA. In clinical studies most commonly used tissue preservation method is formalin fixation, however, it chemically alters the nucleotides inducing nucleic acid fragmentation and decrease RNA quality (Butler et al. 2016) for this reason it is not suitable for LCM studies. Very limited or no study attempted to optimize fixative solvent system for the lung tissue specimens in LCM studies.

The objective of this study was to optimize an appropriate tissue fixation solvent for the lung tissue to support clinical setup as well our ongoing pulmonary research hypothesizing the role of sex hormones in lung physiology. The lung is comprised of multiple cells, so it is very crucial to identify the particular cell type responsible for altered physiology especially in the airway. In this, we utilized lung specimens from control and mixed allergen (MA)-induced asthmatic C57BL/6/J mouse model to optimize the RNA integrity. We used Carnoy's solution or paraformaldehyde solution as fixative to determine the RNA integrity and optimized Carnoy's solution is the best solvent for fixing tissue to perform LCM studies which provided good quality RNA for subsequent gene expression analysis. Selection of proper solvent system for tissue fixation is very important because RNA is a very labile biomolecule, and easily degraded, especially during LCM studies, where very low amounts of tissue material are

collected. It is required to put special cautions to preserve RNA yield and integrity, which are decisive for downstream genomic studies. Solution used for tissue fixation in the study did not alter the tissue morphology and often is used in tissues subjected to LCM studies (Parlato et al. 2002). In the initial study, we found very low RNA integrity number (RIN) in LCM-isolated cells from paraformaldehyde solution-fixed mouse lung. PCR was performed in LCM-isolated cells from paraformaldehyde-fixed lung tissue and found highly variable CT value (ranges from 34 to 40) for s16 indicating the lower quality of RNA (Schroeder et al. 2006). Considering this fact, we are not able to get reliable qRT-PCR data with cells isolated from paraformaldehyde-fixed lung tissue.

RNA integrity number (RIN) represents the ratio of ribosomal subunit 26S and 18S, which is measured at 1–10 scale. The higher RIN number represents a good quality of RNA. Ribosomal RNA is used as a standard for measuring the RNA quality and our study showed that using Carnoy's solution for lung tissue fixation we got highest RNA yield and integrity (Kang et al. 2017; Kim et al. 2003; Shibutani et al. 2000). Further, to confirm the effectiveness of isolated RNA, we evaluated cellular gene expression in murine lung tissue by using different cell types like airway smooth muscle cells, epithelial cells, and granulomatous foci under normal and pathological condition. Higher expression of α -SMA and fibronectin in airway smooth muscle cells concord with earlier reports suggesting increased airway remodeling genes in asthmatic condition. Whereas, RNA isolated from epithelium tissue showed epithelium marker E-cadherin and increased TNF- α expression in granulomatous foci upon MA exposure suggests quality and integrity of isolated RNA was optimum to evaluate downstream effects.

We optimized our protocol to reduce the impact of several events that affect the quality and quantity of RNA in the lung tissue using LCM technique. With a small amount of RNA, we showed a difference in cellular genes as per pathophysiological condition by qRT-PCR, which shows the sensitivity of the optimized method. In the present study, we describe an optimized protocol using the Carnoy's solution as a fixative solution for lung tissue to isolate RNA from ASM cells,




Fig. 4 Different cellular gene expression in LCM-isolated mouse lung tissue. Change in expression of different cellular genes are observed in accordance with treatment shows effectiveness of LCM and integrity of RNA. **a** In ASM cell expression of α -smooth muscle actin (α -SMA), fibronectin, and E-cadherin is significantly changed. **b** Airway epithelium tissue showed increased expression of E-cadherin (epithelial marker), whereas decreased in the α -SMA expression in MA-treated group than the control. **c** Cell from granulomatous foci showed high expression of TNF α gene in MA-treated group compared to PBS. Data represented as mean \pm SEM of at least six mice per group; *** p < 0.001, * p < 0.05 PBS vs MA

epithelial, and granulomatous foci. The fixative solution has a major impact on RNA quality and integrity; Carnoy's solution provides better RNA quality and integrity compared to

paraformaldehyde in different lung cell types. Good quality RNA can provide a more accurate evaluation of cell-specific downstream analysis, which offers the opportunity for a more precise diagnostic and therapeutic approach in translational lung research.

Author contributions JL, RP, NSA, and SV have conceptualized the study. JL, RP, and NSA have performed the experimental procedures. JL, RP, NSA, and SV have contributed in data analysis and manuscript preparation. SV revised final draft and approved for submission.

Funding information Supported by NIH grants R01 HL123494 (Venkatachalem). Additional support in part from ND EPSCoR with NSF #1355466 and NDSU RCA Activity.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Amini P, Ettlin J, Opitz L, Clementi E, Malbon A, Markkanen E (2017) An optimised protocol for isolation of RNA from small sections of laser-capture microdissected FFPE tissue amenable for next-generation sequencing. *BMC Mol Biol* 18:22
- Anderson SD (2010) Indirect challenge tests: airway hyperresponsiveness in asthma: its measurement and clinical significance. *Chest* 138:25s–30s
- Antanaviciute A, Daly C, Crinnion LA, Markham AF, Watson CM, Bonthron DT, Carr IM (2015) GeneTIER: prioritization of candidate disease genes using tissue-specific gene expression profiles. *Bioinformatics* 31:2728–2735
- Ardekani AM, Akhondi MM, Sadeghi MR (2008) Application of genomic and proteomic technologies to early detection of cancer. *Arch Iran Med* 11:427–434
- Arthur G (2016) Albert Coons: harnessing the power of the antibody. *Lancet Respir Med* 4:181–182
- Bousquet J, Lacoste JY, Chanez P, Vic P, Godard P, Michel FB (1996) Bronchial elastic fibers in normal subjects and asthmatic patients. *Am J Respir Crit Care Med* 153:1648–1654
- Brioschi M, Eligini S, Crisci M, Fiorelli S, Tremoli E, Colli S, Banfi C (2014) A mass spectrometry-based workflow for the proteomic analysis of in vitro cultured cell subsets isolated by means of laser capture microdissection. *Anal Bioanal Chem* 406:2817–2825
- Brown AL, Smith DW (2009) Improved RNA preservation for immunolabeling and laser microdissection. *RNA* 15(12):2364–2374
- Butler AE, Matveyenko AV, Kirakossian D, Park J, Gurlo T, Butler PC (2016) Recovery of high-quality RNA from laser capture microdissected human and rodent pancreas. *J Histotechnol* 39:59–65
- Corces MR, Buenrostro JD, Wu B, Greenside PG, Chan SM, Koenig JL, Snyder MP, Pritchard JK, Kundaje A, Greenleaf WJ, Majeti R, Chang HY (2016) Lineage-specific and single-cell chromatin accessibility charts human hematopoiesis and leukemia evolution. *Nat Genet* 48:1193
- Cox ML, Schray CL, Luster CN, Stewart ZS, Korytko PJ, KN MK, Paulauskis JD, Dunstan RW (2006) Assessment of fixatives, fixation, and tissue processing on morphology and RNA integrity. *Exp Mol Pathol* 80:183–191
- Dillon D, Zheng K, Costa J (2001) Rapid, efficient genotyping of clinical tumor samples by laser-capture microdissection/PCR/SSCP. *Exp Mol Pathol* 70:195–200

- Domazet B, MacLennan GT, Lopez-Beltran A, Montironi R, Cheng L (2008) Laser capture microdissection in the genomic and proteomic era: targeting the genetic basis of cancer. *Int J Clin Exp Pathol* 1: 475–488
- Emmert-Buck MR, Bonner RF, Smith PD, Chuaqui RF, Zhuang Z, Goldstein SR, Weiss RA, Liotta LA (1996) Laser capture microdissection. *Science* 274:998–1001
- Fatkin D, Seidman CE, Seidman JG (2014) Genetics and disease of ventricular muscle. *Cold Spring Harb Perspect Med* 4:a021063
- Gautam V, Singh A, Singh S, Sarkar AK (2016) An efficient LCM-based method for tissue specific expression analysis of genes and miRNAs. *Sci Rep* 6:21577
- Hines EA, Sun X (2014) Tissue crosstalk in lung development. *J Cell Biochem* 115:1469–1477
- Kang L, George P, Price DK, Sharakhov I, Michalak P (2017) Mapping genomic scaffolds to chromosomes using laser capture microdissection in application to Hawaiian picture-winged drosophila. *Cytogenet Genome Res* 152:204–212
- Kerk NM, Ceserani T, Tausta SL, Sussex IM, Nelson TM (2003) Laser capture microdissection of cells from plant tissues. *Plant Physiol* 132:27
- Kerman IA, Buck BJ, Evans SJ, Akil H, Watson SJ (2006) Combining laser capture microdissection with quantitative real-time PCR: effects of tissue manipulation on RNA quality and gene expression. *J Neurosci Methods* 153:71–85
- Kim JO, Kim HN, Hwang MH, Shin HI, Kim SY, Park RW, Park EY, Kim IS, Van Wijnen AJ, Stein JL, Lian JB, Stein GS, Choi JY (2003) Differential gene expression analysis using paraffin-embedded tissues after laser microdissection. *J Cell Biochem* 90:998–1006
- Lander ES (1996) The new genomics: global views of biology. *Science* 274:536–539
- Luz D, Ribeiro U Jr, Chassot C, De Salles Collet e Silva F, Ceconello I, Corbett C (2008) Carnoy's solution enhances lymph node detection: an anatomical dissection study in cadavers. *Histopathology* 53:740–742
- MacArthur DG, Manolio TA, Dimmock DP, Rehm HL, Shendure J, Abecasis GR, Adams DR, Altman RB, Antonarakis SE, Ashley EA, Barrett JC, Biesecker LG, Conrad DF, Cooper GM, Cox NJ, Daly MJ, Gerstein MB, Goldstein DB, Hirschhorn JN, Leal SM, Pennacchio LA, Stamatoyannopoulos JA, Sunyaev SR, Valle D, Voight BF, Winckler W, Gunter C (2014) Guidelines for investigating causality of sequence variants in human disease. *Nature* 508:469
- O'Byrne PM, Inman MD (2003) Airway hyperresponsiveness. *Chest* 123: 411s–416s
- Oellrich A, Smedley D (2014) Linking tissues to phenotypes using gene expression profiles. *Database (Oxford)* 2014:bau017
- Parlato R, Rosica A, Cucurullo V, Mansi L, Macchia P, Owens JD, Mushinski JF, De Felice M, Bonner RF, Di Lauro R (2002) A preservation method that allows recovery of intact RNA from tissues dissected by laser capture microdissection. *Anal Biochem* 300: 139–145
- Pereira MA, Dias AR, Faraj SF, Cirqueira CdS, Tomitao MT, Carlos Nahas S, Ribeiro Jr U, de Mello ES (2015) Carnoy's solution is an adequate tissue fixative for routine surgical pathology, preserving cell morphology and molecular integrity. *Histopathology* 66:388–397
- Santoro S, Lopez ID, Lombardi R, Zauli A, Osiceanu AM, Sorosina M, Clarelli F, Peroni S, Cazzato D, Marchi M, Quattrini A, Comi G, Calogero RA, Lauria G, Martinelli Boneschi F (2018) Laser capture microdissection for transcriptomic profiles in human skin biopsies. *BMC Mol Biol* 19:7
- Schroeder A, Mueller O, Stocker S, Salowsky R, Leibler M, Gassmann M, Lightfoot S, Menzel W, Granzow M, Ragg T (2006) The RIN: an RNA integrity number for assigning integrity values to RNA measurements. *BMC Mol Biol* 7:3
- Shibutani M, Uneyama C, Miyazaki K, Toyoda K, Hirose M (2000) Methacarn fixation: a novel tool for analysis of gene expressions in paraffin-embedded tissue specimens. *Lab Invest* 80:199
- Takahashi H, Kamakura H, Sato Y, Shiono K, Abiko T, Tsutsumi N, Nagamura Y, Nishizawa NK, Nakazono M (2010) A method for obtaining high quality RNA from paraffin sections of plant tissues by laser microdissection. *J Plant Res* 123:807–813
- Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, Sutton GG, Smith HO, Yandell M, Evans CA, Holt RA, Gocayne JD, Amanatides P, Ballew RM, Huxon DH, Wortman JR, Zhang Q, Kodira CD, Zheng XH, Chen L, Skupski M, Subramanian G, Thomas PD, Zhang J, Gabor Miklos GL, Nelson C, Broder S, Clark AG, Nadeau J, McKusick VA, Zinder N, Levine AJ, Roberts RJ, Simon M, Slayman C, Hunkapiller M, Bolanos R, Delcher A, Dew I, Fasulo D, Flanigan M, Florea L, Halpern A, Hannenhalli S, Kravitz S, Levy S, Mobarry C, Reinert K, Remington K, Abu-Threideh J, Beasley E, Biddick K, Bonazzi V, Brandon R, Cargill M, Chandramouliswaran I, Charlab R, Chaturvedi K, Deng Z, Francisco VD, Dunn P, Eilbeck K, Evangelista C, Gabrielian AE, Gan W, Ge W, Gong F, Gu Z, Guan P, Heiman TJ, Higgins ME, Ji R-R, Ke Z, Ketchum KA, Lai Z, Lei Y, Li Z, Li J, Liang Y, Lin X, Lu F, Merkulov GV, Milshina N, Moore HM, Naik AK, Narayan VA, Neelam B, Nusskern D, Rusch DB, Salzberg S, Shao W, Shue B, Sun J, Wang ZY, Wang A, Wang X, Wang J, Wei M-H, Wides R, Xiao C, Yan C, Yao A, Ye J, Zhan M, Zhang W, Zhang H, Zhao Q, Zheng L, Zhong F, Zhong W, Zhu SC, Zhao S, Gilbert D, Baumhueter S, Spier G, Carter C, Cravchik A, Woodage T, Ali F, An H, Awe A, Baldwin D, Baden H, Barnstead M, Barrow I, Beeson K, Busam D, Carver A, Center A, Cheng ML, Curry L, Danaher S, Davenport L, Desilets R, Dietz S, Dodson K, Doup L, Ferriera S, Garg N, Gluecksmann A, Hart B, Haynes J, Haynes C, Heiner C, Hladun S, Hostin D, Houck J, Howland T, Ibegwam C, Johnson J, Kalush F, Kline L, Koduru S, Love A, Mann F, May D, McCawley S, McIntosh T, McMullen I, Moy M, Moy L, Murphy B, Nelson K, Pfannkoch C, Pratts E, Puri V, Qureshi H, Reardon M, Rodriguez R, Rogers Y-H, Romblad D, Ruhlfel B, Scott R, Sitter C, Smallwood M, Stewart E, Strong R, Suh E, Thomas R, Tint NN, Tse S, Vech C, Wang G, Wetter J, Williams S, Williams M, Windsor S, Winn-Deen E, Wolfe K, Zaveri J, Zaveri K, Abril JF, Guigó R, Campbell MJ, Sjolander KV, Karlak B, Kejariwal A, Mi H, Lazareva B, Hatton T, Narechania A, Diemer K, Muruganujan A, Guo N, Sato S, Bafna V, Istrail S, Lippert R, Schwartz R, Walenz B, Yooseph S, Allen D, Basu A, Baxendale J, Blick L, Caminha M, Carnes-Stine J, Caulk P, Chiang Y-H, Coyne M, Dahlke C, Mays AD, Dombroski M, Donnelly M, Ely D, Esparham S, Fosler C, Gire H, Glanowski S, Glasser K, Glodek A, Gorokhov M, Graham K, Gropman B, Harris M, Heil J, Henderson S, Hoover J, Jennings D, Jordan C, Jordan J, Kasha J, Kagan L, Kraft C, Levitsky A, Lewis M, Liu X, Lopez J, Ma D, Majoros W, McDaniel J, Murphy S, Newman M, Nguyen T, Nguyen N, Nodell M, Pan S, Peck J, Peterson M, Rowe W, Sanders R, Scott J, Simpson M, Smith T, Sprague A, Stockwell T, Turner R, Venter E, Wang M, Wen M, Wu D, Wu M, Xia A, Zandieh A, Zhu X (2001) The sequence of the human genome. *Science* 291:1304
- Yarova PL, Stewart AL, Sathish V, Britt RD, Thompson MA, Lowe AP, Freeman M, Aravamudan B, Kita H, Brennan SC (2015) Calcium-sensing receptor antagonists abrogate airway hyperresponsiveness and inflammation in allergic asthma. *Sci Trans Med* 7:284ra2e60–2284ra260
- Zaidi S, Choi M, Wakimoto H, Ma L, Jiang J, Overton JD, Romano-Adesman A, Bjornson RD, Breitbart RE, Brown KK, Carriero NJ, Cheung YH, Deanfield J, DePalma S, Fakhro KA, Glessner J, Hakonarson H, Italia MJ, Kaltman JR, Kaski J, Kim R, Kline JK, Lee T, Leipzig J, Lopez A, Mane SM, Mitchell LE, Newburger JW, Parfenov M, Pe'er I, Porter G, Roberts AE, Sachidanandam R, Sanders SJ, Seiden HS, State MW, Subramanian S, Tikhonova IR, Wang W, Warburton D, White PS, Williams IA, Zhao H, Seidman JG, Brueckner M, Chung WK, Gelb BD, Goldmuntz E, Seidman CE, Lifton RP (2013) De novo mutations in histone-modifying genes in congenital heart disease. *Nature* 498:220–223