

Construction Sequences and Certifying 3-connectivity

Jens M. Schmidt

Received: 1 September 2009 / Accepted: 1 September 2010 / Published online: 11 September 2010
© Springer Science+Business Media, LLC 2010

Abstract Tutte proved that every 3-vertex-connected graph G on more than 4 vertices has a *contractible edge*. Barnette and Grünbaum proved the existence of a *removable edge* in the same setting. We show that the sequence of contractions and the sequence of removals from G to K_4 can be computed in $O(|V|^2)$ time by extending Barnette’s and Grünbaum’s theorem. As an application, we derive a certificate for the 3-vertex-connectivity of graphs that can be easily computed and verified.

Keywords Algorithms and data structures · Construction sequence · 3-connected · Certifying algorithm · Tutte contraction · Removable edges

1 Introduction

For a given set O of operations and a given set B of base graphs, a *construction sequence* of a graph G is a sequence of operations in O that constructs G when being applied to a base graph in B . For 3-vertex-connected (we say 3-connected) graphs G , we focus on construction sequences with $B = \{K_4\}$ and for which O consists either of inverse contractions (*Tutte’s construction sequence*) or inverse removals (*Barnette’s and Grünbaum’s construction sequence*). These two construction sequences are the inverse of the sequence of contractions and the sequence of removals, respectively. We will define contractions, removals and their inverse operations in Sect. 2.

Inductively defined constructions of graph classes are important, because the set of operations used in the constructions can often be exploited to prove properties on these graph classes. For 3-connected graphs, the existence theorems on contractible

This research was supported by the Deutsche Forschungsgemeinschaft within the research training group “Methods for Discrete Structures” (GRK 1408) and is an extended version of [13].

J.M. Schmidt (✉)

Dept. of Computer Science, Freie Universität Berlin, Berlin, Germany
e-mail: jens.schmidt@inf.fu-berlin.de

and removable edges yield such inductively defined constructions. Although these existence theorems are used frequently in graph theory [14, 15, 18], we are not aware of any computational results to find the sequence of contractions or removals, except for sequences where contractions and removals are allowed to intermix [1]. Moreover, efficient algorithms are unlikely to be derived from the existence proofs as they, e.g., in the case of Barnette and Grünbaum, depend heavily on adding longest paths, which are NP-hard to find. The main contribution of this paper is a structural result about the existence of Barnette's and Grünbaum's construction sequence and, based on that result, a simple algorithm to compute such a sequence in time $O(|V|^2)$. In addition, we show that Barnette's and Grünbaum's construction sequence can be transformed in linear time to the sequence of contractions, obtaining a close connection between these two sequences and a simple quadratic time algorithm for computing Tutte's construction sequence. Both algorithms do not rely on the 3-connectivity test of Hopcroft and Tarjan [6], which runs in linear time but is rather involved.

The concept of *certifying algorithms*, which give a small and easy-to-verify certificate of correctness along with their output, was initiated by Blum and Kannan [3] and developed further by Mehlhorn et al. [8–10]. While being important for program verification, certifying algorithms often provide new insights into a problem, which can lead to new techniques. For that reason they are a major goal for problems on which the known fast solutions are complicated and difficult to implement. Testing a graph on 3-connectivity is such a problem. Yet, surprisingly little work has been devoted to certify 3-connectivity, although a sophisticated linear-time recognition algorithm (not giving an easy-to-verify certificate) is known for over 35 years [6, 19, 20]. In fact, we are aware of only one certifying algorithm (in the sense of Mehlhorn et al.) for that problem, which runs in quadratic time, but is quite involved [1]. Using construction sequences, we give a simple alternative solution with running time $O(|V|^2)$ that performs essentially DFS-traversals and show that the used certificate is easy-to-verify in time $O(|E|)$.

We first recapitulate well-known results on the existence of construction sequences in Sects. 2.1 and 2.2 and point out how the sequence of contraction can be obtained from Barnette's and Grünbaum's sequence in linear time. Sections 2.3 and 3 cover the main idea for the existence result that we use for computing Barnette's and Grünbaum's construction sequence. The end of Sect. 3 deals with the representation of construction sequences. Section 4 shows how to use construction sequences for a certifying 3-connectivity test.

2 Construction Sequences

Let $G = (V, E)$ be a finite graph with $n := |V|$, $m := |E|$, $V(G) = V$ and $E(G) = E$. A graph is *connected* if there is a path between any two vertices and *disconnected* otherwise. For $k \geq 1$, a graph is *k-vertex-connected* if $n > k$ and deleting every $k - 1$ vertices leaves a connected graph. We will write *k-connected* throughout the paper when referring to *k-vertex-connectivity*. A vertex (resp. a pair of vertices) that leaves a disconnected graph upon deletion is called a *cut vertex* (resp. a *separation pair*). Note that *k-connectivity* does not depend on parallel edges or self-loops. From now

on, we assume for simplicity that our input graph $G = (V, E)$ is simple, although all results can be extended to multigraphs. A path leading from vertex v to vertex w is denoted by $v \rightarrow w$. For a vertex v in a graph, let $N(v) = \{w \mid vw \in E\}$ denote its set of neighbors and $deg(v)$ its degree. For a graph G , let $\delta(G)$ be the minimum degree of its vertices.

A *subdivision* of a graph G is a graph that replaces each edge in $E(G)$ by a path of length at least one. Conversely, we want a function that returns the original graph without subdivided edges. If $deg(v) = 2$ for a vertex v in a graph G , let $smooth_v(G)$ be the graph obtained from G by deleting v followed by adding an edge between its neighbors; we say v is *smoothed*. Otherwise, let $smooth_v(G) = G$. Let $smooth(G)$ be the graph obtained by smoothing every vertex in G . For an edge $e \in E$, let $G \setminus e$ denote the graph obtained from G by deleting e . Let K_n be the complete graph on n vertices.

The following are well-known corollaries of Menger’s theorem [11].

Lemma 1 (Fan Lemma) *Let v be a vertex in a graph G that is k -connected with $k \geq 1$ and let A be a set of at least k vertices in G with $v \notin A$. There are k internally vertex-disjoint paths P_1, \dots, P_k from v to distinct vertices $a_1, \dots, a_k \in A$ such that for each of these paths $V(P_i) \cap A = a_i$.*

Lemma 2 (Expansion Lemma [21]) *Let G be a k -connected graph. The graph obtained by adding a new vertex v joined to at least k vertices in G is still k -connected.*

2.1 Tutte’s Characterization and Its Inverse

Although G is simple, contractions cannot always avoid parallel edges in intermediate graphs. E. g., consider a cycle with an additional vertex connected to all cycle vertices by an edge. This is a *wheel graph*, and the edges adjacent to the non-cycle vertex are called *spokes*. The contraction of any edge that is not a spoke in a wheel graph will create a parallel edge. That is why we define contractions to preserve graphs to be simple. *Contracting* an edge $e = xy$ in a graph deletes e , merges vertices x and y , and replaces every set of parallel edges by a single edge. An edge e is called *contractible* if contracting e results in a 3-connected graph.

A *vertex splitting* takes a vertex v of a 3-connected graph, replaces v by two vertices x and y with an edge between them and replaces every former edge uv that was incident to v with either the edge ux, uy or both such that $|N(x)| \geq 3$ and $|N(y)| \geq 3$ in the new graph. Vertex splitting as defined here is therefore the exact inverse of contracting a contractible edge that has end vertices of degree ≥ 3 .

Theorem 3 (Corollary of Tutte [17]) *The following statements are equivalent:*

- A simple graph G is 3-connected*
- \iff *There exists a sequence of contractions from G to K_4 on contractible edges $e = xy$ with $|N(x)| \geq 3$ and $|N(y)| \geq 3$* (1)
- \iff *There exists a construction sequence from K_4 to G using vertex splittings* (2)

We describe next a straightforward $O(n^2)$ algorithm to compute (1) for a graph G on more than 4 vertices. First, we decrease the number of edges to $O(n)$ in G by applying the following algorithm due to Nagamochi and Ibaraki.

Theorem 4 (Nagamochi, Ibaraki [12]) *Let G be a connected graph and $k \in \mathbb{N}$. There is an $O(m)$ time algorithm computing a spanning subgraph of G that has at most $k(n - 1)$ edges and is k -connected (resp. k -edge-connected) if and only if G is k -connected (resp. k -edge-connected). Moreover, if G is k -connected (resp. k -edge-connected), the spanning subgraph contains a vertex of degree k .*

This algorithm preserves the 3-connectivity or respectively, the non 3-connectivity of G . Moreover, if G is 3-connected, the resulting graph contains a vertex of degree 3 and by a result of Halin [5], every vertex of degree 3 is incident to a contractible edge e . We get e by subsequently contracting each of the three incident edges and testing the resulting graph with the algorithm of Hopcroft and Tarjan [6] on 3-connectivity. Iteration of both subroutines gives us the whole contraction sequence in $O(n^2)$ time. However, the Hopcroft-Tarjan test is difficult to implement and we will give a much simpler algorithm that is capable of computing both characterizations later. In both approaches, we use the algorithm of Theorem 4 to preprocess the input graph in advance. As it allows for a very simple implementation, we do not certify this preprocessing step.

2.2 Barnette’s and Grünbaum’s Characterization and Its Inverse

The Barnette and Grünbaum operations (*BG-operations*) consist of the following operations on a 3-connected graph (see Figs. 1(a)–(c)).

- (a) add an edge xy (possibly a parallel edge)
- (b) subdivide an edge ab by vertex x and add the edge xy for $y \notin \{a, b\}$
- (c) subdivide two distinct, non-parallel edges by vertices x and y , respectively, and add the edge xy

In all three cases, let xy be the edge that was *added* by the BG-operation.

Theorem 5 (Barnette and Grünbaum [2], Tutte [18]) *A graph G is 3-connected if and only if G can be constructed from K_4 using BG-operations.*

Theorem 5 was proven in this notation by Barnette and Grünbaum [2], but implicitly described in a theorem about *nodal connectivity* by Tutte [18, Theorem 12.65].




Fig. 1 The three operations of Barnette and Grünbaum

If not stated otherwise, every construction sequence uses only BG-operations. Let a BG-operation be *basic*, if it does not create parallel edges and let a construction sequence be *basic*, if it only uses basic BG-operations.

Like in Theorem 3, we want the inverse of a BG-operation. Let *removing* the edge $e = xy$ of a graph be the operation of deleting e followed by smoothing x and y . An edge $e = xy$ in G is called *removable*, if removing e yields a 3-connected graph. We show that removing a removable edge $e = xy$ with $|N(x)| \geq 3$, $|N(y)| \geq 3$ and $|N(x) \cup N(y)| \geq 5$ is exactly the inverse of a BG-operation.

Theorem 6 *The following statements are equivalent:*

- (3) A simple graph G is 3-connected
- \iff There exists a sequence of removals from G to K_4 on removable edges $e = xy$ with $|N(x)| \geq 3$, $|N(y)| \geq 3$ and $|N(x) \cup N(y)| \geq 5$
- \iff There exists a construction sequence from K_4 to G using BG-operations
- \iff There exists a basic construction sequence from K_4 to G using BG-operations

Proof Theorem 5 establishes (3) \iff (5). Moreover, the proof of Theorem 5 in [2] implicitly shows that on simple graphs basic operations suffice. Thus, only the equivalence for (4) remains. We first prove (6) \implies (4) and then (4) \implies (5).

BG-operations operate by definition on 3-connected graphs, this holds in particular for the ones in sequence (6). Let G' be the graph obtained by a basic BG-operation in (6) that adds the edge $e = xy$. The operation can clearly be undone by removing e in G' . Since BG-operations preserve 3-connectivity with Theorem 5, $|N(x)| \geq 3$ and $|N(y)| \geq 3$ hold in G' .

It remains to show that $|N(x) \cup N(y)| \geq 5$ in G' . If $|N(x)| \geq 4$ or $|N(y)| \geq 4$, $|N(x) \cup N(y)| \geq 5$ follows, since x and y are neighbors and no self-loops exist. Thus, let $|N(x)| = |N(y)| = 3$. Having $N(x) \setminus \{y\} \neq N(y) \setminus \{x\}$ yields $|N(x) \cup N(y)| \geq 5$ as well, so let $N(x) \setminus \{y\}$ and $N(y) \setminus \{x\}$ contain the same two vertices a and b . If $|V(G')| > 4$, a or b must be adjacent to a vertex c that is neither adjacent to x nor y . But then $\{a, b\}$ is a separation pair, contradicting the 3-connectivity of G' . On the other hand, $|V(G')| = 4$ is only possible when operation (a) was performed as BG-operation, since operations (b) and (c) create new vertices. This gives a contradiction, as (a) is not a basic operation on K_4 .

We prove (4) \implies (5). Let G' be the graph containing a removable edge $e = xy$ that is removed in (4). Note that G' can have parallel edges due to previous removals but no self-loops. The removal can be undone by one of the three BG-operations. On smoothing of e , we count how many end vertices are deleted. This is either 0, 1, or 2. If no end vertex is deleted, removing e just deletes e which is inverted by operation (a). If exactly one end vertex, say $x \in V(G')$, is deleted, let f be the edge in which x was smoothed. Then (b) can be applied, because $y \notin f$ (see Fig. 2(a)) since




Fig. 2 Cases that would fail when undoing a removal

otherwise x would have had only 2 neighbors in G' , contradicting the assumption $|N(x)| \geq 3$.

If both end vertices, x and y , are deleted, let f_1 and f_2 be the edges in which x and y were smoothed, respectively. Operation (c) can only be applied if f_1 and f_2 are neither identical nor parallel. But $f_1 = f_2$ would again contradict $|N(x)| \geq 3$ in G' (see Fig. 2(b)), and f_1 being parallel to f_2 would contradict $|N(x) \cup N(y)| \geq 5$ in G (see Fig. 2(c)), since in that case $N(x) \cup N(y)$ consists only of x , y and the two vertices $f_1 \cap f_2$. □

We show that Barnette’s and Grünbaum’s characterization is algorithmically at least as powerful as Tutte’s by giving a simple linear time transformation. Lemma 7 allows us to focus on computing BG-operations only.

Lemma 7 *Every construction sequence using BG-operations can be transformed in linear time to the sequence (1) of contractions.*

Proof We transform every BG-operation in reverse order of the construction sequence to 0, 1 or 2 contractions each. Operation (a) yields no contraction while operation (b) yields the contraction of exactly one part of the subdivided edge (either xa or xb in Fig. 1). For an operation (c), let $e = ab$ and $f = vw$ be the edges that are subdivided with x and y . Both edges share at most one vertex; w.l.o.g. let $a = v$ be that vertex if it exists. We contraction the edges xb and yw in arbitrary order. In all cases, contractions inverse BG-operations except for the added edge xy , which is left over. But additional edges do not harm the 3-connectivity of the graph nor subsequent contractions. Thus, we have found a contraction sequence to K_4 unless the first contraction in the case of an operation (c) yields a graph H that is not 3-connected. Let H' be the 3-connected graph after the second contraction of the same operation (c). Then H can be obtained from H' by applying operation (b) and therefore is 3-connected. □

2.3 Identifying Intermediate Graphs with Subdivisions in G

Let $K_4 = G_0, G_1, \dots, G_z = G$ be the 3-connected graphs obtained in a construction sequence \mathcal{Q} to a simple 3-connected graph G using the basic BG-operations C_0, \dots, C_{z-1} . We can reverse \mathcal{Q} by starting with G and removing the added edges of BG-operations in reverse order. Suppose we would delete the added edge of every C_i instead of removing it and treat emerging paths containing interior vertices of degree




Fig. 3 The graphs G_0, \dots, G_z and S_0, \dots, S_z of a construction sequence of G . On graphs S_i , the *dashed edges and vertices* are in G but not in S_i and vertices depicted in *black* are *real vertices*. For example, the path $C_0 = e \rightarrow h \rightarrow g$ is a *BG-path* for S_0 , yielding S_1 . The *links* of S_1 are the paths C_0 , $a \rightarrow b \rightarrow c$ and the single edges $ae, ef, fc, cd, da, fg, gd$

2 as (topological) edges in G_i (see Fig. 3). Then iteratively paths are deleted instead of edges being removed and we obtain the sequence of subdivisions $G = S_z, \dots, S_0$ in G with S_0 being a K_4 -subdivision. This leads to the following proposition.

Proposition 8 *Let \mathcal{Q} be a construction sequence from a graph G_0 to G using BG-operations. Then G contains a subdivision of G_0 that is specified by \mathcal{Q} .*

In particular, Proposition 8 yields with Theorem 5 that every 3-connected graph contains a subdivision of K_4 (Theorem of J. Isbell [2]). Each S_i is a subdivision of G_i . Conversely, $G_i = \text{smooth}(S_i)$ for all $0 \leq i \leq z$, since smoothing a graph is precisely the inverse operation of subdividing a graph without vertices of degree two. The vertices x in S_i with $\text{deg}(x) \geq 3$ are called *real vertices*, because they correspond to vertices in G_i . Real vertices have at least 3 neighbors in G_i , because G_i is 3-connected.

Note that in non-basic construction sequences G_i can have parallel edges, although S_i is always simple.

Definition 9 Let the *links* of each S_i be the unique paths in S_i with only their end vertices being real. Let two links be *parallel* if they share the same end vertices. Then a *BG-path* for S_i is a path $P = x \rightarrow y$ in G with the following properties:

1. $S_i \cap P = \{x, y\}$.
2. If a link of S_i contains x and y , the end vertices of that link are x and y .

3. If x and y are inner vertices of links L_x and L_y of S_i , respectively, L_x and L_y are not parallel.

The links of S_i partition $E(S_i)$ because S_i is 2-connected, has therefore minimum degree two and is not a cycle. It is easy to see that every BG-path for S_i corresponds to a BG-operation on G_i and vice versa. We will exploit this duality in the next section.

In general, construction sequences are not bound to start with K_4 . Titov and Kelmans [7, 16] extended Theorem 5 by proving the existence of a construction sequence even when starting with an arbitrary 3-connected graph G_0 instead of K_4 , as long as a subdivision of G_0 is contained in G . This is a generalization of Theorem 5, since every 3-connected graph contains a K_4 -subdivision by Proposition 8.

Theorem 10 [7, 16] *Let G_0 be a 3-connected graph. A simple graph G is 3-connected and contains a subdivision of G_0 if and only if G can be constructed from G_0 using basic BG-operations.*

3 Prescribing Subdivisions

If G is 3-connected, the 3-connected base graph G_0 of the construction sequences of Theorem 5 and 10 corresponds to a subdivision $H \subset G$ of G_0 by Proposition 8. The proofs of Theorems 5 and 10 show only for a very special subdivision H in G that a construction sequence from H to G using BG-paths exists. In fact, the G_0 -subdivision containing the maximum number of edges in G is chosen for both theorems. The construction sequence is then obtained by adding longest BG-paths. Unfortunately, computing these depends heavily on solving the longest paths problem, which is known to be NP-hard even for 3-connected graphs [4].

Suppose we choose some subdivision H of G_0 in advance; we say that H is *prescribed*. Is it possible to strengthen Theorems 5 and 10 to start a construction sequence using BG-paths with H ? Such a result would provide an efficient computational approach to construction sequences, since it allows us to search the neighborhood of H in G for BG-paths, yielding a new prescribed subdivision of a 3-connected graph.

However, when restricted to basic operations it is not possible to prescribe H , as the minimal counterexample in Fig. 4 shows: Consider the graph G consisting of $H := K_4$ depicted in black with an additional vertex x connected to three vertices of H . Then every BG-path for H will create a parallel link, which is a path of length two having x as its middle vertex, although G is simple. But what if we drop the condition that construction sequences have to be basic? The following theorem shows that at this expense we can indeed start a construction sequence from any prescribed subdivision.

Fig. 4 Every possible BG-operation adds a parallel edge to the *black* subgraph





Fig. 5 The case $H \neq \text{smooth}(H)$. Dashed edges are in $E(G) \setminus E(H)$, arrows depict the BG-path $x' \rightarrow y'$



Theorem 11 Let G be a 3-connected graph and $H \subset G$ with H being a subdivision of a 3-connected graph. There is a BG-path for H in G . Moreover, for every link L of H of length at least 2 there is a parallel link (maybe L itself) that contains an inner vertex on which a BG-path for H starts.

Proof We distinguish two cases.

– $H \neq \text{smooth}(H)$.

Then links of length at least 2 exist in H and we pick an arbitrary one of them, say $T = a \rightarrow b$. Let x be an inner vertex of T and let I be the union of inner vertices of all parallel links of T . We show that there is a vertex in I on which a BG-path for H starts. By the 3-connectivity of G , the graph $G \setminus \{a, b\}$ is connected. Since H contains at least four vertices, there exists a path $P = x \rightarrow y$ in $G \setminus \{a, b\}$ with $y \in V(H) \setminus I$ (see Fig. 5). The path P has the Property 9.2. Let x' be the last vertex in P that is contained in I and let y' be the first vertex in P that is contained in $V(H) \setminus I$. Then the subpath $x' \rightarrow y'$ of P has Properties 9.1–9.3 and is a BG-path for H .

– $H = \text{smooth}(H)$.

Then H consists only of real vertices and since $H \neq G$, there is a vertex in $V(G) \setminus V(H)$ or an edge in $E(G) \setminus E(H)$. At first, assume that there is a vertex $x \in V(G) \setminus V(H)$. Then, by the 2-connectivity of G and Fan Lemma 1 we can find a path $P = y_1 \rightarrow x \rightarrow y_2$ with no other vertices in H than y_1 and y_2 . For P the Properties 9.1–9.3 hold, because every link in H is an edge. Now suppose that $V(G) = V(H)$ and let e be an edge in $E(G) \setminus E(H)$. Then e must be a BG-path for H , since both end vertices are real. □

In Theorem 11, non-basic operations can only occur in the case $H = \text{smooth}(H)$ when a path through a vertex of $V(G) \setminus V(H)$ is chosen. Although we cannot avoid that, it is possible to obtain a basic construction by augmenting the BG-operations with a fourth operation (d), which can be seen as combination of operations (a) and (b):

(d) connect a new vertex to three distinct vertices

Operation (d) preserves 3-connectivity with Lemma 2 and is basic, because each new edge ends on the new vertex. Whenever we encounter a vertex in $V(G) \setminus V(H)$ in Theorem 11, we know by Fan Lemma 1 and the 3-connectivity of G that there are three internally vertex-disjoint paths to real vertices in H with all inner vertices being in $V(G) \setminus V(H)$. Adding these paths to H is called an *expand* operation and corresponds to operation (d) in the smoothed graph. This gives the following result.

Theorem 12 *Let G be a simple graph and let H be a subdivision of a 3-connected graph. Then*

$$\begin{aligned}
 &G \text{ is 3-connected and } H \subseteq G \\
 \iff &\delta(G) \geq 3 \text{ and there exists a construction sequence from } H \text{ to } G \quad (7) \\
 &\text{using BG-paths}
 \end{aligned}$$


$$\begin{aligned}
 \iff &\delta(G) \geq 3 \text{ and there exists a basic construction sequence from } H \text{ to } G \quad (8) \\
 &\text{using BG-paths and the expand operation}
 \end{aligned}$$

Proof Let G be 3-connected and $H \subseteq G$. Then $\delta(G) \geq 3$ holds and if $H = G$, the desired construction sequences are empty and exist. If $H \subset G$, we can apply Theorem 11 iteratively with or without the additional expand operation and the construction sequences exist as well. For the sufficiency part, both construction sequences imply $H \subseteq G$, since only paths are added to construct G . Additionally, G must be 3-connected, as adding BG-paths to each S_i preserves S_{i+1} to be a subdivision of a 3-connected graph with Theorem 5, and $\delta(G) \geq 3$ ensures that the last subdivision G of a 3-connected graph is 3-connected itself. \square

A straightforward algorithm to compute Barnette’s and Grünbaum’s construction sequence of a 3-connected graph is to search iteratively for removable edges. But in contrast to the algorithm in Sect. 2.1 that computes contractible edges, this approach only leads to an $O(n^3)$ algorithm. The reason for the additional factor of n is that not all vertices with degree 3 must have an incident removable edge (see Fig. 6 for a counterexample on 9 vertices) and we have to try every edge in the worst case. Computing BG-paths instead of BG-operations allows us to obtain better running times. For this aim, we need to represent construction sequences.

An obvious representation of a construction sequence \mathcal{Q} would be to store the graph $G_0 = \text{smooth}(H)$ and in addition every BG-operation, which gives the se-

Fig. 6 A 3-connected graph having a vertex x of degree 3 with no incident edge being removable. Removing each incident edge of x results in the black separation pair



quence $G_0, \dots, G_z = G$. Unfortunately, the graphs G_i are not necessarily subgraphs of G_{i+1} , so we have to take care of relabeled edges when specifying each operation.

Whenever an edge e is subdivided as part of an operation (b) or (c), we specify it by its index in G_i followed by assigning new indices to the new degree-two vertex and one of the two new separated edge parts in G_{i+1} . The other edge part keeps the index of e .

Similarly, on operations (a) and (b), real end vertices of the added edge are specified by their indices in G_i . We assign a new index to the added edge in G_{i+1} , too. Finally, we have to impose the constraint that G_z is not just isomorphic but identical to G , meaning that vertices and edges of G_z and G are labeled by exactly the same indices, since otherwise we would have to solve the graph isomorphism problem to check that \mathcal{Q} really constructs G .

On the other hand, the identification of G_i with a subdivision S_i in G allows us to represent \mathcal{Q} without indexing issues: We just store $S_0 \subset G$ and the BG-paths C_0, \dots, C_{z-1} . Hence, we can represent each construction sequence \mathcal{Q} of G in the following two ways.

- *Edge representation:* Represent \mathcal{Q} by G_0 and a sequence of BG-operations, along with specifying new and old indices for each operation, such that G_z and G are labeled the same.
- *Path representation:* Represent \mathcal{Q} by S_0 and BG-paths C_0, \dots, C_{z-1} .

Both representations refer to the same sequence of graphs G_0, \dots, G_z and are linear in the graph size. Assuming the uniform cost model, this size is $\Theta(m)$, as the uniform cost model is independent on the size of the numbers processed; in particular, the space amount of each index is 1 instead of $O(\log n)$. The next lemma states that it does not matter which of the two representations we compute.

Lemma 13 *The edge and path representations of a construction sequence \mathcal{Q} can be transformed into each other in $O(m)$ time. Moreover, the representation computed is a unique representation of \mathcal{Q} .*

Proof Let G_0 and a sequence of BG-operations along with their specified indices on edges and vertices be given. If an operation O' subdivides an edge e' , we define $\beta(e', O')$ to be the edge that gets a new index. Let e be the added edge of an operation in \mathcal{Q} . Exploiting the duality of BG-paths and BG-operations, the edge e corresponds to a BG-path C , which will be subdivided by inserting $|C| - 1$ vertices in the construction sequence. To compute the BG-path C from e , we have to keep track of the at most $|C| - 1$ operations that subdivide e and glue the subdivided parts back together.

Whenever an operation $O \in \mathcal{Q}$ subdivides e , we store a pointer at $\beta(e, O)$ to e . Moreover, on all edges f that point to e and are subdivided by an operation O'' , we store a pointer at $\beta(f, O'')$ to e . In both cases, we append $\beta(e, O)$ (resp. $\beta(f, O'')$) to a list stored on the edge e . Therefore, we keep track of all new edges $\beta(e, O)$ and $\beta(f, O'')$ that subdivided C . Eventually, we get all the edges in which C got subdivided by augmenting the list of e with e itself. Hence, we have computed the set of edges that C consists of. Since G_z has the same labeling as G , the indices of e and all other edges in C are still contained in G .

The set of edges is not necessarily in the order of appearance in C , but this can be easily fixed in time $O(|C|)$ by temporarily storing the incidence information of every vertex in C and extracting the BG-path C from a degree-one vertex. In order to compute S_0 , we analogously maintain pointers for each edge of G_0 and get the links of S_0 . Since the links of S_0 together with C_0, \dots, C_{z-1} partition $E(G) \setminus E(S_0)$, the running time is $O(m)$.

Conversely, let S_0 and the sequence C_0, \dots, C_{z-1} of BG-paths be given. We remove BG-paths in reversed order from G by deleting their edge (there is only one edge left this way, the one added in the corresponding BG-operation) followed by smoothing their end vertices. Therefore, we pass through the graph sequence G_z, \dots, G_0 and get G_0 . If both end vertices of the BG-path $C_i = a \rightarrow b$ are real after deleting ab , we can keep their index and construct the corresponding BG-operation (a).

Otherwise, let a have degree 2 after deleting ab and let e and f be its incident edges. When a is smoothed, we can assign the lowest index of e and f to the new edge. Thus, all indices that are necessary for constructing the operation (b) can be found in constant time. If additionally b has degree 2, the same procedure constructs operation (c). It remains to show that always unique representations of \mathcal{Q} are computed. The path representation with BG-paths is by definition unique. In contrast, edge representations can vary in their indices. However, picking the incident edge with lowest index before smoothing a vertex creates a unique representation, since all edge indices of G are given. □

If G is simple, the construction sequences (7) and (8) can be transformed into each other efficiently.


Lemma 14 *For simple graphs G , the construction sequences (7) and (8) can be transformed into each other in $O(m)$.*

Proof With Lemma 13, we can assume that the construction sequence (7) is given in the path representation. We will rearrange the order of BG-paths to generate basic operations. For each BG-path P , its position in the construction sequence and a pointer to the first BG-path $F(P)$ that ends at an inner vertex of P (if that path exists) is stored. We define the position of each link of S_0 as 0. Performing a bucket sort on the lower end vertices of each BG-path (lower in any given total order on $V(G)$) followed by a stable bucket sort on the remaining end vertices gives a list of BG-paths sorted in lexicographic order of the end vertex. This list can be used to efficiently group paths that have the same end vertices.

Let R_{ab} be the set of all BG-paths and links of S_0 having end vertices a and b . We apply the following rule: If a path $P \in R_{ab}$ has length one and does not have the first position of all paths in R_{ab} , we append it to the end of the construction sequence and remove it from R_{ab} . This does not harm the construction sequence, since a and b were already real and P has no inner vertices.

The path with the first position in R_{ab} cannot lead to a non-basic operation. We look at all other paths $P \in R_{ab}$, which are possibly non-basic, but must contain an inner vertex w that is an end vertex of the subsequent BG-path $F(P) = v \rightarrow w$.

Fig. 7 No expand operation can be formed



Without harming the construction sequence, P can be moved to the position of $F(P)$, since a and b were already real and no inner vertex of P is part of a BG-path before $F(P)$ is applied. If v is real at the point in time when $F(P)$ is applied, we can glue P and $F(P)$ together to an expand operation, which is basic due to its new vertex w . Otherwise, v is an inner vertex of a link (see Fig. 7) and P and $F(P)$ can be replaced with the two BG-paths $v \rightarrow a$ and $b \rightarrow w$. Both BG-paths are basic, since they contain end vertices of degree 2.

Conversely, the three internally vertex-disjoint paths of each expand operation can be easily split into two BG-paths, possibly inducing non-basic operations. □

4 Certifying and Testing 3-connectivity in $O(n^2)$


We use construction sequences in the path representation as a certificate for the 3-connectivity of graphs. This leads to a new, certifying method for testing graphs on being 3-connected. The total running time of this method is $O(n^2)$. This is dominated by the time needed for finding the construction sequence and every improvement made there will automatically result in a faster 3-connectivity test. The input graph is a multigraph and does not have to be biconnected nor connected. We follow the steps:

- Apply the linear-time algorithm of Nagamochi and Ibaraki to the input graph G' in order to get a graph $G = (V, E)$ where the number of edges is in $O(n)$.
- Try to compute a K_4 -subdivision in G in $O(n)$.
 - Success: Let S_0 be the K_4 -subdivision.
 - Failure: Return a separation pair or cut vertex.
- Try to compute a construction sequence from prescribed S_0 to G in $O(n^2)$.
 - Success: Return the construction sequence.
 - Failure: Return a separation pair or cut vertex.

The graph G output by Nagamochi and Ibaraki is 3-connected if and only if the input graph G' is 3-connected. We first describe how to find a K_4 -subdivision by one Depth First Search (DFS), which as a byproduct eliminates self-loops and parallel edges and sorts out graphs that are not connected or have vertices with degree less than 3.

Lemma 15 *Let G be a graph on at least 4 vertices. There is a simple DFS-based algorithm that computes either a K_4 -subdivision or a separation pair in G in time $O(n + m)$.*

Fig. 8 Finding a K_4 -subdivision. Dashed edges depict (possibly empty) paths, arcs depict backedges



Proof Let T be a DFS-tree of G and let a (resp. b) be the vertex in T that is visited first (resp. second). We can assume that both a and b have exactly one child in T , respectively, as otherwise a and b form a separation pair. We choose two arbitrary neighbors c and d of a that are different from b (see Fig. 8). W.l.o.g., let d be visited later by the DFS than c . Let i be the least common ancestor of c and d in T . Then $i \neq b$, as b has exactly one child in T . Since $d \neq i$ holds, let j be the child of i that is contained in the path $i \rightarrow d$ in T .

If G is 3-connected, we can find a backedge e that starts on a vertex z in the subtree rooted at j and ends on an inner vertex z' of $a \rightarrow i$ in time $O(n)$. If e does not exist, a and i form a separation pair. Otherwise, let k be the nearest ancestor of z contained in the path $j \rightarrow d$ in T . Each of the three backedges ac , ad and e close a cycle when added to T , resulting in six internally vertex-disjoint paths connecting the vertices in $\{a, z'\}$, $\{z', i\}$, $\{i, k\}$, $\{k, a\}$, $\{z', k\}$ and $\{a, i\}$, respectively. Thus, we have found a K_4 -subdivision with real vertices a, z', i and k . \square

We now show how to carry out the last step of the algorithm. Let H be the computed K_4 -subdivision. In order to find the construction sequence, we use the path representation and try to find iteratively BG-paths along the lines of Theorem 11.

Lemma 16 *Let H be a subdivision of a 3-connected graph that is contained in a 3-connected graph G with $m = O(n)$. There is an algorithm that computes a BG-path for H in time $O(n)$.*

Proof We compute the links of H , assign an index to every link and store this index on each of the inner vertices of that link in $O(n)$ total time. Moreover, we maintain a pointer on that index that points to the end vertices of the link. It remains to show how to find a BG-path along the lines of Theorem 11.

In case $H \neq \text{smooth}(H)$, we pick an arbitrary vertex x of degree two. Let $T = a \rightarrow b$ be the link that contains x and let W be the set of vertices $V(H) \setminus V(T)$ minus all vertices in parallel links of T . We compute the path $P = x \rightarrow y'$ by temporarily deleting a and b and performing a DFS from x that stops on the first vertex $y' \in W$. We can check whether a vertex lies in a parallel link of T in constant time by comparing the end vertices of its containing link with a and b . Thus, the subpath $x' \rightarrow y'$ with x' being the last vertex contained in T or in a parallel link of T is a BG-path and can be found efficiently. Similarly, in case $H = \text{smooth}(H)$ we delete temporarily all edges in $E(H)$ and start a DFS from a vertex $x \in V(H)$ that has

an incident edge in the remaining graph. The traversal is stopped on the first vertex $y \in V(H) \setminus \{x\}$ and the path $x \rightarrow y$ is then the desired BG-path. \square

Each time we have found a BG-path, a new index is assigned to it and stored on each of its inner vertices. We also store a pointer on this index that points to both end vertices of the BG-path. These update operations can be carried out in $O(n)$ time as well. Iterating the procedure, a construction sequence can be found in time $O(n^2)$ if the input graph G is 3-connected.

By Theorem 12, if G is not 3-connected, no construction sequence can exist. In that case, it remains to show that we can always find a separation pair or cut vertex. For some subdivision $H \subset G$ of a 3-connected graph, the DFS starting at vertex x must fail to find a new BG-path. If $H \neq \text{smooth}(H)$, the end vertices of the link that contains x must form a separation pair. Otherwise, $H = \text{smooth}(H)$ and x must be a cut vertex due to Theorem 11. Thus, if G is not 3-connected, the algorithm returns either a separation pair or a cut vertex.

Theorem 17 *The construction sequences (7) and (8) can be computed in $O(n^2)$ and establish a certifying 3-connectivity test with the same running time.*

This raises the following open question.

Question 18 Is there a certifying algorithm with running time $o(n^2)$ that computes a certificate for 3-connectivity of at most linear size?

4.1 Verifying the Construction Sequence


We could validate the certificate by transforming the path representation to the edge representation using Lemma 13 and checking the validity of the BG-operations by comparing indices, but there is a more direct way. First, it can be checked in linear time that all BG-paths C_0, \dots, C_{z-1} are paths in G and that these paths partition $E(G) \setminus E(S_0)$. We try to remove the BG-paths C_{z-1}, \dots, C_0 from G in that order (i.e., we delete the paths followed by smoothing its end vertices). If the certificate is valid, this is well defined since all removed BG-paths are edges. On the other hand, we can detect longer BG-paths $|C_i| \geq 2$ before their removal. In that case, the certificate is not valid, since the inner vertices of C_i are not attached to BG-paths C_j , $j > i$.

We verify that every removed $C_i = ab$ corresponds to a BG-operation by using Definition 9 of BG-paths, and start with checking that a and b lie in our current subgraph for Condition 9.1.

Conditions 9.2 and 9.3 can now be checked in constant time: Consider the situation immediately after the deletion of ab , but before smoothing a and b . Then all links in our subgraph are single edges, except possibly the ones containing a and b as inner vertices.

Therefore, Condition 9.2 is not met for C_i if a is a neighbor of b and at least one of the vertices a and b has degree two (see Fig. 9 for possible configurations). Condition 9.3 is not met if $N(a) = N(b)$ and both a and b have degree

Fig. 9 Cases where Condition 9.2 fails when $a \in N(b)$



two. Both conditions can be checked in constant time. Note that encountering BG-paths $C_{z-1}, C_{z-2}, \dots, C_i$ does not necessarily imply that the current subgraph is 3-connected, since a path C_j with $j < i$ and being no BG-path might occur later.

It remains to validate that the graph after removing all BG-paths equals K_4 . This can be done in constant time by checking it on being simple and having exactly 4 vertices of degree three.

Theorem 19 *The sequences (4)–(8) can be checked on validity in time linearly dependent on their length.*

Acknowledgement We want to thank the anonymous reviewers for their detailed and helpful comments on the original version of this paper.

References

1. Albroscheit, S.: Ein Algorithmus zur Konstruktion gegebener 3-zusammenhängender Graphen (in German). Diploma thesis, Freie Universität Berlin (2006)
2. Barnette, D.W., Grünbaum, B.: On Steinitz’s theorem concerning convex 3-polytopes and on some properties of 3-connected graphs. In: Many Facets of Graph Theory. Lecture Notes in Mathematics, pp. 27–40. Springer, Berlin (1969)
3. Blum, M., Kannan, S.: Designing programs that check their work. In: Proceedings of the 21st Annual ACM Symposium on Theory of Computing (STOC’89), New York, pp. 86–97 (1989)
4. Garey, M.R., Johnson, D.S., Tarjan, R.E.: The planar Hamiltonian circuit problem is NP-complete. SIAM J. Comput. **5**(4), 704–714 (1976)
5. Halin, R.: Zur Theorie der n-fach zusammenhängenden Graphen. Abh. Math. Semin. Univ. Hamb. **33**(3), 133–164 (1969)
6. Hopcroft, J.E., Tarjan, R.E.: Dividing a graph into triconnected components. SIAM J. Comput. **2**(3), 135–158 (1973)
7. Kelmans, A.K.: Graph expansion and reduction. In: Algebraic Methods in Graph Theory, Szeged, Hungary, vol. 1, pp. 317–343 (1978)
8. Kratsch, D., McConnell, R.M., Mehlhorn, K., Spinrad, J.P.: Certifying algorithms for recognizing interval graphs and permutation graphs. SIAM J. Comput. **36**(2), 326–353 (2006). Preliminary version in SODA 2003, pp. 158–167
9. Mehlhorn, K., Näher, S.: From algorithms to working programs: On the use of program checking in LEDA. In: Proceedings of the 23rd International Symposium on Mathematical Foundations of Computer Science (MFCS’98), pp. 84–93 (1998)
10. Mehlhorn, K., Näher, S., Seel, M., Seidel, R., Schilz, T., Schirra, S., Uhrig, C.: Checking geometric programs or verification of geometric structures. Comput. Geom. Theory Appl. **12**(1–2), 85–103 (1999)

11. Menger, K.: Zur allgemeinen Kurventheorie. *Fund. Math.* **10**, 96–115 (1927)
12. Nagamochi, H., Ibaraki, T.: A linear-time algorithm for finding a sparse k -connected spanning subgraph of a k -connected graph. *Algorithmica* **7**(1–6), 583–596 (1992)
13. Schmidt, J.M.: Construction sequences and certifying 3-connectedness. In: Proceedings of the 27th International Symposium on Theoretical Aspects of Computer Science (STACS'10), Nancy, France, pp. 633–644 (2010)
14. Thomassen, C.: Kuratowski's theorem. *J. Graph Theory* **5**(3), 225–241 (1981)
15. Thomassen, C.: Reflections on graph theory. *J. Graph Theory* **10**(3), 309–324 (2006)
16. Titov, V.K.: A constructive description of some classes of graphs. Ph.D. Thesis, Moscow (1975)
17. Tutte, W.T.: A theory of 3-connected graphs. *Indag. Math.* **23**, 441–455 (1961)
18. Tutte, W.T.: Connectivity in graphs. In: *Mathematical Expositions*, vol. 15. University of Toronto Press, Toronto (1966)
19. Vo, K.-P.: Finding triconnected components of graphs. *Linear Multilinear Algebra* **13**, 143–165 (1983)
20. Vo, K.-P.: Segment graphs, depth-first cycle bases, 3-connectivity, and planarity of graphs. *Linear Multilinear Algebra* **13**, 119–141 (1983)
21. West, D.B.: *Introduction to Graph Theory*. Prentice Hall, New York (2001)