

Extending Convex Partial Drawings of Graphs

Tamara Mchedlidze¹ · Martin Nöllenburg¹ ·
Ignaz Rutter¹

Received: 10 February 2014 / Accepted: 11 June 2015 / Published online: 24 June 2015
© Springer Science+Business Media New York 2015

Abstract Given a plane graph G (i.e., a planar graph with a fixed planar embedding and outer face) and a biconnected subgraph G' with a fixed planar straight-line convex drawing, we consider the question whether this drawing can be extended to a planar straight-line drawing of G . We characterize when this is possible in terms of simple necessary conditions, which we prove to be sufficient. This also leads to a linear-time testing algorithm. If a drawing extension exists, one can be computed in the same running time.

Keywords Extension of a partial drawing · Fixed cycle · Fixed inner face · Convex shape · Straight-line drawing · Linear-time algorithm

1 Introduction

The problem of extending a partial drawing of a graph to a complete one is a fundamental problem in graph drawing that has many applications, e.g., in dynamic graph drawing and graph interaction. This problem has been studied most in the planar setting and often occurs as a subproblem in the construction of planar drawings.

A preliminary version of this work appeared in the Proceedings of the 21st International Symposium on Graph Drawing (GD 2013) [8].

✉ Tamara Mchedlidze
mched@iti.uka.de
Martin Nöllenburg
noellenburg@kit.edu
Ignaz Rutter
rutter@kit.edu

¹ Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

The earliest example of such a drawing extension result are so-called Tutte embeddings. In his seminal paper “How to Draw a Graph” [12], Tutte showed that any triconnected planar graph admits a convex drawing with its outer vertices fixed to an arbitrary convex polygon. The strong impact of this fundamental result is illustrated by its, to date, more than 850 citations and the fact that it received the “Best Fundamental Paper Award” from GD’12. The work of Tutte has been extended in several ways. In particular, it has been strengthened to only require polynomial area [5], even in the presence of collinear points [3]. Hong and Nagamochi extended the result to show that triconnected graphs admit convex drawings when their outer vertices are fixed to a star-shaped polygon [6]. For general subdrawings, the decision problem of whether a planar straight-line drawing extension exists is NP-hard [10]. Pach and Wenger [9] showed that every subdrawing of a planar graph that fixes only the vertex positions can be extended to a planar drawing with $O(n)$ bends per edge and that this bound is tight. The drawing extension problem has also been studied in a topological setting, where edges are represented by non-crossing curves. In contrast to the straight-line variant, it can be tested in linear time whether a drawing extension of a given subdrawing exists [1]. Moreover, there is a characterization via forbidden substructures [7]. In case the topology of a straight-line drawing of a subgraph H allows a drawing extension, Timothy et al. [4] showed that there exists an extension with at most $72|V(H)|$ bends per edge.

In this paper, we study the problem of finding a planar straight-line drawing extension of a plane graph for which an arbitrary biconnected subgraph has been fixed in form of a planar straight-line convex drawing. The question of whether the internal faces of the subgraph can be completed have been extensively studied [3, 5, 12]. We fill the gap by studying what happens with the outer face. In particular, we study the problem of finding a planar straight-line drawing extension of a plane graph for which an arbitrary cycle has been fixed to a convex polygon. It is easy to see that a drawing extension does not always exist in this case; see Fig. 1. Let G be a plane graph and let C be a simple cycle of G drawn as a convex polygon Γ_C in the plane. The following two simple conditions are clearly necessary for the existence of a drawing extension: (i) C has no chords that must be embedded outside of C and (ii) for every vertex v with neighbors on C that must be embedded outside of C there exists a placement of v outside Γ_C such that the straight-line drawing of the graph induced by C and v is plane and bounded by the same cycle as in G . We show in this paper that these two conditions are in fact sufficient. Both conditions can be tested in linear time, and if they are satisfied, a corresponding drawing extension can be constructed within the same time bound (Theorem 4). This result implies a linear time algorithm to test the extendability of a planar straight-line convex drawing of a biconnected subgraph (Theorem 5).

Our paper starts with some necessary definitions (Sect. 2) and useful combinatorial properties (Sect. 3). Our main result is Theorem 4, the proof of which has two steps. We first show in Sect. 4 that the conditions are sufficient if Γ_C is one-sided (i.e., it has an edge whose incident inner angles are both less than 90°). Afterward, we show in Sect. 5 that, for an arbitrary convex polygon Γ_C , we can place the neighborhood of C in such a way that the drawing is planar, and such that the boundary C' of its outer face is a one-sided polygon $\Gamma_{C'}$. Moreover, our construction ensures that the remaining graph satisfies the conditions for extendability of $\Gamma_{C'}$. The general result then follows directly from the one-sided case.

2 Definitions and a Necessary Condition

2.1 Plane Graphs and Subgraphs

A graph $G = (V, E)$ is *planar* if it has a drawing Γ in the plane \mathbb{R}^2 without edge crossings. Drawing Γ subdivides the plane into connected regions called *faces*; the unbounded region is the *outer* and the other regions are the *inner* faces. The boundary of a face is called *facial cycle*, and *outer cycle* for the outer face. The cyclic ordering of the edges around each vertex of Γ together with the description of the outer face of Γ characterize a class of drawings with the same combinatorial properties, which is called an *embedding* of G . A graph G with a planar embedding is called *plane graph*. A *plane subgraph* H of G is a subgraph of G together with a planar embedding that is the restriction of the embedding of G to H .

Let G be a plane graph and let C be a simple cycle of G . Cycle C is called *strictly internal*, if it does not contain any vertex of the outer face of G . A chord of C is called *outer* if it lies outside C in G . A cycle without outer chords is called *outerchordless*. The *subgraph of G inside C* is the plane subgraph of G that is composed of the vertices and edges of C and all vertices and edges of G that lie inside C .

2.2 Connectivity

A graph G is *k-connected* if the removal of any set of at most $k - 1$ vertices of G does not disconnect the graph. For $k = 2, 3$ a k -connected graph is also called *biconnected* and *triconnected*, respectively. An internally triangulated plane graph is triconnected if and only if there is no edge connecting two non-consecutive vertices of its outer cycle (see, for example, [2]).

2.3 Star-Shaped and One-Sided Polygons

Let Π be a polygon in the plane. Two points inside or on the boundary of Π are mutually *visible*, if the open straight-line segment connecting them belongs to the interior of Π . The *kernel* $K(\Pi)$ of polygon Π is the set of all the points inside Π from which all vertices of Π are visible. We say that Π is *star-shaped* if $K(\Pi) \neq \emptyset$. We observe that, with our definition of mutual visibility, a star-shaped polygon ensures that its kernel has positive area.

A convex polygon Π with k vertices is called *one-sided*, if there exists an edge e (i.e., a line segment) of Π such that the orthogonal projection to the line supporting e maps all polygon vertices onto segment e . Then e is called the *base edge* of Π . Without loss of generality let $e = (v_1, v_k)$ and v_1, \dots, v_k be the clockwise ordered sequence of vertices of Π .

2.4 Extension of a Drawing

Let G be a plane graph and let H be a plane subgraph of G . Let Γ_H be a planar straight-line drawing of H . We say that Γ_H is *extendable* if drawing Γ_H can be completed to a




Fig. 1 **a** A botanical flower with petals and stamen. (Image courtesy of Pearson Scott Foresman). **b** The convex polygon Γ_C of cycle C is drawn in black. Vertices $w_{i,j}$ and $w_{l,k}$ are petals of C with $w_{l,k} < w_{i,j}$, vertices w and w' are stamens of v_i . Petal $w_{l,k}$ is realizable, whereas petal $w_{i,j}$ is not, since vertex $w_{i,j}$ cannot be placed in the plane without changing the embedding or intersecting C

planar straight-line drawing Γ_G of the plane graph G . Then Γ_G is called an *extension* of Γ_H . A planar straight-line drawing of G is called *convex*, if every face of G (including the outer face) is represented as a convex polygon.

The following theorem by Hong and Nagamochi [6] shows the extendability of a prescribed star-shaped outer face of a plane graph.

Theorem 1 (Hong and Nagamochi [6]) *Every drawing of the outer face of a triconnected planar graph G as a star-shaped polygon can be extended to a convex drawing of G . Such a drawing can be computed in linear time.*

2.5 Petals and Stamens

Let G be a plane graph, and let P_{uv} be a path in G between vertices u and v . Its subpath from vertex a to vertex b is denoted by $P_{uv}[a, b]$. Let C be a simple cycle of G , and let v_1, \dots, v_k be the vertices of C in clockwise order. Given two vertices v_i and v_j of C , we denote by $C[v_i, v_j]$ the subpath of C encountered when we traverse C clockwise from v_i to v_j . Assume that C is represented by a convex polygon Γ_C in the plane. We say that a vertex v_i , $1 \leq i \leq k$ of Γ_C is *flat*, if $\angle v_{i-1}v_i v_{i+1} = \pi$. Throughout this paper, we assume that convex polygons do not have flat vertices.

A vertex $w \in V(G) \setminus V(C)$ adjacent to at least two vertices of C and lying outside C in G , is called a *petal* of C ; see Fig. 1. Consider the plane subgraph G' of G induced by the vertices $V(C) \cup \{w\}$. Vertex w appears on the boundary of G' between two vertices of C , i.e., after some $v_i \in V(C)$ and before some $v_j \in V(C)$ in clockwise order. To indicate this fact, we will denote petal w by $w_{i,j}$. Edges $(w_{i,j}, v_i)$ and $(w_{i,j}, v_j)$ are called the *outer edges* of petal $w_{i,j}$. The subpath $C[v_i, v_j]$ of C is called the *base* of the petal $w_{i,j}$. A vertex v_f is *internal to petal $w_{i,j}$* , if it appears on C after v_i and before v_j in clockwise order. A petal $w_{i,i+1}$ is called *trivial*. A vertex of $V(G) \setminus V(C)$ adjacent to exactly one vertex of C is called a *stamen* of C .

Let v be a petal of C and let u be either a petal or a stamen of C , we say that u is *nested* in v , and denote this fact by $u < v$, if u lies in the cycle delimited by the base

and the outer edges of petal v . For two stamens u and v , neither $u < v$ nor $v < u$. So for each pair of stamens or petals u and v we have either $u < v$, or $v < u$, or none of these. This relation $<$ is a partial order. A petal or a stamen u of C is called *outermost* if it is maximal with respect to $<$.

2.6 Necessary Petal Condition

Let again G be a plane graph and let C be an outerchordless cycle of G represented by a convex polygon Γ_C in the plane. Let $w_{i,j}$ be a petal of C . Let G' be the plane subgraph of G , induced by the vertices $V(C) \cup \{w_{i,j}\}$. We say that $w_{i,j}$ is *realizable* if there exists a planar drawing of G' which is an extension of Γ_C . This gives us the necessary condition that Γ_C is *extendable only if each petal of C is realizable*. In the rest of the paper we prove that this condition is also sufficient.

3 Combinatorial Properties of Graphs and Petals

In this section, we derive several properties of petals in graphs, which we use throughout the construction of the drawing extension in the remaining parts of this paper. Proposition 1 allows us to restrict our attention to maximal plane graphs for which the given cycle C is strictly internal. The remaining lemmas are concerned with the structure of the (outermost) petals of C in such a graph.

Proposition 1 *Let G be a plane graph on n vertices and let C be a simple outerchordless cycle of G . There exists a plane supergraph G' of G with $O(n)$ vertices such that (i) G' is maximal plane, (ii) there are no outer chords of C in G' , (iii) each petal of G' with respect to C is either trivial or has the same neighbors on C as in G , and (iv) C is strictly internal to G . Graph G' can be constructed within $O(n)$ time.*

Proof First, we arbitrarily triangulate the graph in the interior of C . Clearly, this creates neither outer chords nor new petals. For triangulating the graph outside of C , we have to be more careful to avoid the creation of outer chords and potentially unrealizable petals. We proceed in three steps. First, we ensure that, for each edge e of C , the incident face outside of C is a triangle. In this phase, we create new petals, all of which are realizable. Second, we ensure that all faces incident to vertices of C are triangles without introducing any new petals. After this step, all faces incident to vertices of C are triangles and C is strictly internal. We then triangulate the remaining faces arbitrarily. In the following, we describe these steps in more detail and argue their correctness.

In the first step, we create, for each edge e of C , a new vertex v_e that is adjacent to the endpoints of e and embed it in the face incident to e outside of C (refer to Fig. 2a). Note that each such vertex v_e is a trivial petal, and it is hence realizable. Clearly, after performing this operation for each edge e of C , every face incident to an edge of C is a triangle. Note that, besides the new trivial petals, we do not create or modify any other petals. Moreover, we introduce no chords of C since any new edge is incident to a new vertex. Clearly, this step can be performed in $O(|C|)$ time.




Fig. 2 **a** Graph G is depicted by *black*, cycle C is *bold*. Gray vertices are added during the first step of the augmentation. **b** Example of an augmentation during the second step. Face f is considered, which is depicted by *dashed lines*. Vertex v_f is added to lie inside f and connected to u , v and w . As a result v has no other incident non-triangular face

For the second step, we traverse all faces incident to vertices of C that are not triangles. Let f be such a face that is incident to some vertex v of C ; see Fig. 2b. Notice that f can be either an inner or the outer face of G . Let (v, u) and (v, w) be the two edges incident to v that bound f . Note that (v, u) and (v, w) do not belong to C since, according to step 1, f would be a triangle in this case. Hence, since there are no outer chords of C , u and w are not in C . We create a new vertex v_f , connect it to u , v and w , and embed it in f . This reduces the number of non-triangular faces incident to v by 1. Note that this operation neither produces a petal nor a chord of C . After treating all faces incident to vertices of C in this way, all inner faces incident to vertices of C are triangles and C is strictly internal. This step is accomplished within time proportional to the number of non-triangular faces incident to the vertices of C , i.e., in $O(n)$ time.

In the third step, we triangulate the remaining faces arbitrarily. None of the edges added in this step is incident to a vertex of C . Hence, this produces neither petals nor chords. Finally, if C is not strictly internal in the resulting graph, there is at most one vertex of C on the outer face. (To see this, observe that the outer face is a triangle, and if two of its vertices belong to C , the edge between them would be an outer chord, contradicting the properties of our construction.) If this case occurs, we add an additional vertex into the outer face and connect it to all three vertices of the outer face. Afterwards C is strictly internal. Again this can neither produce a petal nor an outer chord.

The resulting graph G' is triangulated and each petal of C in G' is either trivial or has the same neighbors on C as in G . Obviously, the procedure adds only $O(n)$ vertices and runs in $O(n)$ time. This concludes the proof. \square

Lemma 1 *Let G be an n -vertex triangulated planar graph with a strictly internal outerchordless cycle C . Then the following statements hold. (i) Each vertex of C that is not internal to an outermost petal is adjacent to two outermost petals. (ii) There is a simple cycle C' whose interior contains C and that is formed by the outermost stamens and petals of C . Finally, cycle C' can be found in $O(n)$ time.*




Fig. 3 **a** Illustration for the proof of Lemma 1. Cycle C is shown by **bold black curve**. **b** Illustration for the proof of Lemma 2. Cycle C is depicted by **bold black**. Three paths from u to v are shown by **bold grey curves**

Proof For (i), observe that each edge of C is incident to a triangle outside of C , whose tip then is a petal. Thus, every vertex of C is adjacent to at least two petals p_1 and p_2 such that none of them contains the other one; see Fig. 3a. For a vertex v of C that is not contained in the base of an outermost petal, this implies that there exist distinct outermost petals p'_1 and p'_2 with $p_1 \prec p'_1$ and $p_2 \prec p'_2$. Since v is not internal to any of them, it must be incident to one of the outer edges of each of them.

For (ii), let v be a vertex of C and let u and w be two stamens or petals that are adjacent in the circular ordering of neighbors around v ; refer again to Fig. 3a. Then (v, u) and (v, w) together bound a face, which must be a triangle, implying that u and w are adjacent. Applying this argument to all adjacent pairs of outermost petals or stamens around C yields the claimed cycle in G that traverses exactly the outermost petals and stamens. Obviously, this construction can be carried out in $O(n)$ time. \square

The following lemma is an auxiliary lemma that is only used in the proof of Lemma 3.

Lemma 2 *Let G be an n -vertex triangulated planar graph with a strictly internal outerchordless cycle C . Let $u \in C$ be internal to an outermost petal v . Then there exists a chordless path from u to v that contains no other vertices of C , which can be found in $O(n)$ time.*

Proof If u is adjacent to v , there is nothing to prove. Otherwise, let v_1 and v_2 be the first vertex on C to the left and right of u , respectively, that are adjacent to v ; see Fig. 3b. Let C' denote the cycle consisting of $C[v_1, v_2]$ and v together with two edges (v, v_1) and (v, v_2) . The graph consisting of C' and all edges and vertices embedded inside C' is inner-triangulated and chordless, and hence triconnected. Thus, by Menger’s theorem, the graph contains three vertex-disjoint paths from u to v , which can be constructed in $O(n)$ time [11]. The middle one of these paths cannot contain any vertices of C . We obtain the claimed path from the middle path by iteratively short-cutting it using chords. This procedure can be accomplished in time linear in the number of vertices in the path. \square




Fig. 4 Illustration for the proof of Lemma 3. Cycle C is solid black, cycle C_{shell} is solid blue. Edges incident to petals are dashed. The identified paths are bold gray. **a** Case 1a: Vertex b is contained in the counterclockwise path from x to y . **b–c** Case 1b: Vertex b is contained in the clockwise path from x to y (Color figure online)

Lemma 3 *Let G be an n -vertex maximal planar graph with a strictly internal out-erchordless cycle C . Let u and v be two adjacent vertices on C that are not internal to the same petal. Then there exists a third vertex w of C such that there exist three chordless disjoint paths from u , v and w to the vertices of the outer face of G such that none of them contains other vertices of C . Such paths can be constructed in $O(n)$ time.*

Proof Let C_{shell} denote the cycle going through the outermost petals and stamens of C , which exists by Lemma 1. It follows from Menger’s theorem that there exists a set $A = \{a, b, c\}$ of three vertices on C_{shell} that have disjoint paths to the outer face, such that none of them contains any other vertices of C_{shell} . Our goal is to identify vertex w and to connect the vertices u , v , w by vertex-disjoint paths to the vertices of A such that together we obtain the claimed paths. Assume without loss of generality that v is the immediate successor of u on C in counterclockwise order.

We distinguish cases based on the number of outermost petals.

Case 1: There are exactly two outermost petals x and y . Since u and v are not both internal to the same petal, we can assume that u is in the base of petal x and v is in the base of petal y . Note further that one of u and v must be contained in the bases of both petals, and hence adjacent to them; see Fig. 4. Without loss of generality, we assume that v is. The case that u is adjacent to both of them is completely symmetric.

We consider the subpaths of C_{shell} from x to y . At least one of them contains a vertex of A in its interior. If one of them contains all three of them in its interior, let b denote the middle one, otherwise, let b denote an arbitrary one. Let a and c denote the vertices of $A \setminus \{b\}$ that occur before and after b in counterclockwise direction along C_{shell} .

We distinguish cases based on whether b is contained in the clockwise or in the counterclockwise path from x to y .

Case 1a: Vertex b is contained in the counterclockwise path from x to y . See Fig. 4a for an illustration. It then follows that b is a stamen of v . We choose w as an arbitrary vertex in the base of petal y such that it is distinct from u and v (note that this is always possible since C has length at least 3).




Fig. 5 Illustration for the proof of Lemma 3. Cycle C is solid black, cycle C_{shell} is solid blue. Edges incident to petals are dashed. The identified paths are bold gray. **a** Case 2a: **b–c** Case 2b (Color figure online)

Now the claimed paths can be constructed as follows. For u , take the path from u to x (Lemma 2), and from there along C_{shell} to a , avoiding b . For v , take the edge (v, b) . For w , take the path from w to y (Lemma 2), and from there along C_{shell} to c , avoiding b . It follows from the choice of b that these paths are disjoint.

Case 1b: Vertex b is contained in the clockwise path from x to y . Then b must be a stamen that is, by construction, not adjacent to v .

If b is not adjacent to u , choose b 's unique neighbor on C as w ; see Fig. 4b. The paths are constructed as follows. For u , take the path from u to x , as guaranteed by Lemma 2, and from there to c , avoiding b . For v , take the edge (v, y) and from there to a , avoiding b . For w , take the edge (w, b) . It follows from the choice of b that the constructed paths are vertex disjoint.

Otherwise b is adjacent to u ; see Fig. 4c. Then choose w as a vertex that lies on C counterclockwise between u and v . Such a vertex exists, since C contains at least three vertices. The paths are constructed as follows. For u , we take the edge (u, b) . For v , we take the edge (v, x) and the path from x along C_{shell} to c , avoiding b . For w , we take the path from w to y (Lemma 2) and from there to a , avoiding b . Again, it follows from the choice of b that the constructed paths are vertex disjoint.

Case 2: There are at least three outermost petals of C . See Fig. 5 for the illustration of this case. By the conditions of the Lemma, vertices u and v are not internal to the same petal. Since u and v are adjacent, at most one of them is internal to a petal. Without loss of generality assume that v is not internal to a petal, and let x and y denote the two outermost petals adjacent to v , such that u is in the base of petal x . One of the two subpaths of C_{shell} connecting x and y contains in its interior a vertex of A . If there are three vertices of A in the interior of one of these paths, let b denote the middle one, otherwise, let b denote an arbitrary one. The vertices of A before and after it in counterclockwise direction along C_{shell} are a and c , respectively.

We distinguish cases based on whether b is contained in the clockwise or in the counterclockwise path from x to y along C_{shell} .

Case 2a: Vertex b is in the counterclockwise path from x to y along C_{shell} . Then, b is a stamen adjacent to v by the choice of x and y ; see Fig. 5a. Let w be any vertex adjacent to y but distinct from v . Note that $w \neq u$ by the assumption that

there are at least three petals. We construct the following paths. For u , we apply Lemma 2 to construct a path from u to x and then follow C_{shell} to a , avoiding b . For v , we take the edge vb . Finally, for w , we take the edge wy and then traverse C_{shell} to c , avoiding b . Note that the paths are disjoint by the choice of b .

Case 2b: Vertex b is in the clockwise path from x to y along C_{shell} . Then, b is a petal or a stamen but, by definition, not adjacent to v . If b is adjacent to a vertex that is distinct from u as well, then we choose this vertex as w ; see Fig. 5b. For w , we take as path the edge wb . We use the path from u to x to c (along C_{shell}) and the path from v to y to a as above.

If b is adjacent to u (Fig. 5c), then pick w as a vertex adjacent to y but distinct from u and v (such a vertex exists since petal y has at least two neighbors on C , and since there are at least three petals). Then the paths are ub , the path from v to x to c and from w to y to a . As above the paths can be chosen such that they are disjoint.

In all cases, if a path contains chords, we iteratively short-cut it by using chords. In this way we obtain the three chordless paths. The time complexity of the construction is bounded by: an application of Lemma 1, which takes $O(n)$ time; an application of Menger's theorem, which can be accomplished in time linear in the number of vertices [11]; and finally, a constant number of applications of Lemma 2, each of which takes linear time. Thus, we infer that the overall construction takes $O(n)$ time. \square

4 Extension of a One-Sided Polygon


Let G be a plane graph, and let C be a simple outerchordless cycle, represented by a one-sided polygon Γ_C . In this section, we show that if each petal of C is realizable, then Γ_C is extendable to a straight-line drawing of G . This result serves as a tool for the general case, which is shown in Sect. 5.

The drawing extension we produce preserves the outer face, i.e., if the extension exists, then it has the same outer face as G . It is worth mentioning that, if we are allowed to change the outer face, the proof becomes rather simple, as the following theorem shows.

Theorem 2 *Let G be a maximal plane graph and let C be an outerchordless cycle of G , represented in the plane by a one-sided polygon Γ_C . Then, for a suitable choice of the outer face of G , the drawing Γ_C is extendable.*

Proof Let (v_1, v_k) be the base edge of Γ_C . Edge (v_1, v_k) is incident to two faces of G , to a face f_{in} inside C and to a face f_{out} outside C . We select f_{out} as the outer face of G . With this choice, edge (v_1, v_k) is on the outer face of G . Let v be the third vertex of this face. We place the vertex v far enough from Γ_C , so that all vertices of Γ_C are visible from v . Thus, we obtain a planar straight-line drawing of the subgraph G_v induced by the vertices $V(C) \cup \{v\}$ such that each face is represented by a star-shaped polygon. Each subgraph of G inside a face of G_v is triconnected, and therefore, we can complete the existing drawing to a straight-line planar drawing of G , by Theorem 1. \square

Fig. 6 Illustration of Fact 1. Points of the set S_{pt} are black. Neither segments \overline{ab} and \overline{cd} intersect each other, nor do they intersect a segment between two consecutive points of S_{pt}



In the rest of the section we show that extendability of Γ_C can be efficiently tested, even if the outer face of G has to be preserved. The following simple geometric fact will be used in the proof of the result of this section; see Fig. 6 for an illustration.

Fact 1 *Let $pqrt$ be a convex quadrilateral and let o be the intersection of its diagonals. Let S_{pt} be a one-sided convex polygon with base \overline{pt} , that lies inside triangle Δopt . Let \overline{ab} and \overline{cd} be such that $b, d \in S_{pt}$, ordered clockwise as t, d, b, p and $a, c \in \overline{qr}$, ordered as q, a, c, r . Then, neither \overline{ab} and \overline{cd} intersect each other, nor do they intersect a segment between two consecutive points of S_{pt} .*

We are now ready to prove the main result of this section.

Theorem 3 *Let G be a n -vertex plane graph and C be a simple outerchordless cycle of G , represented in the plane by a one-sided polygon Γ_C . If every petal of C is realizable, then Γ_C is extendable. If it exists, such an extension can be constructed in $O(n)$ time.*

Proof For the following construction we need to ensure that G is a maximal plane graph and cycle C is strictly internal. If this is not the case, we apply Proposition 1, to complete G to a maximal plane graph, where C is strictly internal. For simplicity of notation, we refer to the new graph as G . Notice that the application of Proposition 1 only introduces trivial, and therefore realizable, petals. Moreover, no outer chord is introduced. After the construction of the extension of C , we remove the vertices and edges added by the application of Proposition 1.

Let v_1, \dots, v_k be the clockwise ordering of the vertices of C , so that (v_1, v_k) is the base of Γ_C . We rotate Γ_C so that (v_1, v_k) is horizontal. Let a, b, c be the vertices of the outer face of G in clockwise order; see Fig. 7. By Lemma 3, there exists a vertex v_j , $1 < j < k$, such that there exist chordless disjoint paths between v_1, v_j, v_k , and the vertices a, b, c , respectively. Denote these paths by P_{v_1a}, P_{v_jb} and P_{v_kc} . Some vertices of P_{v_1a} and P_{v_kc} are possibly adjacent to each other, as well as to the boundary of C . Depending on these adjacencies, we show how to draw the paths P_{v_1a}, P_{v_kc} and how to place vertex b , so that the graph induced by these vertices and cycle C is drawn with star-shaped faces. Then, the drawing of G can be completed by applying Theorem 1.

Let v_i be the topmost vertex of Γ_C . It can happen that there are two adjacent topmost vertices v_i and v_{i+1} . However, v_{i-1} and v_{i+2} have a smaller y -coordinate, since Γ_C does not contain flat vertices. In the following, we assume that v_i and v_{i+1} have the same y -coordinate. The case when v_i is unique can be seen as a special case where $v_i = v_{i+1}$. Without loss of generality assume that $i + 1 \leq j \leq k - 1$, the case where $2 \leq j \leq i$ is treated symmetrically. Notice that the presence of the path P_{v_jb} ensures that edges between vertices of P_{v_1a} and P_{v_kc} can only lie in the interior




Fig. 7 Illustration for the proof of Theorem 3. Edges between $C[v_1, v_i]$ and $P_{v_1a}[v_1, w'] \cup \{v_k\}$ are gray. Edges between $C[v_{i+1}, v_j]$ and $P_{v_1a}[w, a]$ are dashed




Fig. 8 Illustration for the proof of Theorem 3. For space reasons lines are shown by curves

of the cycle delimited by these paths and edges (v_1, v_k) and (a, c) ; refer to Fig. 7. Consider a vertex of P_{v_1a} that is a petal of C . The base of such a petal cannot contain edge (v_{k-1}, v_k) , since this would cause a crossing with P_{v_kc} . Moreover, if the base of this petal contains edge (v_1, v_k) , then it cannot contain any edge (v_f, v_{f+1}) for $i \leq f < j$, since otherwise this petal would not be realizable. Thus a vertex of P_{v_1a} is either adjacent to v_k or to a vertex v_f , where $i + 1 \leq f \leq j$, but not both. It is worth mentioning that a vertex of P_{v_1a} cannot be adjacent to any v_f , $j + 1 \leq f \leq k - 1$, since such an adjacency would cause a crossing either with P_{v_jb} or with P_{v_kc} .

Let s , l and l_c be three distinct lines through v_j that lie clockwise between the slopes of edges (v_{j-1}, v_j) and (v_j, v_{j+1}) ; see Fig. 8. Such lines exist since Γ_C does not contain flat vertices. Let l_i be the line through v_i with the slope of (v_{i-1}, v_i) . Let l_a be the half-line originating at an internal point of (v_1, v_k) towards $-\infty$, slightly rotated counterclockwise from the horizontal position, so that it crosses l_i . Let q denote the

intersection point of ℓ_a and ℓ_i . Let p be any point on ℓ_a to the left of q . Let ℓ'_a be the line through p with the slope of ℓ . By construction of lines s , ℓ and ℓ_c , line ℓ'_a crosses s above the polygon Γ_C at point p_a and line ℓ_c below this polygon at point p_c .

Let G' be the plane subgraph of G induced by the vertices of C , P_{v_1a} , and P_{v_kc} . The outer cycle of G' consists of edge (a, c) and a path P_{ac} between vertices a and c .

Claim 1 *The vertices of P_{v_1a} and P_{v_kc} can be placed on lines ℓ_a , ℓ'_a and ℓ_c such that the resulting straight-line drawing of G' is planar; path P_{ac} is represented by an x -monotone polygonal chain, and the inner faces of G' are star-shaped polygons.*

The vertices of P_{v_1a} will be placed on line ℓ_a between points p and q and on line ℓ'_a above point p_a . The vertices of P_{v_kc} will be placed on ℓ_c below p_c . In order to place the vertices, we need to understand how the vertices of P_{v_1a} are adjacent to vertices of C . As we travel on P_{v_1a} from v_1 to a , we first meet all vertices adjacent to v_1, \dots, v_i and then all vertices adjacent to v_{i+1}, \dots, v_j , since G is a planar graph. Let w be the first vertex of P_{v_1a} adjacent to v_f , $i + 1 \leq f \leq j$, and let w' be the vertex preceding w on P_{v_1a} . We place vertices of $P_{v_1a}[v_1, w']$, in the order they appear in the path, on line ℓ_a , between q and p , in increasing distance from v_1 . We place all vertices of $P_{v_1a}[w, a]$ on ℓ'_a above p_a in increasing distance from p . We draw the edges between the vertices of C and P_{v_1a} . Notice that vertex w might not exist, since it might happen that none of the vertices of P_{v_1a} is adjacent to v_f , $i + 1 \leq f \leq k$. In this case all vertices of P_{v_1a} are placed on line ℓ_a , between q and p .

In the following, we show that the constructed drawing is planar. Notice that the quadrilateral formed by the points w, a, v_j, v_{i+1} is convex, by the choice of line ℓ'_a and the positions of vertices w and a on it. Also, notice that the points of vertices v_{i+1}, \dots, v_j form a one-sided polygon with base segment $\overline{v_{i+1}v_j}$, which lies in the triangle Δov_jv_{i+1} , where o is the intersection of $\overline{v_{i+1}a}$ and $\overline{v_jw}$. Thus, by Fact 1, the edges connecting $C[v_{i+1}, v_j]$ and $P_{v_1a}[w, a]$ do not cross each other. By applying Fact 1, we can also prove that edges connecting $P_{v_1a}[v_1, w']$ with $C[v_1, v_i]$, cross neither each other, nor Γ_C . Recalling that vertices of $P_{v_1a}[v_1, w']$ can be also adjacent to v_k , we notice that these edges also do not cross Γ_C , by the choice of line ℓ_a . Finally, path P_{v_1a} is chordless, and therefore the current drawing is planar. Notice that the subpath of $P_{a,c}$ that has already been drawn is represented by an x -monotone chain. We next draw the vertices of P_{v_kc} . We observe that in the already constructed drawing path P_{v_1a} taken together with edge (v_1, v_k) is represented by an x -monotone chain, each point of which is visible from any point below the line ℓ'_a . This means that any point below line ℓ'_a , can be connected by a straight-line segment to the vertices $V(P_{v_1a}) \cup \{v_k\}$ without creating any crossing either with P_{v_1a} or with (v_1, v_k) . We also notice that any of the vertices v_j, \dots, v_k can be connected to a point of ℓ_c , without intersecting Γ_C . Recall that p_c denotes the intersection point of ℓ_c and ℓ'_a . Thus we place the vertices of P_{v_kc} on the line ℓ_c , below ℓ'_a , in increasing distance from point p_c . Applying Fact 1 we can prove that the edges induced by $\{v_j, \dots, v_k\} \cup V(P_{v_kc})$ are drawn without crossings. Edges between P_{v_kc} and P_{v_1a} cross neither P_{v_1a} , nor (v_1, v_k) by the choice of lines ℓ_c and ℓ'_a .

We have constructed a planar straight-line drawing of G' . We notice that path P_{ac} is drawn as an x -monotone polygonal chain. We also notice that the faces of G' , created when placing vertices of P_{v_1a} (resp. P_{v_kc}) are star-shaped and have their

kernels arbitrarily close to the vertices of P_{v_1a} (resp. P_{v_kc}). This concludes the proof of the claim.

We now extend the constructed drawing of G' to a drawing of G . Notice that vertex b is possibly adjacent to some of the vertices of P_{ac} . Thus, placing b at an appropriate distance above P_{ac} , the edges between b and P_{ac} can be drawn straight-line without intersecting P_{ac} and therefore no other edge of G' . The faces created when placing b are star-shaped and have their kernels arbitrarily close to b . We finally apply Theorem 1.

The time complexity of the described construction is determined by: an application of Proposition 1, after which the number of vertices in the graph is still $O(n)$; an application of Lemma 3; the construction of a constant number of lines; a constant number of line-line intersections; a linear number of vertices and edges traversed and drawn on the plane; and finally, by multiple applications of Theorem 1, each of which requires time linear in the size of the subgraph it affects.

5 Main Theorem

Let G be a maximal plane graph and C be a strictly internal simple outerchordless cycle of G , represented by an arbitrary convex polygon Γ_C in the plane. In Theorem 4 we prove that it is still true that if each petal of C is realizable, then Γ_C is extendable. Before stating and proving Theorem 4, we introduce notation that will be used through this section.

Recall that v_1, \dots, v_k denote the vertices of C . Let $w_{i,j}$ be an outermost petal of C in G . Let ℓ_i (resp. ℓ_j) be a half-line originating at v_i (resp. v_j) and passing through v_{i+1} (resp. v_{j-1}); see Fig. 9a. Since $w_{i,j}$ is realizable, lines ℓ_i and ℓ_j intersect. Denote by $\text{apex}(w_{i,j})$ their intersection point and by $\text{cone}(w_{i,j})$ the subset of \mathbb{R}^2 that is obtained by the intersection of the half-planes defined by ℓ_i and ℓ_j , not containing Γ_C . It is clear that any internal point of $\text{cone}(w_{i,j})$ is appropriate to draw $w_{i,j}$ so that the plane subgraph of G induced by $V(C) \cup \{w_{i,j}\}$ is crossing-free. For consistency, we also define $\text{cone}(w)$ and $\text{apex}(w)$ of an outer stamen w of C as follows. Assume that w is adjacent to $v_i \in C$. Then $\text{cone}(w) \subset \mathbb{R}^2$ is the union of the half-planes defined by lines of edges (v_{i-1}, v_i) and (v_i, v_{i+1}) , that do not contain Γ_C . We set $\text{apex}(w) = v_i$.

Let P (resp. S) denote the set of outermost petals (resp. stamens) of C in G . By Lemma 1, there exists a cycle C_{shell} in G that contains exactly $P \cup S$. Let G_{shell} denote the plane subgraph of G induced by the vertices of C and C_{shell} ; see Fig. 9b. Let C'_{shell} denote the outer cycle of G_{shell} . We denote the graph consisting of C , C'_{shell} and edges between them by G'_{shell} . Each petal or stamen of C , say w , that belongs to C_{shell} but not to C'_{shell} , belongs to a face of G'_{shell} . We denote this face by $\text{shell}(w)$. We categorize the faces of G_{shell} as follows. The faces that lie inside cycle C are called *faces of C* . The faces that are bounded only by C_{shell} and its chords, are called *faces of C_{shell}* . Notice that each face of C_{shell} is a triangle. Notice that a face of G_{shell} that is comprised by two consecutive edges adjacent to the same vertex of C (not belonging to C), is a triangle, and contains no vertex of $G \setminus G_{\text{shell}}$, since both facts would imply that the taken edges are not consecutive. Finally, faces bounded by a subpath of C and two edges adjacent to the same petal, are called *petal faces*. The plane subgraph of G




Fig. 9 **a** Vertex $w_{i,j}$ is the petal of C with base $C[v_i, v_j]$. Point $\text{apex}(w_{i,j})$ is red, region $\text{cone}(w_{i,j})$ is gray. Vertex w is a stamen of v_{j+1} . The union of the half-planes defined by lines of edges (v_j, v_{j+1}) and (v_{j+1}, v_{j+2}) is $\text{cone}(w)$. Point $\text{apex}(w)$ is blue. **b** Graph G_{shell} . Polygon Γ_C is black. Cycle C_{shell} is bold. Cycle C'_{shell} is blue. Graph G'_{shell} is comprised by blue, red and black edges. Vertices of B are squares (Color figure online)

inside a petal face is triangulated and does not have a chord connecting two vertices of its outer face, and therefore is triconnected. Thus we have the following

Observation 1 *Each vertex of $G \setminus G_{\text{shell}}$ either lies in a face of C , or in a face that is a triangle, or in a petal face, or outside C'_{shell} . Each subgraph of G inside a petal face is triconnected.*

Theorem 4 *Let G be an n -vertex plane graph and let C be a simple outerchordless cycle of G , represented by a convex polygon Γ_C in the plane. The drawing Γ_C is extendable to a straight-line drawing of G if and only if each petal of C is realizable. Both, testing the condition and the construction of an extension can be accomplished in $O(n)$ time.*

Proof The condition that each petal of C is realizable is clearly necessary. Next we show that it is also sufficient.

Assume that each petal of C is realizable. Similarly to the proof of Theorem 3, we first complete G to a maximal plane graph, such that cycle C becomes strictly internal, by applying Proposition 1. The new maximal plane graph (for simplicity of notation denoted also by G) contains no outer chord and only realizable petals, since the newly added petals are trivial. When the construction of the extension of C is completed, we simply remove the vertices and edges added by Proposition 1.

We first show how to draw the graph G'_{shell} . Afterward we complete it to a drawing of G_{shell} . Our target is to represent C'_{shell} as a one-sided polygon, so that Theorem 3 can be applied for the rest of G that lies outside C'_{shell} .

Claim 2 *Polygon Γ_C can be extended to a straight-line drawing of graph G'_{shell} where its outer face C'_{shell} is represented by a one-sided polygon such that each petal of C'_{shell} is realizable. Moreover, C'_{shell} contains in its interior all points of $\{\text{apex}(w) \mid w \in C_{\text{shell}}\}$.*




Fig. 10 **a** Construction of drawing of graph G_{shell} shown in Fig. 9b. Apex points are gray, points of B are black squares. Polygon Π is bold gray, lines $\{\ell(w) \mid w \in C'_{\text{shell}}\}$ are dashed gray. **b** Choice of the base for the one-sided polygon representing C'_{shell} . Angle of petal w is large, thus none of the edges in the base of w is chosen as the base edge. Edge $e = (u, v)$ is a possible choice for the base edge

Proof Recall that P (resp. S) denotes the set of outermost petals (resp. stamens) of C in G . Let B denote the set of vertices of C , to which stamens $S \cap C'_{\text{shell}}$ are adjacent; refer to Fig. 9b. By construction of the apex points, the set $\{\text{apex}(w) \mid w \in P \cap C'_{\text{shell}}\} \cup B$ is in convex position, and we denote by Π its convex hull; refer to Fig. 10a. Polygon Π may be degenerate, and may contain only a single vertex or a single edge. We treat these cases separately to complete the construction of the drawing of the graph G'_{shell} .

Case 1: Polygon Π is non-degenerate. Let p be a point inside Π . Let $\ell(w)$ denote a half-line from p through w , where w is a vertex of Π . If we order the constructed half-lines around p , any two consecutive lines have between them an angle less than π . If $w \in B$, we substitute $\ell(w)$ by the same number of slightly rotated lines as the number of stamens of C'_{shell} adjacent to w , without destroying the order; refer to Fig. 10a. Thus, for each $w \in C'_{\text{shell}}$, a line $\ell(w)$ is defined. Notice that, for any $w \in P \cap C'_{\text{shell}}$, line $\ell(w)$ passes through $\text{apex}(w)$, and the infinite part of $\ell(w)$ lies in $\text{cone}(w)$. Thus, for any position of w on a point of $\ell(w) \cap \text{cone}(w)$, edges between C and w do not cross Γ_C . For a stamen $w \in S \cap C'_{\text{shell}}$, line $\ell(w)$ crosses $\text{cone}(w)$ very close to $\text{apex}(w)$, and its infinite part lies in $\text{cone}(w)$. Thus, similarly, for any position of w on a point of $\ell(w) \cap \text{cone}(w)$, edges between C and w do not cross Γ_C .

We next decide which edge of C'_{shell} to “stretch”, i.e., which edge will serve as the base edge of the one-sided polygon representing C'_{shell} . In order to be able to apply Theorem 3, this one-sided polygon must be such that each petal of C'_{shell} is realizable. Thus we choose the base edge e of C'_{shell} as follows; refer to Fig. 10b. Let w be a petal of C'_{shell} and let (w, p) , (w, q) be the outer edges of petal w . Consider the lines $\ell(p)$ and $\ell(q)$, and the angle between them that contains $\text{apex}(w)$, we call it *angle of* petal w . Observe that there exists at most one petal (called *large petal*) of C'_{shell} whose angle is greater than π . Let $e = (u, v)$ be any edge of C'_{shell} such that (1) e does not belong to the base of a large petal (such an edge exists,




Fig. 11 Construction of Case 1. Corresponding G_{shell} is shown in Fig. 9b

since there are no double edges in G) and (2) if e belongs to the base of some petal, then at least one of u, v is adjacent to an outermost petal of C'_{shell} . The edge $e = (u, v)$, identified in the previous paragraph, will serve as the base of the one-sided polygon, which we are going to construct next. Recall that $\ell(u)$ and $\ell(v)$ are consecutive in the sequence of half-lines we have constructed. Let κ be a circle around Γ_C that contains in the interior the polygon Π and the set of points $\{\text{apex}(w) \mid w \in C_{shell}\}$. Let ℓ be a half-line bisecting the angle between $\ell(u)$ and $\ell(v)$ (refer to Fig. 11). Let λ be a parabola with ℓ as axis of symmetry and the center of κ as focus. We position and parametrize λ such that it does not cross ℓ and κ .

With this placement of λ , each half-line $\ell(w)$, $w \in \Pi$, crosses λ , moreover, intersections with $\ell(u)$ and $\ell(v)$ are on different branches of λ and are the last ones as we walk on λ from its origin to infinity. Let Π' be the convex polygon comprised by the intersection points of lines $\{\ell(w) \mid w \in V(C'_{shell})\}$ with λ . Notice that Π' is a one-sided polygon with the base edge $e = (u, v)$. We scale Π' with respect to the center of κ so that Π' contains the circle κ in the interior. The minimum scale factor is determined by the segment of Π' that is closest to the center of κ . As a result, for each $w \in C$, $\text{cone}(w) \cap \Pi'_{in} \neq \emptyset$, where Π'_{in} denotes the interior of Π' .

We next show that each petal of $C_{shell'}$ is realizable by the choice of edge e . Thus, if e does not belong to the base of a petal of C'_{shell} , then all petals of C'_{shell} are realizable. Otherwise, $e = (u, v)$ belongs to the base of an outermost petal w of C'_{shell} , and is chosen such that either (u, w) or (v, w) exists and represents an outer edge of w . Recall that w is not a large petal, thus, the angle between the lines $\ell(u)$ and $\ell(v)$ is at most π . Since Π' is a one-sided polygon with the base edge (u, v) , petal w is realizable. Since w is an outermost petal of C'_{shell} , we infer that all the remaining petals containing e in their base are also realizable. This concludes the proof of the claim in the non-degenerate case.

Case 2: Polygon Π is degenerate and has two vertices. Notice that in this case cycle C'_{shell} contains at most two petals, since each petal of C'_{shell} is a vertex of Π . Assume that C'_{shell} contains two petals, w and w' . Since C'_{shell} contains at least three vertices (G does not have double edges), there would be at least one stamen w'' in C'_{shell} . However, the vertex of C , to which w'' is adjacent, together with w




Fig. 12 a–b Case 2: Polygon Π is degenerated and has two vertices. c Case 3: Polygon Π is degenerated and has a single vertex, and C'_{shell} contains three or more stamens of C adjacent to the same vertex

and w' , contribute three vertices to Π , which is a contradiction to the assumption that Π has two vertices. Thus, in this case C'_{shell} contains at most one petal of C . In the following we consider two subcases, first when C'_{shell} contains one petal, and second when it contains no petal.

Case 2.a: C'_{shell} contains a petal w of C (Fig. 12a). Notice that there are at least two stamens of C in C'_{shell} , since otherwise C'_{shell} would contain only two vertices, and therefore there would be a double edge in G . Moreover, all stamens of C that are in C'_{shell} are adjacent to the same vertex v_i of C . Let s_1, \dots, s_k be the stamens adjacent to v_i , appearing clockwise in C'_{shell} . Let $\ell(s_k)$ be a line through v_i that does not cross Γ_C ; see Fig. 13a. There exists a half-line $\ell(w)$ in $\text{cone}(w)$ that does not cross $\ell(s_k)$. Let $\ell(s_1)$ be a half-line through v_i , lying in the half-plane defined by $\ell(s_k)$ not containing Γ_C , slightly rotated clockwise. Let $\ell(s_2), \dots, \ell(s_{k-1})$ be lines through v_i that lie clockwise between $\ell(s_1)$ and $\ell(s_k)$. The angle between two consecutive half-lines $\ell(s_1), \dots, \ell(s_k), \ell(w)$ is less than π . Moreover, each of the half-lines has its infinite part in the corresponding cone. Moreover, for any position of w (resp. s_i) on a point of $\ell(w) \cap \text{cone}(w)$ (resp. $\ell(s_i) \cap \text{cone}(s_i)$), edges between C and w (resp. s_i) do not cross Γ_C . Thus, we can choose base edge e and then apply a construction using circle κ and parabola λ identical to the case when polygon Π is not degenerate.

Case 2.b: C'_{shell} does not contain any petal of C (Fig. 12b). In this case C'_{shell} contains at least three stamens of C , and all of them are adjacent to two vertices of C , say v_i and v_j , otherwise Π would contain more than two vertices. Without loss of generality assume that at least two stamens of C are adjacent to v_i . Let them be s_1, \dots, s_k . Let t_1, \dots, t_p be the stamens adjacent to v_j . The construction of half-lines $\ell(s_1), \dots, \ell(s_k), \ell(t_1), \dots, \ell(t_p)$ can be done analogous to Case 2.a, where line $\ell(w)$ is substituted by a set of closely placed lines $\ell(t_1), \dots, \ell(t_p)$. Since the angle between two consecutive half-lines $\ell(s_1), \dots, \ell(s_k), \ell(t_1), \dots, \ell(t_p)$ is less than π , we proceed again as in the non-degenerate case.

Case 3: Polygon Π is degenerated and has a single vertex (Fig. 12c). We first notice that in this case C'_{shell} does not contain any petal of C . This is because C'_{shell} contains at least one stamen of C , adjacent to a vertex v_i of C . Thus a petal together with v_i would contribute two vertices to Π . Therefore, C'_{shell} contains only stamens of C . These stamens are adjacent to a single vertex of C , say v_i ,




Fig. 13 **a** Construction in Case 2.a. **b** Construction in Case 3. **c** If C'_{shell} contains only two stamens of C , then a vertex of C , v_i , belongs to the outer cycle of G

since otherwise Π would contain more than one vertex. If there are only two stamens adjacent to v_i (see Fig. 13c), then either there is a double edge in G , or v_i belongs to the outer cycle of G . The latter can not be the case, since C is a strictly internal cycle of G , as ensured in the beginning of the proof. So C'_{shell} contains three or more stamens s_1, \dots, s_k of C , and they are all adjacent to v_i . Fig. 13b illustrates the construction of lines $\ell(s_1), \dots, \ell(s_k)$ in this last case. Notice that the angles between two consecutive half-lines $\ell(s_1), \dots, \ell(s_k)$ can be made less than π , since v_i is not flat. Thus we can again proceed as in the non-degenerate case. This concludes the proof of the claim.

Let Γ'_{shell} be the constructed drawing of G'_{shell} . Recall that each petal or stamen w of C , that does not belong to C'_{shell} , lies in a face of G'_{shell} , denoted by $shell(w)$. Let $\Gamma_{shell(w)}$ denote the polygon representing face $shell(w)$ in Γ'_{shell} . By construction, $cone(w) \cap \Gamma_{shell(w)} \neq \emptyset$. We next explain how to extend the drawing of G'_{shell} to the drawing of G_{shell} . For each edge (u, v) of C'_{shell} , we add a convex curve, lying close enough to this edge inside Γ'_{shell} . More precisely, we consider the perpendicular bisector of the edge (u, v) and place a point o on this line, so that o is closer to (u, v) than any apex point inside $\Gamma'_{shell(w)}$. We then construct a parabola with origin o passing through the vertices u and v . Let μ be the closed curve that is formed by the union of the parts of all these parabolas (one for each edge of C'_{shell}) that lie inside C'_{shell} . We notice that we have placed them so that all the points of $\{apex(w) \mid w \in C\}$ are still in the interior of μ . Thus μ is intersected by all the sets $cone(w)$, for each $w \in C$. We place each vertex w of $C_{shell} \setminus C'_{shell}$ on $\mu \cap cone(w)$ in the order they appear in C_{shell} . Since all edges induced by C_{shell} lie outside of C_{shell} , and both endpoints of such an edge are placed on a single convex curve, they can be drawn straight without intersecting each other, or other edges of G_{shell} . Thus, we have constructed a planar extension of Γ_C to a drawing of G_{shell} , call it Γ_{shell} .

Recall the definitions of faces of C , faces of C_{shell} and petal faces from the beginning of this section. The faces of C appear in Γ_{shell} as convex polygons. The faces of C_{shell} are triangles, and the petal faces of G_{shell} are star-shapes whose kernel is close to the

corresponding petal. By Observation 1, each vertex of $G \setminus G_{\text{shell}}$ either lies in a face of C , or in a face that is a triangle, or in a petal face, or outside C'_{shell} . Moreover, a subgraph of G inside a petal face is triconnected. Thus, by multiple applications of Theorem 1, we can extend the drawing of G_{shell} to a straight-line planar drawing of the subgraph of G inside C'_{shell} .

Finally, notice that in the constructed drawing of G_{shell} each petal of its outer cycle, i.e., C'_{shell} , is realizable (Claim 2). Moreover, by construction of G_{shell} , C'_{shell} has no outer chords. Thus, we can apply Theorem 3, to complete the drawing of G lying outside C'_{shell} .

The time complexity of the construction is determined by the following basic operations: an application of Proposition 1; the construction of a constant number of subgraphs of linear size, namely C_{shell} , G_{shell} , C'_{shell} and G'_{shell} ; a constant number of traversals of these subgraphs, for computing cones, apex points, polygon Π' and the actual drawing of these subgraphs; a linear number line-line intersections to construct the apex points and the cones; the construction of a linear number of lines to host the vertices of C'_{shell} ; a traversal of the neighborhood of C'_{shell} to determine which edge to “stretch” for the one-sided representation of C'_{shell} ; a linear number of parabola-line intersections for the placement of vertices of C'_{shell} ; the construction of a number of parabolas to form the curve μ , which takes time proportional to the number of apex points plus the size of C'_{shell} , which is linear in total; multiple applications of Theorem 1, each of which runs in time linear in the number of vertices it places; and finally, an application of Theorem 3. Thus, we conclude that the overall time complexity of the construction is $O(n)$. \square

We conclude with the following general statement that follows from Theorem 4 and one of the known algorithms that constructs drawing of a planar graph with a prescribed outer face (e.g. [5, 12] or Theorem 1).

Theorem 5 *Let G be a plane graph and H be a biconnected plane subgraph of G . Let Γ_H be a straight-line convex drawing of Γ_H . Γ_H is extendable to a planar straight-line drawing of G if and only if the outer cycle of H is outerchordless and each petal of the outer cycle of H is realizable.*

6 Conclusions

In this paper, we have studied the problem of extending a given convex drawing of a cycle of a plane graph G to a planar straight-line drawing of G . We characterized the cases when this is possible in terms of two simple necessary conditions, which we proved to also be sufficient.

It can be easily seen that the drawing area depends on the input. Thus, if C has a petal $w_{i,j}$, such that the edges (v_i, v_{i+1}) , (v_{j-1}, v_j) of C (refer to Fig. 9a) are almost parallel, the area required for the drawing becomes arbitrarily large. Another situation which blows up the area is the following. Consider a cycle of length three which surrounds C and has one common vertex v_i with C . If additionally v_i is almost flat in C , i.e., edges (v_{i-1}, v_i) and (v_i, v_{i+1}) have almost the same slope, then the three-cycle needs to be arbitrarily long in order to contain C . Thus, our main open question is

whether simple conditions forbidding such situations are sufficient to ensure that the area of the final drawing is polynomial. For our construction we have repeatedly used Theorem 1 [6], which also does not ensure that the area is polynomial. Thus, in order to answer our open question it also has to be investigated when a straight-line drawing inside a fixed star-shaped polygon has polynomial area.

Finally, as an extension of our research, it would be interesting to investigate whether more involved necessary conditions are sufficient for more general shapes of a cycle, for instance star-shaped polygons.

Acknowledgments Martin Nöllenburg received financial support by the Concept for the Future of KIT (Grant YIG 10-209). Ignaz Rutter was supported by a fellowship within the Postdoc-Program of the German Academic Exchange Service (DAAD). Part of this work was done within GRADR – EUROGIGA project no. 10-EuroGIGA-OP-003.

References

1. Angelini, P., Di Battista, G., Frati, F., Jelínek, V., Kratochvíl, J., Patrignani, M., Rutter, I.: Testing planarity of partially embedded graphs. In: 21st Annual ACM-SIAM Symposium on Discrete Algorithms (SODA'10), pp. 202–221. SIAM (2010)
2. Avis, D.: Generating rooted triangulations without repetitions. *Algorithmica* **16**, 618–632 (1996)
3. Chambers, E.W., Eppstein, D., Goodrich, M.T., Löffler, M.: Drawing graphs in the plane with a prescribed outer face and polynomial area. *J. Gr. Algorithms Appl.* **16**(2), 243–259 (2012)
4. Chan, T.M., Frati, F., Gutwenger, C., Lubiw, A., Mutzel, P., Schaefer, M.: Drawing partially embedded and simultaneously planar graphs. In: *Graph Drawing*, vol. 8871 of *Lecture Notes in Computer Science*, pp. 25–39. Springer, Berlin (2014)
5. Duncan, C.A., Goodrich, M.T., Kobourov, S.G.: Planar drawings of higher-genus graphs. *J. Gr. Algorithms Appl.* **15**(1), 7–32 (2011)
6. Hong, S.-H., Nagamochi, H.: Convex drawings of graphs with non-convex boundary constraints. *Discret Appl Math* **156**(12), 2368–2380 (2008)
7. Jelínek, V., Kratochvíl, J., Rutter, I.: A Kuratowski-type theorem for planarity of partially embedded graphs. *Comput. Geom. Theory Appl.* **46**(4), 466–492 (2013)
8. Mchedlidze, T., Nöllenburg, M., Rutter, I.: Drawing planar graphs with a prescribed cycle. In: Wismath S., Wolff A. (eds.) *Proceedings of the 21st International Symposium Graph Drawing (GD'13)*, vol. 8242 of *LNCS*, pp. 316–327. Springer, Berlin (2013)
9. Pach, J., Wenger, R.: Embedding planar graphs at fixed vertex locations. *Gr. Comb.* **17**(4), 717–728 (2001)
10. Patrignani, M.: On extending a partial straight-line drawing. *Int. J. Found. Comput. Sci.* **17**(5), 1061–1070 (2006)
11. Ripphausen-Lipa, H., Wagner, D., Weihe, K.: The vertex-disjoint menger problem in planar graphs. *SIAM J. Comput.* **26**(2), 331–349 (1997)
12. Tutte, W.T.: How to draw a graph. *Proc. Lond. Math. Soc.* **13**(3), 743–768 (1963)