

## Laparoscopic repair of incisional hernia: Outcomes of 100 consecutive cases comprising 25 wall defects larger than 15 cm

Giovanni Carlo Ferrari · Angelo Miranda ·  
Stefano Di Lernia · Fabio Sansonna · Carmelo Magistro ·  
Dario Maggioni · Ildo Scandroglio · Andrea Costanzi ·  
Maurizio Franzetti · Raffaele Pugliese

Published online: 6 March 2008  
© Springer Science+Business Media, LLC 2008

### Erratum to: Surg Endosc DOI 10.1007/s00464-007-9707-9

(1) On 1st page of article, right-hand column, 4th paragraph, 7th line, change “(>20 cm)” to “(>15 cm)”.

(2) On 2nd page, Table 1 title, change “according to Chevrel [9]” to “modified from Chevrel [8, 9]”.

(3) On 6th page, left-hand column, 7th line from end of Discussion, change “since Chevrel defined as “giant” only hernias larger than 20 cm [29]” to “since Bernard considered as “giant” hernias with surface of 182 cm<sup>2</sup> [19].”

---

The online version of the original article can be found under doi:  
[10.1007/s00464-007-9707-9](https://doi.org/10.1007/s00464-007-9707-9).

---

G. C. Ferrari · A. Miranda · S. D. Lernia · F. Sansonna (✉) ·  
C. Magistro · D. Maggioni · I. Scandroglio · A. Costanzi ·  
M. Franzetti · R. Pugliese  
Surgery and Videolaparoscopy Department, Niguarda Hospital,  
Milan, Piazza Ospedale Maggiore 3, 20162 Milano, Italy  
e-mail: fabios55@libero.it