

Thoracic sympathectomy: a review of current indications

Moshe Hashmonai¹ · Alan E. P. Cameron² · Peter B. Licht³ · Chris Hensman⁴ ·
Christoph H. Schick⁵

Received: 5 March 2015 / Accepted: 16 June 2015 / Published online: 27 June 2015
© Springer Science+Business Media New York 2015

Abstract

Background Thoracic sympathetic ablation was introduced over a century ago. While some of the early indications have become obsolete, new ones have emerged. Sympathetic ablation is being still performed for some odd indications thus prompting the present study, which reviews the evidence base for current practice.

Methods The literature was reviewed using the PubMed/Medline Database, and pertinent articles regarding the indications for thoracic sympathectomy were retrieved and evaluated. Old, historical articles were also reviewed as required.

Results and conclusions Currently, thoracic sympathetic ablation is indicated mainly for primary hyperhidrosis, especially affecting the palm, and to a lesser degree, axilla and face, and for facial blushing. Despite modern pharmaceutical, endovascular and surgical treatments, sympathetic ablation has still a place in the treatment of very selected cases of angina, arrhythmias and cardiomyopathy. Thoracic sympathetic ablation is indicated in several painful conditions: the early stages of complex regional pain syndrome, erythromelalgia, and some pancreatic and other painful abdominal pathologies. Although

ischaemia was historically the major indication for sympathetic ablation, its use has declined to a few selected cases of thromboangiitis obliterans (Buerger's disease), microemboli, primary Raynaud's phenomenon and Raynaud's phenomenon secondary to collagen diseases, paraneoplastic syndrome, frostbite and vibration syndrome. Thoracic sympathetic ablation for hypertension is obsolete, and direct endovascular renal sympathectomy still requires adequate clinical trials. There are rare publications of sympathetic ablation for primary phobias, but there is no scientific basis to support sympathetic surgery for any psychiatric indication.

Keywords Sympathectomy · Thoracoscopy · Hyperhidrosis

The indications for sympathetic ablation have evolved since Alexander first performed this operation in 1889 to treat epilepsy [1]. Jonnesco [2] and Jaboulay [3] extended its application to the treatment of exophthalmic goitre. François-Franck [4] reported its use for the treatment of glaucoma, idiotism and hyperthyroidism. These indications became obsolete long ago. However, even currently, sympathetic surgery is performed and reported for some odd indications. The present review was undertaken in order to examine the evidence base for the indications currently in use. The older open methods have given way to endoscopic techniques, but this review is concerned with the indications for sympathectomy, techniques of which will not be discussed in detail.

Methods

Search of the literature was performed in the PubMed/Medline Database for the years 1990–2014 (the period in which the endoscopic approach of sympathetic ablation

On behalf of the International Society of Sympathetic Surgery.

✉ Moshe Hashmonai
hasmonai@inter.net.il

¹ Faculty of Medicine, Technion-Israel Institute of Technology, PO Box 359, 30952119 Zikhron Ya'akov, Haifa, Israel

² Nuffield Hospital, Ipswich, UK

³ Department of Cardiothoracic Surgery, Odense University Hospital, Odense, Denmark

⁴ Specialist Surgeon, Melbourne, Australia

⁵ German Hyperhidrosiscenter, Munich, Germany

supplanted open surgery). To identify the indications in current use, an initial search was made using as terms “Indications” AND “Thoracic Sympathectomy OR Sympathicotomy OR Sympathetic ablation” NOT “Renal Sympathectomy”. During the complete retrieval process, some additional rare indications were identified and included in the present review. This was followed by individual searches in the Database for each indication, using as terms the pathology, e.g. “Palmar hyperhidrosis” AND “Thoracic Sympathectomy OR Sympathicotomy OR Sympathetic ablation”. To focus on the subject, exclusion of similar pathologies was added to the search items (e.g. for the search of palmar hyperhidrosis “NOT Axillary hyperhidrosis” was added). Review articles and clinical trials were sorted out. Selection of studies was made in three steps: (a) by the relevance of the title; (b) by scrutinizing the abstracts of articles selected in the first stage; and (c) by scrutinizing the full text of articles selected in the second stage. All review articles and clinical trials were scrutinized according to the same three steps. Older publications were retrieved as appropriate.

Results

The search of the literature for current indications of thoracic sympathetic ablation yielded 68 articles. Step one selection (by titles) restricted the list to 31 and a further selection (step 2, by abstracts) restricted the list to 14 publications. The following current indications were identified in current use for performing thoracic sympathectomy: hyperhidrosis—palmar, axillary and facial (craniofacial); blushing; cardiac indications; pain conditions; ischaemia; and psychiatric conditions. For hypertension, renal sympathectomy has supplanted thoracic sympathetic ablation. The figures of retrieved articles for each indication are listed in Table 1. A paucity of clinical trials is noticeable. Randomized trials and reviews based on randomized trials are extremely rare. Articles pertinent to such studies are outlined in the discussion.

Discussion

Hyperhidrosis

Hyperhidrosis is excessive secretion from the eccrine sweat glands [5]. Most cases are primary, of unknown aetiology, but a few may be secondary to other disorders [6]. Hyperhidrosis can cause severe social, emotional and occupational handicaps from an early age. Many treatments have been proposed [7, 8], but the only one which may abolish palmar hyperhidrosis is sympathetic ablation [8].

The first sympathetic ablation for hyperhidrosis was performed by Kotzareff [9] and reported in 1920. Since that publication, thoracic sympathetic ablation became widely used for palmar overperspiration, and later extended for axillary and facial hyperhidrosis.

Primary palmar hyperhidrosis

Primary palmar hyperhidrosis (PHH) is currently the major indication for upper thoracic sympathectomy. An exhaustive review in 2000 [8] has shown that resection of the T2 and T3 ganglia invariably results in palmar anhidrosis. It has been proven that quality of life of patients with PHH is substantially improved by sympathetic ablation [10], and improved quality of life remains stable in the long term [11]. Furthermore, according to patients' subjective evaluation, the worse pre-sympathetic ablation quality of life resulted in the best quality of life estimation following the procedure [12]. Nevertheless, surgery is still controversial because of the risk of compensatory hyperhidrosis—excessive sweating developing in areas unaffected by the sympathetic ablation—often resistant to treatment [13], and which may be very debilitating [14]. Performing a pre-operative chemical sympathetic block was suggested as a method to evaluate the probability of developing compensatory hyperhidrosis following surgical sympathectomy [15]. To reduce the degree and occurrence of compensatory hyperhidrosis, a multitude of technical variations have been suggested [13], but published recommendations are often contradictory. There is an enormous bulk of literature dealing with this question. While many studies advocate that lowering the level of ablation (from T2 to T3 or T4) [16–18] or restricting the extent of ablation reduce the severity and incidence of compensatory hyperhidrosis [19], other studies contradict these claims [16, 20–22]. Two recent reviews published in 2011 underline this controversy. In a “consensus for the treatment of hyperhidrosis” based on a review of randomized studies, a group of authors suggested ablation at the upper level of the third rib (a presumably T3 ablation) as the best procedure for PHH [23]. Another simultaneously published systematic review and meta-analysis of the literature suggested that T3 ablation alone resulted in a significantly higher degree of compensatory sweating than T2–T4 sympathectomy [24], recommending the latter approach. Although there is no agreement, it appears that lowering the level of sympathetic ablation and limiting its extent reduces compensatory hyperhidrosis, but the amount of achieved reduction in palmar sweating is lower as well [25–27], and the percentage of recurrences of PHH is increased [13]. Due to the risk of compensatory hyperhidrosis and other late side effects [28], and possible early complications [29–31], sympathetic ablation should be restricted to those whose

Table 1 Results of the literature search

Indication	Articles obtained by the search			Articles retrieved by:	
	Total #	Reviews	Clinical trials	Title	Abstract
Hyperhidrosis					
Palmar	502	36	61	65	55
Axillary	67	14	8	18	9
Facial (Craniofacial)	164	20	12	17	12
Blushing	22	4	2	13	13
Cardiac indications					
Angina	51	6	4	22	10
Arrhythmias					
Ventricular tachycardia	58	8	2	5	2
Long QT syndrome	61	17	0	10	6
Cardiomyopathy	41	4	5	5	3
Pain conditions					
Complex regional pain syndrome	45	4	1	18	9
Erythromelalgia	11	2	0	2	1
Pancreatic and splanchnic pain	43	9	2	10	5
Ischaemia					
Atherosclerosis	7	1	0	2	2
Thromboangiitis obliterans	37	2	0	4	3
Raynaud's phenomenon	77	9	4	9	3
Scleroderma	34	8	2	8	4
Rheumatoid arthritis	7	0	0	1	1
Lupus erythematosus	4	0	0	1	0
Paraneoplastic syndrome	3	1	0	2	1
Microemboli	1	0	0	1	1
Frostbite	9	4	1	4	3
Vibration syndrome	4	1	0	3	3
Hypertension	1	0	0	0	0
Psychiatric indications (phobias)	8	2	1	6	6

quality of life is severely affected. The surgeon should be well aware of the various procedures and their expected results, and discuss the operation with the patient in order to obtain a truly informed consent for the operation.

Failures of sympathetic ablation for PHH may be due either to an incomplete or inadequate ablation of the correct sympathetic segment (occurring early after the initial operation), or to a true recurrence (developing months or years after the initial operation), probably due to nerve regeneration or sprouting of nerve fibres from the injured sympathetic chain [32]. Resympathectomy has been reported to be beneficial for both types of recurrent overperspiration [33, 34]. However, both surgeons and patients should be aware that repeat ablation may increase the occurrence of compensatory hyperhidrosis [34, 35].

There is a multitude of non-surgical procedures in the treatment of PHH [8, 36]. None of them secures permanent cure. Therefore, their use may be attempted in milder cases of PHH. However, for patients with disabling PHH,

sympathetic ablation remains the only method of treatment which may produce permanent and complete relief of palmar overperspiration and should be regarded as first-line treatment [37]. Due to the potential side effects of sympathectomy, patients should be well informed of these limitations before embarking on the procedure.

Axillary hyperhidrosis

Axillary sweating may accompany PHH but isolated axillary hyperhidrosis (AHH) is also encountered. Severe isolated AHH also affects the quality of life and impinges upon the daily social and working interactions of the patient. Sympathetic ablation may alleviate the condition. However, in the era of open surgery, AHH was not considered as a primary indication for sympathectomy, excision of the axillary skin being a simpler and effective procedure [38, 39], despite the drawback of poor healing and an unsightly scar.

The advent of endoscopic surgery simplified the operative procedure and AHH became one of the indications for sympathetic ablation. It has been shown that sympathectomy for AHH should be at a lower level of ablation than for PHH [40–42], going down to T4–T5 [40, 41], and in 2008 satisfactory results have been published in a randomized controlled study [42]. However, analysis of one of the largest published series [43] has shown that, compared to the results for palmar hyperhidrosis, the results for AHH are worse: the percentage of totally satisfied patients was lower, the percentage of dissatisfied patients was double, and the percentage of those regretting the operation was 4.5 times higher for AHH than for PHH.

Restricting the indication for sympathetic surgery for AHH is also due to the existence of several local non-surgical and surgical options which may alleviate excessive perspiration. Local treatments include antiperspirants, iontophoresis and Botox-A injections. Non-surgical local treatment of the perspiring area by microwaves has been reported to achieve substantial reduction in axillary sweat [44]. Local surgery includes subcutaneous curettage, either with a curette [45] or by suction [46], which has also been stated to be successful in a high percentage of cases. Successful treatment of AHH by excision of the axillary skin, either partial [47] or complete [48], has also been advocated. A recent comparative study has shown that local excision yields superior results than sympathetic ablation for AHH [49], practically avoiding the sequel of compensatory hyperhidrosis.

When the axilla is an isolated area of overperspiration, local treatment may be sufficient and satisfactory. Sympathetic ablation for AHH should be therefore restricted only to patients whose AHH is concomitant with PHH. This restricted indication seems to be at present the gold standard for sympathetic ablation in cases of AHH.

Facial hyperhidrosis

The reported incidence of facial hyperhidrosis varies tremendously. Some publications claim it is very frequent while others hardly mention it. One dermatological study reported that facial hyperhidrosis accounted for 5 % of patients who seek help for overperspiration [50], and another report stated that facial hyperhidrosis was the most frequently affected site [51]. In a sympathetic surgical study, facial hyperhidrosis (including blushing) was the indication in 10 % of 650 procedures [52], while in an Asian study, facial hyperhidrosis was seen in just one of every 40 patients who suffered from palmar hyperhidrosis [53]. Facial hyperhidrosis should be distinguished from Frey's Syndrome, a frequent sequela of parotidectomy, in which gustatory stimuli result in facial sweating or blushing, and for which sympathetic surgery should not be considered.

Several, non-surgical treatment modalities have been suggested for facial hyperhidrosis, including oral medications (glycopyrrolate and clonidine), topical medications and botulinum toxin injections [36].

Due to the possibility of developing Horner's syndrome and compensatory hyperhidrosis, sympathetic ablation is considered the last-line of treatment [36]. Yet, and although rarely reported, thoracoscopic sympathetic ablation for isolated facial hyperhidrosis is effective [36, 43, 52–56]. Ablation is directed at the higher levels of the sympathetic chain, which is considered necessary to achieve an effect in the face. The most common surgical intervention is R2 (rib) sympathectomy, but reports include R2–R3 sympathectomy [43], G2 (ganglion) ramicotomy [44], G2 ablation [52] and even clipping of the lower part of the stellate ganglion without causing any Horner's syndrome [56].

Blushing

Normally, reddening of the face is part of a physiologic thermoregulatory response to hyperthermia that results from increased cutaneous blood flow caused by transient vasodilatation. This may also be triggered by multiple conditions which have been reviewed extensively [57]. Blushing is triggered by emotion and should be distinguished from flushing which appears as a symptom of underlying somatic disease. In severe cases, blushing is often associated with a feeling of embarrassment, and some individuals may experience so much distress that they develop a phobia about it [58]. The exact nature of this vasomotor phenomenon is unclear and the prevalence is unknown, but the incidence varies with geographical location. However, due to the emotional nature of the phenomenon, psychotherapeutic approaches have been considered [59] in addition to local [60] and systemic pharmacotherapies [61]. In a recent review of available treatments [62], the authors considered sympathetic ablation to be the most appropriate treatment of blushing.

Thoracoscopic sympathetic ablation for facial blushing was first described in 1985 [63], but the number of papers on this subject is much lower compared to those for primary hyperhidrosis [43, 52, 64–85]. In Scandinavia, surgical treatment of blushing accounted for approximately half of all sympathectomies [77, 85], whereas sympathetic ablation for this indication is quite rare in Asia [68]. The reason could be either a different cultural attitude to the problem or, possibly, a true constitutional difference between populations.

There is no consensus on the extent or level of sympathetic ablation for isolated facial blushing, which has been targeted at R1–R3 [65], R2 [52, 73], R2–R3 [18, 25, 63, 75, 76], R2–R4 [70], R2–R5 [64, 71] or R3–R4 [18]. However,

a recent randomized clinical study [85] comparing R2 versus R2–R3 sympathetic ablation did not show any significant difference in local effect.

Results from the literature indicate that sympathetic ablation for facial blushing is effective, but meticulous and critical patient selection is essential and thorough information about the high risk of side effects is crucial. In general, complications and side effects following sympathetic ablation for isolated facial blushing are similar to those seen after sympathetic surgery for primary hyperhidrosis [29]. Compensatory sweating is the most commonly reported side effect, seen in almost all patients treated for facial blushing [18, 43, 44], and gustatory sweating is observed post-operatively in one-third of patients [36]. However, contrary to sympathetic surgery for palmar and axillary hyperhidrosis, the risk of Horner's syndrome is increased when sympathetic ablation is performed for blushing, because the sympathetic chain is approached at the level of the second rib, which is considered essential to achieve an effect in the face. Blushing must be of major concern to the patient, enough to tolerate a substantial amount of compensatory sweating after the operation as well as gustatory sweating. It should also be kept in mind that upper chest or neck blushing as well as a slowly emerging and long-lasting facial flushing respond poorly to sympathetic surgery.

Cardiac indications

Sympathetic surgery has now very little place in the management of cardiac conditions, but a few indications remain.

Angina

The possible use of sympathetic ablation to treat angina was suggested by François-Franck [4], based on his studies and the understanding of the afferent sympathetic pathways. Cervical sympathectomy for cardiac indications was performed as early as the 1920s with positive results [86]. Although sympathetic ablation results in vasodilatation, the effect of this procedure on pain is attributed to interruption of significant pain pathways from the heart and not to improvement of cardiac perfusion [87, 88].

With the advent of coronary surgery and endovascular procedures, one might expect that sympathectomy for angina would become obsolete, but sporadic cases of non-compliant coronary artery spasm successfully treated by T2 to T4 sympathetic ablation are being published, remaining a useful tool in the treatment of the very selected cases of refractory angina [66, 67, 89]. A preliminary stellate sympathetic block may confirm the efficacy of the procedure prior to surgery [90]. However, the haemodynamic

instability of these patients remains a concern when considering surgery.

Arrhythmias

In the late 1970s and early 1980s, Schwartz [91] experimentally demonstrated the influence of the autonomic nervous system on cardiac electric activity. The role of the left and right cardiac sympathetic nerves has also been elucidated by the addition of right cardiac sympathetic ablation to left cardiac sympathetic denervation in managing cardiac dysrhythmia [92]. Presently, the role of the autonomic nervous system in modulating cardiac electrophysiology and arrhythmogenesis [93] is well established and justifies sympathetic ablation for the treatment of arrhythmias in appropriate conditions.

1. Ventricular tachycardia: The medical treatment of arrhythmias is by beta-blockers or by implanted cardioverter-defibrillators for refractory cases. Yet, in a small percentage of patients, ventricular tachycardia refractory to traditional therapies (so-called electrical storm) is encountered [94]. For decades, it has been accepted that the sympathetic nervous system has profound and recurrent effects on cardiac electric activity [95, 96]. Based on this concept, Estes and Izlar [97] performed bilateral stellectomy in the 1960s, in an attempt to treat ventricular arrhythmias. In 1968, Zipes et al. [98] reported a successful case of bilateral stellectomy in conjunction with atrial pacing in a patient who had malignant ventricular arrhythmias. Sympathetic ablation, performed in the rare cases with ventricular tachycardia irresponsive to conventional therapy, with substantial success was reported as recently as 2012 [92] and 2014 [99]. Bilateral ablation may give superior results than left sympathectomy alone [99].

Two recent publications, one animal [100] and one human study [101], examined the effect of renal denervation on ventricular arrhythmias. Due to the relative ease in performing endovascular renal denervation, when indicated, this technique may possibly supplant in the near future the use of thoracic sympathectomy in the treatment/prevention of ventricular tachycardia.

2. Long QT syndrome (LQTS): LQTS is a hereditary disorder characterized by a prolonged QT interval and a polymorphic ventricular tachycardia, leading to severe events: syncope, cardiac arrest, and sudden death [102]. There are eight forms of congenital LQTS in which gene mutations cause potassium, sodium and calcium ion channel disorders [103, 104]. In a recent review, the treatment of LQTS was delineated to

consist of β -adrenergic blocking agents, cardioverter-defibrillator implantation and sympathetic ablation [105]. Moss and McDonald [106] in 1971 reported their pioneer work of successfully treating long QT syndrome with left cardiac sympathetic ablation. In humans, left cardiac sympathectomy normalizes the prolonged QT interval, reduces QT dispersion and thereby diminishes the probability of malignant arrhythmias [104]. In 2004, a group of 147 patients with LQTS who underwent sympathectomy were reported [107]. Almost half of them were free of symptoms post-operatively and on long-term follow-up. In the remaining individuals, the occurrence of cardiac events decreased substantially. These results illustrate the value of sympathetic surgery for LQTS.

Cardiomyopathy

The treatment of cardiomyopathy is medical. Patients with heart failure due to cardiomyopathy are not good candidates for surgery, yet sympathectomy has been attempted in order to improve their cardiac performance. In a study of patients with dilated cardiomyopathy, a high thoracic sympathetic ablation significantly reduced the left ventricular cavity, boosting the ejection fraction, and thus improving the left ventricular diastolic function [108]. The same results were obtained in a controlled study published in 2010, and improvement persisted at 6-month follow-up [109]. A later study, further confirmed that left ventricular function in patients with dilated cardiomyopathy is improved by sympathectomy [110]. Thus, it may be concluded that sympathetic blockade has a place as an adjunctive therapy in selected patients with severe dilated cardiomyopathy.

Pain conditions

Sympathetic ablation may be indicated for a few painful conditions.

Complex regional pain syndrome

This is an out of proportion, bizarre, and poorly understood painful response triggered by trauma to a limb. It causes allodynia (the perception of a non-painful stimulus as painful) and hyperpathia (an exaggerated pain response to a painful stimulus). Sympathetic dysfunction, usually overactivity, becomes evident. Varying trophic changes can follow and eventually the condition may lead to irreversible severe incapacity.

Synonyms include causalgia, post-traumatic sympathetic dystrophy, reflex dystrophy, Sudek's atrophy, post-

traumatic dystrophy, shoulder-hand syndrome, post-traumatic pain syndrome, sympathetic neurovascular dystrophy, post-traumatic spreading neuralgia, to mention just a few [111]. However, because the role of the autonomous nervous system is obscure in the pathogenesis of these disorders, the term "Complex regional pain syndrome" (CRPS) is preferred. CRPS is subdivided into two groups: I—following tissue trauma and II—following trauma to a nerve [112]. The pathogenesis is obscure. A regional inflammatory response to the initial trigger has been implicated [113, 114]. Clinical findings suggest that patients with CRPS have autonomic dysfunction within the central nervous system [115].

There is very little good evidence in the literature to guide treatment of CRPS. Early recognition and a multidisciplinary approach to management seem important in obtaining a good outcome. Treatments aimed at pain reduction and rehabilitation of limb function form the mainstay of therapy. Bisphosphonate-type compounds and steroids have been used as pharmacotherapy, with some beneficial effect [116]. Emerging data indicate a possible role of inflammation in the overall pathophysiology and have led to treatment with anti-inflammatory medications [117] and steroids. Numerous other oral drugs, including muscle relaxants, benzodiazepines, antidepressants, anticonvulsants, and opioids, have been reported on anecdotally [116]. Behavioural therapy, physiotherapy or occupational therapy is also reported in the armamentarium of CRPS treatment, possessing a potential beneficial effect [118]. Invasive treatments, including nerve blocks, intraspinally administered drugs and neuromodulatory therapies like spinal cord stimulation have all been attempted, but no definite treatment has been established [116].

Due to the possibility of sympathetic overactivity, sympathetic ablation has been used to treat CRPS as well. This concept has been validated in an animal study aimed to determine the effect of sympathectomy on allodynia and heat hyperalgesia produced by a prior L5 and L6 spinal nerve ligation [119]. In a human study, overactivity of the sympathetic system has been demonstrated as well [120]. Yet, two systematic reviews [121, 122] detected only uncontrolled studies and reports of personal experience, and some authors advocated that sympathectomy should be avoided in the treatment of CRPS [123]. Nevertheless, the use of sympathetic ablation in patients with CRPS is repeatedly published. Chemical sympathetic blocks have been successful [124] as well as permanent sympathetic denervation achieved by surgery [120, 125–130]. Even in cases with block failure, sympathectomy may still succeed [102]. Good reported results are as high as 94 % [126]. A meta-analysis of 1528 collected cases of causalgia [131] published in 2003 considered most cases to be cured by

surgical sympathectomy. However, it seems necessary to perform the ablation early in the course of CRPS as with time it becomes independent of sympathetic activity [132]. The level of sympathetic ablation performed has varied between the stellate ganglion down to T7 [100], but more recently the stellate ganglion has not been included, and the usual level in use was restricted from T2 only [129] to T2–T3/T4 [128]. Finally, it may be of interest to report an anecdotal case in which sympathetically maintained pain which recurred after sympathectomy was relieved by the addition of contralateral sympathectomy, raising the question of crossover sympathetic innervation [133].

Until systematic controlled studies are performed, the present data allows cautious use of timely sympathetic ablation in properly selected patients.

Erythromelalgia

Erythromelalgia is a rare syndrome of unknown aetiology, characterized by episodes of burning pain, redness, oedema and increased temperature in the upper, lower or both extremities [134]. SCN9A gene mutation has been identified in some but not in all patients [135]. Patients with erythromelalgia without SCN9A gene mutation seem to be more refractory to treatment [135]. Erythromelalgia may be primary or secondary to several conditions, and microvascular arteriovenous shunting has been implied [136]. Several drugs have been used in its treatment, aspirin being foremost [136]. A neurophysiologic study has shown another pathology: distal small-fibre neuropathy with selective involvement of cutaneous sympathetic fibres [137]. Based on this finding, sympathetic ablation has been used in the treatment of erythromelalgia, but the reported cases are scanty. Both failure [138] and success [134, 135, 139] have been reported.

Pancreatic pain and other painful abdominal pathologies

Sympathetic ablation has been reported as beneficial for abdominal pain. The main indications are pancreatic pathologies (cancer or chronic inflammation) [140], but also adrenal metastases [141], and other abdominal malignancies as validated in a recent prospective multicentre trial [142]. Sympathetic ablation has been used as well in the treatment of intractable abdominal pain due to other pathologies, such as portal vein thrombosis [143].

Splanchnicectomy is the usual procedure [140, 141, 143–149]. The technique includes transection of the post-ganglionic sympathetic nerve branches which unite to create the greater and the lesser splanchnic nerves. However, addition of sympathetic chain ablation has also been reported. Kusano et al. [144] performed a T5 to T9 chain resection/electrocoagulation, whereas Lönroth et al. [143]

divided the sympathetic chain and splanchnic branches from T4 down to T10–11. Although a bilateral approach has been advocated [140, 145, 146], some authors [144–149] recommended a unilateral approach, according to the location of the abdominal pain, reserving the bilateral approach only for diffuse abdominal pain. If pain recurred, a contralateral additional splanchnicectomy was performed. The endoscopic approach is invariably used. The timing of the operation in relation to other treatments seems to be important. Howard et al. [146] were impressed that splanchnicectomy appeared to work best in patients who have had no prior operative or endoscopic intervention. Isa et al. [150], in a systematic review published in 2014, reported that pre-operative opioid use was associated with a worse outcome after thoracoscopic splanchnicectomy in patients with chronic pancreatitis. Similarly, Amr and Makharita [142], in their prospective randomized multicentre study of patients with inoperable abdominal or pelvic cancer, found that early sympathetic ablation obtained better results than in patients operated on after opioid treatment had failed.

When performed for pain due to malignancy, the results are good because survival is short. When performed for chronic pancreatitis, results of splanchnicectomy are variable. Short-term results are almost invariably good [141, 146, 149]. Studies with long-term follow-up reported variable outcomes. Stone and Chauvin [149], in their series of alcoholic pancreatitis patients, reported persistent relief with a mean follow-up of 16 months. However, in this study, a bilateral truncal vagotomy had been added to the sympathetic procedure. The role of additional vagal ablation has not been elucidated, and further appropriate studies are required. Kusano et al. [144] reported pain relief at a 13.7-month mean follow-up in two-thirds of patients only. In the study of Buscher et al. [145], only a 50 % long-term persistent pain relief was reported, whereas Maher et al. [148] reported only temporary improvement in most patients.

Ischaemia

In the past, sympathectomy was used extensively for ischaemic disorders. Mathieu Jaboulay performed the first sympathetic ablation for an ischaemic foot ulcer in 1898 by perivascular denudation [151]. René Leriche, in 1913, extended the use of this procedure [152]. In 1924, Royle [153] performed the first lumbar sympathectomy for spastic paralysis, whereas Diez [154] was the first to use this procedure for ischaemic lesions of the limbs. Adson [155] in 1925 extended the indications to include vasospastic disorders.

Ischaemia results from several obstructive and spastic disorders.

Obstructive ischaemia

Atherosclerosis and thromboangiitis obliterans (TAO also called Buerger's disease) are the main pathologies causing ischaemia, the former affecting primarily the leg, whereas the latter having a tendency to involve the hands as well in its advanced stages. Atherosclerosis may limit walking due to insufficient muscular oxygenation during exercise (intermittent claudication) or if more severe may cause tissue loss. Several studies which have elucidated the pathophysiological effect of sympathetic ablation on muscular perfusion [156–167] have shown that sympathectomy is not beneficial in intermittent claudication. Although rare publications still report on sympathectomies performed for atherosclerosis causing digital ulcerations [168], success is limited and far inferior to that for Raynaud's phenomenon. The present state is that sympathectomy is not indicated in the treatment of atherosclerotic ischaemia.

TAO is a non-atherosclerotic segmental vasculitis affecting the small- and medium-sized arteries and veins of the extremities and is strongly associated with smoking [169, 170]. Upper limb sympathetic ablation has been successfully performed for intractable pain, ulceration or gangrene [171–174]. However, in a recent analysis of 344 cases, the duration of effect was limited, especially if smoking continued [175].

Whatever the source, microemboli may induce digital ischaemia and be unamenable to embolectomy. In such instances, sympathectomy may prove to be beneficial. Whatever the source of emboli [176–180], the treatment of all these lesions should be directed to the primary pathology, reserving additional sympathetic ablation for the cases in which digital ischaemia persists. A preliminary sympathetic chemical block may indicate likelihood of the successful sympathetic ablation [177].

Vasoconstrictive ischaemia

Vasoconstrictive ischaemia may occur in a series of pathologies: primary Raynaud's phenomenon, scleroderma, rheumatoid arthritis, lupus erythematosus, paraneoplastic syndrome and vibration syndrome.

Raynaud's phenomenon is a paroxysmal arterial or arteriolar vasospastic disorder predominantly afflicting the fingers, characterized by digital ischaemia induced by cold or emotional stress [181]. Raynaud's phenomenon may be primary, or secondary to other underlying diseases (collagen diseases, paraneoplastic syndrome, frostbite and vibration syndrome). When present, treatment should be directed to the primary pathology, as well as prevention of cold exposure and medications such as vasodilator drugs and calcium channel blockers [181]. In advanced stages, when persistent ischaemia, ulceration and gangrene

develop upper thoracic sympathectomy may alleviate ischaemic pain, maximize tissue preservation and limit amputation [182, 183]. However, the effect of sympathetic ablation was found to be temporary in the majority of cases [182, 183]. This was recently reconfirmed in a "summary of evidence" study [184]. Therefore, an initial attempt to obtain sympathetic block by phenol injections [185] may be beneficial, leaving the sympathetic chain intact for a repeated ablation for recurrent symptoms. Reoperative digital sympathectomy has been proven to be successful [186].

In scleroderma, half of all patients develop digital ischaemia. Initial treatment is medical, including prevention of exposure to cold, stress and trauma; smoking cessation, and oral medications like vasodilating agents, statins, endothelin receptor antagonists and phosphodiesterase-5 inhibitors [187]. If digital ulcers develop, intravenous prostacyclins may help [187]. Upper thoracic sympathetic ablation is occasionally performed for scleroderma with critical digital ischaemia [188, 189]. In the majority of reports for this indication, a periarterial [167] or digital [190, 191] approach is described. The reason is probably due to the additional surgical orthopaedic procedures required, which are performed by hand orthopaedic surgeons. The results are good on short term and may persist or deteriorate according to the progression of the basic disease [167]. There are no studies comparing digital versus upper thoracic sympathectomy. If an additional procedure in the hand is required, performing a local sympathectomy is logical. However, the perivascular digital approach requires an individual operation for each involved finger, whereas an upper thoracic sympathetic ablation affects all digits and, logically, should be the procedure of choice in cases with multiple affected fingers.

Digital ischaemia has been described in rheumatoid arthritis and lupus erythematosus (two collagen diseases), and sympathetic ablation has been reported to be beneficial [167]. One sign of the paraneoplastic syndrome is digital ischaemia. In one such reported case with severe bilateral Raynaud's phenomenon, bilateral sympathectomy permanently relieved ischaemia of the hands until demise of the patients after 10 months [192].

The first description of the pathophysiology of frostbite was given by Larrey, Surgeon-General of the imperial French armies, who participated in the 1812 disastrous retreat from Moscow of the Grande Armée [193]. Frostbite damage develops through three major mechanisms: direct cellular injury, hypoxia, and the release of vasoactive and toxic by-products [194]. The current literature contains work both discrediting [195] and supporting [196–198] the effect of sympathetic ablation in frostbite injuries. According to these studies, sympathetic ablation in the immediate post-thawing period may be detrimental by

increasing oedema and accelerating tissue destruction [199]. However, if delayed until 24–48 h, the opposite effect has been observed. Pain and paraesthesia are also improved [200]. The beneficial effect of sympathectomy in the long term has been confirmed in a recent study [201].

Hand-arm vibration syndrome is one of the most common occupational disorders in the industrialized world [202]. Symptoms include paraesthesiae and pain, unrelated to episodes of “white fingers” [203]. In advanced cases, vasospasm and ischaemic changes including gangrenous lesions have been observed [202, 204]. However, unlike the case of other primary pathologies and Raynaud’s phenomena, there is a striking paucity of studies advocating sympathetic ablation in the treatment of vibration-induced ischaemic lesions. Probably, this is because simply avoiding vibration improves symptoms [202].

Hypertension

The treatment of hypertension by sympathetic ablation was initiated in the 1930s [205], first by the transthoracic approach and later at the infradiaphragmatic level [206]. The efficacy of sympathetic ablation in the treatment of hypertension was confirmed in the study of Smithwick and Thompson [207] who in 1953 compared 1266 hypertensive patients following thoracolumbar sympathectomy with 467 hypertensive controls and observed 5-year mortality rates of 19 and 54 %, respectively. Of those who survived surgery, 45 % had significantly lower blood pressure afterwards, and the therapeutic effect of sympathectomy lasted 10 years or more. With the advent of modern effective antihypertensive drugs with minimal side effects, sympathectomy was largely abandoned in the 1960s [208]. However, there is a small percentage (but substantial number) of patients with resistant hypertension. For these patients, renal sympathectomy remains a useful treatment. Open periarterial stripping of the renal vessels was used in the past to obtain renal sympathetic denervation [209]. With the advent of endovascular procedures and the possibility to ablate the adventitia by intraluminal radiofrequency [210], renal sympathetic ablation became a relatively simple procedure and gained popularity. The long-term efficacy of the procedure remains to be determined. At present, it is not considered to be a routine procedure for resistant hypertension, a better design and future randomized studies being required [211]. The thoracic approach for sympathetic ablation to treat hypertension is obsolete.

Psychiatric indications

As odd as it may sound, sympathetic ablation for a psychiatric indication has been performed, not in the early

years of surgery, but as late as this century [212]. In this article, the authors reviewed their experience with sympathetic ablation in the treatment of blushing, excessive sweating and “social phobia with and without physical symptoms”. A social phobia or, as it should be better called, social withdrawal or anxiety may be secondary to a physical problem, like palmar hyperhidrosis or axillary overperspiration [213]. It has been shown that hyperhidrotics show higher stress levels in social interactions and more depressive symptoms compared to control subjects [214]. Social withdrawal may also be secondary to blushing [81, 215, 216]. In these cases, it is understandable that curing the underlining condition, patients’ social withdrawal will also be reduced. The reverse, namely overperspiration, being secondary to social anxiety disorders is also possible [217]. However, treating a primary phobia by sympathetic surgery has no scientific basis in the literature.

A search in the Medline under the terms “Social Phobia and Sympathetic Ablation” has yielded only five publications dealing specifically with the subject, all by the same group of authors [218–222]. A scrupulous review of these publications leads to the following two conclusions. Firstly, the authors’ classification is based on an hypothesis of neural pathways, but there is no objective proof for it being correct. Secondly, there is no objective proof that it is correct to use sympathetic ablation in the treatment of a primary phobia/anxiety. The fact that all treatments have failed is no justification to treat a primary phobia by sympathetic surgery [219, 221]. In their article [219], the authors claim to have operated upon 45 people for a social phobia and performed a follow-up on 20 patients. However, all they present in the article is a single case report! Performing sympathetic ablation for a primary social phobia remains equivalent to performing a scientifically yet unproven human experiment. There is no psychiatric indication for sympathetic surgery.

Conclusions

Thoracic sympathetic ablation is currently the only curative modality for primary palmar hyperhidrosis. Although serious sequelae may ensue, patients with severe symptoms should be offered this possibility, provided they receive a detailed information regarding side effects. Presently, there are several effective non-surgical and surgical modalities of treating axillary hyperhidrosis. As the results of sympathectomy for axillary hyperhidrosis are inferior to those for palmar hyperhidrosis, sympathetic ablation should be reserved only for cases in which the operation is performed primarily for concomitant palmar hyperhidrosis. Sympathetic ablation should be reserved as the last line of

treatment for facial overperspiration and blushing. Compensatory hyperhidrosis, the development of excessive sweating in areas of the body unaffected anatomically by the sympathetic denervation, and to a lesser degree gustatory sweating, are the major drawbacks of sympathetic surgery for primary hyperhidrosis, which requires its strictly judicious use, for these cases. Thoracic sympathetic ablation has a place in some rare and strictly selected cases of angina, arrhythmias and cardiomyopathy unresponsive to standard medication. Sympathetic ablation may be beneficial if performed in the early stages of complex regional pain syndrome, and some painful abdominal pathologies, in particular pancreatic or other malignancies. The use of sympathetic ablation for ischaemia has substantially declined, but still has a place in thromboangiitis obliterans and selected cases of microemboli, primary Raynaud's phenomenon, Raynaud's phenomenon secondary to collagen diseases, paraneoplastic syndrome and frostbite. In dubious cases, a preliminary chemical sympathetic block may suggest what benefit could be expected from the procedure. Since the advent of endovascular renal sympathetic ablation, thoracic sympathetic procedures for hypertension became obsolete. Sympathetic ablation is being used by a narrow group of surgeons for cases of primary phobias, but it should be emphasized that there is no scientific basis to perform sympathetic surgery for any psychiatric indication.

Compliance with Ethical Standards

Disclosures Moshe Hashmonai, Alan E. P. Cameron, Peter B. Licht, Chris Hensman and Christoph H. Schick have no conflicts of interest or financial ties to disclose.

References

- Alexander W (1889) The treatment of epilepsy. *YJ Pentland, Edinburgh*, pp 27–106
- Jonnesco T (1896) Resectia totala si bilaterala a simpaticulu cervical in cazuri de epilepsie si gusa exophthalmica. *Romania Med* 4:479–496
- Jaboulay M (1900) La régénération du goitre extirpé dans la maladie de Basedow et la section du sympathique cervical dans cette maladie. In: Martin E (ed) *Chirurgie du Grand Sympathique et du Corps Thyroïde (Les Différents Goïtres)*. Articles originaux et observations réunis, A Storck & Cie, Lyon, pp 3–5
- François-Franck M (1899) Signification physiologique de la résection du sympathique dans la maladie de Basedow, l'épilepsie, l'idiotie et le glaucome. *Bull Acad Med Paris* 41:565–594
- Sato K, Kang WH, Saga K, Sato KT (1989) Biology of the sweat glands and their disorders II. Disorders of sweat gland function. *J Am Acad Dermatol* 20:713–726
- Sato K (1991) Hyperhidrosis. *JAMA* 265:651
- Connolly M, de Berker D (2003) Management of primary hyperhidrosis. A summary of the different treatment modalities. *Am J Clin Dermatol* 4:681–697
- Hashmonai M, Kopelman D, Assalia A (2000) The treatment of primary palmar hyperhidrosis: a review. *Surg Today* 30: 211–218
- Kotzareff A (1920) Résection partielle du tronc sympathique cervical droit pour hyperhidrose unilatérale (régions faciale, cervicale, thoracique et brachiale droites). *Rev Med Suisse Rom* 40:111–113
- Panhofer P, Zacherl J, Jakesz R, Bischof G, Neumayer C (2006) Improved quality of life after sympathetic block for upper limb hyperhidrosis. *Br J Surg* 93:582–586
- Panhofer P, Gleiss A, Eilenberg WH, Jakesz R, Bischof G, Neumayer C (2013) Long-term outcomes after endorhachic sympathetic block at the T4 ganglion for upper limb hyperhidrosis. *Br J Surg* 100:1471–1477
- Wolosker N, Yazbek G, de Campos JR, Munia MA, Kauffman P, Jatene FB, Puech-Leão P (2010) Quality of life before surgery is a predictive factor for satisfaction among patients undergoing sympathectomy to treat hyperhidrosis. *J Vasc Surg* 51: 1190–1194
- Hashmonai M (2014) Surgical options for the treatment of primary palmar hyperhidrosis: consensus and controversies. In: Huddle JR (ed) *Hyperhidrosis: causes, treatment options and outcomes*. Nova Science Publishers Inc, Hauppauge, pp 111–135
- Hashmonai M, Shein M (1994) Upper thoracic sympathectomy—open approaches. In: Paterson-Brown S, Garden J (eds) *Principles and practice of surgical laparoscopy*. WB Saunders, London, pp 587–603
- Miller DL, Force SD (2008) Temporary thoracoscopic block for hyperhidrosis. *Ann Thorac Surg* 85:1211–1214
- Gunn TM, Davis DM, Speicher JE, Rossi NP, Parekh KR, Lynch WR, Iannettoni MD (2014) Extended level of sympathetic chain removal does not increase the incidence of severity of compensatory hyperhidrosis after endoscopic thoracic sympathectomy. *J Thorac Cardiovasc Surg* 148:2673–2676
- Ishy A, de Campos JR, Wolosker N, Kauffman P, Tedde ML, Chiavoni CR, Jatene FB (2011) Objective evaluation of patients with palmar hyperhidrosis submitted to two levels of sympathectomy: T3 and T4. *Interact CardioVasc Thorac Surg* 12:545–548
- Reisfeld R (2006) Sympathectomy for hyperhidrosis: should we place the clamps at T2–T3 or T3–T4? *Clin Auton Res* 16:384–389
- Li X, Tu YR, Lin M, Lai FC, Chen JF, Dai ZJ (2008) Endoscopic thoracic sympathectomy for palmar hyperhidrosis: a randomized control trial comparing T3 and T2–4 ablation. *Ann Thorac Surg* 85:1747–1751
- Cai S, Huang S, An J, Li Y, Weng Y, Liao H, Chen H, Liu L, He J, Zhang J (2014) Effect of lowering or restricting sympathectomy levels on compensatory hyperhidrosis. *Clin Auton Res* 24:143–149
- Katara AN, Domino JP, Cheah WK, So JB, Ning C, Lomanto D (2007) Comparing T2 and T2–T3 ablation in thoracoscopic sympathectomy for palmar hyperhidrosis; a randomized control trial. *Surg Endosc* 21:1768–1771
- Yano M, Kiriya M, Fukai I, Sasaki H, Kobayashi Y, Mizuno K, Haneda H, Suzuki E, Endo K, Fujii Y (2005) Endoscopic thoracic sympathectomy for palmar hyperhidrosis: efficacy of T2 and T3 ganglion resection. *Surgery* 138:40–45
- Cerfolio RJ, de Campos JRM, Bryant AS, Connery CP, Miller DL, DeCamp MM, McKenna RJ, Krasna MJ (2011) The society of thoracic surgeons expert consensus for the surgical treatment of hyperhidrosis. *Ann Thorac Surg* 91:1642–1648
- Deng B, Tan QY, Jiang YG, Zhao YP, Zhou JH, Ma Z, Wang RW (2011) Optimization of sympathectomy to treat palmar

- hyperhidrosis: the systemic review and meta-analysis of studies published in the last decade. *Surg Endosc* 25:1893–1901
25. Panhofer P, Ringhofer C, Gleiss A, Jakesz R, Prager M, Bischof G, Neumayer C (2014) Quality of life after sympathetic surgery at the T4 ganglion for primary hyperhidrosis: clip application versus diathermy cut. *Int J Surg* 12:1478–1483
 26. Baumgartner FJ, Reyes M, Sarkisyan GG, Iglesias A, Reyes E (2011) Thoracoscopic sympathectomy for disabling hyperhidrosis: a prospective randomized comparison between two levels. *Ann Thorac Surg* 92:2015–2019
 27. Liu Y, Yang J, Liu J, Yang F, Jiang G, Li J, Huang Y, Wang J (2009) Surgical treatment of primary palmar hyperhidrosis: a prospective randomized study comparing T3 and T4 sympatotomy. *Eur J Cardiothorac Surg* 35:398–402
 28. Furlan AD, Mailis A, Papagapiou M (2000) Are we paying a high price for surgical sympathectomy? A systematic literature review of late complications. *J Pain* 1:245–257
 29. Cameron AE (2003) Specific complications and mortality of endoscopic thoracic sympathectomy. *Clin Auton Res* 13(Suppl 1):I31–I35
 30. Ojimba TA, Cameron AE (2004) Drawbacks of endoscopic thoracic sympathectomy. *Br J Surg* 91:264–269
 31. Gossot D, Kabiri H, Caliandro R, Debrosse D, Girard P, Grunenwald D (2001) Early complications of thoracoscopic sympathectomy: a prospective study of 940 procedures. *Ann Thorac Surg* 71:1116–1119
 32. Kao MC (2001) Transthoracic endoscopic sympathectomy. *Surg Endosc* 15:1365
 33. Lin TS, Fang HY, Wu CY (2000) Repeat transthoracic endoscopic sympathectomy for palmar and axillary hyperhidrosis. *Surg Endosc* 14:134–136
 34. Freeman RK, van Woerkom JM, Vyverberg A, Ascioti AJ (2009) Reoperative endoscopic sympathectomy for persistent or recurrent palmar hyperhidrosis. *Ann Thorac Surg* 88:412–417
 35. Licht PB, Clausen A, Ladegaard L (2010) Resympathectomy. *Ann Thorac Surg* 89:1087–1090
 36. Walling HW, Swick BL (2011) Treatment options for hyperhidrosis. *Am J Clin Dermatol* 12:285–295
 37. Baumgartner FJ, Bertin S, Konecny J (2009) Superiority of thoracoscopic sympathectomy over medical management for the palmo-plantar subset of severe hyperhidrosis. *Ann Vasc Surg* 23:1–7
 38. Ellis H (1979) Transaxillary sympathectomy in the treatment of hyperhidrosis of the upper limb. *Am Surg* 45:546–551
 39. Moran KT, Brady MP (1991) Surgical management of primary hyperhidrosis. *Br J Surg* 78:279–283
 40. Gossot D, Debrosse D, Grunenwald D (2000) Sympathectomie thoracique pour hyperhidrose axillaire isolée. (full text in English on www.e2med.com/ad) *Ann Dermatol Venereol* 127:1065–1067
 41. Hsu CP, Shia SE, Hsia JY, Chuang CY, Chen CY (2001) Experiences in thoracoscopic sympathectomy for axillary hyperhidrosis and osmidrosis: focusing on the extent of sympathectomy. *Arch Surg* 136:1115–1117
 42. Munia MAS, Wolosker N, Kaufmann P, Milanes de Campos JR, Puech-Leão P (2008) Sustained benefit lasting one year from T4 instead of T3–T4 sympathectomy for isolated axillary hyperhidrosis. *Clinics* 63:771–774
 43. Rex LO, Drott C, Claes G, Göthberg G, Dalman P (1998) The Borås experience of endoscopic thoracic sympathectomy for palmar, axillary, facial hyperhidrosis and facial blushing. *Eur J Surg* 164(Suppl 580):23–26
 44. Hong HC, Lupin M, O’Shaughnessy KF (2012) Clinical evaluation of a microwave device for treating axillary hyperhidrosis. *Dermatol Surg* 38:728–735
 45. Rompel R, Peros I, Petres J (1997) Subcutaneous curettage for the treatment of axillary hyperhidrosis. *Eur J Dermatol* 7:43–46
 46. Wollina U, Köstler E, Schönlebe J, Haroske G (2008) Tumescence suction curettage versus minimal skin resection with subcutaneous curettage of sweat glands in axillary hyperhidrosis. *Dermatol Surg* 34:709–716
 47. Weaver PC (1970) Axillary skin excision as a treatment for axillary hyperhidrosis. *Postgrad Med J* 46:422–424
 48. Stenquist B (1985) Axillary hyperhidrosis: a simple surgical procedure. *J Dermatol Surg Oncol* 11:388–391
 49. Heidemann E, Licht PB (2013) A comparative study of thoracoscopic sympathectomy versus local surgical treatment for axillary hyperhidrosis. *Ann Thorac Surg* 95:264–268
 50. Walling HW (2011) Clinical differentiation of primary from secondary hyperhidrosis. *J Am Acad Dermatol* 64:690–695
 51. Benohanian A, Solish N (2004) Hyperhidrosis: sweating out the details. *Can J Cont Med Ed* 16:77–81
 52. Reisfeld R, Nguyen R, Pnini A (2000) Endoscopic thoracic sympathetic ablation for treatment of essential hyperhidrosis syndrome: experience with 650 patients. *Surg Laparosc Endosc Percutan Tech* 10:5–10
 53. Kao MC, Chen YL, Lin JY, Hsieh CS, Tsai JC (1996) Endoscopic sympathectomy treatment for craniofacial hyperhidrosis. *Arch Surg* 131:1091–1094
 54. Kim DH, Paik HC, Lee DY (2004) Comparative analysis of T2 selective division of rami-communicantes (ramicotomy) with T2 sympathetic clipping in the treatment of craniofacial hyperhidrosis. *Eur J Cardiothorac Surg* 26:396–400
 55. Lin TS, Fang HY (2000) Transthoracic endoscopic sympathectomy for craniofacial hyperhidrosis: analysis of 46 cases. *J Laparoendosc Adv Surg Tech A* 10:243–247
 56. Lin TS, Chou MC (2002) Needleoscopic thoracic sympathetic block by clipping for craniofacial hyperhidrosis: an analysis of 28 cases. *Surg Endosc* 16:1055–1058
 57. Izikson L, English JC 3rd, Zirwas MJ (2006) The flushing patient: differential diagnosis, workup, and treatment. *J Am Acad Dermatol* 55:193–208
 58. Scholing A, Emmelkamp PM (1996) Treatment of fear of blushing, sweating, or trembling. Results at long-term follow-up. *Behav Modif* 20:338–356
 59. Dijk C, Buwalda FM, de Jong PJ (2012) Dealing with fear of blushing: a psychoeducational group intervention for fear of blushing. *Clin Psychol Psychother* 19:481–487
 60. Drummond PD, Minosora K, Little G, Keay W (2013) Topical ibuprofen inhibits blushing during embarrassment and facial flushing during aerobic exercise in people with a fear of blushing. *Eur Neuropsychopharmacol* 23:1747–1753
 61. Pelissolo A, Moukheiber A (2013) Open-label treatment with escitalopram in patients with social anxiety disorder and fear of blushing. *J Clin Psychopharmacol* 33:695–698
 62. Garcia Tirado FJ, Rieger Reyes C (2013) Tiene tratamiento la eretofobia? *Aten Primaria* 45:279–281
 63. Wittmoser R (1985) Treatment of sweating and blushing by endoscopic surgery. *Acta Neurochir (Wien)* 74:153–154
 64. Yilmaz EN, Dur AH, Cuesta MA, Rauwerda JA (1996) Endoscopic versus transaxillary thoracic sympathectomy for primary axillary and palmar hyperhidrosis and/or facial blushing: 5-year experience. *Eur J Cardiothorac Surg* 10:168–172
 65. Drott C, Claes G, Olsson-Rex L, Dalman P, Fahlén T, Göthberg G (1998) Successful treatment of facial blushing by endoscopic transthoracic sympathectomy. *Br J Dermatol* 138:639–643
 66. Krasna MJ, Jiao X, Sonett J, Gamliel Z, King K (2000) Thoracoscopic sympathectomy. *Surg Laparosc Endosc Percutan Tech* 10:314–318
 67. Drott C, Claes G, Rex L, Dalman P, Göthberg G, Fahlén T (2001) Long-term effects after surgery for hand sweating and facial blushing. Patients are satisfied in spite of troublesome side-effects. *Lakartidningen* 98:1766–1772

68. Ueyama T, Matsumoto Y, Abe Y, Yuge O, Iwai T (2001) Endoscopic thoracic sympathectomy in Japan. *Ann Chir Gynaecol* 90:200–202
69. Drott C, Claes G, Rex L (2002) Facial blushing treated by sympathetic denervation—longlasting benefits in 831 patients. *J Cosmet Dermatol* 1:115–119
70. Rajesh YS, Pratap CP, Woodyer AB (2002) Thoracoscopic sympathectomy for palmar hyperhidrosis and Raynaud's phenomenon of the upper limb and excessive facial blushing: a five year experience. *Postgrad Med J* 78:682–684
71. Lardinois D, Ris HB (2002) Minimally invasive video-endoscopic sympathectomy by use of a transaxillary single port approach. *Eur J Cardiothorac Surg* 21:67–70
72. Sterodimas A, Nicolaou M, Paes TR (2003) Successful use of Botulinum toxin-A for the treatment of neck and anterior chest wall flushing. *Clin Exp Dermatol* 28:592–594
73. Neumayer C, Zacherl J, Holak G, Jakesz R, Bischof G (2003) Experience with limited endoscopic thoracic sympathetic block for hyperhidrosis and facial blushing. *Clin Auton Res* 13(Suppl 1):152–157
74. Yuraitis M, Jacob CI (2004) Botulinum toxin for the treatment of facial flushing. *Dermatol Surg* 30:102–104
75. Saetre T, Flørenes T (2005) Thoracoscopic sympathectomy in the treatment of palmar hyperhidrosis and facial blushing. *Tidsskr Nor Laegeforen* 125:2795–2797
76. Adair A, George ML, Camprodon R, Broadfield JA, Rennie JA (2005) Endoscopic sympathectomy in the treatment of facial blushing. *Ann R Coll Surg Engl* 87:358–360
77. Licht PB, Ladegaard L, Pilegaard HK (2006) Thoracoscopic sympathectomy for isolated facial blushing. *Ann Thorac Surg* 81:1863–1866
78. Nicolaou M, Paes T, Wakelin S (2006) Blushing: an embarrassing condition, but treatable. *Lancet* 367:1297–1299
79. Chou SH, Kao EL, Lin CC, Huang M (2006) The outcome of ganglion clipping in hyperhidrosis and blushing. *Clin Auton Res* 16:240–242
80. Malmivaara A, Kuukasjärvi P, Autti-Ramo I, Kovanen N, Mäkelä M (2007) Effectiveness and safety of endoscopic thoracic sympathectomy for excessive sweating and facial blushing: a systematic review. *Int J Technol Assess Health Care* 23:54–62
81. Fibla JJ, Molins L, Mier JM, Vidal G (2009) Effectiveness of sympathetic block by clipping in the treatment of hyperhidrosis and facial blushing. *Interact CardioVasc Thorac Surg* 9:970–972
82. Jadresic E, Suárez C, Palacios E, Palacios F, Matus P (2011) Evaluating the efficacy of endoscopic thoracic sympathetic ablation for generalized social anxiety disorder with blushing complaints: a comparison with sertraline and no treatment—Santiago de Chile 2003–2009. *Innov Clin Neurosci* 8:24–35
83. Coveliers H, Atif S, Rauwerda J, Wisselink W (2011) Endoscopic thoracic sympathectomy: long-term results for treatment of upper limb hyperhidrosis and facial blushing. *Acta Chir Belg* 111:293–297
84. Smidfelt K, Drott C (2011) Late results of endoscopic thoracic sympathectomy for hyperhidrosis and facial blushing. *Br J Surg* 98:1719–1724
85. Licht PB, Pilegaard HK, Ladegaard L (2012) Sympathectomy for isolated facial blushing: a randomized clinical trial. *Ann Thorac Surg* 94:401–405
86. Ransohoff JL (1925) Cervical sympathectomy for angina pectoris. *Ann Surg* 81:585–590
87. Pather N, Partab P, Singh B, Satyapal KS (2003) The sympathetic contributions to the cardiac plexus. *Surg Radiol Anat* 25:210–215
88. Galiñanes M, Loubani M, Sensky PR, Hassouna A, Cherryman GR, Leverment JN, Samani NJ (2004) Efficacy of transmyocardial laser revascularization and thoracic sympathectomy for the treatment of refractory angina. *Ann Thorac Surg* 78:122–128
89. Yoshida K, Inoue T, Hirakawa N, Node K (2008) Endoscopic sympathectomy as a novel strategy for vasospastic angina refractory to medical treatments. *J Cardiol* 52:49–52
90. Abbate A, Hamza M, Cassano AD, Melchior R, Roberts C, Grizzard J, Shah K, Hastillo A, Kasirajan V, Crea F, Lanza GA, Vetrovec GW (2012) Sympathectomy as a treatment for refractory coronary artery spasm. *Int J Cardiol* 161:e7–e9
91. Schwartz PJ, Malliani A (1975) Electrical alternation of the T-wave: clinical and experimental evidence of its relationship with the sympathetic nervous system and with the long Q-T syndrome. *Am Heart J* 89:45–50
92. Ajjjola OA, Lellouche N, Bourke T, Tung R, Ahn S, Mahajan A, Shivkumar K (2012) Bilateral cardiac sympathetic denervation for the management of electrical storm. *J Am Coll Cardiol* 59:91–92
93. Shen MJ, Zipes DP (2014) Role of the autonomic nervous system in modulating cardiac arrhythmias. *Circ Res* 114:1004–1021
94. Methangkool E, Chua JH, Gopinath A, Shivkumar K, Mahajan A (2014) Anesthetic considerations for thoracoscopic sympathetic ganglionectomy to treat ventricular tachycardia storm: a single-center experience. *J Cardiothorac Vasc Anesth* 28:69–75
95. Sosunov EA, Anyukhovskiy EP, Gainullin RZ, Plotnikov A, Danilo P Jr, Rosen MR (2001) Long-term electrophysiological effects of regional cardiac sympathetic denervation of the neonatal dog. *Cardiovasc Res* 51:659–669
96. Gilmour RF Jr (2001) Life out of balance: the sympathetic nervous system and cardiac arrhythmias. *Cardiovasc Res* 51:625–626
97. Estes EH Jr, Izlar HL Jr (1961) Recurrent ventricular tachycardia. A case successfully treated by bilateral cardiac sympathectomy. *Am J Med* 31:493–497
98. Zipes DP, Festoff B, Schaal SF, Cox C, Sealy WC, Wallace AG (1968) Treatment of ventricular arrhythmia by permanent atrial pacemaker and cardiac sympathectomy. *Ann Intern Med* 68:591–597
99. Vaseghi M, Gima J, Kanaan C, Ajjjola OA, Marmureanu A, Mahajan A, Shivkumar K (2014) Cardiac sympathetic denervation in patients with refractory ventricular arrhythmias or electrical storm: immediate and long-term follow-up. *Heart Rhythm* 11:360–366
100. Linz D, Wirth K, Ukena C, Mahfoud F, Pösch J, Linz B, Böhm M, Neuberger HR (2013) Renal denervation suppresses ventricular arrhythmias during acute ventricular ischemia in pigs. *Heart Rhythm* 10:1525–1530
101. Remo BF, Preminger M, Bradfield J, Mittal S, Boyle N, Gupta A, Shivkumar K, Steinberg JS, Dickfeld T (2014) Safety and efficacy of renal denervation as a novel treatment of ventricular tachycardia storm in patients with cardiomyopathy. *Heart Rhythm* 11:541–546
102. Schwartz PJ (2013) Practical issues in the management of the long QT syndrome: focus on diagnosis and therapy. *Swiss Med Wkly* 143:w13843
103. Shimizu W (2005) The long QT syndromes: therapeutic implications of a genetic diagnosis. *Cardiovasc Res* 67:347–356
104. Wang LX (2003) Role of left cardiac sympathetic denervation in the management of congenital long QT syndrome. *J Postgrad Med* 49:179–181
105. Schwartz PJ, Ackerman MJ (2013) The long QT syndrome: a transatlantic clinical approach to diagnosis and therapy. *Eur Heart J* 34:3109–3116
106. Moss AJ, McDonald J (1971) Unilateral cervicothoracic sympathetic ganglionectomy for the treatment of long QT interval syndrome. *N Engl J Med* 285:903–904

107. Schwartz PJ, Priori SG, Cerrone M, Spazzolini C, Odero A, Napolitano C, Bloise R, De Ferrari GM, Klersy C, Moss AJ, Zareba W, Robinson JL, Hall WJ, Brink PA, Toivonen L, Epstein AE, Li C, Hu D (2004) Left cardiac sympathetic denervation in the management of high-risk patients affected by the long-QT syndrome. *Circulation* 109:1826–1833
108. Xiu CH, Liu FQ, Wang HQ, Wang X, Qu RH (2006) Effects of cardiac sympathetic blockade on left ventricular diastolic function in patients with dilated cardiomyopathy and severe heart failure. *Zhonghua Yi Xue Za Zhi* 86:1170–1173
109. Pêgo-Fernandes PM, Moreira LF, Souza GE, Bacal F, Bocchi EA, Stolf NA, Jatene FB (2010) Endoscopic left sympathetic blockade in the treatment for dilated cardiomyopathy. *Arq Bras Cardiol* 95:685–690
110. Guo W, Liu F, Fu L, Qu R, Wang G, Zhang C (2012) Effects of high thoracic epidural sympathetic blockade for the treatment of severe chronic heart failure due to dilated cardiomyopathy. *Acta Cardiol* 67:533–539
111. Thompson JE, Garrett WV (1991) Extremity causalgia and mimocausalgia. In: Ernst CB, Stanley JC (eds) *Current therapy in vascular surgery*, 2nd edn. BC Decker Inc, Philadelphia, pp 202–205
112. Low PA, Engstrom JW (2012) Disorders of the autonomic nervous system. In: Longo DL, Fauci AS, Kasper DL, Hauser SL, Jameson JL, Loscalzo J (eds) *Harrison's principles of internal medicine*, 18th edn. McGraw Hill, New York, pp 3351–3360
113. Tracey DJ, Walker JS (1995) Pain due to nerve damage: are inflammatory mediators involved? *Inflamm Res* 44:407–411
114. Goris RJA (1998) Reflex sympathetic dystrophy: model of a severe regional inflammatory response syndrome. *World J Surg* 22:197–202
115. Wasner G, Heckmann K, Maier C, Baron R (1999) Vascular abnormalities in acute reflex sympathetic dystrophy (CRPS I): complete inhibition of sympathetic nerve activity with recovery. *Arch Neurol* 56:613–620
116. Rowbotham MC (2006) Pharmacologic management of complex regional pain syndrome. *Clin J Pain* 22:425–429
117. Sharma A, Williams K, Raja SN (2006) Advances in treatment of complex regional pain syndrome: recent insights on a perplexing disease. *Curr Opin Anaesthesiol* 19:566–572
118. Albazaz R, Wong YT, Homer-Vanniasinkam S (2008) Complex regional pain syndrome: a review. *Ann Vasc Surg* 22:297–306
119. Kim SH, Na HS, Sheen K, Chung JM (1993) Effects of sympathectomy on a rat model of peripheral neuropathy. *Pain* 55:85–92
120. Freedman M, Greis AC, Marino L, Sinha A, Hestenburg J (2014) Complex regional pain syndrome: diagnosis and treatment. *Phys Med Rehabil Clin N Am* 25:291–303
121. Mailis A, Furlan A (2003) Sympathectomy for neuropathic pain. *Cochrane Database Syst Rev* 2:CD002918 [Update in *Cochrane Database Syst Rev* 2010(7):CD002918]
122. Straube S, Derry S, Moore RA, Cole P (2013) Cervico-thoracic or lumbar sympathectomy for neuropathic pain and complex regional pain syndrome. *Cochrane Database Syst Rev* 9:CD002918
123. Quisel A, Gill JM, Witherell P (2005) Complex regional pain syndrome: which treatments show promise? *J Fam Pract* 54:599–603
124. Agarwal-Kozlowski K, Lorke DE, Habermann CR, Schulte am Esch J, Beck H (2011) Interventional management of intractable sympathetically mediated pain by computed tomography-guided catheter implantation for block and neuroablation of the thoracic sympathetic chain: technical approach and review of 322 procedures. *Anaesthesia* 66:699–708
125. Baron R, Maier C (1996) Reflex sympathetic dystrophy: skin blood flow, sympathetic vasoconstrictor reflexes and pain before and after surgical sympathectomy. *Pain* 67:317–326
126. Mockus MB, Rutherford RB, Rosales C, Pearce WH (1987) Sympathectomy for causalgia. Patient selection and long-term results. *Arch Surg* 122:668–672
127. Olcott C, Eltherington LG, Wilcosky BR, Shoor PM, Zimmerman JJ, Fogarty TJ (1991) Reflex sympathetic dystrophy—the surgeon's role in management. *J Vasc Surg* 14:488–495
128. Bandyk DF, Johnson BL, Kirkpatrick AF, Novotney ML, Back MR, Schmacht DC (2002) Surgical sympathectomy for reflex sympathetic dystrophy syndromes. *J Vasc Surg* 35:269–277
129. Singh B, Moodley J, Shaik AS, Robbs JV (2003) Sympathectomy for complex regional pain syndrome. *J Vasc Surg* 37:508–511
130. Golubev VG, Krupatkin AI, Zeinalov VT, Merkulov MV, Kuz'michev VA (2008) New facilities in the management of complex regional pain upper limb syndrome with thoracoscopic sympathectomy. *Vestn Ross Akad Med Nauk* 8:52–55
131. Hassantash SA, Afrakhteh M, Maier RV (2003) Causalgia: a meta-analysis of the literature. *Arch Surg* 138:1226–1231
132. Schwartzman RJ (2000) New treatments for reflex sympathetic dystrophy. *New Engl J Med* 343:654–656
133. Valley MA, Rogers JN, Gale DW (1995) Relief of recurrent upper extremity sympathetically-maintained pain with contralateral sympathetic blocks: evidence for crossover sympathetic innervation? *J Pain Symptom Manag* 10:396–400
134. Shiga T, Sakamoto A, Koizumi K, Ogawa R (1999) Endoscopic thoracic sympathectomy for primary erythromelalgia in the upper extremities. *Anesth Analg* 88:865–866
135. Zhang L, Wang WH, Li LF, Dong GX, Zhao J, Luan JY, Sun TT (2010) Long-term remission of primary erythromelalgia with R1150 W polymorphism in SCN9A after chemical lumbar sympathectomy. *Eur J Dermatol* 20:763–767
136. Ljubojević S, Lipozencić J, Pustisek N (2004) Erythromelalgia. *Acta Dermatovenerol Croat* 12:99–105
137. Sandroni P, Davis MD, Harper CM, Rogers RS 3rd, Harper CM Jr, Rogers RS 3rd, Ofallon WM, Rooke TW, Low PA (1999) Neurophysiologic and vascular studies in erythromelalgia: a retrospective analysis. *J Clin Neuromuscul Dis* 1:57–63
138. Gaur S, Koroscil T (2009) Late-onset erythromelalgia in a previously healthy young woman: a case report and review of the literature. *J Med Case Rep* 3:106
139. Nakajima Y, Koizumi K, Hirata T, Hirai K, Sakamoto A, Shimizu K (2004) Successful thoracoscopic sympathectomy for primary erythromelalgia in the upper extremities. *Jpn J Thorac Cardiovasc Surg* 52:524–526
140. Ihse I, Zoucas E, Gyllstedt E, Lillo-Gil R, Andrén-Sandberg Å (1999) Bilateral thoracoscopic splanchnicectomy: effects on pancreatic pain and function. *Ann Surg* 230:785–790
141. Lang-Lazdunski L, Le Pimpec-Varthes F, Riquet M (2002) Videothoracoscopic splanchnicectomy for intractable pain from adrenal metastasis. *Ann Thorac Surg* 73:1290–1292
142. Amr YM, Makharrata MY (2014) Neurolytic sympathectomy in the management of cancer pain-time effect: a prospective, randomized multicenter study. *J Pain Symptom Manage* 48:944–956
143. Lönroth H, Hyltander A, Lundell L (1997) Unilateral left-sided thoracoscopic sympathectomy for visceral pain control: a pilot study. *Eur J Surg* 163:97–100
144. Kusano T, Miyazato H, Shiraiishi M, Yamada M, Matsumoto M, Muto Y (1997) Thoracoscopic thoracic splanchnicectomy for chronic pancreatitis with intractable abdominal pain. *Surg Laparosc Endosc* 7:213–218
145. Buscher HCJL, Jansen JBMJ, van Dongen R, Bleichrodt RP, van Goor H (2002) Long-term results of bilateral thoracoscopic splanchnicectomy in patients with chronic pancreatitis. *Br J Surg* 89:158–162
146. Howard TJ, Swofford JB, Wagner DL, Sherman S, Kehman GA (2002) Quality of life after bilateral thoracoscopic

- splanchnicectomy: long-term evaluation in patients with chronic pancreatitis. *J Gastrointest Surg* 6:845–852
147. Leksowski K (2001) Thoracoscopic splanchnicectomy for the relief of pain due to chronic pancreatitis. A pilot study. *Surg Endosc* 15:592–596
 148. Maher JW, Johlin FC, Heitshusen D (2001) Long-term follow-up of thoracoscopic splanchnicectomy for chronic pancreatic pain. *Surg Endosc* 15:706–709
 149. Stone HH, Chauvin EJ (1990) Pancreatic denervation for pain relief in chronic alcohol associated pancreatitis. *Br J Surg* 77:303–305
 150. Issa Y, Ali UA, Bouwense SAW, van Santvoort HC, van Goor H (2014) Preoperative opioid use and the outcome of thoracoscopic splanchnicectomy in chronic pancreatitis: a systemic review. *Surg Endosc* 28:405–412
 151. Jaboulay M (1899) Le traitement de quelques troubles trophiques du pied et de la jambe par la dénudation de l'artère fémorale et la distension des nerfs vasculaires. *Lyon Med* 91:467–468
 152. Leriche R (1913) De l'élongation et de la section des nerfs périsvasculaires dans certains syndromes douloureux d'origine artérielle et dans quelques troubles trophiques. *Lyon Chir* 10:378–382
 153. Royle ND (1924) The treatment of spastic paralysis by sympathetic ramisection. *Surg Gynecol Obstet* 39:701–720
 154. Diez J (1924) Un nuevo método de simpatectomia periférica para el tratamiento de las afecciones tróficas y gangrenosas de los miembros. *Bol Soc Cir Buenos Aires* 8:792–806
 155. Adson AW, Brown GE (1925) Treatment of Raynaud's disease by lumbar ramisection and ganglionectomy and perivascular sympathetic neurectomy of the common iliacs. *JAMA* 84:1908–1910
 156. Bernard C (1852) Sur les effets de la section de la portion céphalique du grand sympathique. *Comptes Rend Soc Biol* 4:168–170
 157. Gaskell WH (1878) Further researches on the vasomotor nerves of ordinary muscles. *J Physiol* 1:262–302, 426/3–15
 158. Lee BY, Ostrander LE, Thoden WR, Madden JL (1987) Effect of lumbar sympathectomy on muscle blood flow: distribution of perfusion measured by hydrogen clearance in skeletal muscle. *J Rehabil Res Dev* 24:1–8
 159. Wright CJ (1975) Effect of sympathetic inhibition on maximal muscle blood flow in the normal and atherosclerotic human leg. *Can J Surg* 18:431–435
 160. Rex L, Claes G, Drott C, Pegenius G, Elam M (1998) Vasomotor and sudomotor function in the hand after thoracoscopic transection of the sympathetic chain: implications for choice of therapeutic strategy. *Muscle Nerve* 21:1486–1492
 161. Fraunhofer S, Henzler M, Strotzer M, von Sömmogy S (1991) Measuring regional blood flow in evaluating CT-controlled lumbar sympathetic neurolysis in arterial occlusive disease. *Vasa Suppl* 33:160–161
 162. Lombardi D, Milone F, Barone G (1978) Critical considerations on paradoxical gangrene after sympathetic denervation. *Minerva Chir* 33:709–718
 163. Beaconsfield P (1954) A. Effect of exercise on muscle blood flow in normal and sympathectomized limbs. B. Collateral circulation before and after sympathectomy. *Ann Surg* 140:786–795
 164. Prokhorov GG (1988) The blood supply and oxygen balance of the different tissues of the extremities following experimental sympathectomy. *Fiziol Zh Im I M Secherova* 74:1447–1453
 165. Kreutzer W, Schenk WG (1972) Hemodynamic responses to lumbar sympathectomy. An experimental study of changes in blood flow and limb oxygen consumption in the dog with acute or chronic arterial obstruction. *J Cardiovasc Surg (Torino)* 13:532–537
 166. Brothers TE, Wakefield TW, Jacobs LA, Lindenauer SM (1993) Effects of lumbar sympathectomy on canine transcuteaneous oxygen tension. *Surgery* 113:433–437
 167. Hartzell TL, Makhni EC, Sampson C (2009) Long-term results of periarterial sympathectomy. *J Hand Surg Am* 34:1454–1460
 168. Shl Karimov, Berkinov UB, Krotov NF, Ganiev DA (2011) The experience with thoracoscopic sympathectomy for the treatment of distal lesions of limb arteries. *Vestn Ross Akad Med Nauk* 1:18–22
 169. Piazza G, Creager MA (2010) Thromboangiitis obliterans. *Circulation* 121:1858–1861
 170. Ketha SS, Cooper LT (2013) The role of autoimmunity in thromboangiitis obliterans (Buerger's disease). *Ann NY Acad Sci* 1285:15–25
 171. Ishibashi H, Hayakawa N, Yamamoto H, Nishikimi N, Yano T, Nimura Y (1995) Thoracoscopic sympathectomy for Buerger's disease: a report on the successful treatment of four patients. *Surg Today* 25:180–183
 172. Komori K, Kawasaki K, Okazaki J, Eguchi D, Mawatari K, Okadome K, Sadanaga N (1995) Thoracoscopic sympathectomy for Buerger's disease of the upper extremities. *J Vasc Surg* 22:344–345
 173. Lee AD, Agarwal S, Sadhu D (2006) A 7-year experience with thoracoscopic sympathectomy for critical upper limb ischemia. *World J Surg* 30:1644–1647
 174. Wong J, Lam ST, Ong GB (1978) Buerger's disease: a review of 105 patients. *Aust N Z J Surg* 48:382–387
 175. Ates A, Yekeler I, Ceviz M, Erkut B, Pac M, Basoğlu A, Koçak H (2006) One of the most frequent vascular diseases in north-eastern of Turkey: thromboangiitis obliterans or Buerger's disease (experience with 344 cases). *Int J Cardiol* 111:147–153
 176. Haimovici H (1982) Arterial thromboembolism of the upper extremity associated with the thoracic outlet syndrome. *J Cardiovasc Surg (Torino)* 23:214–220
 177. Urschel HC Jr (1993) Dorsal sympathectomy and management of thoracic outlet syndrome with VATS. *Ann Thorac Surg* 56:717–720
 178. Urschel HC Jr, Razzuk MA (1998) Neurovascular compression in the thoracic outlet. Changing management over 50 years. *Ann Surg* 228:609–617
 179. Banis JC Jr, Rich N, Whelan TJ Jr (1977) Ischemia of the upper extremity due to noncardiac emboli. *Am J Surg* 134:131–139
 180. Sakakibara K, Matsumoto M, Motohashi S, Suzuki S (2013) Recurrent refractory arterial thromboembolism in a patient with anti-phospholipid antibody syndrome. *Intern Med* 52:1145–1146
 181. Landry GJ (2013) Current medical and surgical management of Raynaud's syndrome. *J Vasc Surg* 57:1710–1716
 182. Matsumoto Y, Ueyama T, Endo M, Sasaki H, Kasashima F, Abe Y, Kosugi I (2002) Endoscopic thoracic sympathectomy for Raunaud's phenomenon. *J Vasc Surg* 36:57–61
 183. de Trafford JC, Lafferty K, Potter CE, Roberts VC, Cotton LT (1988) An epidemiological survey of Raynaud's phenomenon. *Eur J Vasc Surg* 2:167–170
 184. Coveliers HME, Hoexum F, Nederhoed JH, Wisselink W, Rauwerda JA (2011) Thoracic sympathectomy for digital ischemia: a summary of evidence. *J Vasc Surg* 54:273–277
 185. Dondelinger RF, Kourdzil JC (1987) Percutaneous phenol block of the upper thoracic sympathetic chain with computed tomography guidance. A new technique. *Acta Radiol* 28:511–515
 186. Dorafshar AH, Seitz IA, Zachary L (2009) Reoperative digital sympathectomy in refractory Raynaud's phenomenon. *Plastic Reconstr Surg* 123:36e–38e
 187. Chung L (2007) Therapeutic options for digital ulcers in patients with systemic sclerosis. *J Dtsch Dermatol Ges* 5:460–465

188. Touati Y, Mathieu JP, Houel F, Imbert P (1982) Current indications for superior dorsal sympathectomy. A propos of 84 cases. *J Mal Vasc* 7:187–192
189. Balas P, Tripolitis A (1974) Upper thoracic sympathectomy on systemic scleroderma. *Vasc Surg* 8:234–237
190. Ward WA, van Moore A (1995) Management of finger ulcers in scleroderma. *J Hand Surg Am* 20:868–872
191. Bogoch ER, Gross DK (2005) Surgery of the hand in patients with systemic sclerosis: outcomes and considerations. *J Rheumatol* 32:642–648
192. Maier C, Baron R, Loose R, Schröder D (1994) Endoscopische transthorakale Sympathektomie bei paraneoplastische Raynaud-Syndrom. *Dtsch Med Wschr* 119:1162–1166
193. Larrey DJ (1817) *Mémoires de Chirurgie Militaire et Campagnes*. Tome IV. J Smith, Paris, pp 134–140
194. Bracker MD (1992) Environmental and thermal injury. *Clin Sports Med* 11:419–436
195. Bouwman DL, Morrison SOTR, Lucas CE, Ledgerwood AM (1980) Early sympathetic blockade for frostbite—is it of value? *J Trauma* 20:744–749
196. Shumacker HB Jr, Kilman JW (1964) Sympathectomy in the treatment of frostbite. *Arch Surg* 89:575–584
197. Leriche R (1940) A propos des gelures et de leur traitement immédiat par l’infiltration lombaire. *Presse Med* 48:75
198. Golding MR, Dejong P, Sawyer PN, Hennigar GR, Wesolowski SA (1963) Protection from early and late sequelae of frostbite by regional sympathectomy: mechanism of “cold sensitivity” following frostbite. *Surgery* 53:303–308
199. Murphy JV, Banwell PE, Roberts AHN, McGrouther DA (2000) Frostbite: pathogenesis and treatment. *J Trauma* 48:171–178
200. Taylor MS (1999) Lumbar epidural sympathectomy for frostbite injuries of the feet. *Mil Med* 164:566–567
201. Khan MI, Tariq M, Rehman A, Zafar A, Sheen SN (2008) Efficacy of cervicothoracic sympathectomy versus conservative management in patients suffering from incapacitating Raynaud’s syndrome after frost bite. *J Ayub Med Coll Abbottabad* 20:21–24
202. Chetter IC, Kent PJ, Kester RC (1998) The hand arm vibration syndrome: a review. *Cardiovasc Surg* 6:1–9
203. Goldsmith PC, Molina FA, Bunker CB, Terenghi G, Leslie TA, Fowler CJ, Polak JM, Dowd PM (1994) Cutaneous nerve fibre depletion in vibrating white finger. *J Roy Soc Med* 87:377–381
204. Stoyneva Z, Lyapina M, Tzvetkov D, Vodenicharov E (2003) Current pathophysiological views on vibration-induced Raynaud’s phenomenon. *Cardiovasc Res* 57:615–624
205. Findley FM (1936) Hypertension: its surgical approach. *Calif Western Med* 45:334–340
206. Adson AW, Craig WM, Brown GE (1936) Surgery in its relation to hypertension. *Surg Gynecol Obstet* 62:314–331
207. Smithwick RH, Thompson JE (1953) Splanchnicectomy for essential hypertension: results in 1266 cases. *JAMA* 152:1501–1504
208. Cohen DL, Townsend RR (2011) Renal denervation revisited: promising treatment for resistant hypertension. *J Clin Hypertens* 13:931–932
209. Harris SH (1935) Renal sympathectomy: its scope and limitations. *Proc R Soc Med* 28:1497–1510
210. Krum H, Schlaich M, Whitbourn R, Sobotka PA, Sadowski J, Bartus K, Kapelak B, Walton A, Sievert H, Thambar S, Abraham WT, Esler M (2009) Catheter-based renal sympathetic denervation for resistant hypertension: a multicentre safety and proof-of-principle cohort study. *Lancet* 373:1275–1281
211. Lobo MD, de Belder MA, Cleveland T, Collier D, Dasgupta I, Deanfield J, Kapil V, Knight C, Matson M, Moss J, Paton JF, Poulter N, Simpson I, Williams B, Caulfield MJ (2015) Joint UK societies’ consensus statement on renal denervation for resistant hypertension. *Heart* 101:10–16
212. Rantanen T, Telaranta T (2013) Long-term results of endoscopic sympathetic block using the Lin-Telaranta classification. *Surg Endosc* 27:3860–3864
213. Bragança GM, Lima SO, Pinto Neto AF, Marques LM, Melo EV, Reis FP (2014) Evaluation of anxiety and depression prevalence in patients with primary severe hyperhidrosis. *An Bras Dermatol* 89:230–235
214. Gross KM, Schote AB, Schneider KK, Schulz A, Meyer J (2014) Elevated social stress levels and depressive symptoms in primary hyperhidrosis. *PLoS ONE* 9(3):e92412
215. Callejas MA, Grimalt R, Mejía S, Peri JM (2012) Results of video-assisted sympathectomy for blushing. *Actas Dermosifiliogr* 103:525–531
216. Sciuchetti FJ, Ballabio D, Corti F, Romano F, Benenti C, Costa Angeli M (2006) Endoscopic thoracic sympathectomy by clamping in the treatment of social phobia: the Monza experience. *Minerva Chir* 61:417–420
217. Davidson JR, Foa EB, Connor KM, Churchill LE (2002) Hyperhidrosis in social anxiety disorders. *Prog Neuropsychopharmacol Biol Psychiatry* 26:1327–1331
218. Telaranta T (1998) Treatment of social phobia by endoscopic thoracic sympathectomy. *Eur J Surg* 164(Suppl 580): 27–32
219. Pohjavaara P, Telaranta T, Väisänen E (2001) Endoscopic sympathetic block—new treatment of choice for social phobia? *Ann Chir Gynaecol* 90:177–184
220. Väisänen E, Pohjavaara P, Telaranta T (2001) Sympathetic surgery for psychiatry. Case report. *Ann Chir Gynaecol* 90:187–188
221. Lin CC, Telaranta T (2001) Lin-Telaranta classification: the importance of different procedures for different indications in sympathetic surgery. *Ann Chir Gynaecol* 90:161–166
222. Pohjavaara P, Telaranta T, Väisänen E (2003) The role of the sympathetic nervous system in anxiety: is it possible to relieve anxiety with endoscopic sympathetic block? *Nord J Psychiatry* 57:55–60