

Histology of *Quercus ilex* roots during infection by *Phytophthora cinnamomi*

Miguel Ángel Redondo¹ · Ana Pérez-Sierra² · Paloma Abad-Campos^{3,4} ·
Lilian Torres⁴ · Alejandro Solla⁵ · José Reig-Armiñana⁶ · Francisco García-Breijo^{4,6}

Received: 19 February 2015 / Revised: 6 August 2015 / Accepted: 12 August 2015 / Published online: 3 September 2015
© Springer-Verlag Berlin Heidelberg 2015

Abstract

Key message The speed of infection of *Quercus ilex* by *Phytophthora cinnamomi* is influenced by the method of inoculation used, and structural changes in the host do not differ depending on whether primary or secondary roots are infected.

Abstract This study aimed to elucidate the infection process of the invasive pathogen *Phytophthora cinnamomi* on primary and secondary roots of 2-month-old *Quercus ilex* seedlings. To test if different methods of inoculation lead to different changes in the host caused by the pathogen, the root system of plants was either immersed into a suspension of *P. cinnamomi* zoospores, or placed in direct

contact with agar plugs colonized by *P. cinnamomi* mycelium. Histology of root sections obtained every 24 h for 10 days revealed similar changes in the structure of cells and tissues of the host irrespective of the inoculation method used. However, the immersion method resulted in a delay in the colonization of the host, different aerial symptoms, and the formation of different reproductive structures of the pathogen. Emerging secondary and tertiary roots and sites where secondary or tertiary roots were about to emerge were identified as main entry points. Hyphae in the xylem tissues were more frequently found in secondary than in primary roots, but in both types of roots the phloem was the most important pathway of colonization. For the first time in the interaction between *Q. ilex* and *P. cinnamomi*, transmission electron microscopy was used to describe degradation of the host cell walls, pit penetration and extrahaustorial matrix. Haustoria development during intracellular growth and hyphal aggregations (stromata) caused no damage to the host cell walls indicating hemibiotrophic parasitism.

Communicated by W. Osswald.

Electronic supplementary material The online version of this article (doi:10.1007/s00468-015-1275-3) contains supplementary material, which is available to authorized users.

✉ Miguel Ángel Redondo
miguel.angel.redondo@slu.se

- ¹ Department of Forest Mycology and Plant Pathology, Swedish University of Agricultural Sciences, Uppsala, Sweden
- ² Forest Research, Alice Holt Lodge, Farnham, Surrey, UK
- ³ Instituto Agroforestal Mediterráneo, Universitat Politècnica de València, Valencia, Spain
- ⁴ Departamento de Ecosistemas Agroforestales, Escuela Técnica Superior de Ingeniería Agronómica y del Medio Natural, Universitat Politècnica de València, Valencia, Spain
- ⁵ Ingeniería Forestal y del Medio Natural, Universidad de Extremadura, Plasencia, Spain
- ⁶ Laboratorio de Anatomía Vegetal “Julio Iranzo”, Jardí Botànic de la Universitat de València, Valencia, Spain

Keywords Cell structure · Histological alterations · Histopathology · Microscopy · Pathogenesis · Invasive pathogen

Introduction

Research increasingly points to root health as the key for future sustainability of managed and natural forest ecosystems. Primary roots provide trees with stability and determine the ability of trees to explore deep water. Fine secondary roots, responsible for the majority of absorption of water and nutrients supplied to the crown, are important elements of plant health and are indicators of tree vitality

(Corcobado et al. 2013; Laliberté et al. 2015; McConnell and Balci 2015). Species of the genus *Phytophthora* are among the organisms causing most serious damage to fine roots of trees. These fungus-like eukaryotes which are taxonomically classified as oomycetes cause devastating diseases in an extensive range of tree hosts (Tsao 1990). The oomycete *Phytophthora cinnamomi* is very effective at damaging roots of plants and for this reason causes broad and economically important impacts in forestry, horticulture, and in the nursery industry (Hardham 2005). Based on scientific and economic importance *P. cinnamomi* was recently included in the list of the 10 most relevant oomycetes in molecular plant pathology (Kamoun et al. 2015).

Quercus ilex is the most abundant tree in Spain and one of the commonest tree species in the Mediterranean basin (Ruiz de la Torre 2006). *Quercus ilex* forests are under serious threat mainly because of the lack of regeneration (Pulido et al. 2013), episodic events of drought (Corcobado et al. 2014) and decline caused by several species of *Phytophthora* (Corcobado et al. 2010; de Camilo-Alves et al. 2013; Pérez-Sierra et al. 2013). Tree mortality caused by *P. cinnamomi* is especially relevant in the ‘dehesa’ or ‘montado’ agro-silvo-pastoral systems, and in forests where this pathogen concurs with other abiotic or biotic stressors (de Camilo-Alves et al. 2013; Corcobado et al. 2014; Linaldeddu et al. 2014).

Seed germination is a critical step in natural forest regeneration processes. In soils infested with *P. cinnamomi*, acorns of *Q. ilex* do not germinate (Rodríguez-Molina et al. 2002; Martín-García et al. 2015). If acorns escape from infection, seedlings will be exposed to a second challenge coinciding at a time when conditions are most favourable for *P. cinnamomi* infections, i.e. mean maximum temperatures around 25 °C and soil moisture contents above 30 % vol (Corcobado et al. 2013). At this time of the year, occurring during May in southern Spain, seedlings are about 2 months old, and radicle elongation represents a crucial step for seedling survival (Cubera et al. 2012). Any limitations to the positive geotropic extension of the primary root during germination may have a significant impact on the ability of the tree to tolerate soil drought and compete with herbaceous plants in summer (Cubera et al. 2012). No information about the infection process of the primary roots of *Q. ilex* trees by *P. cinnamomi* at this critical age is available in the literature.

Artificial inoculations of young trees with *Phytophthora* spp. performed under controlled conditions have allowed scientists to describe functional and structural changes of woody plants occurring during the infection process (Blaschke 1994; Brummer et al. 2002; Oh and Hansen 2007; Horta et al. 2010; Rytönen et al. 2013; Dalio et al. 2014; Oßwald et al. 2014). Particularly in holm oak

(*Quercus ilex*), *P. cinnamomi* is able to trigger a generalized dysfunction in water relations of seedlings (Robin et al. 2001). The reduction of stomatal conductance of *Q. ilex* seedlings after infection by *P. cinnamomi* may result from changes in concentrations of hormones in host tissues which would be responsible for stomatal closure (Cahill et al. 1986; Robin et al. 2001). Reduction of predawn water potential of *Q. ilex* leaves could be caused by water stress resulting from root loss (Maurel et al. 2001; Robin et al. 2001). Besides altering water relations, *P. cinnamomi* is able to reduce N and P contents of leaves and modify the performance of *Q. ilex* seedlings after infection (Maurel et al. 2001; Pérez-Sierra et al. 2013; Corcobado et al. 2014).

Histology of secondary roots of 6-month-old *Q. ilex* seedlings revealed alterations of the structure of the tissues when trees were inoculated with *P. cinnamomi* (Ruiz-Gómez et al. 2012, 2015). Changes in the host included cell wall thickening, an increase in the number and size of intercellular spaces in the central cylinder, accumulation of phenolic compounds in the middle lamella and of pectic substances which sealed pits of the xylem vessels, and the formation of lignitubers (Ruiz Gómez et al. 2015). Lignitubers have been recently described as structures in response to *P. cinnamomi* infection able to isolate cells from penetrating hyphae, even though these encasements alone do not prevent colonization of the host by the pathogen (Crone et al. 2013a; Jung et al. 2013). Structures of the pathogen included chlamydospores forming inside the cortex cells, and stromata within the vascular cylinder (Ruiz Gómez et al. 2015). Stromata are hyphae which aggregate randomly inside the host (Willettts 1997) and are thought to be involved in the long-term survival of *P. cinnamomi* in natural environments (Crone et al. 2013a, b; Jung et al. 2013). Haustoria, i.e. specialized hyphae capable of penetrating the host cell and absorbing nutrients from the host cytoplasm, were not confirmed in secondary roots of *Q. ilex* infected by *P. cinnamomi* (Ruiz-Gómez et al. 2012). The authors suggested further electronic microscopy studies to better characterize these structures. In the present work, different microscopy techniques (including electronic microscopy) were used to study primary and secondary roots of *Q. ilex* seedlings after being inoculated with *P. cinnamomi* by two methods: root immersion into a suspension of motile zoospores and direct contact of *P. cinnamomi* mycelium with the roots. It was hypothesized that (1) different methods of *P. cinnamomi* inoculation will lead to different changes in the host caused by the pathogen, (2) different inoculation methods will change the timing of infection, and (3) the anatomic differences between primary and secondary roots will imply differences regarding infection process and structure formation in these two types of roots.

Materials and methods

Plant and pathogen material

All histology assessments were carried out in roots of *Q. ilex* seedlings 2 months after germination (Fig. 1SIa). Acorns were collected directly from the crown of a single tree located in Malpartida de Plasencia, Extremadura, SW Spain (39°58'N 6°5'W, 443 m above sea level). Seeds were surface-disinfested (5 min under a 10 % NaCl solution), germinated in individual free-draining containers containing sterile vermiculite (pH \approx 7) inside a growth chamber at 25 °C (60–70 % relative humidity, 12 h photoperiod) and seedlings were irrigated with sterile water (Horta et al. 2010).

A single A2 mating type isolate of *P. cinnamomi* (PS-694) which has been shown to be highly virulent in *Q. ilex* (Pérez-Sierra et al. 2013) was used. The isolate was preserved in the fungal culture collection maintained at the Instituto Agroforestal Mediterráneo, Universitat Politècnica de València (Spain), in sterile soil extract and oatmeal agar slants (Sigma-Aldrich, Steinheim, Germany) at 14 °C in the dark.

Inoculations

Inoculations consisted of two different methods, (1) immersion of the root system of seedlings into a liquid containing a suspension of *P. cinnamomi* zoospores, and (2) direct contact of the root system of seedlings with agar plugs colonized by mycelium of *P. cinnamomi*. For the first inoculation method, 1 L autoclaved glass flasks containing V8 juice agar (2 g CaCO₃, 200 mL Campbell's V8 juice and 15 g agar in 800 mL distilled water) were used. Hyphal tips were transferred onto the V8 juice agar surface and allowed to grow for 5 days at 25 °C. Then flasks were filled with non-sterile soil extract (Jung et al. 1999) and incubated for 72 h at room temperature for sporangia production. To encourage zoospore release, the flasks were cold-shocked at 4 °C for 30 min. After this period, the liquid suspension was adjusted to approximately 10³ motile zoospores mL⁻¹ and the whole root system of *Q. ilex* seedlings was immersed into the suspension and incubated for 10 days at 25 °C (Fig. 1SIb). For non-infected controls, the root system of *Q. ilex* seedlings was immersed in non-colonized V8 medium flasks filled with non-sterile soil extract. Plants of *Q. ilex* do not show substantial changes in their root system even if exposed to waterlogging during 2.5 months (Corcobado et al. 2014).

For the second inoculation method, seedlings were removed from vermiculite and the root system was washed in sterile water. Inoculations were made by placing 2 cm²

V8 agar plugs containing actively growing mycelium in contact with five secondary roots and the primary root at distances of 0.5–1 cm from their tips (Fig. 1SI c, d) (Horta et al. 2010). For non-infected controls, non-colonized V8 agar plugs were placed in contact with five secondary roots and the primary root. To avoid desiccation, agar plugs and root tips were covered with a cotton ball moistened with sterile water. In total, 20 seedlings per inoculation method were used plus 12 additional non-inoculated seedlings which were used as controls. The whole infection process was maintained at 25 °C, coinciding with the temperature of optimal plant growth and maximum radicle damage of *Q. ilex* by *P. cinnamomi* (Martín-García et al. 2015).

Sampling and microscopy

Root segments 1–1.5 cm in length and 1–2 mm diameter were sampled from primary and secondary roots every 24 h over 10 days. Each day, two plants per inoculation method were harvested and about 20 necrotic roots segments per plant were sampled. Before necrotic lesions were observable, root segments from about 1 cm above the root tips were randomly harvested. For pathogen re-isolation segments from the whole root ball were used (see below). Segments were prepared and sectioned differently depending on the microscopy technique used.

For Light Microscopy (LM), samples were fixed with 5 % glutaraldehyde–paraformaldehyde in 0.02 M Phosphate Buffer Saline (PBS) (pH 7.2) for 12 h at 4 °C (Casano et al. 2011), then rinsed in buffer and later dehydrated in graded series of ethanol and embedded in Spurr's resin (ref. 14300; Electron Microscopy Science, Hatfield, PA, USA) according to manufacturer's instructions (<http://www.emsdiasum.com/microscopy/technical/datasheet/14300.aspx>). Semi-thin sections 2 μ m thick were obtained using a diamond knife DIATOME Histo 45° (DIATOME, Hatfield, PA, USA) and a Nova LKB Bromma Ultramicrotome. Sections were stained with 0.5 % aqueous toluidine blue and observed under an Olympus Provis AX-70 light microscope (Olympus Corp., Japan). Images were obtained through an Infinity 2 CCD (Lumenera Corp., Ottawa, ON, Canada) digital camera and processed by Lumenera AnalySIS software. In total, 150–200 root segments were examined through LM.

For Fluorescence Microscopy (FM), samples were fixed with formaldehyde (40 %)-glacial acetic acid-ethanol (75 %) (5:5:90 vol) and rehydrated by immersing them in distilled water for 20 min. Fresh sections 25–30 μ m thick were obtained with a freezing Leica CM1325 microtome, and then stained with calcofluor white and 10 M KOH (1:1 vol). An Olympus U-ULS 100 HG epifluorescence system with U-MWU (excitation filter 330–385 nm, dichroic mirror 400 nm, barrier filter 420 nm) and U-MWBV

(excitation filter 400–440 nm, dichroic mirror 455 nm, barrier filter 475 nm) cubes was used. Pictures were obtained and processed as indicated above. In total, 40–50 root segments were examined through FM.

For Low-Temperature Scanning Electron Microscopy (LTSEM), samples were fixed and rehydrated as for fluorescence microscopy. Root segments were then frozen with liquid nitrogen, left for sublimation for 15 min and gold covered for 30 s. Root surface observations were carried out by a JEOL JSM 5410 microscope (JEOL USA Inc, Peabody, MA, USA), and pictures were obtained with an Olympus MegaView III camera and processed by Olympus analySIS getIT software (Olympus Corp., Japan). In total, 40–50 root segments were examined through LTSEM.

For Transmission Electron Microscopy (TEM), samples were fixed as indicated for light microscopy, then washed three times with 0.02 M PBS (pH 7.4) for 15 min and fixed again with 2 % osmium tetroxide (OsO₄) in 0.01 M PBS (pH 7.4) for 2 h at room temperature. After washing in buffer, samples were dehydrated and embedded as indicated for light microscopy. Ultrathin sections 80 nm thick were made with a diamond knife (mod. DIATOME ultra 45°; DIATOME, Hatfield, PA, USA), mounted on copper grids of 100 mesh, and then stained with 10 % uranyl acetate and 0.1 % lead citrate using the “Synaptek Grid-Stick Kit” (EMS; <http://www.ems-diasum.com/microscopy/technical/data-sheet/71175.aspx>). Sections were observed at 80 kV under the JEOL JEM-1010 microscope (JEOL USA Inc, Peabody, MA, USA). Images were obtained using an Olympus MegaView III camera and processed by Olympus analySIS getIT software (Olympus Corp., Japan). In total, 40–50 root segments were examined through TEM.

Re-isolation of *Phytophthora cinnamomi*

Re-isolation of *P. cinnamomi* was made from roots inoculated by the two methods. One seedling per day and inoculation method was used. The primary root and two or three secondary roots of each seedling were collected, cut into 6 mm segments, and segments grouped in two clusters of four segments depending if they were proximal or distal from the point of attachment. Over the duration of the experiment about 450 fine root segments were plated in total onto selective CMA-PARBPH agar medium (Pérez-Sierra et al. 2013). After 2–3 days of incubation at 22 °C in the dark, the presence of *P. cinnamomi* in these segments was examined.

Results

Phytophthora cinnamomi was re-isolated with the same success rate (about 50 %) from the primary and secondary roots, independently of the inoculation method used. Re-

isolations of *P. cinnamomi* from the primary root were more successful in distal rather than in proximal segments (70 vs. 34 %, respectively). Re-isolation success from secondary root segments, however, was similar in distal and proximal segments (about 50 %).

Inoculations of seedlings in contact with agar plugs colonized by *P. cinnamomi* produced a faster colonization of plant tissues than inoculations consisting of dipping the roots into a suspension of *P. cinnamomi* zoospores. Figure 1 summarizes the timeline of events observed during infection of *Q. ilex* roots by *P. cinnamomi*. Roots in contact with mycelium showed visible necrosis 24 h after inoculation whereas roots immersed into the zoospore suspension showed necrosis after 3 days post-inoculation (dpi) (Fig. 1). Furthermore, plants inoculated by the root-mycelium contact showed aerial symptoms earlier than plants inoculated by the dip method (5 and 7 dpi, respectively) and symptoms differed depending on the method (leaf necrosis vs wilting of apical leaves, respectively) (Fig. 3SI). Most abundant necrotic lesions were observed in the fine lateral roots and occasionally in the cell elongation zone of the primary root. *P. cinnamomi* hyphae, root necrosis and aerial symptoms were not observed in the control seedlings (Fig. 2SI).

At 1 dpi, LTSEM images from sections of immersed roots revealed zoospore encystment and germination, and penetration of germinative tubes into the rhizodermis (Fig. 2). At 3 dpi, hyphae colonizing the cortical parenchyma of immersed root sections were observed (Fig. 3a, FM), and at 4 dpi haustorial-like structures (Fig. 3b, FM) and cell collapse in the cortex were first detected (Fig. 3c, LM). At 5 dpi, hyphae reached the phloem and vascular cambium, and phloem degradation started (Fig. 3d, LM). Hyphae in the xylem and cambium degradation were observable after 10 days of infection (Fig. 4, LM).

In roots inoculated by direct mycelium contact, fungal penetration started immediately (Fig. 1), and at 1 dpi hyphae were detected in the cortex (Fig. 5, FM). Regions of secondary and tertiary root emergence were major infection entry points for *P. cinnamomi* (Fig. 6b, LM). Fluorescence microscopy allowed observation of numerous hyphae growing inside parenchyma cells of cortex and their intercellular spaces (Fig. 5a, b, FM). In roots inoculated by direct mycelium contact hyphae were first detected in the xylem 3 dpi (Fig. 6a, LM), together with haustorial-like structures observed in the phloem (Fig. 5c, FM). In secondary roots inoculated by direct mycelium contact hyphae approached the primary root via the vascular tissue (Fig. 6b, LM) and cortex. Within the first days after inoculation, damage in root tissues included collapse of cortex cells at 2 dpi but especially obvious was the degradation of the phloem (3 dpi; Fig. 6c, d, LM). Cambium degradation




Fig. 1 Sequence of events of early infection of *Quercus ilex* roots by *Phytophthora cinnamomi* after inoculation **a** through root immersion into a zoospore suspension or **b** by direct contact of roots with

mycelium. Above and below the axes pathogen advance and tissue damages are indicated, respectively




Fig. 2 Low-temperature scanning electron microscopy image showing encystment of *Phytophthora cinnamomi* zoospores (Z) on a secondary *Quercus ilex* root 1 dpi via root immersion into a zoospore suspension. On the left side, a germ tube (GT) allows the pathogen to penetrate the rhizodermis. Bar = 10 μm

started 7 dpi, at the time when the phloem cells were completely distorted (Fig. 6d, LM). Even though *P. cinnamomi* hyphae were observed in the xylem, vessels were rarely altered. Interestingly, hyphae in the xylem tissues of primary roots were very scarce in comparison to the abundant hyphae observed in the xylem of secondary roots.

Over the rhizodermis of primary roots inoculated by the dip method, sporangia were observed 6 dpi (Fig. 7a–c, LM). Zoospores released from these sporangia were also observed 6 dpi. Inoculation of seedlings by root-mycelium contact resulted in the formation of hyphal aggregations after 6 days and chlamydozoospores after 7 days in the cortex of secondary roots (Fig. 7d, LM). Hyphal aggregations were of different shapes but of similar size (Fig. 7e–i, LM). Gametangia or oospores were not observed inside the host.

Ultrathin sections observed in transmission electron microscopy allowed detection of abundant vesicles inside some *P. cinnamomi* hyphae (Fig. 8a, TEM). Electron-dense material was observed surrounding hyphae when contacting host cells (Fig. 8b, TEM), surrounding hyphae




Fig. 3 Fluorescence micrographs **a**, **b** and light microscopy images **c**, **d** of primary and secondary *Quercus ilex* roots inoculated by immersion into a zoospore suspension of *Phytophthora cinnamomi* zoospores. Slides were stained with calcofluor white and toluidine blue, respectively. **a** Cross section of primary root 3 dpi with abundant hyphae (H) growing through the cortical parenchyma (C). Bar = 40 μ m. **b** Cross section of primary root 4 dpi with hypha

(H) penetrating a cortical parenchyma cell (C) through a haustorial-like structure (Ha). Bar = 20 μ m. **c** Cross section of primary root 4 dpi with hypha (H) growing beyond the cork cambium (CC) in the intercellular spaces of the secondary cortex (C), and causing cellular collapses (CoC). Sta = starch. Bar = 20 μ m. **d** Longitudinal section of a primary root 5 dpi with hypha (H) perforating a phloem cell (Ph). Bar = 20 μ m

inside host cells (Fig. 8c, TEM) and a papilla was formed at the opposite part of the wall where the hyphae contacted the cell (Fig. 8d, TEM). Electron-dense material depositions were also observed as a matrix around haustorial-like structures when penetrating the host (Fig. 9a–c, TEM), but no extrahaustorial membrane was observed. Interestingly, very little if any electron-dense material was deposited around the hyphal aggregations, and the cell walls of neighbouring cells surrounding the hyphal aggregations did not show degradation (Fig. 7e–i, LM). On the contrary, neighbouring cells surrounding hyphae had their walls degraded (Fig. 8b, c, TEM), and their plasmalemma was frequently detached (Figs. 8d, 9d, TEM). At the entrance points of hyphae penetration, cell walls were strongly degraded by the pathogen (Figs. 9a, b, 10a, b, TEM). Occasionally, *P. cinnamomi* hyphae seemed to oppress the host cell walls (Fig. 9d, TEM) or even folding them at the entrance points (Fig. 10d, TEM). Occasionally, cell wall invaginations surrounded the penetrating hyphae forming a lignituber-like structure (Fig. 9c, TEM).

Discussion

Within 10 days of infection *P. cinnamomi* was able to colonize all tissues of the primary and the secondary roots of 2-month-old *Q. ilex* seedlings. Except for the xylem, severe alteration of the structure of cells was observed in the cortex, phloem and vascular cambium tissues independent of the inoculation method used. The lignin-based composition of the secondary walls of the xylem may explain why the xylem is so resistant to *P. cinnamomi* and other microorganisms (Hatakka 2005). Damage was particularly noticeable in the less lignified but more nutrient-rich phloem as reported in previous studies (Oh and Hansen 2007).

Based on results, different methods of *P. cinnamomi* inoculation lead to similar changes in *Q. ilex* tissues. However, the method of inoculation used influenced the speed of the infection process and the type of aerial symptoms and asexual spores produced. It should be acknowledged that differences in the maintenance of plants




Fig. 4 Light microscopy sections of *Quercus ilex* primary roots inoculated by immersion into a suspension of *Phytophthora cinnamomi* zoospores. Slides were stained with *toluidine blue*. **a** Cross section 10 dpi with hyphae (H) growing through the vascular cambium and phloem (Ph), causing cellular collapse and cell degradation in the vascular cambium (VC*). Xy = xylem. Bar = 20 μm . **b** Longitudinal section 10 dpi with a hypha (H) growing through the xylem vessels (Xy). No damage in the xylem was observed. Bar = 20 μm

between the two inoculation methods used might have influenced the results. Nevertheless, previous research showed that depending on the inoculation method, the symptoms shown by seedlings and their final mortality differed significantly (Hansen et al. 2005; Haque and Diez 2012; Rytönen et al. 2013). Each host may show a different infection process (Cahill et al. 1989) depending on the species of *Phytophthora* and the inoculation method used, as indicated by the significant *Phytophthora* spp. \times inoculation method interaction reported by Haque and Diez (2012). Using the dip method with a suspension of *P. cinnamomi* chlamydospores, at 3 dpi all tissues of secondary *Q. ilex* roots were colonized, but first aerial symptoms consisted of leaf discoloration by 14 dpi (Ruiz Gómez et al. 2015). Root dip into a zoospore suspension mimics the process that occurs in nature, probably better than the mycelium-root contact method. In a leaf inoculation study, higher infection rates and more real symptoms occurred using a zoospore suspension as an inoculum




Fig. 5 Fluorescence micrographs of *Quercus ilex* secondary roots inoculated by direct contact of mycelium of *Phytophthora cinnamomi*. Slides were stained with *calcofluor white*. **a** Longitudinal section 1 dpi showing hyphae (H) growing through cortical cells (C). Rh = rhizodermis. Bar = 40 μm . **b** Cross section 3 dpi showing hyphae (H) growing inside cortical parenchyma cells (C) and through intercellular spaces. Bar = 20 μm . **c** Cross section 3 dpi showing abundant hyphae (H) growing in the phloem (Ph). A haustorial-like structure (Ha) was also observable. En = endodermis. C = cortical parenchyma. Bar = 20 μm

source compared to mycelial agar plugs (Hansen et al. 2005).

Infection started immediately by direct hyphal penetration through the epidermal cells, and occurred after 1 day if zoospores were used. This conformed to results of Rytönen et al. (2013) but contrasted to other studies reporting




Fig. 6 Light microscopy images of *Quercus ilex* secondary roots inoculated by direct contact of mycelium of *Phytophthora cinnamomi*. Slides were stained with *toluidine blue*. **a** Longitudinal section 3 dpi with hyphae (H) growing through the xylem (Xy) and no cell damage observed. *Bar* = 40 μm . **b** Detail of a longitudinal section of a lateral root branching point (general view inserted in the upper right) with hyphae (H) growing towards the primary root

through the vascular tissue. *Bar* = 40 μm . **c** Cross section 3 dpi with many hyphae (H) located inside phloem cells (Ph) causing its degradation. Some haustoria-like structures can be observed (Ha). No hyphae were visible in the cortex (C) and xylem (Xy). *En* = endodermis. *Bar* = 20 μm . **d** Cross section 7 dpi with abundant hyphae (H) growing through the vascular tissue and endodermis. Completely degraded phloem (Ph) is observable. *Bar* = 20 μm

immediate penetration of *P. cinnamomi* if zoospores were the source of inoculum (Cahill et al. 1989; Hardham 2005). By immersion of *Q. ilex* roots for 10 min into a chlamydo-spore suspension (25×10^3 IU mL⁻¹), infection of secondary roots was also successful 1 dpi (Ruiz-Gómez et al. 2012, 2015). This variation might be due to differences in typology of zoospores suspension and inoculum localization between different studies. In the present work, and in the work performed by Rytönen et al. (2013), a non-sterile zoospore suspension was used, and the possible presence of bacteria in this suspension during the infection process might affect zoospores survival and infectiveness. In contrast, other work using axenic solutions of zoospores as a source of inoculum reported immediate penetration (Cahill et al. 1989; Hardham 2005). During root segments selection at early stages of infection some of the penetration spots might have been missed because necrotic areas were absent and sample harvesting was performed at random along the root. To avoid the uncertainty of missing

infection sites using zoospore inoculum Cahill et al. (1989) and Hardham (2005) applied the zoospores in well-localized areas, which were later used in the histological studies. No appressoria, reported for other *Phytophthora* species (Judelson and Blanco 2005; Oh and Hansen 2007), were observed here. Two main infection courts were identified: the emerging secondary or tertiary roots and sites where the developing secondary or tertiary roots were about to emerge. This result could be explained by the fragmentation and discontinuity of tissues in the region immediately surrounding the site of root emergence (Fahn 1990; O'Gara et al. 2015).

Once the epidermis was penetrated, the majority of hyphae observed in the cortical tissue were growing in the intercellular spaces. At this time, extracellular enzymes of *Phytophthora* may be involved in the degradation of the host cell walls including the middle lamella (Brummer et al. 2002; Hardham 2005). Haustoria developed when intracellular growth occurred, preceded by the




Fig. 7 Light microscopy sections of *Phytophthora cinnamomi*-infected secondary *Quercus ilex* roots stained with *toluidine blue*, in which asexual reproductive and survival structures were observed. **a** Longitudinal section of immersion-inoculated secondary root section, 6 dpi, with several sporangia (Sp) developing on rhizodermis (Rh). Some zoospores (Z) have been released. C = cortex parenchyma. Bar = 40 μm . **b, c** Detail of irregular **b** and common shaped

c sporangia developing on the root surface. Bar = 20 μm . **d** Chlamydospores (Chl) developing in the intercellular spaces of cortical parenchyma of mycelia contact inoculated secondary root 7 dpi, cross section. Bar = 20 μm . **e, f, g, h, i** Hyphal aggregations (stromata) observed 6–10 days after inoculation by the mycelia-root contact method. Bar = 20 μm

accumulation of electron-dense material in the plant cell wall at the site of future invasion, as observed in other studies (Brummer et al. 2002). Hyphae surrounding cortical cells probably caused the release of phenol/tannin-like compounds (Cahill et al. 1989; Horta et al. 2010) or toxins which might be the cause of the necrosis observed on the primary and secondary roots starting from 3 and 1 dpi, respectively. The increase of the intercellular spaces of the cortex tissue and the increase in cell damage in the secondary cortical region of the primary roots in comparison

to those in control plants may explain the detachment of the bark from roots, as observed by the authors of this work.

Irrespective of the inoculation method and the type of roots used, 2 days after cortex colonization *P. cinnamomi* hyphae were detected in the phloem. The images confirm that the phloem is the most important pathway for the vertical colonization of *P. cinnamomi* within the plant. Evidence of this movement is shown in Fig. 6b, c where numerous hyphae were detected in the phloem but not in




Fig. 8 Transmission electron microscopy examinations of *Phytophthora cinnamomi*-infected primary and secondary *Quercus ilex* roots. **a** Hypha (H) with abundant vesicles (V). Bar = 400 nm. **b** Hyphae growing in the intercellular spaces (H) and electron-dense material (EdM) deposition surrounding the hyphae. The cell wall close to the hyphae is degraded (CW*). Xy = xylem. Bar = 2 μ m. **c** Hyphae (H) inside a xylem vessel (Xy) with electron-dense material (EdM)

deposition surrounding the hyphae. The cell wall in contact with the hyphae is slightly degraded (CW*). Bar = 2 μ m. **d** Hyphae growing in the intercellular space (H), and electron-dense material as a papilla (P) is deposited in the opposite part of the cell wall. Neighbouring cell plasmalemma is detached (PI*). CW = cell wall. Sta = starch. Bar = 2 μ m

the cortex. In consequence, *P. cinnamomi* is able to easily pass through sieve plates connecting sieve elements. Sieve occlusion, usually occurring by callose deposition around the sieve pores within minutes after damage (Mullendore et al. 2010), was not observed to stop phloem colonization by *P. cinnamomi* (Fig. 3d). Hyphae colonizing the xylem of primary roots were rarely observed, in agreement with previous studies (Brummer et al. 2002; Oh and Hansen 2007), but it was frequent in secondary roots. With the exception of this difference, the infection process did not differ between primary and secondary roots.

Complete destruction of the phloem continued 2 or 3 days after the hyphae first reached this tissue, which confirms observations of infection by other *Phytophthora* species (Oh and Hansen 2007), and of other hosts infected by *P. cinnamomi* (Cahill et al. 1989). Lignitubers were observed here as a response to *P. cinnamomi* infection and

are known to form as a general plant response to isolate cells from penetrating hyphae even though these encasements alone do not completely prevent infection (Crone et al. 2013a; Jung et al. 2013). Tyloses were not detected in accordance to Ruiz Gómez et al. (2015) but in contrast to other studies (Cahill et al. 1989; Blaschke 1994). If occurring, tyloses in *Q. ilex* may form in response to *P. cinnamomi* after 10 days of infection.

Five days after penetration, *P. cinnamomi* produced sporangia in the rhizodermis (Fig. 7). On other hosts, sporangia production by *P. cinnamomi* was reported on infected cortex tissue 48–72 h after inoculation (Cahill et al. 1989; Hardham 2005). In secondary roots of 6-month-old *Q. ilex* seedlings inoculated by *P. cinnamomi*, sporangia formation was not reported, but mature chlamydo spores were observed in the cortex 14 dpi (Ruiz-Gómez et al. 2012, 2015). Differences in timing of asexual




Fig. 9 Transmission electron microscopy examinations of *Phytophthora cinnamomi*-infected *Quercus ilex* roots. **a** Hypha in the intercellular space (H) penetrating a host cell. At the entrance point the cell wall is degraded (CW*) and extra-haustorial matrix of electron-dense material was deposited (EdM) around the hypha (H). Bar = 800 nm. **b** Hypha (H) moving from a protoxylem cell to another (H). Bar = 2 μ m. **c** Hypha growing in the intercellular space (H) and penetrating a host cell (H). Cell wall invagination as a

haustorial encasement or lignituber-like structure (CWi). At the final part of the invagination, the hypha bypass the cell wall and electron-dense matrix is deposited (EdM). Bar = 2 μ m. **d** Three hyphae growing inside a cell (H) and surrounded by abundant electron-dense material (EdM). The one in the left seems to push the wall (PCW) of the neighbouring cell. The plasmalemma of the host cells is detached (PI*). Bar = 2 μ m

spore formation between studies may occur due to differences in host species, type of root, and method of inoculation used, and probably the temperature, age of tissues and concentration of infective units, all of which will also contribute to make results not comparable. Usually 24 h after formation (Thomas Jung, personal communication) sporangia mature and their papillae dissolve allowing release of motile zoospores ready to start new cycles of infection. This rapid cycle, occurring here within 6 days may explain why *P. cinnamomi* is so successful in causing sudden death of adult *Q. ilex* trees during early summer and early autumn in southern Spain and Portugal (de Camilo-Alves et al. 2013) and impedes natural regeneration. Numerous cycles of spore formation, rapid infection and fine roots destruction occurring before symptom expression

explain the difficulties associated with disease management.

Infection process of *Q. ilex* by *P. cinnamomi* at the ultrastructural level has never been described. In the current study, transmission electron microscopy images showed for the first time in the *Q. ilex*–*P. cinnamomi* interaction, plasmalemma detachment and wall degradation of host cells, hyphae penetration through pits, and depositions of electron-dense material as papillae attached to the cell wall or as an extra-haustorial matrix surrounding the penetrating hyphae. Localized reinforcement of the cell wall through deposition of papillae at sites of pathogen detection appears to be a common component of the pattern-triggered immunity of plants (Underwood 2012). The nature of this dense material is unknown (Oßwald et al.




Fig. 10 Transmission electron microscopy examinations of *Phytophthora cinnamomi*-infected *Quercus ilex* roots. **a** Hypha (H) penetrating a host cell and haustorial-like structure (Ha) with electron-dense material depositions (EdM). Electron-dense material is also deposited as a papilla in the inner side of cell wall (P). **b** Selective enlargement from (a), showing the entrance point in which the cell wall is degraded (CW*), and electron-dense material is

deposited (EdM). Mitochondrion (M). Hyphae (H), Bar = 200 nm. **c** Hypha (H) penetrating a host cell through a pit. Edm = electron-dense material. Bar = 2 μm. **d** Selective enlargement from (c), showing the entrance point in which the pit wall appears to be mechanically deformed by the hyphae (H) and folded (PW). These folds expose the pit membrane (PM). Mitochondrion (M). Bar = 200 nm

2014) though some authors suggested it is formed by phenolic-like compounds and/or callose (Hardham 2001; Brummer et al. 2002; Horta et al. 2010; Underwood 2012). During later stages of infection, oomycete haustoria are often encased by a double-layered callose-containing membrane structure (Lu et al. 2012), not observed here. The vesicles observed inside *P. cinnamomi* hyphae (Fig. 8a) were reported to be involved in the release of cell wall-degrading enzymes by the pathogen (Hardham 2005).

Besides hyphae proliferating through the host tissue, hyphal aggregations varied in density and appearance. These formations called stromata or ‘singular stroma’ were recently reported in secondary roots of *Q. ilex* (Ruiz Gómez et al. 2015) but are novel in primary roots of this tree species. Typically stromata were confined to one cell, but it was also observed that emerging hyphae penetrated new root cells in close proximity to form more stromata

(Fig. 7h, i) in accordance to what was observed in herbaceous hosts (Crone et al. 2013a). Stromata have been described as morphologically variable formations in regard to size, compactness, and degree of differentiation (Willetts 1997). Functionally, it is suggested that due to the hyphal density of the stromata their capacity to store nutrients acquired from the host material is significant, resulting in the high production of mycelium and spores when conditions are favourable for germination (Willetts 1997). In consequence, stromata act as survival propagules (Willetts 1997; Crone et al. 2013a, b; Jung et al. 2013), and this explains why *P. cinnamomi* is such a persistent pathogen under *Q. ilex* forests. The fact that neighbouring host cells close to stromata did not show damage, necrosis or thickening of walls points to the hypotheses that through these structures the pathogen (1) impedes recognition by the host, (2) suppresses the host’s defence gene expression, and

(3) accumulates nutrients from the host to be invested into the formation of chlamydo spores and oospores when conditions become stressful, therefore successfully assuring pathogen persistence.

Haustorium-like structures were not observed in other interactions (Oh and Hansen 2007; Rytönen et al. 2013), indicating that not all *Phytophthora* species are hemibiotrophic. The presence of haustoria, i.e. structures involving feeding relationships between the host and the pathogen (Judelson and Blanco 2005), and the formation of numerous chlamydo spores indicate that *P. cinnamomi* may temporarily behave as a biotrophic pathogen (Crone et al. 2013a; Ruiz Gómez et al. 2015). It was recently postulated that a necrotrophic mode of *P. cinnamomi* growth would not provide enough nutrients to produce the large numbers of chlamydo spores (Crone et al. 2013a). The biotrophic mode of growth with nutrient acquisition aided through haustoria, followed by formation of dense masses of hyphae in stromata may be a prerequisite to the formation of the high numbers of chlamydo spores (Crone et al. 2013a).

Conclusions

Artificial inoculations may not mimic the real interactions between hosts and pathogens in nature. Nevertheless, inoculation methods used under controlled environments may avoid interactions with external variables and make results comparable between experiments. Irrespective of the inoculation method used, similar structural changes in *Q. ilex* roots were observed, although dipping the root system of plants into a zoospore inoculum suspension caused a more delayed colonization, different aboveground symptoms, and the formation of different reproductive structures of the pathogen in comparison to placing the root system of plants in contact to mycelium. This study reports no differences in the initial infection of primary and secondary roots, and provides valuable data of how *P. cinnamomi* colonizes *Q. ilex* at an ontogenic stage critical for tree regeneration. The pathogen rapidly reaches the phloem, which is the most important pathway of vertical colonization of *P. cinnamomi* within the plant. *Quercus ilex* trees exhibited a variety of directed responses to the pathogen, including general wall thickening, cell collapse, encasement of hyphae, and deposition of electron-dense materials around hyphae. This is the first time transmission electron microscopy has been used to describe cell structures occurring during infection of *Q. ilex* by *P. cinnamomi*. This work will improve further investigation of the pathogenic pathways and survival of *P. cinnamomi* within host tissues, as well as detailed plant pathogenic interactions within the roots. Moreover, results about the

timing of production of propagules in fine roots and the time lag between infection and appearance of visible symptoms will be useful for further epidemiological studies.

Author contribution statement Miguel Ángel Redondo: Experiment design, *P. cinnamomi* inoculation, sampling and histological procedures, elaboration of microscopy images, interpretation of results, and manuscript writing. Ana Pérez-Sierra: Experiment design, *Q. ilex* seedlings growing, *P. cinnamomi* inoculation and sampling. Paloma Abad-Campos: Experiment design, *P. cinnamomi* inoculation and sampling. Lilian Torres: *P. cinnamomi* inoculation, sampling and histological procedures. Elaboration of microscopy images, and interpretation of results. Alejandro Solla Acorn: recollection, interpretation of results and manuscript writing. José Reig-Armiñana: Supervision of histological procedures, microscopy images elaboration, and interpretation of results. Francisco García-Breijo: Supervision of histological procedures, supervision of microscopy images elaboration, interpretation of results and manuscript writing.

Acknowledgments This research was financially supported by the Project AGL2011-30438, by the Vicerrectorado de Investigación from the Polytechnic University of Valencia, and by the “Julio Irazo” laboratory from the Botanic Garden of Valencia. The authors deeply appreciate the help of the staff from both institutions, particularly the valuable contribution of Nuria Cebrián Gómez. Additionally, the staff from the microscopy sections from Polytechnic University of Valencia and University of Valencia have provided us with valuable help. We are grateful to the two anonymous reviewers for the valuable comments in an earlier version of this manuscript.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

- Blaschke H (1994) Decline symptoms on roots of *Quercus robur*. Eur J For Pathol 24:386–398. doi:10.1111/j.1439-0329.1994.tb00832.x
- Brummer M, Arend M, Fromm J et al (2002) Ultrastructural changes and immunocytochemical localization of the elicitor quercinin in *Quercus robur* L. roots infected with *Phytophthora quercina*. Physiol Mol Plant Pathol 61:109–120. doi:10.1006/pmpp.2002.0419
- Cahill DM, Weste GM, Grant BR (1986) Changes in cytokinin concentrations in xylem extrudate following infection of *Eucalyptus marginata* Donn ex Sm with *Phytophthora cinnamomi* Rands. Plant Physiol 81:1103–1109
- Cahill D, Legge B, Weste GM (1989) Cellular and histological changes induced by *Phytophthora cinnamomi* in a group of plant species ranging from fully susceptible to fully resistant. Phytopathology 79:417–424. doi:10.1094/Phyto-79-417
- Casano LM, del Campo EM, García-Breijo FJ et al (2011) Two Trebouxia algae with different physiological performances are ever-present in lichen thalli of Ramalina farinacea. Coexistence versus competition? Environ Microbiol 13:806–818. doi:10.1111/j.1462-2920.2010.02386.x
- Corcobado T, Cubera E, Pérez-Sierra A et al (2010) First report of *Phytophthora gonapodyides* involved in the decline of *Quercus ilex* in xeric conditions in Spain. New Dis Rep 22:33

- Corcobado T, Cubera E, Moreno G, Solla A (2013) *Quercus ilex* forests are influenced by annual variations in water table, soil water deficit and fine root loss caused by *Phytophthora cinnamomi*. *Agric For Meteorol* 169:92–99. doi:10.1016/j.agrformet.2012.09.017
- Corcobado T, Cubera E, Juárez E et al (2014) Drought events determine performance of *Quercus ilex* seedlings and increase their susceptibility to *Phytophthora cinnamomi*. *Agric For Meteorol* 192–193:1–8. doi:10.1016/j.agrformet.2014.02.007
- Crone M, McComb JA, O'Brien PA, Hardy GESJ (2013a) Survival of *Phytophthora cinnamomi* as oospores, stromata, and thick-walled chlamydospores in roots of symptomatic and asymptomatic annual and herbaceous perennial plant species. *Fungal Biol* 117:112–123. doi:10.1016/j.funbio.2012.12.004
- Crone M, McComb JA, O'Brien PA, Hardy GESJ (2013b) Assessment of Australian native annual/herbaceous perennial plant species as asymptomatic or symptomatic hosts of *Phytophthora cinnamomi* under controlled conditions. *For Pathol* 43:245–251. doi:10.1111/efp.12027
- Cubera E, Moreno G, Solla A, Madeira M (2012) Root system of *Quercus suber* L. seedlings in response to herbaceous competition and different watering and fertilisation regimes. *Agrofor Syst* 85:205–214. doi:10.1007/s10457-012-9492-x
- Dalio RJD, Fleischmann F, Humez M, Osswald W (2014) Phosphite protects *Fagus sylvatica* seedlings towards *Phytophthora plurivora* via local toxicity, priming and facilitation of pathogen recognition. *PLoS One* 9:e87860. doi:10.1371/journal.pone.0087860
- de Camilo-Alves C, de Sampaio P, da Clara MIE, de Almeida Ribeiro NMC (2013) Decline of mediterranean oak trees and its association with *Phytophthora cinnamomi*: a review. *Eur J For Res* 132:411–432. doi:10.1007/s10342-013-0688-z
- Fahn A (1990) Plant anatomy. Pergamon Press, Oxford
- Hansen EM, Parke JL, Sutton W (2005) Susceptibility of Oregon forest trees and shrubs to *Phytophthora ramorum*: a comparison of artificial inoculation and natural infection. *Plant Dis* 89:63–70. doi:10.1094/PD-89-0063
- Haque MMU, Diez JJ (2012) Susceptibility of common alder (*Alnus glutinosa*) seeds and seedlings to *Phytophthora alni* and other *Phytophthora* species. *For Syst* 21:313–322. doi:10.5424/fs/2012212-02267
- Hardham AR (2001) The cell biology behind *Phytophthora* pathogenicity. *Australas Plant Pathol* 30:91–98. doi:10.1071/AP01006
- Hardham AR (2005) *Phytophthora cinnamomi*. *Mol Plant Pathol* 6:589–604. doi:10.1111/j.1364-3703.2005.00308.x
- Hatakka A (2005) Biodegradation of lignin. *Biopolym Online*. doi:10.1002/3527600035.bpol1005
- Horta M, Caetano P, Medeira C et al (2010) Involvement of the β -cinnamomin elicitor in infection and colonisation of cork oak roots by *Phytophthora cinnamomi*. *Eur J Plant Pathol* 127:427–436. doi:10.1007/s10658-010-9609-x
- Judelson HS, Blanco FA (2005) The spores of *Phytophthora*: weapons of the plant destroyer. *Nat Rev Microbiol* 3:47–58. doi:10.1038/nrmicro1064
- Jung T, Cooke DEL, Blaschke H et al (1999) *Phytophthora quercina* sp. nov., causing root rot of European oaks. *Mycol Res* 103:785–798. doi:10.1017/S0953756298007734
- Jung T, Colquhoun IJ, Hardy GESJ (2013) New insights into the survival strategy of the invasive soilborne pathogen *Phytophthora cinnamomi* in different natural ecosystems in Western Australia. *For Pathol* 43:266–288. doi:10.1111/efp.12025
- Kamoun S, Furzer O, Jones JDG et al (2015) The top 10 oomycete pathogens in molecular plant pathology. *Mol Plant Pathol* 16:413–434. doi:10.1111/mpp.12190
- Laliberté E, Lambers H, Burgess TI, Wright SJ (2015) Phosphorus limitation, soil-borne pathogens and the coexistence of plant species in hyperdiverse forests and shrublands. *N Phytol* 206:507–521. doi:10.1111/nph.13203
- Linaldeddu BT, Scanu B, Maddau L, Franceschini A (2014) *Diplodia corticola* and *Phytophthora cinnamomi*: the main pathogens involved in holm oak decline on Caprera Island (Italy). *For Pathol* 44:191–200. doi:10.1111/efp.12081
- Lu Y-J, Schornack S, Spallek T et al (2012) Patterns of plant subcellular responses to successful oomycete infections reveal differences in host cell reprogramming and endocytic trafficking. *Cell Microbiol* 14:682–697. doi:10.1111/j.1462-5822.2012.01751.x
- Martín-García J, Solla A, Corcobado T et al (2015) Influence of temperature on germination of *Quercus ilex* in *Phytophthora cinnamomi*, *P. gonapodyides*, *P. quercina* and *P. psychrophila* infested soils. *For Pathol* 45:215–223. doi:10.1111/efp.12159
- Maurel M, Robin C, Capron G, Desprez-Loustau M-L (2001) Effects of root damage associated with *Phytophthora cinnamomi* on water relations, biomass accumulation, mineral nutrition and vulnerability to water deficit of five oak and chestnut species. *For Pathol* 31:353–369. doi:10.1046/j.1439-0329.2001.00258.x
- McConnell ME, Balci Y (2015) Fine root dynamics of oak saplings in response to *Phytophthora cinnamomi* infection under different temperatures and durations. *For Pathol* 45:155–164. doi:10.1111/efp.12150
- Mullendore DL, Windt CW, As HV, Knoblauch M (2010) Sieve tube geometry in relation to phloem flow. *Plant Cell Online* 22:579–593. doi:10.1105/tpc.109.070094
- O'Gara E, Howard K, McComb J et al (2015) Penetration of suberized periderm of a woody host by *Phytophthora cinnamomi*. *Plant Pathol* 64:207–215. doi:10.1111/ppa.12244
- Oh E, Hansen EM (2007) Histopathology of infection and colonization of susceptible and resistant Port-Orford-Cedar by *Phytophthora lateralis*. *Phytopathology* 97:684–693. doi:10.1094/PHTO-97-6-0684
- Oßwald W, Fleischmann F, Rigling D et al (2014) Strategies of attack and defence in woody plant–*Phytophthora* interactions. *For Pathol* 44:169–190. doi:10.1111/efp.12096
- Pérez-Sierra A, López-García C, León M et al (2013) Previously unrecorded low-temperature *Phytophthora* species associated with *Quercus* decline in a Mediterranean forest in eastern Spain. *For Pathol* 43:331–339. doi:10.1111/efp.12037
- Pulido F, McCreary D, Cañellas I et al (2013) Oak regeneration: ecological dynamics and restoration techniques. In: Campos P, Huntsinger L, Oviedo Pro JL (eds) Mediterranean oak woodland working landscapes. Springer, Dordrecht, pp 123–144
- Robin C, Capron G, Desprez-Loustau ML (2001) Root infection by *Phytophthora cinnamomi* in seedlings of three oak species. *Plant Pathol* 50:708–716. doi:10.1046/j.1365-3059.2001.00643.x
- Rodríguez-Molina MC, Torres-Vila LM, Blanco-Santos A et al (2002) Viability of holm and cork oak seedlings from acorns sown in soils naturally infected with *Phytophthora cinnamomi*. *For Pathol* 32:365–372. doi:10.1046/j.1439-0329.2002.00297.x
- Ruiz de la Torre J (2006) Flora mayor. Organismo Autónomo Parques Nacionales, Dirección General para la Biodiversidad, Madrid
- Ruiz Gómez FJ, Navarro-Cerrillo RM, Sánchez-Cuesta R, Pérez-de-Luque A (2015) Histopathology of infection and colonization of *Quercus ilex* fine roots by *Phytophthora cinnamomi*. *Plant Pathol* 64:605–616. doi:10.1111/ppa.12310
- Ruiz-Gómez FJ, Sánchez-Cuesta R, Navarro-Cerrillo RM, Pérez-de-Luque A (2012) A method to quantify infection and colonization of holm oak (*Quercus ilex*) roots by *Phytophthora cinnamomi*. *Plant Methods* 8:39. doi:10.1186/1746-4811-8-39
- Rytönen A, Lilja A, Werres S et al (2013) Infectivity, survival and pathology of Finnish strains of *Phytophthora plurivora* and *Ph. pini* in Norway spruce. *Scand J For Res* 28:307–318. doi:10.1080/02827581.2012.756926

- Tsao PH (1990) Why many phytophthora root rots and crown rots of tree and horticultural crops remain undetected? EPPO Bull 20:11–17. doi:[10.1111/j.1365-2338.1990.tb01174.x](https://doi.org/10.1111/j.1365-2338.1990.tb01174.x)
- Underwood W (2012) The plant cell wall: a dynamic barrier against pathogen invasion. Front Plant Sci 3:85. doi:[10.3389/fpls.2012.00085](https://doi.org/10.3389/fpls.2012.00085)
- Willets HJ (1997) Morphology, development and evolution of stromata/sclerotia and macroconidia of the Sclerotiniaceae. Mycol Res 101:939–952. doi:[10.1017/S0953756297003559](https://doi.org/10.1017/S0953756297003559)