

STRUCTURAL RESULTS FOR PLANAR SETS
WITH MANY SIMILAR SUBSETS

BERNARDO M. ÁBREGO, GYÖRGY ELEKES*,
SILVIA FERNÁNDEZ-MERCHANT

Received November 23, 2000

Consider a k -element subset P of the plane. It is known that the maximum number of sets similar to P that can be found among n points in the plane is $\Theta(n^2)$ if and only if the cross ratio of any quadruplet of points in P is algebraic [3], [9].

In this paper we study the structure of the extremal n -sets A which have cn^2 similar copies of P . As our main result we prove the existence of large lattice-like structures in such sets A . In particular we prove that, for n large enough, A must contain m points in a line forming an arithmetic progression, or $m \times m$ lattices, when P is not cocyclic or collinear. On the other hand we show that for cocyclic or collinear sets P , there are n -element sets A with $c_P n^2$ copies of P and without $k \times k$ lattice subsets.

1. Introduction

We identify the plane with the field of complex numbers \mathbb{C} . For $A, B \subseteq \mathbb{C}$ and $z, w \in \mathbb{C}$ we denote $zA = \{za : a \in A\}$, and $A + w = \{a + w : a \in A\}$. Also, we say that A and B are similar, and we write $A \sim B$, if there are complex numbers w and $z \neq 0$ such that $B = zA + w$ (i.e., we do not allow reflections).

For every pattern set P and finite set $A \subset \mathbb{C}$ we define

$$S_P(A) = |\{X \subseteq A : X \sim P\}|.$$

Mathematics Subject Classification (2000): 52C10

* Supported by Hungarian National Foundation for Scientific Research (OTKA), Grants No. T 42750 and T 47056.

It is a natural question, posed many times by Erdős and Purdy [4]–[6], to determine or estimate the following function

$$(1) \quad S_P(n) = \max_{|A|=n} S_P(A);$$

i.e., the maximum number of subsets similar to a given pattern P that can be found among n points in the plane. In [3] Erdős and the second author started the study of this function. They noticed that $S_P(n) \leq n(n-1)$ (a set similar to P is determined by the location of any pair of reference points). They also proved that $S_P(n) \geq cn^{2-b\log^{-a}n}$ for some constants $a, b, c > 0$ depending only on P , and $S_P(n) = \Theta(n^2)$ when P is an algebraic set or $|P|=3$. Later, Laczkovich and Ruzsa [9] proved that $S_P(n) = \Theta(n^2)$ if and only if the cross ratio among every quadruplet of distinct elements in P is algebraic. Recall that the cross ratio of a quadruplet (a, b, c, d) of different complex numbers is given by

$$(a; b; c; d) = \frac{(c-a)(d-b)}{(d-a)(c-b)}.$$

Even though this settles the order of magnitude for a big class of sets P , it is not known if $\lim_{n \rightarrow \infty} S_P(n)/n^2$ exists for any non-trivial sets P , e.g., triangles.

In this article we take a qualitative approach to the problem. For a pattern set P satisfying $S_P(n) = \Theta(n^2)$, we say that A is a (P, c) -rich set if $S_P(A) \geq c|A|^2$. Our objective is to find structural properties about (P, c) -rich sets. To accomplish this, we introduce in Section 2 the notion of a $G_P(m)$ set. Then we state our main result, Theorem 1, which asserts that every sufficiently large (P, c) -rich set must contain a $G_P(m)$ set. The importance of this result is seen through its corollaries. We prove, for example, the existence of large arithmetic progressions or lattice structures (when P is not cocyclic or collinear) among (P, c) -rich sets. Other corollaries include the fact that the related function $S_P^{gen}(n)$, where the maximum in (1) is restricted to sets in general position, satisfies $S_P^{gen}(n) = o(n^2)$; and the existence of regular triangle lattices among sufficiently large extremal sets of the function $S_T(n)$, where T is an equilateral triangle.

For the sake of continuity, we postpone the proof of Theorem 1 to Section 4, where we introduce the reader to some necessary Number Theory results by Freiman [7], Balog, Szemerédi [2], Laczkovich, Ruzsa [9], and Fürstenberg, Katznelson [8].

Finally, in Section 3, we look closely to the case when P is a cocyclic or a collinear set. For this case we construct (P, c) -rich sets contained in at most

$|P|$ lines. From this construction we characterize cocyclic or collinear sets as the only sets P for which there are arbitrarily large (P, c) -rich sets avoiding $|P| \times |P|$ -lattice structures.

2. P -generated sets and the main result

We start this section by introducing the concept of a $G_P(m)$ set which will be essential for our main result. From now on $P = \{p_1, p_2, \dots, p_k\}$ will be the pattern set, and for every positive integer m , $I_m = \{j \in \mathbb{Z} : |j| \leq m\}$ and $[m] = \{j \in \mathbb{Z} : 1 \leq j \leq m\}$.

We say that A is a P -generated set of parameter m , for short $G_P(m)$ set, if for some triplet of points in P , say p_1, p_2, p_3 , there exist complex numbers u, v , and $z \neq 0$ such that $A = \bigcup_{j=1}^k L_j$ where

$$L_j = \frac{p_1 - p_j}{p_1 - p_3}(u + zI_m) + \frac{p_2 - p_j}{p_2 - p_3}(v + zI_m).$$

(We suggest depicting the special case $u = v = 0, z = 1$; all others are similar.)

A $G_P(m)$ set has the property that whenever we ‘place’ a similar copy of P in such a way that $p_1 \in L_1$ and $p_2 \in L_2$, then $p_j \in L_j$ for all $j \geq 3$ (hence the term ‘generated’). We make this precise in the following proposition.

Proposition 1. *For $\alpha, \beta \in I_m$ and $1 \leq j \leq k$ let*

$$a_{j,\alpha,\beta} = \frac{p_1 - p_j}{p_1 - p_3}(u + z\alpha) + \frac{p_2 - p_j}{p_2 - p_3}(v + z\beta) \in L_j.$$

If $a_{1,\alpha,\beta} \neq a_{2,\alpha,\beta}$ the set $\{a_{1,\alpha,\beta}, a_{2,\alpha,\beta}, a_{3,\alpha,\beta}, \dots, a_{k,\alpha,\beta}\}$ is similar to P .

Proof. For fixed α and β we have

$$\begin{aligned} a_{j,\alpha,\beta} &= \frac{p_1}{p_1 - p_3}(u + z\alpha) + \frac{p_2}{p_2 - p_3}(v + z\beta) + \left(\frac{u + z\alpha}{p_3 - p_1} + \frac{v + z\beta}{p_3 - p_2}\right)p_j \\ &= u_{\alpha,\beta} + z_{\alpha,\beta}p_j \end{aligned}$$

where $u_{\alpha,\beta}$ and $z_{\alpha,\beta}$ do not depend on j and $z_{\alpha,\beta} \neq 0$ since, by assumption $a_{1,\alpha,\beta} \neq a_{2,\alpha,\beta}$. ■

By definition L_1, L_2 and L_3 are arithmetic progressions of lengths $2m+1, 2m+1$ and $4m+1$ respectively. On the other hand, for $j \geq 4$ the set L_j is generated by $\frac{p_1 - p_j}{p_1 - p_3}z$ and $\frac{p_2 - p_j}{p_2 - p_3}z$. It turns out that these vectors are parallel if and only if $\{p_1, p_2, p_3, p_j\}$ lie on a circle or a line. Hence L_j is entirely contained in a line whenever p_1, p_2, p_3 , and p_j are cocyclic or collinear, and it is a $(2m+1) \times (2m+1)$ lattice otherwise.

If P is a triangle and A is a $G_P(m)$ set corresponding to the values $u = v = 0$ and $z = 1$, then by the above proposition we have $S_P(A) \geq (2m + 1)^2 - 1$. Also, by our previous remark, $n = |A| = 8m + 1$. Thus A is an n -element set with at least $n^2/16$ triplets similar to P . This is a minor improvement over the $n^2/18$ construction given in [3]; which can in fact be seen as a proper subset of A .

The following theorem is the main result of the paper. It describes the structure of sets in the plane with a large number of subsets similar to P by showing the existence of large $G_P(m)$ subsets. The proof of the theorem will be postponed to Section 4. Before that, we will explore some interesting geometric consequences.

Theorem 1. *For every $c > 0$ and m positive integer there is a threshold $N_0 = N_0(c, m)$ with the following property.*

Every (P, c) -rich set with $n > N_0$ elements must contain a $G_P(m)$ set.

Proof. See Section 4. ■

Corollary 1. *If $S'_P(n, m)$ denotes the maximum number of subsets similar to P , among all n -element subsets of the plane with no m points on a line, then for every fixed m we have $S'_P(n, m) = o(n^2)$.*

Proof. Follows directly from Theorem 1 and the fact that any $G_P(m)$ set has m collinear points. ■

As a further corollary we also have $S_P^{gen}(n) = o(n^2)$, where this last function is the maximum number of subsets similar to P among n -sets in general position (no 3 on a line no 4 on a circle).

The next corollary states the existence of large lattices among (P, c) -rich sets when P is not cocyclic or collinear.

Corollary 2. *If P is a finite set and $\{p_1, p_2, p_3, p_4\} \subseteq P$ is not a cocyclic or a collinear set then for every $c > 0$ and m positive integer there is a threshold $N_0 = N_0(c, m)$ with the following property.*

Every (P, c) -rich set with $n > N_0$ elements must contain a $(2m+1) \times (2m+1)$ lattice with generators x and y satisfying $x/y = (p_1; p_2; p_3; p_4)$.

Proof. Assume without loss of generality that $p_1 = 0$ and $p_2 = 1$. Suppose Q is a (P, c) -rich set with $|Q| = n > N_0$, where N_0 is given by Theorem 1. Hence Q must contain a $G_P(m)$ set, so in particular Q contains the set

$$\begin{aligned} L_4 &= \frac{p_4}{p_3}(u + zI_m) + \frac{1 - p_4}{1 - p_3}(v + zI_m) \\ &= \left(\frac{p_4}{p_3}u + \frac{1 - p_4}{1 - p_3}v \right) + \frac{p_4}{p_3}zI_m + \frac{1 - p_4}{1 - p_3}zI_m. \end{aligned}$$

Let $x = \frac{1-p_4}{1-p_3}z$, $y = \frac{p_4}{p_3}z$. Notice that $x/y = (0; 1; p_3; p_4)$ and since $\{p_1, p_2, p_3, p_4\}$ is not a cocyclic set then $(0; 1; p_3; p_4) \notin \mathbb{R}$, and consequently x and y are linearly independent vectors. Thus L_4 is the required lattice. ■

There are n -sets with cn^2 triples determining equilateral triangles, and containing no 3×3 lattices (e.g. any $G_P(m)$ set with n points and P an equilateral triangle). But large lattices are unavoidable whenever $c > 1/6$, more precisely we have the following.

Corollary 3. *Let T be an equilateral triangle. For m positive integer and $\varepsilon > 0$ there is a threshold $N_1 = N_1(\varepsilon, m)$ with the following property.*

Every set A with $n > N_1$ elements for which $S_T(A) \geq \left(\frac{1}{6} + \varepsilon\right)n^2$ must contain a $(2m + 1) \times (2m + 1)$ regular triangle lattice.

Proof. Let T be the triangle $\{0, 1, e^{i\pi/3}\}$ and R the rhomb $\{0, 1, e^{i\pi/3}, e^{-i\pi/3}\}$. Suppose A is an n -element set satisfying $S_T(A) \geq \left(\frac{1}{6} + \varepsilon\right)n^2$. For $i = 1, 2$ define X_i as the number of pairs in A that are the vertices of exactly i equilateral triangles with vertices in A . Observe that according to our definitions $S_T(A) = \frac{1}{3}X_1 + \frac{2}{3}X_2$ and $S_R(A) = X_2$, hence

$$X_1 + 2X_2 \geq \left(\frac{1}{2} + 3\varepsilon\right)n^2$$

and using the trivial inequality $X_1 + X_2 < (1/2)n^2$ we conclude that

$$S_R(A) = X_2 > 3\varepsilon n^2.$$

Hence by [Corollary 2](#), for $n > N_1 = N_0(3\varepsilon, m)$ we have that A contains a $(2m + 1) \times (2m + 1)$ lattice with generators x and y satisfying $x/y = (0; 1; e^{i\pi/3}; e^{-i\pi/3}) = -e^{i\pi/3}$, so the lattice is in fact a regular triangle lattice. ■

As a further corollary we can remark that in [1] it was proved that $S_T(n) \geq \left(\frac{1}{3} - \frac{\sqrt{3}}{4\pi}\right)n^2 + O(n^{3/2})$, so the extremal sets for the function $S_T(n)$ contain large regular lattices.

When the set P is cocyclic [Theorem 1](#) cannot guarantee a lattice substructure, but in that case we can always obtain a set of k concurrent lines with many points.

Corollary 4. *A set of arithmetic progressions in \mathbb{C} is said to be concurrent if the lines containing the progressions are concurrent. If P is a cocyclic set with k elements then for every $c > 0$ and m positive integer there is a threshold $N_0 = N_0(c, m)$ with the following property.*

Every (P, c) -rich set with $n > N_0$ elements must contain a set of k concurrent arithmetic progressions each of size $2m + 1$.

Proof. Let $g_j = (p_1; p_2; p_3; p_j)$ for $j \geq 2$. Assume again without loss of generality that $p_1 = 0$ and $p_2 = 1$. Since P is a cocyclic set then we know that $g_j \in \mathbb{R}$ for all $j \geq 2$. Suppose Q is a (P, c) -rich set with $|Q| = n > N_0$, where N_0 is given by [Theorem 1](#). Thus Q contains a $G_P(m)$ set. Define the lines

$$\begin{aligned} \ell_1 &= \frac{v}{1 - p_3} + \frac{z}{1 - p_3} \mathbb{R} \quad \text{and} \\ \ell_j &= (u + g_j v) + \frac{z p_j}{p_3} \mathbb{R} \quad \text{for } j \geq 2. \end{aligned}$$

Clearly $L_j \subseteq \ell_j$ and every L_j contains a $2m + 1$ arithmetic progression. Now consider the similarity transformation $T : \mathbb{C} \rightarrow \mathbb{C}$ given by $T(w) = \frac{p_3}{z} w - \frac{u}{z}$. By definition we have

$$\begin{aligned} T(\ell_1) &= x + \frac{p_3}{1 - p_3} \mathbb{R} \quad \text{and} \\ T(\ell_j) &= x(1 - p_j) + p_j \mathbb{R} \quad \text{for } j \geq 2, \end{aligned}$$

where $x = \frac{p_3(u+v)-u}{z(1-p_3)}$. A simple calculation shows that $R_j := T(\ell_j) \cap T(\ell_2)$ is given by

$$\begin{aligned} R_1 &= \frac{p_3(1 - \overline{p_3})\overline{x} - \overline{p_3}(1 - p_3)x}{p_3 - \overline{p_3}} \quad \text{and} \\ R_j &= \frac{p_j(\overline{x} + x\overline{p_j}) - \overline{p_j}(x + \overline{x}p_j)}{p_j - \overline{p_j}} \quad \text{for } j \geq 3. \end{aligned}$$

Using the fact that $g_j \in \mathbb{R}$ and $p_j = p_3(p_3 + g_j(1 - p_3))^{-1}$ we obtain for $j \geq 3$

$$R_j = \frac{p_3(1 - \overline{p_3})\overline{x} - \overline{p_3}(1 - p_3)x}{p_3 - \overline{p_3}}.$$

Therefore the set of lines $\{\ell_1, \ell_2, \dots, \ell_k\}$ is concurrent. ▀

3. Cocyclic and collinear patterns

To complement the results from last section we present the following theorem, where for P a cocyclic set, we construct a set contained in k concurrent lines and with many similar copies of P .

The proof of the theorem mimics, in a certain sense, the construction first done in [\[3\]](#). The main difference being the use of the cross ratios $(p_1; p_2; p_3; p_j)$ instead of the value p_j , this allows our set to be contained in a set of at most k lines as opposed to have the grid-like structure described in [\[3\]](#).

Theorem 2. For every k -element cocyclic or collinear set P with $k \geq 3$ and $S_P(n) = \Theta(n^2)$, there is a constant $c = c(P) > 0$ and a collection of lines $\ell_1, \ell_2, \dots, \ell_k$ through the origin satisfying the following property.

For every integer $n \geq k$ there is a n -element set $A_n \subset \bigcup_{j=1}^k \ell_j$ satisfying $S_P(A_n) \geq cn^2$.

Before we see the proof of the theorem let us state the following consequence which characterizes the notion of cocyclicity (or collinearity) for finite sets.

Corollary 5. For every k -element set P satisfying $S_P(n) = \Theta(n^2)$ the following are equivalent.

1. P is cocyclic or collinear.
2. There is a constant $c > 0$ and arbitrarily large (P, c) -rich sets containing no $k \times k$ lattices.

Proof. Trivial for $k \leq 2$. The implication (1) \Rightarrow (2) follows from Theorem 2 and the fact that a set contained in k concurrent lines cannot contain a $k \times k$ lattice. (2) \Rightarrow (1) follows directly from Corollary 2 using any value $m \geq (k - 1)/2$. ■

Proof of Theorem 2. Assume without loss of generality that $p_1 = 0$ and $p_2 = 1$. Let $g_2 = 0$, $g_3 = 1$, and $g_j = (0; 1; p_3; p_j)$ for $4 \leq j \leq k$; note that $g_j \in \mathbb{R}$ since P is cocyclic or collinear. Since $S_P(n) = \Theta(n^2)$, we have that by Laczkovich–Ruzsa Characterization Theorem [9] every g_j is algebraic. Let D be the degree of $L = \mathbb{Q}(g_3, g_4, \dots, g_k)$ over \mathbb{Q} and $\{h_1, h_2, \dots, h_D\}$ be a basis of L over \mathbb{Q} .

We first prove the theorem for an increasing sequence $\{n_m\}$ of values of n . For every $m \geq 1$ consider the following set of real numbers

$$\mathcal{G}_m = \left\{ \sum_{i=1}^D a_i h_i : a_i \in I_m \right\}.$$

Clearly $|\mathcal{G}_m| = (2m + 1)^D$. Let $J_m = \mathcal{G}_m \times \mathcal{G}_m \setminus \{(a, b) \in \mathcal{G}_m \times \mathcal{G}_m : bp_3 - a = 0\}$. For every $(a, b) \in J_m$ define the similarity transformation

$$T_{a,b}(z) = \frac{p_3(a - b)}{p_3 - 1} + \frac{bp_3 - a}{p_3 - 1}z.$$

Let $n_m = \left| \bigcup_{\substack{(a,b) \in J_m \\ 1 \leq j \leq k}} T_{a,b}(p_j) \right|$ and $A_{n_m} = \bigcup_{\substack{(a,b) \in J_m \\ 1 \leq j \leq k}} T_{a,b}(p_j)$. By definition we have

$$(2) \quad T_{a,b}(0) = \frac{p_3}{p_3 - 1}(a - b), \quad \text{and} \quad T_{a,b}(p_j) = p_j(a(1 - g_j) + bg_j) \quad \text{for } j \geq 2.$$




Fig. 1 Examples of the sets A_{n_m} for some sets P .

Let $\ell_1, \ell_2, \dots, \ell_k$ be a collection of lines defined as follows,

$$(3) \quad \ell_1 := \left(\frac{p_3}{p_3 - 1} \right) \mathbb{R}, \text{ and } \ell_j := p_j \mathbb{R} \text{ for } j \geq 2.$$

By construction we have that for every $1 \leq j \leq k$

$$\bigcup_{(a,b) \in J_m} T_{a,b}(p_j) \subseteq \ell_j,$$

thus $A_{n_m} \subset \bigcup_{j=1}^k \ell_j$ and $T_{a,b}(P) \subset A_{n_m}$ for every $(a,b) \in J_m$.

Now, if P is cocyclic then all the lines in (3) are pairwise different which together with (2) gives $T_{a,b}(P) \neq T_{a',b'}(P)$ whenever $(a,b) \neq (a',b')$. On the other hand, if P is collinear we can assume $p_1 = 0 < p_3 < p_4 < \dots < p_k < 1 = p_2$. If $T_{a,b}(P) = T_{a',b'}(P)$ then either $(a,b) = (a',b')$ or by (2) P is symmetric with respect to $1/2$ and $T_{a,b}^{-1} \circ T_{a',b'}(z)$ is a half-turn rotation with center $1/2$. In both cases, for each pair (a,b) there is at most one pair $(a',b') \neq (a,b)$ such that $T_{a,b}(P) = T_{a',b'}(P)$ which means

$$(4) \quad S_P(A_{n_m}) \geq \frac{1}{2} |J_m| \geq \frac{1}{2} (|\mathcal{G}_m|^2 - |\mathcal{G}_m|) \geq \frac{1}{3} |\mathcal{G}_m|^2.$$

Now we will find appropriate bounds for the size of A_{n_m} in terms of $|\mathcal{G}_m|$. To accomplish this we need the following lemma.

Lemma 1. *There are positive integers M, K independent of m so that for every $2 \leq j \leq k$ we have that $(Kg_j)\mathcal{G}_m \subseteq \mathcal{G}_{Mm}$.*

Proof. Let $y \in L = \mathbb{Q}(g_3, g_4, \dots, g_k)$. Express yh_i as a linear combination of $\{h_1, h_2, \dots, h_D\}$ over \mathbb{Q} ,

$$yh_i = \sum_{j=1}^D q_{i,j} h_j.$$

Suppose $q_{i,j} = a_{i,j}/b_{i,j}$ with $a_{i,j}, b_{i,j}$ integers. Let K be the least common multiple of $\{b_{i,j}\}$ and M_0 the maximum of $\{|a_{i,j}|\}$. Then for every $z = \sum_{i=1}^D z_i h_i \in \mathcal{G}_m$ we have

$$Kyz = \sum_{i=1}^D z_i Kyh_i = \sum_{j=1}^D \left(\sum_{i=1}^D \frac{z_i K a_{i,j}}{b_{i,j}} \right) h_j \in \mathbb{Z}(g_3, g_4, \dots, g_k),$$

and

$$\left| \sum_{i=1}^D \frac{z_i K a_{i,j}}{b_{i,j}} \right| \leq \sum_{i=1}^D \left| \frac{z_i K a_{i,j}}{b_{i,j}} \right| \leq KM_0 D m.$$

So by letting $M = KM_0 D$ we infer that $(Ky)\mathcal{G}_m \subseteq \mathcal{G}_{Mm}$. Clearly we can find common values of K and M to make this work simultaneously for $y = g_2, g_3, \dots, g_k$. ■

Besides this lemma we also need the following easy facts for generalized arithmetic progressions of the form \mathcal{G}_m : For any positive integers N, m_1, m_2, m we have that $N\mathcal{G}_m \subseteq \mathcal{G}_{Nm}$ and $\mathcal{G}_{m_1} \pm \mathcal{G}_{m_2} = \mathcal{G}_{m_1+m_2}$. Using these properties we have

$$\left| \bigcup_{(a,b) \in J_m} T_{a,b}(p_1) \right| = |\mathcal{G}_m - \mathcal{G}_m| = |\mathcal{G}_{2m}|$$

(note that $|\mathcal{G}_m| \leq |\mathcal{G}_{2m}| \leq 2^D |\mathcal{G}_m|$), and for $j \geq 2$,

$$\begin{aligned} \left| \bigcup_{(a,b) \in J_m} T_{a,b}(p_j) \right| &= |(1 - g_j)\mathcal{G}_m + g_j\mathcal{G}_m| \leq |K\mathcal{G}_m - Kg_j\mathcal{G}_m + Kg_j\mathcal{G}_m| \\ &\leq |\mathcal{G}_{Km} - \mathcal{G}_{Mm} + \mathcal{G}_{Mm}| = |\mathcal{G}_{(K+2M)m}| \leq (K + 2M)^D |\mathcal{G}_m|, \\ \left| \bigcup_{(a,b) \in J_m} T_{a,b}(p_j) \right| &= |(1 - g_j)\mathcal{G}_m + g_j\mathcal{G}_m| \geq |\mathcal{G}_m|. \end{aligned}$$

So by letting $c_1 = k$ and $c_2 = k(K + 2M)^D$ we have

$$(5) \quad c_1 |\mathcal{G}_m| \leq n_m \leq c_2 |\mathcal{G}_m|.$$

Hence by (4) we conclude that $S_P(A_{n_m}) \geq \frac{1}{3} |\mathcal{G}_m|^2 \geq \frac{1}{3} c_2^{-2} n_m^2$.

To prove the result for arbitrary $n \geq k$ we note that, using (5) and $|\mathcal{G}_m| = (2m + 1)^D$, there is a constant c_3 independent of m such that $n_m < n_{m+1} \leq c_3 n_m$. Thus, if $n_m < n \leq n_{m+1}$ we let A_n be the set A_{n_m} constructed before together with $n - n_m$ extra points in the line ℓ_1 . It is clear that $S_P(A_n) \geq c_4 n^2$ which completes the proof. ■

Remark. If P is collinear all the lines in (3) are the same. If P is cocyclic then all the lines in (3) are different and every similar copy of P lies on a circle passing through the origin.

4. Proof of Theorem 1

The proof of our main result will be based on some deep results of Combinatorial Number Theory, namely those which describe the structure of small sumsets. These results are usually stated for subsets of the integers, but they are in fact true for subsets of any torsion-free abelian group [10], in particular for subsets of \mathbb{C} . The notion of generalized arithmetic progressions (first introduced by Szemerédi [12] in his famous paper) is involved in all of these results. For any d positive integer, $\{n_1, n_2, \dots, n_d\} \subseteq \mathbb{N}$, and $\{z_1, z_2, \dots, z_d\} \subseteq \mathbb{C} \setminus \{0\}$, we call the set

$$\mathcal{G} = \left\{ \sum_{i=1}^d k_i z_i : 0 \leq k_i < n_i \right\}$$

a generalized arithmetic progression of dimension d and parameters $\{n_i\}$ and $\{z_i\}$. In what follows the symbol $\mathcal{G}_{d,n}$ will denote a generalized arithmetic progression of dimension not exceeding d and size n . For short we will use expressions like “there exists a $\mathcal{G}_{d,n}$ ”.

From now on we assume $A, B \subseteq \mathbb{C}$, $|A| = |B| = n$, and $E \subseteq A \times B$. We write $A + B = \{a + b : a \in A, b \in B\}$ and $A +_E B = \{a + b : (a, b) \in E\}$.

Using exponential sums and methods from the geometry of numbers. Freiman obtained the following result that describes the structure of $A + B$ when the cardinality of such set is not much greater than n . Later Ruzsa found a simpler proof [11].

Theorem A (Freiman [7]). *For every $C > 0$ there is a positive constant $c = c(C)$ and a natural number $d = d(C)$ satisfying the following property.*

If $|A + B| \leq Cn$ then $A \cup B$ is contained in a $\mathcal{G}_{d,cn}$.

Balog and Szemerédi found a statistical version of this theorem, by relaxing the assumption that all pairwise sums must be taken into account, and just considering the sums of cn^2 pairs.

Theorem B (Balog, Szemerédi [2]). *For every $C_1, C_2 > 0$ there are positive constants $c_1 = c_1(C_1, C_2)$, $c_2 = c_2(C_1, C_2)$, and a natural number $d = d(C_1, C_2)$ satisfying the following property.*

If $|A +_E B| \leq C_1n$ and $|E| \geq C_2n^2$ then there is a \mathcal{G}_{d,c_1n} satisfying $|A \cap \mathcal{G}_{d,c_1n}| \geq c_2n$ and $|B \cap \mathcal{G}_{d,c_1n}| \geq c_2n$.

Laczkovich and Ruzsa took this theorem a step further by proving the following generalization, which is precisely the result we use for the proof of [Theorem 1](#).

Theorem C (Laczkovich, Ruzsa [9]). *For every $C_1, C_2 > 0$ there are positive constants $c_1 = c_1(C_1, C_2)$, $c_2 = c_2(C_1, C_2)$, and a natural number $d = d(C_1, C_2)$ satisfying the following property.*

If $|A +_E B| \leq C_1n$ and $|E| \geq C_2n^2$ then there is a \mathcal{G}_{d,c_1n} satisfying

$$|E \cap (\mathcal{G}_{d,c_1n} \times \mathcal{G}_{d,c_1n})| \geq c_2n^2.$$

The following lemma, which is a consequence of [Theorem C](#), is used in the proof of [Theorem 1](#). Any proof with explicit bounds of this lemma would in turn provide explicit bounds for the results in this paper.

Lemma 2. *For every $C_1, C_2 > 0$ there is a positive constant $c_3 = c_3(C_1, C_2)$ such that for every positive integer M , there is a threshold function $N = N(C_1, C_2, M)$ satisfying the following property.*

For every $n \geq N$, if $|A +_E B| \leq C_1n$ and $|E| \geq C_2n^2$ then there are arithmetic progressions $\mathcal{G}_1, \mathcal{G}_2$ of length $l \geq M$ and common difference such that

$$|E \cap (\mathcal{G}_1 \times \mathcal{G}_2)| \geq c_3l^2.$$

Proof. By [Theorem C](#), applied to the sets A and B , there are positive constants c_1, c_2 , a positive integer d (all depending on C_1 and C_2 only), and $\mathcal{G} = \mathcal{G}_{d,c_1n}$ a generalized arithmetic progression satisfying $|(\mathcal{G} \times \mathcal{G}) \cap E| \geq c_2n^2$. Suppose \mathcal{G} is given by

$$\mathcal{G} = \left\{ \sum_{i=1}^d k_i z_i : 0 \leq k_i < n_i \right\}$$

with $n_1 \geq n_2 \geq \dots \geq n_d \geq 1$ and $\prod_{i=1}^d n_i = c_1 n$. Let $c_3 = c_2/c_1^2$, $N = M^d/c_1$, and $l = n_1$.

For every $r = (r_2, r_3, \dots, r_d) \in [n_2] \times [n_3] \times \dots \times [n_d]$ define

$$\mathcal{G}_r = \left\{ k_1 z_1 + \sum_{i=2}^d r_i z_i : 0 \leq k_1 < n_1 = l \right\}.$$

Observe that

$$\sum_{r,s \in [n_2] \times [n_3] \times \dots \times [n_d]} |(\mathcal{G}_r \times \mathcal{G}_s) \cap E| \geq |(\mathcal{G} \times \mathcal{G}) \cap E| \geq c_2 n^2$$

so by an averaging argument there are $R_1, R_2 \in [n_2] \times [n_3] \times \dots \times [n_d]$ satisfying

$$|E \cap (\mathcal{G}_{R_1} \times \mathcal{G}_{R_2})| \geq (c_2/c_1^2) n_1^2 = c_3 l^2.$$

Notice that \mathcal{G}_{R_1} and \mathcal{G}_{R_2} are arithmetic progressions with common difference z_1 . If $n \geq N$ then $l^d = n_1^d \geq c_1 n \geq c_1 N = M^d$, i.e., $l \geq M$ and thus $\mathcal{G}_{R_1}, \mathcal{G}_{R_2}$ is the required pair of arithmetic progressions. ■

The other main ingredient for our proof is a two dimensional generalization of the famous Szemerédi Theorem [12] regarding the existence of long arithmetic progressions among subsets of \mathbb{Z} with positive density. The theorem is due to Fürstenberg and Katznelson and was obtained by means of Ergodic Theory [8]. No quantitative proof of this result is known.

Theorem D (Fürstenberg, Katznelson [8]). *For every integer m and constant $c > 0$ there is a threshold function $N_1 = N_1(c, m)$ with the following property.*

For every $n \geq N_1$, if $A \subseteq [n] \times [n]$ and $|A| \geq cn^2$ then A contains an $m \times m$ square sublattice, i.e., a set of the form $(a_1 + b[m]) \times (a_2 + b[m])$ with a_1, a_2, b positive integers.

Finally we are in a position to prove the main theorem.

Proof of Theorem 1. First assume without loss of generality that $p_1 = 0$ and $p_2 = 1$. Suppose Q is a (P, c) -rich set of size n . Let $A = p_3 Q$, $B = (1 - p_3) Q$, and

$$E = \left\{ (a, b) \in A \times B : bp_3 \neq a(1 - p_3) \text{ and } a \frac{p_j}{p_3} + b \frac{1 - p_j}{1 - p_3} \in Q \text{ for all } j \right\}$$

Consider the function $f: E \rightarrow \{X \subset Q: X \sim P\}$ given by

$$\begin{aligned} f(a, b) &= \left\{ a \frac{p_j}{p_3} + b \frac{1 - p_j}{1 - p_3} : 1 \leq j \leq k \right\} \\ &= \left\{ \frac{b}{1 - p_3} + \left(\frac{a}{p_3} - \frac{b}{1 - p_3} \right) p_j : 1 \leq j \leq k \right\}. \end{aligned}$$

Observe that f is well defined since $a/p_3 - b/(1-p_3) \neq 0$ by assumption (i.e., $bp_3 \neq a(1-p_3)$). Let $X \subseteq Q$ satisfy $X \sim P$, thus $X = w + zP$ for some $w, z \in \mathbb{C}$ and $z \neq 0$. Let $a = p_3(w + z)$ and $b = w(1 - p_3)$, notice that $(a, b) \in E$ and $w + zp_j = a \frac{p_j}{p_3} + b \frac{1-p_j}{1-p_3} \in Q$. Thus $f(a, b) = X$, i.e., f is surjective.

Therefore $|E| \geq |\{X \subset Q : X \sim P\}| = S_P(Q) \geq cn^2$, and additionally $|A +_E B| \leq n$ since $A +_E B \subseteq Q$ (see the condition in the definition of E , for $p_j = p_3$). So we have the exact hypothesis of Lemma 2 with $C_1 = 1$ and $C_2 = c$. Let $M = N_1(c_3, 2m + 1)$ and $N_0 = N(1, c, M)$, where N_1 and N are the respective threshold functions of Theorem D and Lemma 2. Therefore, by Lemma 2 there are arithmetic progressions $\mathcal{G}_1, \mathcal{G}_2$ of length $l \geq M$ and common difference such that

$$(6) \quad |E \cap (\mathcal{G}_1 \times \mathcal{G}_2)| \geq c_3 l^2.$$

Suppose $\mathcal{G}_i = \{u_1 + jz : 0 \leq j < l\}$ for $i = 1, 2$. Identify $\mathcal{G}_1 \times \mathcal{G}_2$ with $[l] \times [l]$ and $E \cap (\mathcal{G}_1 \times \mathcal{G}_2)$ with the corresponding subset of $[l] \times [l]$. Now, since $l > M = N_1(c_3, 2m + 1)$ then using (6) as the hypothesis for Theorem D we conclude that $E \cap (\mathcal{G}_1 \times \mathcal{G}_2)$ contains a $(2m + 1) \times (2m + 1)$ square sublattice $S_1 \times S_2$. Suppose $S_i = \hat{u}_i + tzI_m$ for $i = 1, 2$, with t a positive integer and $\hat{u}_1 \in \mathcal{G}_1, \hat{u}_2 \in \mathcal{G}_2$. Since $S_1 \times S_2 \subseteq E$ we have

$$L_j := \frac{p_j}{p_3} S_1 + \frac{1 - p_j}{1 - p_3} S_2 \subset Q \text{ for } 1 \leq j \leq k.$$

Hence $\bigcup_{j=1}^k L_j$ is the required $G_P(m)$ set contained in Q . ■

Remark. We actually proved that Q contains a $G_P(m)$ set for every choice of the triplet p_1, p_2, p_3 .

Acknowledgments. We are grateful to J. Beck and N. Simányi for sharing with us their knowledge on the applications of Ergodic Theory to Combinatorial Number Theory. We also thank the referees for their valuable suggestions to improve the presentation.

References

- [1] B. ÁBREGO and S. FERNÁNDEZ-MERCHANT: On the Maximum Number of Equilateral Triangles I, *Discrete and Computational Geometry* **23** (2000), 129–135.
- [2] A. BALOG and E. SZEMERÉDI: A Statistical Theorem of Set Addition, *Combinatorica* **14** (1994), 263–268.
- [3] GY. ELEKES and P. ERDŐS: Similar Configurations and Pseudogrids, in *Intuitive Geometry*, Colloquia Mathematica Societatis János Bolyai, North Holland, Amsterdam 1994, 85–104.

- [4] P. ERDŐS and G. PURDY: Some Extremal Problems in Geometry, *Journal of Combinatorial Theory* **10** (1971), 246–252.
- [5] P. ERDŐS and G. PURDY: Some Extremal Problems in Geometry III, *Proc. 6th Southeastern Conference in Combinatorics, Graph Theory and Comp.* (1975), 291–308.
- [6] P. ERDŐS and G. PURDY: Some Extremal Problems in Geometry IV, *Proc. 7th Southeastern Conference in Combinatorics, Graph Theory and Comp.* (1976), 307–322.
- [7] G. A. FREIMAN: *Foundations of a Structural Theory of Set Addition*, Translation of Mathematical Monographs vol. 37, American Mathematical Society, Providence, R.I., USA, 1973.
- [8] H. FÜRSTENBERG and Y. KATZNELSON: An ergodic Szemerédi theorem for commuting transformations, *Journal d'Analyse Mathématique* **34** (1978), 275–291.
- [9] M. LACZKOVICH and I. Z. RUZSA: The Number of Homothetic Subsets, *The mathematics of Paul Erdős II*, Springer Verlag 1997, 294–302.
- [10] M. B. NATHANSON: *Additive Number Theory: Inverse Problems and the Geometry of Sumsets*, Springer Verlag 1996.
- [11] I. Z. RUZSA: Generalized Arithmetic Progressions and Sum Sets, *Acta Mathematica Hungarica* **65** (1994), 379–388.
- [12] E. SZEMERÉDI: On sets of integers containing no k elements in arithmetic progression, *Acta Arithmetica* **27** (1975), 199–245.

Bernardo M. Ábrego

*Dept of Mathematics
California State University
18111 Nordhoff St.
Northridge, CA 91330–8313
USA*
bernardo.abrego@csun.edu

György Elekes

*Dept of Computer Science
Eötvös University
Pázmány 1/C
Budapest, H–1117
Hungary*
elekes@cs.elte.hu

Silvia Fernández-Merchant

*Dept of Mathematics
California State University
18111 Nordhoff St.
Northridge, CA 91330–8313
USA*
silvia.fernandez@csun.edu