

Fuzzy closure operators II: induced relations, representation, and examples

R. Bělohávek

Abstract Closure operators (and related structures) are investigated from the point of view of fuzzy set theory. The paper is a follow up to [7] where fundamental notions and result have been established. The present approach generalizes the existing approaches in two ways: first, complete residuated lattices are used as the structures of truth values (leaving the unite interval $[0,1]$ with minimum and other t-norms particular cases); second, the monotony condition is formulated so that it can reflect also partial subthood (not only full subthood as in other approaches). In this paper, we study relations induced by fuzzy closure operators (fuzzy quasiorders and similarities); factorization of closure systems by similarities and by so-called decrease of logical precision; representation of fuzzy closure operators by (crisp) closure operators; relation to consequence relations; and natural examples illustrating the notions and results.

Keywords Closure operator, Fuzzy equivalence, Fuzzy quasiorder, Consequence relation

1

Introduction

This is a follow up to my paper [7]. In [7], closure operators and related structures have been considered from the point of view of fuzzy approach (graded truth approach; with complete residuated lattices taken for the structures of truth values). The aim of this paper is to present further results on fuzzy closure operators.

The organization and the content of the paper are as follows: Sect. 2 recalls the notions and main results of [7]. In Sect. 3, we study some induced (fuzzy) relations: fuzzy quasiorder and equivalence (similarity). We show a way to factorize the complete lattice of closed (w.r.t. to a given fuzzy closure operator) fuzzy sets by an a -cut of a naturally defined similarity relation, parameter a having the role of controlling the coarsity of the factorization.

Another way to factorize the lattice of closed fuzzy sets is by a so-called decrease of logical precision. The factorization processes have natural applications if the structure of closed sets has some natural interpretation and one needs to simplify the structure (as an example, we demonstrate the results on factorization of so-called fuzzy concept lattices). In Sect. 4, we present a natural representation of fuzzy closure operators by (classical) closure operators. Section 5 presents some examples of fuzzy closure operators. In Sect. 6, fuzzy closure operators and consequence relations are briefly discussed.

2

Fuzzy closure operators

Closure operators (and closure systems) play a significant role in both pure and applied mathematics. In the framework of fuzzy set theory, several particular examples of closure operators and systems have been considered (e.g. so-called fuzzy subalgebras, fuzzy congruences, fuzzy topology etc.). Recently, fuzzy closure operators and fuzzy closure systems themselves have been studied, see e.g. [8, 9, 15, 16]. As a matter of fact, a fuzzy set A is usually defined as a mapping from a universe set X into the real interval $[0, 1]$ in the above mentioned works. Therefore, the structure of truth values of the “logic behind” is fixed to $[0, 1]$ equipped with minimum being the operation corresponding to logical conjunction.

A general approach to the study of fuzzy closure operators has been outlined in [7]. Compared to previous approaches, there are basically two points of departure: First, the structure of truth values is assumed to form a complete residuated lattice. Second, the monotonicity condition is defined to mean “if A is almost a subset of B then the closure of A is almost a subset of the closure of B ”.

We now recall basic concepts and results (for proofs and further results we refer to [7]).

Definition 1 A complete residuated lattice is an algebra $\mathbf{L} = \langle L, \wedge, \vee, \otimes, \rightarrow, 0, 1 \rangle$ such that

- (1) $\langle L, \wedge, \vee, 0, 1 \rangle$ is a complete lattice with the least element 0 and the greatest element 1;
- (2) $\langle L, \otimes, 1 \rangle$ is a commutative monoid, i.e. \otimes is commutative, associative, and $x \otimes 1 = x$ holds for each $x \in L$;
- (3) \otimes, \rightarrow form an adjoint pair, i.e.

$$x \otimes y \leq z \quad \text{iff} \quad x \leq y \rightarrow z \quad (1)$$

holds for all $x, y, z \in L$.

R. Bělohávek
Institute for Research and Applications of Fuzzy Modeling,
University of Ostrava, Bráfova 7,
701 03 Ostrava, Czech Republic
e-mail: belohlav@osu.cz

Supported by grant no. 201/99/P060 of the GA ČR and by NATO Advanced Fellowship B for 2000. The author would like to thank to Professor G. Gerla for helpful comments. Part of the paper written during author’s research visit at the Center for Intelligent Systems, State University of New York at Binghamton. Support by the Center and by its director, Professor G.J. Klir, is gratefully acknowledged.

Residuated lattices play the role of structures of truth values in fuzzy logic. Introduced originally in the study of ideal systems of rings [24], residuated lattices have been introduced into the context of fuzzy logic by Goguen [17]. For logical calculi with truth values in residuated lattices (and special types of residuated lattices), basic properties of residuated lattices, and further references we refer to [18–20].

We only recall that the most studied and applied residuated lattices are those defined on the real interval $[0,1]$ (residuated lattices on $[0,1]$ uniquely correspond to left-continuous t-norms). Three most important structures pairs of adjoint operations are the following: the Lukasiewicz one ($a \otimes b = \max(a + b - 1, 0)$), $a \rightarrow b = \min(1 - a + b, 1)$, Gödel one ($a \otimes b = \min(a, b)$, $a \rightarrow b = 1$ if $a \leq b$ and $= b$ else), and product one ($a \otimes b = a \cdot b$, $a \rightarrow b = 1$ if $a \leq b$ and $= b/a$ else). Another important set of truth values is the set $\{a_0 = 0, a_1, \dots, a_n = 1\}$ ($a_0 < \dots < a_n$) with \otimes given by $a_k \otimes a_l = a_{\max(k+l-n, 0)}$ and the corresponding \rightarrow given by $a_k \rightarrow a_l = a_{\min(n-k+l, n)}$. A special case of the latter algebras is the Boolean algebra $\mathbf{2}$ of classical logic with the support $2 = \{0, 1\}$.

A nonempty subset $K \subseteq L$ is called an \leq -filter if for every $a, b \in L$ such that $a \leq b$ it holds that $b \in K$ whenever $a \in K$. An \leq -filter K is called a filter if $a, b \in K$ implies $a \otimes b \in K$. Unless otherwise stated, in what follows we denote by L a complete residuated lattice and by K an \leq -filter in L (both L and K possibly with indices).

An L -set (fuzzy set) [17, 28] A in a universe set X is any map $A : X \rightarrow L$. By L^X we denote the set of all L -sets in X . The concept of an L -relation is defined obviously. Operations on L extend pointwise to L^X , e.g. $(A \vee B)(x) = A(x) \vee B(x)$ for $A, B \in L^X$. Following common usage, we write $A \cup B$ instead of $A \vee B$, etc. Given $A, B \in L^X$, the *subsethood degree* [17] $S(A, B)$ of A in B is defined by $S(A, B) = \bigwedge_{x \in X} A(x) \rightarrow B(x)$. We write $A \subseteq B$ if $S(A, B) = 1$. Analogously, the *equality degree* $E(A, B)$ of A and B is defined by $E(A, B) = \bigwedge_{x \in X} (A(x) \leftrightarrow B(x))$. It is immediate that $E(A, B) = S(A, B) \wedge S(B, A)$. By $\{a_1/x_1, \dots, a_n/x_n\}$ we denote an L -set A for which $A(x) = a_i$ if $x = x_i$ ($i = 1, \dots, n$) and $A(x) = 0$ otherwise.

Recall that a (classical) closure operator on a set X is a mapping $C : 2^X \rightarrow 2^X$ satisfying the following conditions: $A \subseteq C(A)$, if $A_1 \subseteq A_2$ then $C(A_1) \subseteq C(A_2)$, and $C(A) = C(C(A))$, for any $A, A_1, A_2 \in 2^X$. More generally, if \subseteq denotes a partial order, we get the notion of closure operator in an ordered set [10].

Definition 2 An L_K -closure operator (fuzzy closure operator) on the set X is a mapping $C : L^X \rightarrow L^X$ satisfying

$$A \subseteq C(A) \quad (2)$$

$$S(A_1, A_2) \leq S(C(A_1), C(A_2)) \quad (3)$$

whenever $S(A_1, A_2) \in K$

$$C(A) = C(C(A)) \quad (4)$$

for every $A, A_1, A_2 \in L^X$.

If $K = L$, we omit the subscript K and call C an L -closure operator. The set K plays the role of the set of designated truth values. Condition (3) says that the closure preserves also partial subsethood whenever the subsethood degree is designated. Note that for $L = \{0, 1\}$, L_K -closure operators coincide with (classical) closure operators. Note also that for $L = [0, 1]$, $L_{\{1\}}$ -closure operators are precisely fuzzy closure operators studied by Gerla [9, 15, 16].

Definition 3 A system $\mathcal{S} = \{A_i \in L^X \mid i \in I\}$ is called closed under S_K -intersections iff for each $A \in L^X$ it holds that

$$\left(\bigcap_{i \in I, S(A, A_i) \in K} S(A, A_i) \rightarrow A_i \right) \in \mathcal{S}$$

where

$$\begin{aligned} & \left(\bigcap_{i \in I, S(A, A_i) \in K} S(A, A_i) \rightarrow A_i \right) (x) \\ &= \bigwedge_{i \in I, S(A, A_i) \in K} (S(A, A_i) \rightarrow A_i(x)) \end{aligned}$$

for each $x \in X$. A system closed under S_K -intersections will be called an L_K -closure system.

For $K = L$ the subscript will again be omitted. $\mathbf{2}$ -closure systems coincide with closure systems, i.e. systems of sets closed under intersections. In general, being closed under arbitrary intersections is a weaker condition than being closed under S_K -intersections. Closedness under S_K -intersections is, however, equivalent to closedness under intersections of “ K -shifted” L -sets. Let for $a \in L$, $A \in L^X$, denote by $a \rightarrow A$ the L -set defined by $(a \rightarrow A)(x) = a \rightarrow A(x)$.

Theorem 4 ([7]) \mathcal{S} is an L_K -closure system iff for any $a_i \in L$, $i \in I$, it holds $\bigcap_{a_i \in K} (a_i \rightarrow A_i) \in \mathcal{S}$. Therefore, a system \mathcal{S} which is closed under arbitrary intersections is an L_K -closure system iff for each $a \in K$ and $A \in \mathcal{S}$ it holds $a \rightarrow A \in \mathcal{S}$.

The following theorem shows another way to obtain the closure in an L_K -closure system.

Theorem 5 ([7]) Let $\mathcal{S} = \{A_i \in L^X \mid i \in I\}$ be an L_K -closure system. Then for each $A \in L^X$ it holds

$$\bigcap_{i \in I, S(A, A_i) \in K} S(A, A_i) \rightarrow A_i = \bigcap_{i \in I, A \subseteq A_i} A_i .$$

A natural idea is to consider the property “to be closed (w.r.t. a given fuzzy closure operator C)” a graded property. An L -set A can be considered to be “almost closed w.r.t. C ” iff “ A almost equals $C(A)$ ”. This poses a question of whether fuzzy closure systems can be defined as systems of “almost closed” fuzzy sets.

Definition 6 An L -system $\mathbf{S} \in L^X$ is called an L_K -closure L -system in X if for every $A, B \in L^X$ we have

$$\mathbf{S} \left(\bigcap_{A_i \in L^X, S(A, A_i) \in K} (\mathbf{S}(A_i) \otimes \mathbf{S}(A, A_i)) \rightarrow A_i \right) = 1, \quad (5)$$

$$\mathbf{S}(A) \otimes \mathbf{S}(A, B) \otimes \mathbf{S}(B, A) \leq \mathbf{S}(B) \quad (6)$$

$$\text{whenever } \mathbf{S}(B, A) \in K. \quad (7)$$

Remark (1) Note that the L-set

$$\bigcap_{A_i \in L^X, S(A, A_i) \in K} (\mathbf{S}(A_i) \otimes \mathbf{S}(A, A_i)) \rightarrow A_i$$

in X is defined by

$$\left(\bigcap_{A_i \in L^X, S(A, A_i) \in K} (\mathbf{S}(A_i) \otimes \mathbf{S}(A, A_i)) \rightarrow A_i \right)(x) = \bigwedge_{A_i \in L^X, S(A, A_i) \in K} (\mathbf{S}(A_i) \otimes \mathbf{S}(A, A_i)) \rightarrow A_i(x).$$

(2) An L_K -closure L-system is therefore an L-set of L-sets in X . We interpret $\mathbf{S}(A)$ as the degree to which $A \in L^X$ is closed. Condition (6) is naturally interpreted as the requirement that an L-set that is both a subset and a superset of to an “almost closed” L-set is itself “almost closed”.

Let C be an L_K -closure operator in X , \mathcal{S} be an L_K -closure system in X , and \mathbf{S} be an L_K -closure L-system in X . Define operators $C_{\mathcal{S}} : L^X \rightarrow L^X$ and $C_S : L^X \rightarrow L^X$, systems of L-sets $\mathcal{S}_C \subseteq L^X$ and $\mathcal{S}_S \subseteq L^X$, and L-systems of L-sets $\mathbf{S}_C \in L^{L^X}$ and $\mathbf{S}_{\mathcal{S}} \in L^{L^X}$ by

$$C_{\mathcal{S}}(A)(x) = \bigwedge_{i \in I, S(A, A_i) \in K} (\mathbf{S}(A, A_i) \rightarrow A_i(x)) \quad (8)$$

$$(C_S(A))(x) = \bigwedge_{A_i \in L^X, S(A, A_i) \in K} (\mathbf{S}(A_i) \otimes \mathbf{S}(A, A_i)) \rightarrow A_i(x) \quad (9)$$

$$\mathcal{S}_C = \{A \in L^X, A = C(A)\} \quad (10)$$

$$\mathcal{S}_S = \{A \in L^X, \mathbf{S}(A) = 1\} \quad (11)$$

$$\mathbf{S}_C(A) = E(A, C(A)) \quad (12)$$

$$\mathbf{S}_{\mathcal{S}}(A) = E(A, C_{\mathcal{S}}(A)). \quad (13)$$

The situation is depicted in Fig. 1.

Theorem 7 ([2]) Under the above notation, $C_{\mathcal{S}}$ and C_S are L_K -closure operators, \mathcal{S}_C and \mathcal{S}_S are L_K -closure systems, \mathbf{S}_C and $\mathbf{S}_{\mathcal{S}}$ are L_K -closure L-systems, and the diagram in Fig. 1 commutes.




Fig. 1. Commuting diagram of Theorem 7

Each oriented path in the diagram of Fig. 1 defines a mapping (a mapping composed of the mappings represented by the arrows). Commutativity of the diagram in Fig. 1 says that any two mappings corresponding to oriented paths with common starting node and final node are equal. Particularly, we have that the mappings defined by (8)–(13) are pairwise inverse (i.e. we have $C = C_{\mathcal{S}_C}$ and $C = C_{S_S}$, $\mathcal{S} = \mathcal{S}_{C_{\mathcal{S}}}$ and $\mathcal{S} = \mathcal{S}_{S_{\mathcal{S}}}$, and $\mathbf{S} = \mathbf{S}_{C_S}$ and $\mathbf{S} = \mathbf{S}_{\mathcal{S}_S}$); furthermore, we have $C = C_{\mathcal{S}_C}$ etc.

Definition 8 An L_K -Galois connection (fuzzy Galois connection) between the sets X and Y is a pair $\langle \uparrow, \downarrow \rangle$ of mappings $\uparrow : L^X \rightarrow L^Y$, $\downarrow : L^Y \rightarrow L^X$, satisfying

$$\mathbf{S}(A_1, A_2) \leq \mathbf{S}(A_2^{\uparrow}, A_1^{\uparrow}) \quad \text{whenever } \mathbf{S}(A_1, A_2) \in K \quad (14)$$

$$\mathbf{S}(B_1, B_2) \leq \mathbf{S}(B_2^{\downarrow}, B_1^{\downarrow}) \quad \text{whenever } \mathbf{S}(B_1, B_2) \in K \quad (15)$$

$$A \subseteq (A^{\uparrow})^{\downarrow} \quad (16)$$

$$B \subseteq (B^{\downarrow})^{\uparrow}. \quad (17)$$

for every $A, A_1, A_2 \in L^X$, $B, B_1, B_2 \in L^Y$.

If $K = L$ then we again omit the subscript K . Note also that an L_K -Galois connection between X and Y forms a Galois connection between the complete lattices $\langle L^X, \subseteq \rangle$ and $\langle L^Y, \subseteq \rangle$ [10, 21].

Remark Note that Galois connections between sets [10, 21] are just L-Galois connections for $L = 2$.

We will need the following results.

Theorem 9 ([7]) Let C be an L-closure operator, and $Y = \{C(A) | A \in L^X\}$. Then the pair of mappings $\uparrow_C : L^X \rightarrow L^Y$, $\downarrow_C : L^Y \rightarrow L^X$ defined for $A \in L^X$, $B \in L^Y$ and $x \in X$, $A' \in Y$ by

$$A^{\uparrow_C}(A') = \mathbf{S}(A, A')$$

$$B^{\downarrow_C}(x) = \bigwedge_{A \in Y} B(A) \rightarrow A(x)$$

forms an L-Galois connection such that $C = \uparrow_C \downarrow_C$.

Theorem 10 ([2]) For a binary L-relation $I \in L^{X \times Y}$ denote $\uparrow_I : L^X \rightarrow L^Y$ and $\downarrow_I : L^Y \rightarrow L^X$ the mappings defined for $A \in L^X$, $B \in L^Y$ by

$$A^{\uparrow_I}(y) = \bigwedge_{x \in X} (A(x) \rightarrow I(x, y)) \quad (18)$$

$$B^{\downarrow_I}(x) = \bigwedge_{y \in Y} (B(y) \rightarrow I(x, y)). \quad (19)$$

For an L-Galois connection $\langle \uparrow, \downarrow \rangle$ between X and Y denote $I_{\langle \uparrow, \downarrow \rangle}$ the binary L-relation $I \in L^{X \times Y}$ defined for $x \in X$, $y \in Y$ by $I(x, y) = \{1/x\}^{\uparrow}(y) (= \{1/y\}^{\downarrow}(x))$. Then $\langle \uparrow, \downarrow \rangle$ is an L-Galois connection and it holds

$$\langle \uparrow, \downarrow \rangle = \langle \uparrow_{I_{\langle \uparrow, \downarrow \rangle}}, \downarrow_{I_{\langle \uparrow, \downarrow \rangle}} \rangle \quad \text{and} \quad I = I_{\langle \uparrow, \downarrow \rangle}.$$

Induced relations: quasiorder and similarity

An L-relation R on a set X is called

reflexive if $R(x, x) = 1$

symmetric if $R(x, y) = R(y, x)$

K -transitive if $R(x, y) \otimes R(y, z) \leq R(x, z)$

whenever $R(x, y) \in K$ and $R(y, z) \in K$.

An L_K -quasiorder on X is an L-relation on X that is reflexive and K -transitive. An L_K -similarity (or L_K -equivalence) on X is an L-relation on X that is reflexive, symmetric, and K -transitive.

Remark. (1) Clearly, putting $L = 2$ we get the usual (bivalent) notion of a quasiorder and an equivalence relation (no matter what K). Thus, the notions of L_K -quasiorder and L_K -similarity are generalizations of the bivalent notions.

(2) If K is interpreted as the set of sufficiently high (designated) truth values, then K -transitivity means: "if the facts that $\langle x, y \rangle$ belongs to R and $\langle y, z \rangle$ belongs to R are sufficiently true then $\langle x, z \rangle$ also belongs to R (and is sufficiently true in case K is a filter)". For $K = L$ (in which case we omit the subscript K) we get the usual notions of L-quasiorder and L-similarity (called usually fuzzy quasiorder and fuzzy similarity).

Induced L-relations on X

Theorem 11 Let C be an L_K -closure operator on X . Then the relation Q_C in X defined by

$$Q_C(x, y) = C(\{1/x\})(y)$$

is an L_K -quasiorder.

Proof. Since $\{1/x\} \subseteq C(\{1/x\})$, reflexivity follows by

$$Q_C(x, x) = C(\{1/x\})(x) = 1.$$

K -transitivity: Let $Q_C(x, y) \in K$, $Q_C(y, z) \in K$. We have to show $Q_C(x, y) \otimes Q_C(y, z) \leq Q_C(x, z)$. By adjointness and by definition of Q_C we thus have to show that

$$C(\{1/x\})(y) \leq C(\{1/y\})(z) \rightarrow C(\{1/x\})(z)$$

whenever $C(\{1/x\})(y)$, $C(\{1/y\})(z) \in K$. We have $S(\{C(\{1/x\})(y)/\}, C(\{1/x\})) = 1$, therefore, by (3) and using $C(C(\{1/x\})) = C(\{1/x\})$, also $S(C(\{C(\{1/x\})(y)/\}), C(\{1/x\})) = 1$. Furthermore, by $C(\{1/x\})(y) \in K$ and (3),

$$\begin{aligned} C(\{1/x\})(y) &= S(\{1/y\}, \{C(\{1/x\})(y)/y\}) \\ &\leq S(C(\{1/y\}), C(\{C(\{1/x\})(y)/y\})) . \end{aligned}$$

Therefore,

$$\begin{aligned} C(\{1/x\})(y) &\leq S(C(\{1/y\}), C(\{C(\{1/x\})(y)/y\})) \\ &= S(C(\{1/y\}), C(\{C(\{1/x\})(y)/y\})) \\ &\quad \otimes S(C(\{C(\{1/x\})(y)/\}), C(\{1/x\})) \\ &\leq S(C(\{1/y\}), C(\{1/x\})) \\ &\leq C(\{1/y\})(z) \rightarrow C(\{1/x\})(z) . \end{aligned}$$

The proof is complete.

Remark. Note that Q_C actually satisfies a stronger condition than K -transitivity. Namely, as it follows from the proof of Theorem 11, $Q_C(x, y) \otimes Q_C(y, z) \leq Q_C(x, z)$ whenever $Q_C(x, y) \in K$. This property is typical for Pavelka style fuzzy logic (see [22] and also [18]): Take X to be the set of all formulas; put $K = \{1\}$; let C be the operator of syntactic consequence, i.e. for an L-set A of formulas and a formula $x \in X$, let $C(A)(x)$ be the degree of provability of x from A . One easily verifies that C satisfies the above condition stronger than K -transitivity. On the other hand, C does not satisfy the in a sense symmetric condition, i.e. it is not true that if $Q_C(y, z) \in K$ then $Q_C(x, y) \otimes Q_C(y, z) \leq Q_C(x, z)$.

$Q_C(x, y)$ is naturally interpreted as the truth degree to which y belongs to the closure of a singleton containing x . One might wonder what is the relationship between Q_C and $Q_{\mathcal{S}_C}$ defined by

$$Q_{\mathcal{S}_C}(x, y) = \bigwedge_{A \in \mathcal{S}_C, A(x) \in K} A(x) \rightarrow A(y) ,$$

i.e. the truth degree to which it holds that whenever it is sufficiently true that x belongs to some closure then y belongs to that closure as well.

Theorem 12 For any L_K -closure operator C we have $Q_C = Q_{\mathcal{S}_C}$. Therefore, $Q_{\mathcal{S}_C}$ is an L_K -quasiorder.

Proof. On the one hand, $C(\{1/x\})(x) = 1 \in K$ yields

$$\begin{aligned} Q_{\mathcal{S}_C}(x, y) &= \bigwedge_{A \in \mathcal{S}_C, A(x) \in K} A(x) \rightarrow A(y) \\ &\leq C(\{1/x\})(x) \rightarrow C(\{1/x\})(y) \\ &= 1 \rightarrow C(\{1/x\})(y) \\ &= C(\{1/x\})(y) = Q_C(x, y) . \end{aligned}$$

On the other hand, $Q_C(x, y) \leq Q_{\mathcal{S}_C}(x, y)$ is true iff for each $A \in \mathcal{S}_C$ such that $A(x) \in K$ we have $C(\{1/x\})(y) \leq A(x) \rightarrow A(y)$. Applying adjointness twice, the last inequality is equivalent to $A(x) \leq C(\{1/x\})(y) \rightarrow A(y)$ which is true. Indeed, since $A(x) \in K$, (3) gives

$$\begin{aligned} A(x) = 1 \rightarrow A(x) &= \{1/x\}(x) \rightarrow A(x) = S(\{1/x\}, A) \\ &\leq S(C(\{1/x\}), C(A)) = S(C(\{1/x\}), A) \\ &\leq C(\{1/x\})(y) \rightarrow A(y) . \end{aligned}$$

If K is, moreover, a filter in \mathbf{L} , the fact that Q_C is an L_K -quasiorder follows from Theorem 14 and the following statement.

Lemma 13 *Let K be an filter in \mathbf{L} , $\mathcal{S} = \{A_i \in L^X \mid i \in I\}$ be a system of \mathbf{L} -sets. Then the \mathbf{L} -relation $Q_{\mathcal{S}}$ on X given by*

$$Q_{\mathcal{S}}(x, y) = \bigwedge_{i \in I, A_i(x) \in K} A_i(x) \rightarrow A_i(y)$$

is an L_K -quasiorder.

Proof. Reflexivity follows from the fact that $A_i(x) \rightarrow A_i(x) = 1$ and from $\bigwedge \emptyset = 1$. K -transitivity: By definition, we have to show that if

$$\bigwedge_{i \in I, A_i(x) \in K} A_i(x) \rightarrow A_i(y) \in K$$

and

$$\bigwedge_{i \in I, A_i(y) \in K} A_i(y) \rightarrow A_i(z) \in K$$

then

$$\left(\bigwedge_{i \in I, A_i(x) \in K} A_i(x) \rightarrow A_i(y) \right) \otimes \left(\bigwedge_{i \in I, A_i(y) \in K} A_i(y) \rightarrow A_i(z) \right) \leq \left(\bigwedge_{i \in I, A_i(x) \in K} A_i(x) \rightarrow A_i(z) \right),$$

i.e. to show that for each $i \in I$ such that $A_i(x) \in K$ we have

$$\begin{aligned} A_i(x) \otimes \left(\bigwedge_{i \in I, A_i(y) \in K} A_i(y) \rightarrow A_i(z) \right) \\ \leq A_i(z). \end{aligned}$$

This inequality is true. Indeed,

$\bigwedge_{i \in I, A_i(x) \in K} A_i(x) \rightarrow A_i(y) \in K$ and $\bigwedge_{i \in I, A_i(y) \in K} A_i(y) \rightarrow A_i(z) \leq A_i(x) \rightarrow A_i(y)$ gives $A_i(x) \rightarrow A_i(y) \in K$. As also $A_i(x) \in K$, we have $A_i(x) \otimes (A_i(x) \rightarrow A_i(y)) \in K$. From $A_i(x) \otimes (A_i(x) \rightarrow A_i(y)) \leq A_i(y)$ we thus have $A_i(y) \in K$. Therefore, we conclude

$$\begin{aligned} A_i(x) \otimes \left(\bigwedge_{i \in I, A_i(x) \in K} A_i(x) \rightarrow A_i(y) \right) \\ \otimes \left(\bigwedge_{i \in I, A_i(y) \in K} A_i(y) \rightarrow A_i(z) \right) \\ \leq A_i(x) \otimes (A_i(x) \rightarrow A_i(y)) \otimes (A_i(y) \rightarrow A_i(z)) \leq A_i(z), \end{aligned}$$

completing the proof.

Indeed, putting $\mathcal{S} = \mathcal{S}_C$, Lemma 12 yields that $Q_{\mathcal{S}_C}$ is an L_K -quasiorder. Theorem 12 then completes the argument.

Remark. (1) A closer look at the proof of Lemma 15 shows that like Q_C , $Q_{\mathcal{S}}$ satisfies a stronger form of K -transitivity: $Q_{\mathcal{S}}(x, y) \otimes Q_{\mathcal{S}}(y, z) \leq Q_{\mathcal{S}}(x, z)$ whenever $Q_{\mathcal{S}}(x, y) \in K$.

(2) The assumption of Lemma 15 that K be closed w.r.t. \otimes is essential. As a counterexample, consider $I = \{i, j\}$, $X = \{x, y, z\}$, $L = [0, 1]$ equipped with Lukasiewicz structure, $A_i(x) = 0.9$, $A_i(y) = 0.8$, $A_i(z) = 0.7$, $A_j(x) = A_j(y) = A_j(z) = 1$. Taking $K = [0.9, 1]$ which is an \leq -filter not closed w.r.t. \otimes , we have $Q_{\mathcal{S}}(x, y) \otimes Q_{\mathcal{S}}(y, z) = 0.9 \otimes 1 = 0.9 \not\leq 0.8 = Q_{\mathcal{S}}(x, z)$.

It can be easily seen that every L_K -quasiorder on X induces an L_K -similarity E_Q on X by putting $E_Q(x, y) = Q(x, y) \wedge Q(y, x)$. Therefore, for any L_K -closure operator C on X , the \mathbf{L} -relation E_C on X defined by

$$E_C(x, y) = Q_C(x, y) \wedge Q_C(y, x)$$

is an L_K -similarity on X . We say that an L_K -similarity E on X is compatible with $A \in L^X$ if $A(x) \otimes E(x, y) \leq A(y)$ holds for any $x, y \in X$ such that $A(x) \in K$. The condition of compatibility translates verbally to “if it is sufficiently true that x belongs to A and if x and y are similar then y belongs to A as well”.

Theorem 14 *Let C be an L_K -closure operator on X . Then E_C is the largest L_K -similarity on X that is compatible w.r.t. every C -closed \mathbf{L} -set (i.e. w.r.t. every $A \in \mathcal{S}_C$).*

Proof. The fact that E_C is an L_K -similarity on X was established in the above paragraph. Let $x, y \in X$, $A \in \mathcal{S}_C$, $A(x) \in K$. By Theorem 12,

$$\begin{aligned} E_C(x, y) &= Q_{\mathcal{S}_C}(x, y) \wedge Q_{\mathcal{S}_C}(y, x) \leq Q_{\mathcal{S}_C}(x, y) \\ &= \bigwedge_{A \in \mathcal{S}_C, A(x) \in K} A(x) \rightarrow A(y) \leq A(x) \rightarrow A(y), \end{aligned}$$

i.e. $A(x) \otimes E_C(x, y) \leq A(y)$ by adjointness. We proved that E_C is compatible w.r.t. any $A \in \mathcal{S}_C$.

Let E be an L_K -similarity that is compatible w.r.t. any $A \in \mathcal{S}_C$. Take $x, y \in X$ and an $A \in \mathcal{S}_C$ such that $A(x) \in K$. Compatibility of E yields $A(x) \otimes E(x, y) \leq A(y)$, i.e. $E(x, y) \leq A(x) \rightarrow A(y)$. Since x, y , and A were chosen arbitrarily, we conclude

$$E(x, y) \leq \bigwedge_{A \in \mathcal{S}_C, A(x) \in K} A(x) \rightarrow A(y) \leq Q_{\mathcal{S}_C}(x, y).$$

Due to the symmetry of E we finally have

$$E(x, y) \leq Q_{\mathcal{S}_C}(x, y) \wedge Q_{\mathcal{S}_C}(y, x) = E_C(x, y)$$

proving that E_C is the largest L_K -similarity compatible with all $A \in \mathcal{S}_C$.

Factorization of L_K -closure systems by similarity For an L_K -closure operator on X , \mathcal{S}_C is a complete lattice w.r.t. \subseteq . This fact follows directly from the fact that \mathcal{S}_C is closed

w.r.t. arbitrary intersections. For various reasons (e.g. for computational ones), it might not be desirable to distinguish the particular L-sets in X . Rather, it can be advantageous to treat L-sets which are similar in terms of membership degrees of elements of X as if they were the same, i.e. one might desire to perform a kind of abstraction by factorization w.r.t. to a suitable similarity defined on L-sets. A suitable L-similarity relation is described in the following assertion (see e.g. [4]).

Lemma 15 *The L-relation E on X defined by*

$$E(A, B) = \bigwedge_{x \in X} A(x) \leftrightarrow B(x)$$

is an L-similarity on X .

For $A, B \in L^X$, $E(A, B)$ is the truth degree to which it is true that for any $x \in X$, x belongs to A iff x belongs to B . The first observation states that sufficiently high similarity between L-sets is preserved by L_K -closure operators.

Theorem 16 *For an L_K -closure operator C on X and $A, B \in L^X$ we have $E(A, B) \leq E(C(A), C(B))$ whenever $E(A, B) \in K$.*

Proof. It is easy to see that $E(A, B) = S(A, B) \wedge S(B, A)$. Therefore, $E(A, B) \in K$ yields $S(A, B) \in K$ and $S(B, A) \in K$. Applying (3) we get $S(A, B) \leq S(C(A), C(B))$ and $S(B, A) \leq S(C(B), C(A))$, and thus $E(A, B) = S(A, B) \wedge S(B, A) \leq S(C(A), C(B)) \wedge S(C(B), C(A)) = E(C(A), C(B))$.

Intuitively, the complete lattice \mathcal{S}_C can be simplified by putting similar closed L-sets together, i.e. putting together L-sets A and B for which $E(A, B)$ is high. Putting the L-sets together should be compatible w.r.t. the complete lattice structure on \mathcal{S}_C . Recall that for any $a \in L$, an a -cut of E is a (bivalent) relation ${}^a E$ on X defined by $\langle x, y \rangle \in {}^a E$ iff $a \leq E(x, y)$. It is easy to see that ${}^a E$ is always a tolerance on X (i.e. a reflexive and symmetric relation on X). If $\otimes = \wedge$, ${}^a E$ is, moreover, transitive, i.e. an equivalence relation. In general, ${}^a E$ is not transitive. Factorization of a structure by a compatible tolerance relation is, in general, not possible (one needs transitivity so that operations on the factor set can be defined). Surprisingly, Czédli [13] showed a way to factorize lattices by compatible tolerances (for factorization of complete lattices by tolerances see [26]). We now recall the necessary concepts: Let T be a tolerance relation on a support V of a complete lattice $\mathbf{V} = \langle V, \leq \rangle$. T is called compatible if it is preserved under arbitrary infima and suprema, i.e. if $\langle u_i, v_i \rangle \in T$ ($i \in I$) implies $\langle \bigwedge_{i \in I} u_i, \bigwedge_{i \in I} v_i \rangle \in T$ and $\langle \bigvee_{i \in I} u_i, \bigvee_{i \in I} v_i \rangle \in T$. For $v \in V$, denote $v_T = \bigwedge_{\langle v, v' \rangle \in T} v'$ and $v^T = \bigvee_{\langle v, v' \rangle \in T} v'$, and call each set of the form $[v]_T = [v_T, v^T] = \{v' \in V \mid v_T \leq v' \leq v^T\}$ a block of T . Denote $V/T = \{[v]_T \mid v \in V\}$ the set of all blocks of T and call it the factor set of V by T . Introduce a relation \leq_T defined on V/T by $[v]_T \leq [v']_T$ iff $\bigwedge [v]_T \leq \bigwedge [v']_T$ (or, equivalently, iff

$\bigvee [v]_T \leq \bigvee [v']_T$). The following assertion follows from [26].

Theorem 17 *Let C be an L_K -closure operator on X , T be a compatible tolerance relation on $\langle \mathcal{S}_C, \subseteq \rangle$. (1) \mathcal{S}_C/T is the set of all maximal blocks of T , i.e. $\mathcal{S}_C/T = \{B \subseteq \mathcal{S}_C \mid B \times B \subseteq T \text{ \& } ((\forall B' \subset B) B' \times B' \not\subseteq T)\}$. (2) $\langle \mathcal{S}_C/T, \subseteq_T \rangle$ is a complete lattice (factor lattice) where infima and suprema are given by*

$$\bigwedge_{i \in I} [A_i]_T = \left[\left(\bigcap_{i \in I} A_i \right)^T \right]_T \quad \text{and} \quad \bigvee_{i \in I} [A_i]_T = \left[\bigvee_{i \in I} A_i \right]_T.$$

One may easily verify that if T is, moreover, transitive (i.e. a complete congruence on \mathbf{V}), then $\langle \mathcal{S}_C, \subseteq_T \rangle$ is the well-known factor lattice.

To show that the a -cuts ${}^a E$ can be used to factorize \mathcal{S}_C by the above described procedure, we need to verify that ${}^a E$ is compatible w.r.t. \subseteq .

Lemma 18 *Let C be an L_K -closure operator on X . For any $a \in K$, ${}^a E$ is a compatible tolerance relation on the complete lattice $\langle \mathcal{S}_C, \subseteq \rangle$.*

Proof. We show that ${}^a E$ is compatible both with infima and suprema, i.e. we show that $\langle A_i, B_i \rangle \in {}^a E$ ($i \in I$) implies both $\langle \bigwedge_{i \in I} A_i, \bigwedge_{i \in I} B_i \rangle \in {}^a E$ and $\langle \bigvee_{i \in I} A_i, \bigvee_{i \in I} B_i \rangle \in {}^a E$.

Infima: Suppose $\langle A_i, B_i \rangle \in {}^a E$, i.e. $a \leq \bigwedge_{x \in X} (A_i(x) \leftrightarrow B_i(x))$ ($i \in I$). We have to show

$$a \leq \bigwedge_{x \in X} \left(\bigwedge_{i \in I} A_i(x) \leftrightarrow \bigwedge_{i \in I} B_i(x) \right),$$

i.e. to show that for each $i \in I$ we have

$$a \leq \left(\bigwedge_{i \in I} A_i(x) \leftrightarrow \bigwedge_{i \in I} B_i(x) \right).$$

The last inequality is true iff both

$$a \leq (\bigwedge_{i \in I} A_i(x) \rightarrow \bigwedge_{i \in I} B_i(x)) \text{ and}$$

$$a \leq (\bigwedge_{i \in I} B_i(x) \rightarrow \bigwedge_{i \in I} A_i(x))$$

are valid. Due to symmetry we verify only $a \leq (\bigwedge_{i \in I} A_i(x) \rightarrow \bigwedge_{i \in I} B_i(x))$ which is equivalent to $a \otimes \bigwedge_{i \in I} A_i(x) \leq \bigwedge_{i \in I} B_i(x)$. This inequality is true. Indeed, by assumption, $\langle A_i, B_i \rangle \in {}^a E$, i.e.

$$a \leq \bigwedge_{x \in X} (A_i(x) \leftrightarrow B_i(x)),$$

from which it follows $a \otimes A_i(x) \leq B_i(x)$ for any $i \in I, x \in X$. We therefore have

$$a \otimes \bigwedge_{i \in I} A_i(x) \leq \bigwedge_{i \in I} (a \otimes A_i(x)) \leq \bigwedge_{i \in I} B_i(x).$$

Suprema: We have to show $a \leq E(\bigvee_{i \in I} A_i, \bigvee_{i \in I} B_i)$, i.e.

$$a \leq E(C(\bigcup_{i \in I} A_i), C(\bigcup_{i \in I} B_i)).$$

First, observe that (*)

$$a \leq E(\bigcup_{i \in I} A_i, \bigcup_{i \in I} B_i):$$

Indeed, the inequality is true iff both $a \leq S(\bigcup_{i \in I} A_i, \bigcup_{i \in I} B_i)$ and $a \leq S(\bigcup_{i \in I} B_i, \bigcup_{i \in I} A_i)$ hold. Because of symmetry, we verify only the former one:

by definition, $a \leq S(\bigcup_{i \in I} A_i, \bigcup_{i \in I} B_i)$ is true iff

$$a \leq (\bigcup_{i \in I} A_i)(x) \rightarrow (\bigcup_{i \in I} B_i)(x),$$

i.e. iff

$$a \otimes (\bigcup_{i \in I} A_i)(x) \leq (\bigcup_{i \in I} B_i)(x)$$

holds for each $x \in X$. By assumption,

$$\begin{aligned} a \otimes \left(\bigcup_{i \in I} A_i \right) (x) &= \bigvee_{i \in I} (a \otimes A_i(x)) \leq \bigvee_{i \in I} B_i(x) \\ &= \left(\bigcup_{i \in I} B_i \right) (x) \end{aligned}$$

establishing (*).

Now, $a \leq E(\bigcup_{i \in I} A_i, \bigcup_{i \in I} B_i)$ implies both $a \leq S(\bigcup_{i \in I} A_i, \bigcup_{i \in I} B_i)$ and $a \leq S(\bigcup_{i \in I} B_i, \bigcup_{i \in I} A_i)$. Since $a \in K$, (3) implies $a \leq S(C(\bigcup_{i \in I} A_i), C(\bigcup_{i \in I} B_i))$ and $a \leq S(C(\bigcup_{i \in I} B_i), C(\bigcup_{i \in I} A_i))$, i.e.

$$a \leq E\left(C\left(\bigcup_{i \in I} A_i\right), C\left(\bigcup_{i \in I} B_i\right)\right) = E\left(\bigvee_{i \in I} A_i, \bigvee_{i \in I} B_i\right)$$

completing the proof.

Remark. (1) Note that the tolerance relation ${}^a E$ used to factorize \mathcal{S}_C need not to be supplied from the outside. It is determined by selecting an appropriate $a \in K$.

(2) The role of $a \in K$ is to control the granularity of the factorization: since $a \leq b$ implies ${}^b E \subseteq {}^a E$, the rule is “the bigger a , the finer the factorization”. Clearly, for the extreme cases of a , i.e. $a = 0$ (note that ${}^0 E = \mathcal{S}_C \times \mathcal{S}_C$ which is always a compatible relation on \mathcal{S}_C) and $a = 1$ we obtain $\mathcal{S}_C/{}^0 E$ which is a one-element lattice and $\mathcal{S}_C/{}^1 E$ which is a lattice isomorphic to \mathcal{S}_C .

(3) Note also that the fact $a \in K$ is not needed in the proof of compatibility with infima in \mathcal{S}_C .

Factorization of L_K -closure systems by decrease of logical precision

We now mention another way to factorize a system \mathcal{S}_C of all closed L -sets of an L -closure operator C . Let thus C be an L_1 -closure operator on X . It may be the case that one does not need to distinguish close truth values of L_1 . Formally, a kind of a factorization of L_1 is taking place. Assume therefore that there is a complete homomorphism h of L_1 onto L_2 (i.e. a homomorphism which preserves arbitrary infima and suprema). h induces a mapping $h^* : L_1^X \rightarrow L_2^X$ such that for $A \in L_1^X$, $h^*(A) \in L_2^X$ is defined by

$$(h^*(A))(x) = h(A(x))$$

for any $x \in X$. Since there is no danger of misunderstanding, we write simply h instead of h^* .

One may consider h as representing a decrease of logical precision (see [6]): Several truth values from L_1 may collapse into one truth value from L_2 ; instead of $A \in L_1^X$ one may consider $h^*(A) \in L_2^X$ which may be easier to work with, yet sufficiently granular.

Furthermore, h induces a mapping $h(C) : L_2^X \rightarrow L_2^X$ as follows: for $B \in L_2^X$ take any $A \in L_1^X$ such that $h(A) = B$ (such an A always exists due to the fact that h is surjective) and define

$$(h(C))(B) = h(C(A)) .$$

Theorem 19 *Let C be an L_1 -closure operator in X , h be a complete homomorphism of L_1 onto L_2 . Then $h(C)$ is an*

L_2 -closure operator in X and $A \mapsto h(A)$ is a complete lattice homomorphism of \mathcal{S}_C onto $\mathcal{S}_{h(C)}$.

Proof. First, we show that $h(C)$ is defined correctly. In order to do it, take any $A_1, A_2 \in L_1^X$ such that $h(A_1) = h(A_2)$. We need to show $h(C(A_1)) = h(C(A_2))$. It is easy to show that $h(E(A_1, A_2)) = 1$. Since $E(A_1, A_2) \leq E(C(A_1), C(A_2))$, monotonicity of h gives $h(E(C(A_1), C(A_2))) = 1$. As $h(E(C(A_1), C(A_2))) = E(h(C(A_1)), h(C(A_2)))$, we infer $E(h(C(A_1)), h(C(A_2))) = 1$, whence $h(C(A_1)) = h(C(A_2))$. Therefore, $h(C)$ is defined correctly.

Let $\langle \uparrow_c, \downarrow_c \rangle$ be the L -Galois connection between X and \mathcal{S}_C of Theorem 9. Consider an L_1 -relation I between X and \mathcal{S}_C defined for $x \in X$ and $A \in \mathcal{S}_C$ by

$$I(x, A) = A(x) .$$

Let $\langle \uparrow_I, \downarrow_I \rangle$ be the L_1 -Galois connection between X and \mathcal{S}_C according to Theorem 10. It is immediate that $\langle \uparrow_c, \downarrow_c \rangle = \langle \uparrow_I, \downarrow_I \rangle$. We will show that $h(C) = \uparrow_{h(I)} \downarrow_{h(I)}$, i.e. $h(C)$ is a composition of $\uparrow_{h(I)}$ and $\downarrow_{h(I)}$. To this end, take any $A \in L_1^X$. Using repeatedly the fact that h is a complete homomorphism, we get

$$h(A) \uparrow_{h(I)} \downarrow_{h(I)} = \dots = h(A \uparrow_I \downarrow_I) = h(C(A)) = (h(C))(A)$$

proving $h(C) = \uparrow_{h(I)} \downarrow_{h(I)}$. By Theorem 9, $h(C)$ is an L_2 -closure operator in X .

Finally, by [6], $\langle A, A \uparrow_I \rangle \mapsto \langle h(A), h(A \uparrow_I) \rangle$ is a complete lattice homomorphism of $\mathcal{B}_I = \{ \langle A, A \uparrow_I \rangle \mid A \in \mathcal{S}_C \}$ onto $\mathcal{B}_{h(I)} = \{ \langle A, A \uparrow_{h(I)} \rangle \mid A \in \mathcal{S}_{h(C)} \}$ (with the lattice order \leq on \mathcal{B}_I given by $\langle A, A \uparrow_I \rangle \leq \langle B, B \uparrow_I \rangle$ iff $A \subseteq B$, similarly for $\mathcal{B}_{h(I)}$). The fact that $A \mapsto h(A)$ is a complete lattice homomorphism of \mathcal{S}_C onto $\mathcal{S}_{h(C)}$ now directly follows by observing that \mathcal{B}_I is isomorphic to \mathcal{S}_C , and $\mathcal{B}_{h(I)}$ is isomorphic to $\mathcal{S}_{h(C)}$.

$\mathcal{S}_{h(C)}$ is therefore a homomorphic image of \mathcal{S}_C , i.e. a factor lattice of \mathcal{S}_C .

Remark. The fact that $h(C)$ is an L_2 -closure operator can be proved directly (without reference to Theorem 9) as follows: For any $A \in L_1^X$ we have $A \subseteq C(A)$, whence $h(A) \subseteq h(C(A)) = (h(C))(A)$ by monotonicity of h , proving (2). Furthermore, $S(A_1, A_2) \leq S(C(A_1), C(A_2))$ implies

$$\begin{aligned} S(h(A_1), h(A_2)) &= h(S(A_1, A_2)) \leq h(S(C(A_1), C(A_2))) \\ &= S(h(C(A_1)), h(C(A_2))) \\ &= S((h(C))(A_1), (h(C))(A_2)) \end{aligned}$$

proving (3). Finally,

$$\begin{aligned} (h(C))(h(A)) &= h(C(A)) = h(C(C(A))) \\ &= (h(C))(h(C(A))) = (h(C))((h(C))(h(A))) \end{aligned}$$

proving (4).

4

Representation by 2-closure operators

We show that there is a natural one-to-one correspondence between $L_{\{1\}}$ -closure operators on X and special closure operators on $X \times L$. Call a subset $A \subseteq X \times L$ (L -set)-representative if (1) for each $x \in X$ it

holds $\langle x, a \rangle \in A$ and $b \leq a$ implies $\langle x, b \rangle \in A$, and (2) for each $x \in X$ the set $\{a \in L \mid \langle x, a \rangle \in A\}$ has the greatest element.

For any L -set $A \in L^X$ put

$$[A] = \{\langle x, a \rangle \in X \times L \mid a \leq A(x)\} . \quad (20)$$

For any set $A \subseteq X \times L$ put

$$[A] = \left\{ \langle x, a \rangle \in X \times L \mid a = \bigvee_{\langle x, b \rangle \in A} b \right\} . \quad (21)$$

We have immediately the following result.

Lemma 20 *Let $A \in L^X$ be an L -set, $A' \subseteq X \times L$ be a representative set. Then (1) $[A] \subseteq X \times L$ is an representative set, (2) $[A']$ is an L -set such that (3) $A = \llbracket [A] \rrbracket$, $A' = \llbracket [A'] \rrbracket$.*

Definition 21 *A 2-closure operator D on $X \times L$ is called commutative w.r.t. $\llbracket \cdot \rrbracket$ if*

$$\llbracket [D(A)] \rrbracket = D(A) = D(\llbracket [A] \rrbracket) \quad (22)$$

holds for each $A \in X \times L$.

Remark. It is easy to verify that (22) holds iff both $\llbracket [D(A)] \rrbracket \subseteq D(A)$ and $D(\llbracket [A] \rrbracket) \subseteq D(A)$ hold.

For an operator $D : X \times L \rightarrow X \times L$ define an operator $C_D : L^X \rightarrow L^X$ by

$$C_D(A) = \llbracket [D([A])] \rrbracket$$

for $A \in L^X$. For an operator $C : L^X \rightarrow L^X$ define an operator $D_C : X \times L \rightarrow X \times L$ by

$$D_C(A) = \llbracket [C([A])] \rrbracket$$

for $A \in X \times L$.

Theorem 22 *Let C be an $L_{\{1\}}$ -closure operator on X and D be a 2-closure operator on $X \times L$ which is commutative w.r.t. $\llbracket \cdot \rrbracket$. Then (1) D_C is a 2-closure operator on X which is commutative w.r.t. $\llbracket \cdot \rrbracket$, (2) C_D is an $L_{\{1\}}$ -closure operator on X , and (3) $C = C_{D_C}$ and $D = D_{C_D}$.*

Proof. (1) Let $A, A_1, A_2 \subseteq X \times L$. We have

$$A \subseteq \llbracket [A] \rrbracket \subseteq \llbracket [C([A])] \rrbracket = D_C(A) ,$$

proving extensionality of D_C . If $A_1 \subseteq A_2$ then clearly $S(\llbracket [A_1] \rrbracket, \llbracket [A_2] \rrbracket) = 1$, hence $S(C(\llbracket [A_1] \rrbracket), C(\llbracket [A_2] \rrbracket)) = 1$, so

$$D_C(A_1) = \llbracket [C(\llbracket [A_1] \rrbracket)] \rrbracket \subseteq \llbracket [C(\llbracket [A_2] \rrbracket)] \rrbracket = D_C(A_2) ,$$

proving monotonicity of D_C .

$$D_C(D_C(A)) = \llbracket [C(\llbracket [C(\llbracket [A] \rrbracket)] \rrbracket)] \rrbracket = \llbracket [C(C(\llbracket [A] \rrbracket))] \rrbracket = \llbracket [C([A])] \rrbracket = D_C(A) ,$$

proving idempotency. Finally,

$$\llbracket [D_C(A)] \rrbracket = \llbracket [C(\llbracket [A] \rrbracket)] \rrbracket = \llbracket [C([A])] \rrbracket = D_C(A) ,$$

and

$$D_C(\llbracket [A] \rrbracket) = \llbracket [C(\llbracket [A] \rrbracket)] \rrbracket = \llbracket [C([A])] \rrbracket = D_C(A) ,$$

verifying commutativity of D_C .

(2) Let $A, A_1, A_2 \in L^X$. We have

$A = \llbracket [A] \rrbracket \subseteq \llbracket [D(\llbracket [A] \rrbracket)] \rrbracket = C_D(A)$, thus C_D is extensional. If $S(A_1, A_2) = 1$ then $\llbracket [A_1] \rrbracket \subseteq \llbracket [A_2] \rrbracket$, therefore $D(\llbracket [A_1] \rrbracket) \subseteq D(\llbracket [A_2] \rrbracket)$ and thus

$$C_D(A_1) = \llbracket [D(\llbracket [A_1] \rrbracket)] \rrbracket \subseteq \llbracket [D(\llbracket [A_2] \rrbracket)] \rrbracket = C_D(A_2) ,$$

monotonicity of C_D . Using commutativity we further get

$$\begin{aligned} C_D(C_D(A)) &= \llbracket [D(\llbracket [D(\llbracket [A] \rrbracket)] \rrbracket)] \rrbracket = \llbracket [D(D(\llbracket [A] \rrbracket))] \rrbracket \\ &= \llbracket [D(\llbracket [A] \rrbracket)] \rrbracket = C_D(A) , \end{aligned}$$

idempotency of C_D .

(3) For any $A \in L^X$ we have

$C_{D_C}(A) = \llbracket [C(\llbracket [A] \rrbracket)] \rrbracket = C(A)$. For any $A \subseteq X \times L$ we have by commutativity of D that $D_{C_D}(A) = \llbracket [D(\llbracket [A] \rrbracket)] \rrbracket = D(A)$.

Remark. Note that commutativity of D is essential in the foregoing proposition (a counterexample is easy to get).

5 Some examples

In this section we introduce two further properties of L_K -closure operators and show some examples. Call an L -set $A \in L^X$ finite if $\{x \in X \mid A(x) > 0\}$ is a finite set.

Definition 23 *An L_K closure operator C on X is called compact (finitary, or algebraic) if*

$$C(A) = \bigcup \{C(B) \mid B \in L^X, B \subseteq A, B \text{ is finite}\}$$

holds for each $A \in L^X$.

Remark. For $L = 2$ we get the compact closure operators.

Definition 24 *An L_K -closure operator C on X is called topologic if it satisfies*

$$C(A \cup B) = C(A) \cup C(B)$$

for any $A, B \in L^X$.

Remark. For $L = 2$, topologic L_K -closure operators are just closure operators of topologic spaces. The corresponding system \mathcal{S}_C consists of the closed sets of the topology.

Fuzzy subalgebras Let $A = \langle A, F \rangle$ be an algebra, i.e. A is a nonempty set and F is a system of operations on A . An L -set $B \in L^A$ is called an L -subalgebra of A if for each $f : A^n \rightarrow A$ of F and every $a_1, \dots, a_n \in A$, it holds

$$B(a_1) \otimes \dots \otimes B(a_n) \leq B(f(a_1, \dots, a_n)) .$$

Denote by $L\text{-Sub } A$ the set of all L -subalgebras of A .

Remark. For $\otimes = \wedge$, L -subalgebras and their systems are introduced and investigated in [8]. Note that 2-subalgebras coincide with the usual subalgebras.

Theorem 25 *For any algebra $A = \langle A, F \rangle$, $L\text{-Sub } A$ is an $L_{\{1\}}$ -closure system and the corresponding operator*

$C = C_{L\text{-SubA}}$ is an algebraic $L_{\{1\}}$ -closure operator. Moreover, if $\otimes = \wedge$ (i.e. L is an algebra of intuitionistic logic (Heyting algebra)), C is an L -closure operator.

Proof. It is easy to see that $L\text{-SubA}$ is closed under arbitrary intersections, hence $L\text{-SubA}$ is an $L_{\{1\}}$ -closure system and the corresponding C is an $L_{\{1\}}$ -closure operator. It remains to verify the compactness of C . To this end, let $B \in L^A$ and put

$$[B](a) = \bigvee \{ B(a_1)^{|x_1|_t} \otimes \dots \otimes B(a_n)^{|x_n|_t} \mid t \in T_n, \\ a_i \in A, t(a_1, \dots, a_n) = a \}$$

where $\alpha^k = \alpha \otimes \dots \otimes \alpha$ (k -times), T_i denotes the set of all i -ary terms of the type of A , and $|x_i|_t$ denotes the number of occurrences of the variable x_i in t . It is a matter of routine to prove by induction over rank of the term t (defined by $\text{ran}(x_i) = 0$ and $\text{ran}(f(t_1, \dot{s}, t_n)) = 1 + \max\{\text{ran}(t_1), \dot{s}, \text{ran}(t_n)\}$) that $[B] = C(B)$ which implies the compactness of C .

Let $\otimes = \wedge$. We have to prove $S(B_1, B_2) \leq S([B_1], [B_2])$ for every $B_1, B_2 \in L^A$ which holds iff for each $a \in A$ we have $[B_1](a) \wedge S(B_1, B_2) \leq [B_2](a)$. Since \wedge is idempotent we have

$$[B_1](a) \wedge S(B_1, B_2) \quad (23)$$

$$= \bigvee_{t \in T_n, t(a_1, \dots, a_n) = a} (B_1(a_1) \wedge \dots \wedge B_1(a_n)) \wedge S(B_1, B_2)$$

$$= \bigvee_{t \in T_n, t(a_1, \dots, a_n) = a} (B_1(a_1) \wedge S(B_1, B_2) \wedge \dots \wedge B_1(a_n)) \quad (24)$$

$$\wedge S(B_1, B_2) \quad (25)$$

$$\leq \bigvee_{t \in T_n, t(a_1, \dots, a_n) = a} (B_2(a_1) \wedge \dots \wedge B_2(a_n)) = [B_2](a) \quad (26)$$

Remark. Note that in general, $C = C_{L\text{-SubA}}$ is not an L -closure system. As a counterexample consider $L = \{0, \frac{1}{2}, 1\}$ with a Lukasiewicz structure, and a four-element lattice with the support $A = \{a, b, c, d\}$ with the least element a , the greatest element d , and two mutually incomparable elements b and c , i.e. its Hasse diagram is a 45°-rotated square. Let $B_1, B_2 \in L^A$ be given by $B_1(a) = 0, B_1(b) = B_1(c) = B_1(d) = 1, B_2(a) = 0, B_2(b) = B_2(c) = B_2(d) = \frac{1}{2}$. Clearly, B_2 itself is an L -subalgebra of A . On the other hand, $[B_1](a) = 1$ since $B_1(b) \otimes B_1(c) = 1 \leq [B_1](b \wedge c) = [B_1](a)$. We therefore have $S(B_1, B_2) = \frac{1}{2} \not\leq 0 = [B_1](a) \rightarrow [B_2](a) = S([B_1], [B_2])$, i.e. $[] = C$ is not an L -closure operator.

Fuzzy relational closures Let R be an L -relation on the set X , i.e. $R \in L^{X \times X}$. By a reflexive (symmetric, transitive) closure of R it is meant the least L -relation on X which is itself reflexive (symmetric, transitive) and contains R . The reflexive, symmetric, and transitive closure of R is denoted by R^r, R^s , and R^t , respectively. Recall that R is reflexive if $R(x, x) = 1$, symmetric if $R(x, y) = R(y, x)$, and transitive if $R(x, y) \otimes R(y, z) \leq R(x, z)$.

It is immediate that $R^r = R \cup_{x \in X} \{1/\langle x, x \rangle\}$, $R^s = R \cup R^{-1}$ where $R^{-1}(x, y) = R(y, x)$. Since $R \subseteq R^r, R \subseteq R^s, S(R, S) \leq S(R^r, S^r), S(R, S) \leq S(R^s, S^s)$ (both of the

inequalities are easy to verify), and $R^r = R^{rr}, R^s = R^{ss}$, we conclude that both r and s are L -closure operators on $X \times X$. Moreover, by the above description of r and s we conclude that both of them are compact as well as topologic.

To show that $R^t = \bigcup_{i=1}^{\infty} R^i = R \cup R \circ R \cup R \circ R \circ R \cup \dots$ (where $(R \circ S)(x, y) = \bigvee_{z \in X} (R(x, z) \otimes S(z, y))$) it is enough to observe that $R \subseteq \bigcup_{i=1}^{\infty} R^i$; if $R \subseteq S$ and S is transitive then $\bigcup_{i=1}^{\infty} R^i \subseteq S$; and that $\bigcup_{i=1}^{\infty} R^i$ is transitive. Since the two former are evident, we only verify the last condition. $\bigcup_{i=1}^{\infty} R^i$ is transitive iff

$$\left(\bigcup_{i=1}^{\infty} R^i \right) (x, y) \circ \left(\bigcup_{j=1}^{\infty} R^j \right) (y, z) \\ = \left(\bigvee_{i=1}^{\infty} R^i(x, y) \right) \circ \left(\bigvee_{j=1}^{\infty} R^j(y, z) \right) \\ = \bigvee_{i=1}^{\infty} \bigvee_{j=1}^{\infty} R^i(x, y) \circ R^j(y, z) \leq \bigvee_{k=1}^{\infty} R^k(x, z)$$

which holds iff for every i, j we have $R^i(x, y) \circ R^j(y, z) \leq \bigvee_{k=1}^{\infty} R^k(x, z)$. The last statement is true because $R^i(x, y) \circ R^j(y, z) \leq R^{i+j}(x, z)$. It is easy to see that for $K = \{1\}$, the conditions (2)–(4) are satisfied, hence t is an L_K -closure operator on $X \times X$. In general, t is not an L -closure operator (consider $L = \{0, \frac{1}{2}, 1\}$ with Lukasiewicz structure, $X = \{a, b, c\}$, $R(a, b) = R(b, c) = 1, S(a, b) = S(b, c) = \frac{1}{2}$, and $R(x, y) = S(x, y) = 0$ otherwise). t is compact since

$$R^t(x, y) = \bigvee_{i=1}^{\infty} R^i(x, y) \\ = \bigvee_{i=1}^{\infty} \bigvee_{x=z_1, z_2, \dots, z_{i+1}=y} R(z_1, z_2) \otimes \dots \otimes R(z_i, z_{i+1}) \\ = \bigvee_{i=1}^{\infty} \bigvee_{x=z_1, z_2, \dots, z_{i+1}=y} \{R(z_1, z_2)/\langle z_1, z_2 \rangle, \dots, \\ R(z_i, z_{i+1})/\langle z_i, z_{i+1} \rangle\}^t(x, y) .$$

As it is well-known from the classical case ($L = \{0, 1\}$), t is not topologic.

Remark. An easy inspection shows that if L is a Heyting algebra ($\otimes = \wedge$) then t is even an L -closure operator on $X \times X$.

Fuzzy concept lattices By Port-Royal logic [1], a concept is determined by its extent (the collection of all objects which fall under the concept) and its intent (the collection of all attributes which fall under the concept). For instance, the extent of the concept DOG is the collection of all dogs while its intent is the collection of all attributes common to dogs (like “to be a mammal”, “to bark” etc.). Port-Royal theory of concepts has been formalized and developed into a logico-algebraical theory of conceptual data analysis and knowledge representation by Wille et al. [14, 25]. The theory is known as formal concept analysis or theory of concept lattices. The first approach to generalize formal concept lattices from the point of view

of fuzzy logic is [11]. Later on, a general approach to the study of concept lattices from the point of view of fuzzy logic has been pursued, independently, by Pollandt [23] and the present author (see e.g. [2–6]). The theory goes as follows: Let X and Y be non-empty sets interpreted as the set of objects and the set of attributes, respectively, I be an L -relation between X and Y . The triple $\langle X, Y, I \rangle$ is called a (formal) L -context. A (formal) L -concept in $\langle X, Y, I \rangle$ is a pair $\langle A, B \rangle \in L^X \times L^Y$ (i.e. A is an L -set of attributes, B is an L -set of objects) such that B is the L -set of all attributes common to all objects from A , and A is the L -set of all objects sharing all the attributes from B . These verbal conditions translate formally as follows: Let $\uparrow_I : L^X \rightarrow L^Y$ and $\downarrow_I : L^Y \rightarrow L^X$ be defined by (18) and (19). Then $\langle A, B \rangle \in L^X \times L^Y$ is an L -concept in $\langle X, Y, I \rangle$ iff $A^\downarrow = B$ and $B^\uparrow = A$ (one easily verifies that the verbal conditions are expressed exactly by formulas (18) and (19)). The set $\mathcal{B}(X, Y, I) = \{\langle A, B \rangle \in L^X \times L^Y, A^\downarrow = B, B^\uparrow = A\}$ equipped with the partial order \leq defined by

$$\langle A_1, B_1 \rangle \leq \langle A_2, B_2 \rangle \quad \text{iff } A_1 \subseteq A_2 \quad (\text{iff } B_2 \subseteq B_1)$$

is called the L -concept lattice (fuzzy concept lattice) determined by $\langle X, Y, I \rangle$. Fuzzy concept lattice (which is in fact a complete lattice) is the basic derived structure which reveals the conceptual knowledge present in (the input data) $\langle X, Y, I \rangle$ (for more information on the principles of conceptual data analysis see [14]).

Putting in other words, Theorem 10 says that $\langle \uparrow_I, \downarrow_I \rangle$ forms a representative form of L -Galois connection. Theorem 9 implies that the composite mappings $\uparrow_I \downarrow_I : L^X \rightarrow L^X$ and $\downarrow_I \uparrow_I : L^Y \rightarrow L^Y$ are L -closure operators on X and Y , respectively, and that the sets $\{A \in L^X | A = A^{\uparrow_I \downarrow_I}\}$ and $\{B \in L^Y | B = B^{\downarrow_I \uparrow_I}\}$ are dually isomorphic L -closure systems. Moreover, Theorem 9 implies that each L -closure operator on X (L -closure system in X) can be viewed as being of the form $\uparrow_I \downarrow_I$ (or $\downarrow_I \uparrow_I$) (as the set of all extents (or intents) of an L -concept lattice) for some L -context $\langle X, Y, I \rangle$.

We now show an application of so-called Main theorem of L -concept lattices (see [3]) to provide a characterization of lattices of fixed points of L -closure operators. To this end, recall that given a complete lattice $\mathbf{V} = \langle V, \leq \rangle$, a subset $K \subseteq V$ is called infimally (supremally) dense in \mathbf{V} if each $v \in V$ is an infimum (supremum) of some subset of K .

Theorem 26 *A complete lattice $\mathbf{V} = \langle V, \leq \rangle$ is isomorphic to $\langle \mathcal{S}_C, \subseteq \rangle$ for an L -closure operator C in X iff there are mappings $\gamma : X \times L \rightarrow V$ and $\mu : \mathcal{S}_C \times L \rightarrow V$ such that $\gamma(X, L)$ is supremally dense in \mathbf{V} , $\mu(\mathcal{S}_C, L)$ is infimally dense in \mathbf{V} , and $a \otimes b \leq A(x)$ is equivalent to $\gamma(x, a) \leq \mu(A, b)$ for any $a, b \in L, x \in X, A \in \mathcal{S}_C$.*

Proof. The proof follows immediately from [3, Theorem 7], Theorems 9, and 12.

Since $\mathcal{B}(X, Y, I)$ is isomorphic to \mathcal{S}_C for $C = \uparrow_I \downarrow_I$, the above two factorization procedures for \mathcal{S}_C yield immediately factorization procedures for fuzzy concept lattices (see also [4, 6]).

6

Fuzzy closure operators and consequence relations

The concept of a closure operator is an important one from the point of view of logic. Typically, a closure operator C in a given logical calculus arises as follows: for a given set collection A of formulas, the closure $C(A)$ is defined to be the collection of all formulas provable from A . A detailed study of closure operators in the context of two-valued logic can be found in [27]. The situation is analogous in fuzzy logic. Namely, following the seminal work of Pavelka [22], provability degree of a formula from a fuzzy set of formulas is defined in fuzzy logic. Then, a fuzzy closure operator is naturally induced by a fuzzy logical calculus as follows: for a given fuzzy set A of formulas and a given formula φ , the degree to which φ belongs to the closure $C(A)$ of A is defined to be the provability degree of φ from A . For a special structure of truth values (namely, for $L = [0, 1]$ equipped with \min as the connective modeling conjunction, i.e. the standard Gödel algebra [18]), fuzzy closure operators and fuzzy consequence relations have been studied by Chakraborty (see e.g. [12]) and Gerla (see e.g. [15]). However, the study of general fuzzy closure operators and its relations to fuzzy logic is still an open goal (a paper on this topic is in preparation).

Our aim in this section is to present a general result on the relationship between fuzzy closure operators and fuzzy consequence relations. First, we show that each binary L -relation between L^X (the set of all L -sets in a given set X) and X induces in a natural way a fuzzy closure system (and the corresponding fuzzy closure operator). For an L -relation R between L^X and X , and a subset $K \subseteq L$, we say that an L -set $A \in L^X$ is R_K -closed if for any $B \in L^X$ and each $x \in X$ we have

$$S(B, A) \otimes R(B, x) \leq A(x)$$

whenever $S(B, A) \in K$.

Lemma 27 *For any $R \in L^{L^X \times X}$ and $K \subseteq L$, the set \mathcal{S}_R of all L -sets in X that are R_K -closed forms an L_K -closure system.*

Proof. By definition, we have to show that \mathcal{S}_R is closed w.r.t. S_K -intersections, i.e. we have to show that for any $A \in L^X$, $(\bigcap_{B \in \mathcal{S}_R, S(A, B) \in K} S(A, B) \rightarrow B)$ is R_K -closed. Take any $C \in L^X$ such that $S(C, \bigcap_{B \in \mathcal{S}_R, S(A, B) \in K} S(A, B) \rightarrow B) \in K$. We have to show

$$\begin{aligned} & S \left(C, \bigcap_{B \in \mathcal{S}_R, S(A, B) \in K} S(A, B) \rightarrow B \right) \otimes R(C, x) \\ & \leq \bigwedge_{B \in \mathcal{S}_R, S(A, B) \in K} S(A, B) \rightarrow B(x) \end{aligned}$$

is true for any $x \in X$. The last inequality holds iff for any $B \in L^X$ such that $S(A, B) \in K$ we have

$$\begin{aligned} S(A, B) \otimes S\left(C, \bigcap_{B \in \mathcal{S}_R, S(A, B) \in K} S(A, B) \rightarrow B\right) \otimes R(C, x) \\ \leq B(x) \end{aligned}$$

which is clearly true provided both $S(A, B) \otimes S(C, \bigcap_{B \in \mathcal{S}_R, S(A, B) \in K} S(A, B) \rightarrow B) \leq S(C, B)$ and $S(C, B) \in K$ are valid. We have

$$\begin{aligned} S(A, B) \otimes S(C, \bigcap_{B \in \mathcal{S}_R, S(A, B) \in K} S(A, B) \rightarrow B) \\ \leq S(A, B) \otimes S(C, S(A, B) \rightarrow B) \\ = S(A, B) \otimes \bigwedge_{y \in X} (C(y) \rightarrow (S(A, B) \rightarrow B(y))) \\ = S(A, B) \otimes \bigwedge_{y \in X} (S(A, B) \rightarrow (C(y) \rightarrow B(y))) \\ \leq \bigwedge_{y \in X} (S(A, B) \otimes (S(A, B) \rightarrow (C(y) \rightarrow B(y)))) \\ \leq \bigwedge_{y \in X} C(y) \rightarrow B(y) = S(C, B) . \end{aligned}$$

Since $S(A, B) \in K$ and $S(C, \bigcap_{B \in \mathcal{S}_R, S(A, B) \in K} S(A, B) \rightarrow B) \in K$, we conclude $S(C, B) \in K$.

In fact, each L_K -closure system is induced in the way described in Lemma 27 by its corresponding L_K -closure operator: Each L_K -closure operator C on X induces an L -relation $R_C \in L^{X \times X}$ by

$$R_C(A, x) = C(A)(x) .$$

Now, applying Lemma 27 to R_C we get the L_K -closure system corresponding to C :

Lemma 28 For an L_K -closure operator C on X we have $\mathcal{S}_C = \mathcal{S}_{R_C}$.

Proof. Let $A \in \mathcal{S}_C$. We show that A is R_C -closed. Let $S(B, A) \in K$. Then, by (3), $S(B, A) \leq S(C(B), C(A))$, i.e. for each $x \in X$ we have

$$S(B, A) \otimes R_C(B, x) = S(B, A) \otimes C(B)(x) \leq C(A)(x) = A(x) ,$$

whence A is R_C -closed. Conversely, if A is R_C -closed then for any B such that $S(B, A) \in K$ we have

$$S(B, A) \otimes R_C(B, x) \leq A(x) .$$

Putting $B = A$ and considering $S(A, A) = 1 \in K$, we conclude

$$S(B, A) \otimes R_C(B, x) = 1 \otimes R_C(A, x) = C(A)(x) \leq A(x) ,$$

thus $A \in \mathcal{S}_C$.

Recall (see e.g. [15]) that a (bivalent) relation \vdash between 2^X and X is called a consequence relation in (a given set) X if (i) $X \vdash \varphi$ holds for each $\varphi \in X$; (ii) $X \vdash \varphi$ and $X \subseteq Y$ imply $Y \vdash \varphi$; and (iii) $X \cup Y \vdash \varphi$ and $X \vdash \psi$ for any $\psi \in Y$ imply $X \vdash \varphi$.

Definition 29 An L -relation \vdash between L^X and X is called an L_K -consequence relation provided it satisfies

- (i) $A(\varphi) \leq (A \vdash \varphi)$
- (ii) $(A \vdash \varphi) \otimes S(A, B) \leq (B \vdash \varphi)$ whenever $(B \vdash \varphi) \in K$
- (iii) $(\bigwedge_{\varphi \in X} B(\psi) \rightarrow (A \vdash \psi)) \otimes ((A \cup B) \vdash \varphi) \leq (A \vdash \varphi)$ whenever $(\bigwedge_{\varphi \in X} B(\psi) \rightarrow (A \vdash \psi)) \in K$ for any $A, B \in L^X, \varphi \in X$

Remark. Note that condition (iii) may be equivalently replaced by (iii'):

$$\left(\bigwedge_{\varphi \in X} B(\psi) \rightarrow (A \vdash \psi) \right) \otimes (B \vdash \varphi) \leq (A \vdash \varphi)$$

whenever $(\bigwedge_{\varphi \in X} B(\psi) \rightarrow (A \vdash \psi)) \in K$. Indeed, $(B \vdash \varphi) \leq ((A \cup B) \vdash \varphi)$ by (ii), whence (iii) implies (iii'). Conversely, suppose (iii') and take $B' = A \cup B$. By $A(\psi) \leq (A \vdash \psi)$ we have

$$\begin{aligned} \bigwedge_{\psi \in X} ((A \cup B)(\psi) \rightarrow (A \vdash \psi)) \\ = \bigwedge_{\psi \in X} (A(\psi) \rightarrow (A \vdash \psi)) \wedge \bigwedge_{\psi \in X} (B(\psi) \rightarrow (A \vdash \psi)) \\ = \bigwedge_{\psi \in X} (B(\psi) \rightarrow (A \vdash \psi)) . \end{aligned}$$

Therefore,

$$\begin{aligned} \left(\bigwedge_{\psi \in X} (B(\psi) \rightarrow (A \vdash \psi)) \right) \otimes ((A \cup B) \vdash \varphi) \\ = \left(\bigwedge_{\psi \in X} ((A \cup B)(\psi) \rightarrow (A \vdash \psi)) \right) \otimes ((A \cup B) \vdash \varphi) \\ = \left(\bigwedge_{\psi \in X} (B'(\psi) \rightarrow (A \vdash \psi)) \right) \otimes (B' \vdash \varphi) \leq (A \vdash \varphi) \end{aligned}$$

by (iii').

For a mapping $C : L^X \rightarrow L^X$, define an L -relation \vdash_C between L^X and X by

$$(A \vdash_C \varphi) = C(A)(\varphi) .$$

For an L -relation \vdash between L^X and X define a mapping $C_{\vdash} : L^X \rightarrow L^X$ by

$$C_{\vdash}(A)(\varphi) = (A \vdash \varphi) .$$

Theorem 30 Let $C : L^X \rightarrow L^X$ be a mapping, $\vdash \in L^{X \times X}$ be an L -relation. Then (1) C is an L_K -closure operator iff \vdash_C is an L_K -consequence relation; (2) \vdash is an L_K -consequence relation iff C_{\vdash} is an L_K -closure operator; (3) $C = C_{\vdash_C}$ and $\vdash = \vdash_{C_{\vdash}}$.

Proof. Clearly, (3) is true. Therefore, it is sufficient to establish the “ \Rightarrow ”-parts of (1) and (2).

(1) We verify that \vdash_C is an L_K -consequence operator. (i) and (ii) are direct consequences of (2) and (3). We now

verify (iii'), a condition equivalent to (iii) (see Remark following Definition 29):

$$\left(\bigwedge_{\psi \in X} B(\psi) \rightarrow (A \vdash_C \psi) \right) \otimes (B \vdash_C \varphi) \leq (A \vdash_C \varphi)$$

is by definition equivalent to

$$\left(\bigwedge_{\psi \in X} B(\psi) \rightarrow C(A)(\psi) \right) \otimes C(B)(\varphi) \leq C(A)(\varphi)$$

which is true iff

$$\bigwedge_{\psi \in X} B(\psi) \rightarrow C(A)(\psi) \leq C(B)(\varphi) \rightarrow C(A)(\varphi) .$$

The last inequality is valid since $\bigwedge_{\psi \in X} B(\psi) \rightarrow C(A)(\psi) \in K$ implies

$$\begin{aligned} \bigwedge_{\psi \in X} B(\psi) \rightarrow C(A)(\psi) &= S(B, C(A)) \\ &\leq S(C(B), C(C(A))) \leq C(B)(\varphi) \rightarrow C(A)(\varphi) . \end{aligned}$$

(2) We check that C_+ is an L_K -closure operator. (2) and (3) are direct consequences of (i) and (ii). Putting $B = C_+(A)$, (iii') yields

$$\begin{aligned} (C_+(C_+(A))) (\varphi) &= 1 \otimes (C_+(A) \vdash \varphi) \\ &= \bigwedge_{\psi \in X} (B(\psi) \rightarrow (A \vdash \psi)) \otimes (B \vdash \varphi) \leq (A \vdash \varphi) \\ &= C_+(A)(\varphi) . \end{aligned}$$

Remark. Theorem 30 thus establishes a one-to-one correspondence between fuzzy closure operators and fuzzy consequence relations. In [15], Gerla defines graded consequence relation in X as a fuzzy relation (Gerla takes $L = [0, 1]$ and $\otimes = \min$) between 2^X (the power set of X) and X . Gerla then establishes a one-to-one correspondence between graded consequence relations and special fuzzy closure operators (Gerla deals, in our terms, with $L_{\{1\}}$ -closure operators), i.e. not all fuzzy closure operators. As it can be easily seen, the difficulty is in condition (iii) of the definition of L_K -consequence relation: particularly, the condition $(\bigwedge_{\varphi \in X} B(\psi) \rightarrow (A \vdash \psi)) \in K$ is missing in Gerla's definition. Instead, Gerla uses (iii''): $\bigwedge_{\psi \in B} (A \vdash \psi) \otimes ((A \cup B) \vdash \varphi) \leq (A \vdash \varphi)$ where $A, B \in 2^X$ (i.e. are A, B are subsets of X). Clearly, for $A, B \in 2^X$, (iii') is equivalent to $(\bigwedge_{\varphi \in X} B(\psi) \rightarrow (A \vdash \psi)) \otimes ((A \cup B) \vdash \varphi) \leq (A \vdash \varphi)$. Now, it is not true (as Gerla observes) that (iii') is satisfied by \vdash_C for any $L_{\{1\}}$ -closure operator C , i.e. (iii'') is too strong and the above natural relations do not establish a one-to-one correspondence. Theorem 30 show a way to have a one-to-one correspondence, generalizing fully the bivalent case.

References

1. **Arnauld A, Nicole P** (1972) La logique ou l'art de penser. 1662. Also in German: Die Logik oder die Kunst des Denkens. Darmstadt
2. **Bělohlávek R** (1999) Fuzzy Galois connections, Math Logic Quarterly 45(4): 497–504
3. **Bělohlávek R** (2001) Lattices of fixed points of fuzzy Galois connections, Math Logic Quarterly 47(1): 111–116
4. **Bělohlávek R** (2000) Similarity relations in concept lattices, J Logic Comput 10(6): 823–845
5. **Bělohlávek R** (2000) Fuzzy Galois connections and fuzzy concept lattices: from binary relations to conceptual structures. In: Novák V, Perfilieva I (eds) Discovering World With Fuzzy Logic. Physica-Verlag, Heidelberg, New York, pp. 462–494
6. **Bělohlávek R** Logical precision in concept lattices, J Logic Comput (to appear)
7. **Bělohlávek R** Fuzzy closure operators I, J Math Anal Appl (to appear)
8. **Biacino L, Gerla G** (1984) Closure systems and L-subalgebras, Information Sciences 33: 181–195
9. **Biacino L, Gerla G** (1996) An extension principle for closure operators, J Math Anal Appl 198: 1–24
10. **Birkhoff G** (1967) Lattice Theory, 3rd edn. AMS Coll. Publ. 25, Providence, R.I.
11. **Burusco A, Fuentes-Gonzales R** (1994) The study of L-fuzzy concept lattice, Mathware & Soft Computing 3: 209–218
12. **Chakraborty MK** (1994) Graded consequence: further studies, J Appl Non-classical Logics 5: 227–237
13. **Czédli G** (1982) Factor lattices by tolerances, Acta Sci Math (Szeged) 44: 35–42
14. **Ganter B, Wille R** (1999) Formal Concept Analysis. Mathematical Foundations, Springer-Verlag, Berlin
15. **Gerla G** (1996) Graded consequence relations and fuzzy closure operators, J Appl Non-Classical Logics 6: 369–379
16. **Gerla G** Fuzzy Logic Book to appear by Kluwer
17. **Goguen JA** (1967) L-fuzzy sets, J Math Anal Appl 18: 145–174
18. **Hájek P** (1998) Metamathematics of Fuzzy Logic. Kluwer, Dordrecht
19. **Höhle U** (1996) On the fundamentals of fuzzy set theory, J Math Anal Appl 201: 786–826
20. **Novák V, Perfilieva I, Močko J** (1999) Mathematical Principles of Fuzzy Logic. Kluwer, Dordrecht
21. **Ore O** (1944) Galois connections, Trans AMS 55: 493–513
22. **Pavelka J** (1979) On fuzzy logic I, II, III, Zeit Math Log Grungl Math 25: 45–52, 119–134, 447–464
23. **Pollandt S** (1997) Fuzzy Begriffe. Springer-Verlag, Berlin/Heidelberg
24. **Ward M, Dilworth RP** (1939) Residuated lattices, Trans AMS 45: 335–354
25. **Wille R** (1982) Restructuring lattice theory: an approach based on hierarchies of concepts. In: Rival I (ed) Ordered Sets. Reidel, Dordrecht, Boston, pp. 445–470
26. **Wille R** (1985) Complete tolerance relations of concept lattices, In: Eigenthaler G et al (eds), Contributions to General Algebra, Vol. 3. Hölder-Pichler-Tempsky, Wien, pp. 397–415
27. **Wójcicki R** (1988) Theory of Logical Calculi. Basic Theory of Consequence Operations. Kluwer (Synthese Library, Vol. 199), Dordrecht
28. **Zadeh LA** (1965) Fuzzy sets, Inf Control 8(3): 338–353
29. **Zadeh LA** (1971) Similarity relations and fuzzy orderings, Infor Sci 3: 159–176