

# A note on many valued quantum computational logics

Giuseppe Sergioli<sup>1</sup> · Antonio Ledda<sup>1</sup>

Published online: 25 July 2015  
© Springer-Verlag Berlin Heidelberg 2015

**Abstract** The standard theory of quantum computation relies on the idea that the basic information quantity is represented by a superposition of elements of the canonical basis and the notion of probability naturally follows from the Born rule. In this work we consider three valued quantum computational logics. More specifically, we will focus on the Hilbert space  $\mathbb{C}^3$ , we discuss extensions of several gates to this space and, using the notion of effect probability, we provide a characterization of its states.

**Keywords** Quantum computational logics · Qutrits

## 1 Introduction

The usual notion of *uncertainty* seems to be tightly related to an epistemic condition (Lindley 2006). A typical case: a coin was flipped, but a specified knower could not see which side of the coin faced up when it landed. It seems to be generally accepted that uncertainty deals with ignorance: a certain

predicate is uncertain, with respect to a specific knower, if the information available is not sufficient to determine its applicability.

On the other hand, *vagueness* seems to be unrelated with ignorance. Instead, this notion often refers to concepts whose extensions are lacking in clarity (Keefe and Smith 1996; Fine 1975; Williamson 2002). Natural examples of vague concepts are predicates admitting “border-line cases”, in which it is hard to sharply determine whether an object falls completely in the extension of the predicate or not. Rather, an object may possess specific properties to some extent. A successful framework for dealing with vagueness is provided by *many-valued logics* (Goguen 1969; Hájek 1998).

In the microscopic domain, appreciable overlaps between the concepts of uncertainty and vagueness are to mention. A remarkable example is the “Stern–Gerlach experiment” that shows how particles possess an intrinsic angular momentum that can assume certain discrete values only. The experiment is normally conducted using atoms or electrically neutral particles, that are treated as classical spinning dipoles; they will precess in a magnetic field because of the torque that the field exerts on the dipoles. If the magnetic field where the particles move through is not homogeneous, then the force on one end of the dipole of each particle will be slightly greater than the opposing force on the other end, so that there is a net force which deflects the trajectory of the particles. If the particles were classical spinning objects, one would expect that the distribution of their spin angular momentum vectors will be random and the spectrum will be continuous, because each particle will be deflected by a different amount, producing a density distribution on the detector screen. Instead, the particles passing through the Stern–Gerlach apparatus are deflected either up (say  $|0\rangle$ ) or down (say  $|1\rangle$ ) by a specific amount, because—how is well know nowadays—the spectrum of quantum angular momentum is discrete.

---

Communicated by M.L. Dalla Chiara, R. Giuntini, E. Negri and S. Smets.

---

The authors acknowledge the support of MIUR within the FIRB project “Structures and Dynamics of Knowledge 24 and Cognition”, Cagliari: F21J12000140001 and of RAS within the project “Modeling the Uncertainty: Quantum Theory and Imaging Processing”.

---

✉ Giuseppe Sergioli  
antonio.ledda@unica.it  
  
Antonio Ledda  
giuseppe.sergioli@gmail.com

<sup>1</sup> Università di Cagliari, Cagliari, Italy

Formally speaking and by using the common Dirac notation, we describe the state of a particle that is passing through a Stern–Gerlach apparatus as the following superposition.

$|\psi\rangle = a|0\rangle + b|1\rangle$ , where  $|a|^2$  and  $|b|^2$  are the probabilities to detect the particle with spin up or down, respectively. The state  $|\psi\rangle$  is a *pure state*, that represents a maximal piece of information that cannot be increased by any further observation. However, once a state  $|\psi\rangle$  is fixed, by its very nature, it involves an amount of uncertainty, which in this case is a property of the state not related to the observer’s knowledge. A state may also be *mixed* and it can represent a non-maximal piece of information, mathematically represented by a density operator.

We will see that, at the microscopic level, uncertainty and vagueness can be captured under several degrees of freedom rendering the two notions amenable of interactions not available in the classical world.

Useful tools for inquiring into possible interplays of vagueness and uncertainty in the quantum realm are provided by *quantum computational logics*. These logics, investigated by Maria Luisa Dalla Chiara, Roberto Giuntini and other authors, including, for example, the present papers (Bertini and Leporini 2007; Cattaneo et al. 2004a, b; Dalla Chiara et al. 2004, 2005; Giuntini et al. 2011), differ from the well known Birkhoff–von Neumann approach to quantum logic, where propositions ascribing properties are represented by projection operators—or, equivalently, by closed subspaces of a Hilbert space. In quantum computational logics meanings of sentences are no longer represented by projector operators on a Hilbert space, but by means of *quantum information quantities*: qubits, qutrits, density operators. In this framework fuzzy-like structures appear at different levels, and with different status. The aim of this paper is to discuss this further bridge between many-valued and quantum logics.

The paper is organized as follows: in Sect. 2 we provide all the basic notions necessary to render the article self-contained; in Sect. 3 we focus on the concepts of *uncertainty*, *mixedness* and *degrees of truth* as a foundational motivation in the context of many-valued quantum computational logics; in Sect. 4 we discuss some unary quantum gates in the context of standard and many-valued quantum computational logics; in Sect. 5 we introduce the notion of *effect probability*. This notion will be expedient in showing a characterization of mixed states in  $\mathbb{C}^3$ . Finally, we close the paper with some comments.

## 2 Basic notions

For the reader’s convenience, we recall in this section all the basic notions required for a complete understanding of the paper.

Consider the  $n$ -fold tensor product Hilbert space  $\otimes^n \mathbb{C}^d$ , with  $n \geq 1$  and  $d \geq 2$ .

The canonical orthonormal basis  $\mathcal{B}^{(d^n)}$  of  $\otimes^n \mathbb{C}^d$  is defined as follows:

$$\mathcal{B}^{(d^n)} = \left\{ |x_1, \dots, x_n\rangle : x_i \in \left\{ 0, \frac{1}{d-1}, \frac{2}{d-1}, \dots, 1 \right\}, \forall i \in \{1, \dots, n\} \right\}$$

where

- $|x_1, \dots, x_n\rangle$  is an abbreviation for the tensor product  $|x_1\rangle \otimes \dots \otimes |x_n\rangle$ ;
- the vector  $|\frac{i}{d-1}\rangle \in \mathbb{C}^d$  (with  $0 \leq i \leq d-1$ ) is a  $d$ -dimensional column vector with 1 in the  $(i+1)$ th-entry and 0 in all the other  $d-1$  entries.

**Definition 1 (Qudit)** A unit vector in the Hilbert space  $\mathbb{C}^d$  (with  $d \geq 2$ ) is a *qudit*. As a special case, if  $d = 2$  the unit vector is the so called *qubit*, whose extensive expression  $|\psi\rangle = a|0\rangle + b|1\rangle$  (where  $|a|^2 + |b|^2 = 1$ ) was already mentioned in the previous section.

**Definition 2 (Quregister and qumix)** A *quregister* is a unit vector in  $\otimes^n \mathbb{C}^d$  and a *qumix* (or *mixed state*) is a density operator in  $\otimes^n \mathbb{C}^d$ .

So, a vector in  $\otimes^n \mathbb{C}^d$  is a  $n$ -fold tensor product of  $d$ -dimensional vectors. Trivially, qudits are special cases of quregisters.

**Definition 3 (Truth-values of a quregister)** We say that the truth-value of a quregister  $|x_1, \dots, x_{n-1}, \frac{i}{d-1}\rangle \in \otimes^n \mathbb{C}^d$  is  $\frac{i}{d-1}$ , with  $0 \leq i \leq d-1$ .

For, the truth-value of a quregister only depends on its last component. In particular, if  $i = 0$  we say that the register is *false* and if  $i = d-1$  we say that the register is *true*.

Let us remark that the number of different truth-values over the Hilbert space  $\otimes^n \mathbb{C}^d$  is  $d$ , for any value of  $n$ .

**Definition 4 (The truth-value projectors)** A *truth-value projection* on  $\otimes^n \mathbb{C}^d$  is a projector  $P_{\frac{i}{d-1}}^{(d^n)}$  whose range is the closed subspace spanned by the set of all quregisters whose  $n$ th component is  $|\frac{i}{d-1}\rangle$ , where  $P_{\frac{i}{d-1}}^{(d^n)} = I^{(n-1)} \otimes P_{\frac{i}{d-1}}^{(d)}$  and  $0 \leq i \leq d-1$ .

In particular, the *truth-projection* of  $\otimes^n \mathbb{C}^d$  is the projection operator  $P_1^{(d^n)}$  whose range is the closed subspace spanned by the set of all true quregisters of  $\otimes^n \mathbb{C}^d$ .

As an example, let us note that the projector operators  $P_1^{(2^2)} \in \otimes^2 \mathbb{C}^2$  and  $P_1^{(4)} \in \mathbb{C}^4$ , take, respectively, the form:

$$P_1^{(2^2)} = I \otimes P_1^{(2)} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \text{ and}$$

$$P_1^{(4)} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

For any qumix  $\rho \in \otimes^n \mathbb{C}^d$  it is possible to introduce a notion of probability, according with the Born rule, as follows:

**Definition 5** ( $\frac{i}{d-1}$  - probability) Let  $\rho$  be a qumix in  $\otimes^n \mathbb{C}^d$ . The probability (Chiara et al. 2014) that  $\rho$  has the truth-value  $\frac{i}{d-1}$  (with  $0 \leq i \leq d - 1$ ) is defined by as:

$$\Pr_{\frac{i}{d-1}}^{(d)} = \text{tr} \left( P_{\frac{i}{d-1}}^{(d^n)} \rho \right),$$

where  $\text{tr}$  is the trace functional.

From an intuitive point of view,  $\Pr_{\frac{i}{d-1}}^{(d)}$  represents the probability that the information stored by the qumix  $\rho$  is the truth-value  $\frac{i}{d-1}$ .

The unitary evolution of quregisters and qumixes is dictated by quantum logical gates. *Quantum logical gates* (and the quantum operations they naturally induce) are unitary transformations that map quregisters (qumixes) in  $\otimes^n \mathbb{C}^d$  into quregisters (qumixes) in  $\otimes^n \mathbb{C}^d$ . From a foundational perspective, it could be useful to distinguish between

- semiclassical quantum gates: unitary operators that transform basis elements into basis elements;
- genuinely quantum gates: unitary operators that transform basis elements into superposition states.

In the rest of the paper, we will mostly be interested in the Hilbert space  $\mathbb{C}^3$ . We say that a unit vector in  $\mathbb{C}^3$  is a *qutrit*, a density operator (qumix) in  $\mathbb{C}^3$  is a *qutrit-density operator* and a quantum logical gate on  $\mathbb{C}^3$  is a *qutrit-gate*.

### 3 Uncertainty and degrees of truth

#### 3.1 Uncertainty

The concept of uncertainty has been a major topic of discussion for engineers, philosophers and mathematicians working on statistical theories. In particular, in the context of risk analysis (Chernoff and Moses 2012), many scholars categorize uncertainties into two main families: aleatoric and epistemic uncertainty (Kiureghiana and Ditlevsen 2009; Matthies 2007). The first family groups those deriving from

randomness in samples while the second those that stem from a lack of knowledge. Examples of aleatoric uncertainty are the occurrence of an hurricane in the Gulf of Mexico, or the height of an arbitrary individual in a certain population. On the other hand examples of epistemic uncertainty include, for instance, the global effect clouds formation have on the temperature of earth, or the nature of certain earthquake mechanisms (Paté-Cornell 1999). Aleatoric uncertainty comprises unknowns that vary each time the same experiment is performed. According with Kiureghiana and Ditlevsen (2009), aleatoric uncertainty is related to the intrinsic randomness of a phenomenon, and epistemic uncertainty is caused by a lack of knowledge.

In both cases, however, the notion of uncertainty is tightly related to a condition of the modeler. It is a concept that has to deal with *ignorance*: a given predicate is uncertain, relative to a specific modeler, when the available information is not sufficient to determine its applicability (Lindley 2006). However, once we enter the quantum domain, the concept of *uncertainty* assumes a different status, and new degrees of freedom will be available for this concept. Let us recall for a moment the classical two-slits experiment. A coherent light source is placed in front of a screen that contains two parallel slits. The wave nature of light causes the light waves passing through the two slits interfere, producing bright and dark bands on the screen placed behind the slits. The interference phenomenon is formally expressed by a superposition state  $|\psi\rangle = a|0\rangle + b|1\rangle$  where  $|\psi\rangle$  represents the state of a photon before coming up against one of the two slits,  $|0\rangle$  and  $|1\rangle$  represent the states of a photon after it passed through either the first or the second slits, respectively, and  $|a|^2$  and  $|b|^2$  are the respective probabilities. Differently from the classical case, the state  $|\psi\rangle$  is ontologically superposed: before a measurement occurs, the photon is neither in the state  $|0\rangle$  nor in  $|1\rangle$  but in a mere superposition between both. This feature of  $|\psi\rangle$  is unrelated to the ignorance of the modeler and this fact is formally expressed by the unitarity of the state  $|\psi\rangle$ , in fact  $|\psi\rangle$  is said to be a *pure* state. The concepts of *complete knowledge* (or maximal information), *unit vector* and *pure state* are, in this context, one and the same notion.

From a quantum logical point of view also, once we agree on a logical basis, uncertainty is related to proper features of the state only, without any reference to any knowledge of a possible observer (Dalla Chiara et al. 2003). Upon setting  $|0\rangle$  and  $|1\rangle$  to be our choices for the truth values “false” and “true” respectively, the superposed state  $|\psi\rangle = a|0\rangle + b|1\rangle$  expresses a *logical uncertainty* (from now on simply *uncertainty*) between the possible truth values:  $|0\rangle$  and  $|1\rangle$ , with the respective truth-probabilities  $|a|^2$  and  $|b|^2$  (with  $|a|^2 + |b|^2 = 1$ ), in accord with the Born rule.

However, it is not the case that, in general, any superposition state has non-zero truth-probability. Indeed, consider the state  $|\psi'\rangle \in \otimes^2 \mathbb{C}^2$ :

$$|\psi'\rangle = \frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|10\rangle = \left(\frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|1\rangle\right) \otimes |0\rangle. \tag{1}$$

It can be easily verified that  $|\psi'\rangle$  is a superposition state whose truth-probability is 0. On the other hand (due to the non-commutativity of the tensor product), the state  $|\psi''\rangle = |0\rangle \otimes (\frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|1\rangle)$  is both an uncertain and superposed state in  $\otimes^2\mathbb{C}^2$ .

### 3.2 Mixedness

The information a given system provides can be considered along an alternative degree of freedom: its *mixedness*.

In general, a quantum system may not be in a pure state, and the information quantity it describes may not be maximal. This could be due to, e.g., a non-complete efficiency of a preparation procedure, or, in general, to the interaction of the system with the environment, so that decoherence phenomena may arise corrupting the experimenter’s knowledge on the system. On the other hand, there are interesting processes that cannot be represented by unitary evolutions. For instance, consider the case when, at the end of a computational process, a non-unitary operation—a measurement—is applied, and the state of the system collapses into a probability distribution over pure states, namely a *proper mixed state* (mixed state for short) (Timpson and Brown 2005). A mixed state represents a *non-maximal information* on the system, that could be increased by further observations. In this case an evident epistemic feature comes into play. However, even if the concept of mixed state patently involves an observer, the property of “being mixed” should not be regarded as an exclusively epistemic condition devoid of any ontological commitment. In the microscopic context, indeed, any observation substantially modifies a state; and the property of “needing further observation to be completely known” should be considered an ontological feature of a system. As an example, quantum decoherence phenomena (Zurek 2007) consist in the loss of the coherence of the phase angles between the components of a system in a quantum superposition: an amount of information from the system vanishes into the environment (in accord with a sort of *for all practical purposes* pragmatic approach). This loss of the coherence induces a decreasing of the information on the physical system. For this reason, mixed states represent a crucial tool in quantum decoherence theory (Schlosshauer 2007; Zurek 2007).

As a generalization of the unitary case, it is possible for mixed states also to resort a notion of uncertainty that coherently generalizes the concept considered in the case of pure states. We say that a mixed state  $\rho$  represents an uncertain piece of information if its probability value is in the interval

**Table 1**

	Uncertain	Not uncertain
Mixed	$\begin{pmatrix} 1-\lambda & 0 \\ 0 & \lambda \end{pmatrix}$ with $\lambda \in (0, 1)$	$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1/2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1/2 \end{pmatrix}$
Not mixed	$\frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$

(0, 1). Let us remark that, differently from the unitary case, for an arbitrary density operator  $\rho$  the property of being *mixed* is formally equivalent to the fact that

$$\text{Tr}(\rho^2) < 1. \tag{2}$$

This corresponds to the non-unitarity of  $\rho$ : in fact,  $\text{Tr}(\rho^2) = 1$  if and only if  $\rho$  is a pure state. For any density operator  $\rho$  on  $\otimes^n\mathbb{C}^d$ , its (normalized) *linear entropy* SL, defined as  $\frac{d^n}{d^n-1}(1-\text{Tr}(\rho^2))$ , provides a measure of its degree of mixedness, or impurity. Clearly, only when  $\text{SL}(\rho) = 0$ ,  $\rho$  represents a maximal piece of information. In this case  $\rho$  is a pure state.

Although uncertainty and mixedness are proper features of a state, they are independent of each other. The following table exemplifies a few simple cases in which they can be told apart (Table 1).

### 3.3 Degrees of truth

Quantum computational logics, in the interpretation of Maria Luisa Dalla Chiara, Giampiero Cattaneo and other authors, including the present writers (Cattaneo et al. 2004a,b; Dalla Chiara et al. 2003, 2004, 2005), depart from the usual Birkhoff–von Neumann approach, where meanings of sentences are projection operators, or, equivalently, closed subspaces of a Hilbert space.

In this other approach, what really matters are quantum information units: qubits, quregisters, and, more generally, density operators on a given Hilbert space  $\otimes^n\mathbb{C}^2$  (Giuntini et al. 2011). Fuzzy-like structures have appeared in this context which have been extensively studied (Dalla Chiara et al. 2009, 2013; Giuntini et al. 2009). However, as noted in Freytes et al. 2013, this fuzzy behavior is mainly due to probabilistic features concerning uncertainty aspects of a state. Namely, the probabilities of a state to be detected either in the “false” state  $P_0^{(2^n)}$ , or in the “true” state  $P_1^{(2^n)}$ . In fact, the backdrop is an Hilbert space which is a tensor power of the space  $\mathbb{C}^2$ , where only two possible “truth-values” are available:  $|0\rangle$  and  $|1\rangle$ . Within this approach, no other truth values

are allowed. A state may be uncertain, of course; but it would be uncertain with respect to the true and to the false projector.

Under several aspects, this standpoint may be considered unduly restrictive. For, as mentioned in Bertini and Leporini (2007) and Muthukrishnan and Stroud (2000), it is absolutely conceivable to encounter cases in which a physical system may collapse into several states. A definite case in point are qutrits (Goyal et al. 2011; Tarasov 2002), where a state  $|\psi\rangle$  may have probabilities  $a_0, a_{\frac{1}{2}}, a_1$  to be detected in the basis states  $|0\rangle, |\frac{1}{2}\rangle, |1\rangle$ , respectively. Consider, for instance, the states:

$$|\psi\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$$

$$|\psi'\rangle = \left| \frac{1}{2} \right\rangle$$

in  $\mathbb{C}^2$  and  $\mathbb{C}^3$ , respectively. On the one hand,  $|\psi\rangle$  in the first equation above is an uncertain and pure state. However, only truth and false probabilities can be associated with  $|\psi\rangle$ , since  $|\psi\rangle$  is a state in  $\mathbb{C}^2$ .

On the other hand, the state  $|\psi'\rangle$  is neither uncertain, nor mixed: *it is the state*  $|\frac{1}{2}\rangle$ . This other value should not be considered as a superposition of basis elements, because it is on its own another basis element. Indeed, for any  $n$ , in the tensor power  $\otimes^n \mathbb{C}^2$ , the only possible truth values are classical; in  $\mathbb{C}^3$  a new truth-value appears:  $|\frac{1}{2}\rangle$ . The Hilbert space  $\mathbb{C}^4$  provides a neat example. The spaces  $\otimes^2 \mathbb{C}^2$  and  $\mathbb{C}^4$  are clearly isomorphic. Actually they are the same mathematical object. However, from the perspective of quantum computational logics they provide distinct logical semantics.

For instance, the vector

$$|\psi\rangle = |01\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$$

is a true register in  $\otimes^2 \mathbb{C}^2$ , because the last component of the tensor product is  $|1\rangle$ .

On the other hand, this can be easily seen to be not the case in  $\mathbb{C}^4$ , since

$$P_1^{(4)} = |1\rangle\langle 1| = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

where  $|1\rangle \in \mathbb{C}^4$ .

This fact is a consequence of the assumptions we made for the notions of true and false state.

According with Definition 3, it is evident that the outcome of a measurement in  $\otimes^2 \mathbb{C}^2$  is, in terms of truth and false




Fig. 1 .

probabilities, necessarily two-valued, since it is committed to a classical backdrop.

This won't be longer the case if a different notion of truth comes into play. For, in  $\mathbb{C}^4$ , up to the choice of the basis, there are four possible available values:  $|0\rangle, |\frac{1}{3}\rangle, |\frac{2}{3}\rangle, |1\rangle$ .

According with this idea, if we agree on the computational basis, the state

$$|\psi\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$$

would correspond to the state  $|\frac{1}{3}\rangle$ .

This brief observation is expedient to emphasize that the choice of the quantum information units determines the context in which quantum computational logics operate. If we start with a quantum information unit in  $\mathbb{C}^2$ , then every possible quantum computational logic would rely on a classical two-valued setting. On the other hand, if our choice is  $\mathbb{C}^n$ ,  $n > 2$ , new truth-values come into play.

From now on, let us focus on the Hilbert space  $\mathbb{C}^3$ .

A concrete physical system that should be necessarily represented by a state in  $\mathbb{C}^3$  is depicted in Fig. 1. Consider a 3-levels energy system in an excited state  $|\psi_1\rangle$ , whose energy is  $\epsilon_0 + \Delta\epsilon_2$ . Three events—with respective probabilities  $|a_1|^2, |a_2|^2$  and  $|a_3|^2$  (related to the respective gap of energy)—are possible:

1. the system remains in the same state  $|\psi_1\rangle$ ;
2. the system decays in the state  $|\psi_{\frac{1}{2}}\rangle$ , whose respective energy is  $\epsilon_0 + \Delta\epsilon_1$ ;
3. the system collapses to the ground state  $|\psi_0\rangle$ , whose respective energy is  $\epsilon_0$ .

The 3-levels energy system described above can be formally expressed as a qutrit—a unit vector in  $\mathbb{C}^3$ :



$$|\psi\rangle = a_0|\psi_0\rangle + a_{\frac{1}{2}}|\psi_{\frac{1}{2}}\rangle + a_1|\psi_1\rangle.$$

A natural example comes from the 3-levels laser in quantum optics (Broers et al. 1992), that is a special case of the *population inversion phenomenon* that occurs when a system of atoms exists in a non-equilibrium state such that more atoms are in an excited state than in the ground energy level. In this case, the evolution of the state should consider different possible decays from a state to another (different possible energy transactions). The probability of each decay is related with the respective gap of energy. During the decay process, a laser is emitted and its wavelength depends on the corresponding gap of energy.

As we will see in the next section, the observations above induce effective consequences in the extension of the definitions of some standard unary quantum logical gates on the Hilbert space  $\mathbb{C}^3$ .

### 4 Extending the quantum gates

In this section we discuss extensions of several well known quantum gates to the case of qutrits. These constructions exploit the fact that in  $\mathbb{C}^3$ —as well as in  $\otimes^n \mathbb{C}^3$ —the new truth value widens the usual behavior of gates in  $\mathbb{C}^2$ —as well as in  $\otimes^n \mathbb{C}^2$ —along distinct degrees of freedom. In fact, we will see that single gates in  $\mathbb{C}^2$  may admit several extensions in the case of qutrits.

#### 4.1 The negation

*Qubit case* For any  $n \geq 1$ , the *negation* on  $\otimes^n \mathbb{C}^2$  is the unitary operator  $Not^{(2^n)}$  such that, for every element  $|x_1, \dots, x_n\rangle$  of the computational basis  $\mathcal{B}^{(2^n)}$ ,

$$Not^{(2^n)}(|x_1, \dots, x_n\rangle) = |x_1, \dots, x_{n-1}\rangle \otimes |1 - x_n\rangle.$$

We have that:

$$Not^{(2^n)} = \begin{cases} \sigma_x & \text{if } n = 1; \\ I^{(n-1)} \otimes \sigma_x, & \text{otherwise,} \end{cases}$$

where  $\sigma_x := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  is the “first” Pauli matrix.

*Qutrit case* Given the usual basis  $\mathcal{B}^{(3)} = \{|0\rangle, |\frac{1}{2}\rangle, |1\rangle\}$  of  $\mathbb{C}^3$ , it is possible to define a negation  $Not_{|\frac{1}{2}\rangle}^{(3)}$  as expected by

$$Not_{|\frac{1}{2}\rangle}^{(3)}|x\rangle = |1 - x\rangle,$$

where  $x \in \{0, \frac{1}{2}, 1\}$ . We use the subscript  $|\frac{1}{2}\rangle$  to emphasize the fact that  $|\frac{1}{2}\rangle$  is a fixpoint of  $Not_{|\frac{1}{2}\rangle}^{(3)}$ , i.e.  $Not_{|\frac{1}{2}\rangle}^{(3)}|\frac{1}{2}\rangle = |\frac{1}{2}\rangle$ .

We can easily obtain the matrix form  $Not_{|\frac{1}{2}\rangle}^{(3)} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$  such that:

$$Not_{|\frac{1}{2}\rangle}^{(3)} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} c \\ b \\ a \end{pmatrix}.$$

This idea can be easily generalized to the other basis states as follows:

$$Not^{(3)}_{|0\rangle} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \text{ and } Not^{(3)}_{|1\rangle} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Let us remark that, for any  $i \in \{0, \frac{1}{2}, 1\}$ ,  $Not_{|i\rangle}^{(3)} \cdot Not_{|i\rangle}^{(3)} = I^{(3)}$ .

*Remark 1* Note that, given the computational basis  $\mathcal{B}^{(4)} = \{|0\rangle, |\frac{1}{3}\rangle, |\frac{2}{3}\rangle, |1\rangle\}$ , and  $\mathcal{B}^{(2^2)} = \{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$  the Not-like gates they induce are essentially different. Namely,

$$Not_{|\frac{1}{2}\rangle}^{(4)} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

but

$$Not^{(2^2)} = I^{(2)} \otimes Not^{(2)} \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

#### 4.2 The Hadamard gate

*Qubit case* For any  $n \geq 1$ , the *Hadamard gate* on  $\otimes^n \mathbb{C}^2$  is the linear operator  $H^{(2^n)}$  such that for every element  $|x_1, \dots, x_n\rangle$  of the computational basis  $\mathcal{B}^{(2^n)}$ :

$$H^{(2^n)}(|x_1, \dots, x_n\rangle) = |x_1, \dots, x_{n-1}\rangle \otimes \frac{1}{\sqrt{2}} ((-1)^{x_n} |x_n\rangle + |1 - x_n\rangle).$$

We have that

$$H^{(2^n)} = \begin{cases} H & \text{if } n = 1; \\ I^{n-1} \otimes H, & \text{otherwise,} \end{cases}$$

where  $H$  is the *Hadamard* matrix:

$$H = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}.$$

The basic property of  $H^{(2^n)}$  is that, for any  $|\psi\rangle \in \otimes^n \mathbb{C}^2$ :

$$H^{(2^n)}(H^{(2^n)}(|\psi\rangle)) = |\psi\rangle.$$

*Qutrit case* In Anwar et al. (2012), the following extension of the Hadamard gate for qutrits is considered

$$H^{(3)} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \frac{1}{6}(-1 + i\sqrt{3}) & -\frac{1}{6}(1 + i\sqrt{3}) \\ 1 & -\frac{1}{6}(1 + i\sqrt{3}) & \frac{1}{6}(-1 + i\sqrt{3}) \end{pmatrix}$$

as a tool in the framework of distillation protocols for fault tolerant quantum computation-precisely magic state distillation.

We state without proof the main properties of  $H^{(3)}$ :

- Lemma 1**
1. For any  $|\psi\rangle \in \mathcal{B}^{(3)}$ ,  $H^{(3)}|\psi\rangle = a|0\rangle + b|\frac{1}{2}\rangle + c|1\rangle$  s.t  $|a|^2 = |b|^2 = |c|^2 = \frac{1}{3}$ ;
  2.  $H^{(3)}$  is a genuinely quantum gate;
  3.  $H^{(3)} \cdot H^{(3)} = \text{Not}_{|0\rangle}^{(3)} \neq I$ ;
  4. for any density operator<sup>1</sup>  $\rho$  on  $\mathbb{C}^3$ ,  $\text{SL}(\rho) \neq \text{SL}(H^{(3)}(\rho))$ .

As shown in the previous lemma,  $H^{(3)}$  preserves several desirable features of  $H$ : it is a genuinely quantum gate that transforms states in the logical base  $\mathcal{B}^{(3)}$  into superposition states with uniformly distributed probabilities.

Along different lines, other possible extensions of the Hadamard gate are the *square root of the identity gates* (Giuntini et al. 2011):

$$\sqrt{I}_{|0\rangle}^{(3)} = 1 \oplus H_{(\mathbb{C}^2)} = \frac{\sqrt{2}}{2} \begin{pmatrix} \sqrt{2} & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & -1 \end{pmatrix};$$

$$\sqrt{I}_{|\frac{1}{2}\rangle}^{(3)} = \frac{\sqrt{2}}{2} \begin{pmatrix} 1 & 0 & 1 \\ 0 & \sqrt{2} & 0 \\ 1 & 0 & -1 \end{pmatrix};$$

$$\sqrt{I}_{|1\rangle}^{(3)} = H_{(\mathbb{C}^2)} \oplus 1 = \frac{\sqrt{2}}{2} \begin{pmatrix} 1 & 1 & 0 \\ 1 & -1 & 0 \\ 0 & 0 & \sqrt{2} \end{pmatrix}$$

where  $\oplus$  indicates the matrix direct sum.

Some properties of the gates above follow:

- Lemma 2** For any  $i \in \{0, \frac{1}{2}, 1\}$ :

1.  $\sqrt{I}_{|i\rangle}^{(3)}$  is a genuinely quantum qutrit-gate;
2.  $\sqrt{I}_{|i\rangle}^{(3)} \cdot \sqrt{I}_{|i\rangle}^{(3)} = I$ ;

<sup>1</sup> The extensive definition of a density operator  $\rho$  on  $\mathbb{C}^3$  is provided in the next section. When a qutrit quantum gate  $A$  is applied to a density operator  $\rho$  on  $\mathbb{C}^3$ , the evolution of  $\rho$  is given by:  $A\rho A^\dagger$ . Since no danger of confusion will be impending, for the sake of notational simplicity, from now on we write  $A(\rho)$ .

3. for any density operator  $\rho$  on  $\mathbb{C}^3$ ,  $\text{SL}(\rho) = \text{SL}(\sqrt{I}_{|i\rangle}^{(3)}(\rho))$ , as in the qubit case;

4.

$$\begin{aligned} \sqrt{I}_{|0\rangle}^{(3)} \begin{pmatrix} a \\ b \\ c \end{pmatrix} &= \begin{pmatrix} a \\ \frac{1}{\sqrt{2}}(b+c) \\ \frac{1}{\sqrt{2}}(b-c) \end{pmatrix}; \quad \sqrt{I}_{|\frac{1}{2}\rangle}^{(3)} \begin{pmatrix} a \\ b \\ c \end{pmatrix} \\ &= \begin{pmatrix} \frac{1}{\sqrt{2}}(a+c) \\ b \\ \frac{1}{\sqrt{2}}(a-c) \end{pmatrix}; \quad \sqrt{I}_{|1\rangle}^{(3)} \begin{pmatrix} a \\ b \\ c \end{pmatrix} \\ &= \begin{pmatrix} \frac{1}{\sqrt{2}}(a+b) \\ \frac{1}{\sqrt{2}}(a-b) \\ c \end{pmatrix}. \end{aligned}$$

Let us remark that, in the Hilbert space  $\mathbb{C}^3$  the Hadamard gate  $H^{(3)}$  does not behave as a square root of the identity. Instead,  $\sqrt{I}_{|i\rangle}^{(3)}$  is a square root of identity for any  $i \in \{0, \frac{1}{2}, 1\}$ .

### 4.3 The square-root of the negation

*Qubit case* For any  $n \geq 1$ , the square root of the negation (Cattaneo et al. 2009) on  $\otimes^n \mathbb{C}^2$  is the unitary operator  $\sqrt{\text{Not}}^{(2^n)}$  such that, for every element  $|x_1, \dots, x_n\rangle$  of the computational basis  $\mathcal{B}^{(2^n)}$ ,

$$\begin{aligned} \sqrt{\text{Not}}^{(2^n)}(|x_1, \dots, x_n\rangle) &= |x_1, \dots, x_{n-1}\rangle \otimes \frac{1}{2}((1+i)|x_n\rangle \\ &\quad + (1-i)|1-x_n\rangle). \end{aligned}$$

The basic property of  $\sqrt{\text{Not}}^{(2^n)}$  is the following: for any  $|\psi\rangle \in \otimes^n \mathbb{C}^2$ ,

$$\sqrt{\text{Not}}^{(2^n)}(\sqrt{\text{Not}}^{(2^n)}(|\psi\rangle)) = \text{Not}^{(2^n)}(|\psi\rangle).$$

From a logical point of view, therefore, the square root of the negation can be regarded as a kind of “tentative partial negation” that transforms precise pieces of information into maximally uncertain ones. For, we have

$$\text{Pr}_1^{(2)}(\sqrt{\text{Not}}^{(2)}(|0\rangle)) = \frac{1}{2} = \text{Pr}_1^{(2)}(\sqrt{\text{Not}}^{(1)}(|1\rangle)).$$

As noticed in Dalla Chiara et al. (2004), Lemma 17.1.11, this gate possesses no Boolean counterpart.

*Qutrit case* As in the case of the gate  $\text{Not}_{|i\rangle}^{(3)}$ ,  $i \in \{0, \frac{1}{2}, 1\}$ , once we enter in the qutrit world, several possible widening of  $\sqrt{\text{Not}}^{(2)}$  are available. Namely, given the usual basis  $\mathcal{B}^{(3)}$

of  $\mathbb{C}^3$ , it is possible to define the gates  $\sqrt{\text{Not}}_{|0\rangle}^{(3)}$ ,  $\sqrt{\text{Not}}_{|\frac{1}{2}\rangle}^{(3)}$ , and  $\sqrt{\text{Not}}_{|1\rangle}^{(3)}$  as follows:

$$\sqrt{\text{Not}}_{|0\rangle}^{(3)} = 1 \oplus \sqrt{\text{Not}}^{(2)} = \frac{1}{2} \begin{pmatrix} 2 & 0 & 0 \\ 0 & 1+i & 1-i \\ 0 & 1-i & 1+i \end{pmatrix};$$

$$\sqrt{\text{Not}}_{|\frac{1}{2}\rangle}^{(3)} = \frac{1}{2} \begin{pmatrix} 1+i & 0 & 1-i \\ 0 & 2 & 0 \\ 1-i & 0 & 1+i \end{pmatrix};$$

$$\sqrt{\text{Not}}_{|1\rangle}^{(3)} = \sqrt{\text{Not}}^{(3)} \oplus 1 = \frac{1}{2} \begin{pmatrix} 1+i & 1-i & 0 \\ 1-i & 1+i & 0 \\ 0 & 0 & 2 \end{pmatrix}.$$

Similarly to  $\sqrt{I}_{|i\rangle}^{(3)}$ , it can be seen that the gates  $\sqrt{\text{Not}}_{|i\rangle}^{(3)}$  also, act “locally” as genuinely quantum gates on two of the base vectors, leaving unchanged  $|0\rangle, |\frac{1}{2}\rangle, |1\rangle$ , respectively:

$$\sqrt{\text{Not}}_{|0\rangle}^{(3)} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} a \\ \frac{1}{2}[(1+i)b + (1-i)c] \\ \frac{1}{2}[(1-i)b + (1+i)c] \end{pmatrix};$$

$$\sqrt{\text{Not}}_{|\frac{1}{2}\rangle}^{(3)} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} \frac{1}{2}[(1+i)a + (1-i)c] \\ b \\ \frac{1}{2}[(1-i)a + (1+i)c] \end{pmatrix};$$

$$\sqrt{\text{Not}}_{|1\rangle}^{(3)} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} \frac{1}{2}[(1+i)a + (1-i)b] \\ \frac{1}{2}[(1-i)a + (1+i)b] \\ c \end{pmatrix}.$$

Some interesting properties of the gates above are summarized in the following:

**Lemma 3** For any  $i \in \{0, \frac{1}{2}, 1\}$ ,

1.  $\sqrt{\text{Not}}_{|i\rangle}^{(3)}$  is a genuinely quantum qutrit-gate;
2.  $\sqrt{\text{Not}}_{|i\rangle}^{(3)} \cdot \sqrt{\text{Not}}_{|i\rangle}^{(3)} = \text{Not}_{|i\rangle}^{(3)}$ ;
3. for any density operator  $\rho$  on  $\mathbb{C}^3$ ,  $\text{SL}(\rho) = \text{SL}(\sqrt{\text{Not}}_{|i\rangle}^{(3)}(\rho))$  (as in the qubit case).

### 5 Effect characterization of density operators on $\mathbb{C}^3$

As mentioned in Sect. 3.2, when an interaction between a system and the environment comes into play, the state of the system is represented by a qumix. It is well known that Pauli matrices  $\sigma_1, \sigma_2, \sigma_3$  and  $I$  form a basis for the set of density operators on  $\mathbb{C}^2$ , so that an arbitrary density operator  $\rho$  in  $\mathbb{C}^2$  may be represented as

$$\rho = \frac{1}{2}(I + r_1\sigma_1 + r_2\sigma_2 + r_3\sigma_3)$$

where  $r_1, r_2$  and  $r_3$  are real numbers such that  $r_1^2 + r_2^2 + r_3^2 \leq 1$ . The vector  $(r_1, r_2, r_3)$  represents the uniquely determined point in the Bloch sphere associated with  $\rho$ : the Bloch-vector of  $\rho$ . There is a one-to-one correspondence between the space of the length-1 vectors in  $\mathbb{R}^3$  and the space of the density operators in  $\mathbb{C}^2$ .

A representation of this sort can be obtained for any  $n$ -dimensional Hilbert space through generalized Pauli-matrices. In particular, in  $\mathbb{C}^3$ , Pauli matrices are generalized by Gell-Mann matrices (Schlienz and Mahler 1995).

It can be seen that any density operator  $\rho$  in  $\mathbb{C}^3$  (Goyal et al. 2011) can be written as

$$\rho = \frac{1}{3} \left( I + \sqrt{3} \sum_{i=1}^8 r_i \lambda_i \right), \tag{3}$$

where  $r_i$  are real numbers such that  $\sum_{i=1}^8 |r_i|^2 = 1$ .

Let us observe that length-1 vectors in  $\mathbb{R}^8$  and the space of the density operators in  $\mathbb{C}^3$  are not in one-to-one correspondence. Indeed, if  $r_i = 1$  for  $i = 8$  and  $r_i = 0$  otherwise, then the eigenvalues of the corresponding operator  $\rho$  in Eq. (3) would be  $\{\frac{2}{3}, \frac{2}{3}, -\frac{1}{3}\}$ , against the positivity requirement.

In this section, we propose an alternative characterization of density operators in  $\mathbb{C}^3$  through the notion of effect probability.

**Definition 6** Let  $\mathbb{H}$  be a complex Hilbert space that represents the state space of a quantum system  $S$ . The set of effects  $E(\mathbb{H})$  for  $S$  is the set of operators

$$\{A : \forall |\psi\rangle \in \mathbb{H}, 0 \leq \langle \psi | A \psi \rangle \leq 1\}.$$

Following Gudder (1998), an effect represents a yes–no measurements (for example a measurement on a Stern–Gerlach apparatus, whose outputs can only be spin up or spin down (Bush et al. 1995, I.1.2) that may be unsharp (Giuntini and Greulin 1989): sharp measurements are mere idealization, impossible in practice, where measurements are always imprecise to some degree. This unsharpness arises from the interaction between the system and the environment. For example, consider a geiger or a photon counter performing a position measurement on a one-particle quantum system. If the system is completely isolated from the environment and the detector is perfectly accurate, then it clicks if and only if the particle is detected within a certain sensitivity domain  $B \subset R^3$ . However, this situation is a limit condition, not encompassing real situations, where the system interacts with the environment and the measurement device is not perfectly accurate. In this case, an adequate calibration experiment classifies the confidence that the apparatus clicks



when the particle is in  $B$  as a real value in  $(0, 1)$ . According to Gudder (1998), this can be considered an example of a *yes-no measurement* quantum event (the particle is/is not in a certain sensitivity domain  $B$ ) to whose outcomes a probability value can be assigned.

Phenomena of this sort motivated the development of positive operator valued measure (POVM) theory, that generalises standard positive valued measure (PVM) theory (Bush et al. 1995, §1.3). POVM relies on the fact that the physical reality is described as it emerges when investigated by measuring processes, which are to be considered, themselves, as physical processes. Perhaps, the striking difference between PVM and POVM is that, while PVM considers only sharp observables, represented by hermitian operators, and projective measurements, POVM widens this setting to unsharp observables, represented by effects, to which effect-valued measurements, that generalize within this context projective measurements, are associated (Bush et al. 1995, p. 6). Indeed, to any effect valued measurement a probability measurement (effect probability) is naturally associated by the Born rule (Bush et al. 1995, (1.21)):

$$\Pr_{(E)}(\rho) = \text{tr}(E\rho), \tag{4}$$

where  $E$  and  $\rho$  are an effect and a density operator on an Hilbert space  $\mathbb{H}$ , respectively.

In the case of  $\mathbb{C}^3$ , the notion of effect probability turns out to be particularly expedient for our purposes if  $E$  in (4) is the effect:

$$E = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

In fact, using this particular effect probability we can define a three-valued quantum computational logic on  $\mathbb{C}^3$  that consistently generalizes two-valued quantum computational logic on  $\mathbb{C}^2$ . Indeed, for any  $|i\rangle$  in the computational basis,  $\Pr_{(E)}(|i\rangle) = i$ . Moreover,  $\Pr_{(E)}$  is crucial in proving the following:

**Proposition 1** *Let  $\rho$  be a density operator in  $\mathbb{C}^3$  and consider the following set of gates:  $A = \{I, \text{Not}_{|0\rangle}^{(3)}, \sqrt{I}_{|i\rangle}^{(3)}, \sqrt{\text{Not}}_{|i\rangle}^{(3)}\}_{i \in \{0, \frac{1}{2}, 1\}}$ . Then  $\rho$  is uniquely characterized by the following equations:*

1.  $\Pr_{(E)}(\rho) = \frac{1}{6}(3 - \sqrt{3}r_3 - 3r_8)$
2.  $\Pr_{(E)}(\text{Not}_{|0\rangle}^{(3)}\rho) = \frac{1}{2} - \frac{r_3}{\sqrt{3}}$
3.  $\Pr_{(E)}(\sqrt{I}_{|0\rangle}^{(3)}\rho) = \frac{1}{12}(6 - 3\sqrt{3}r_3 - 2\sqrt{3}r_6 - 3r_8)$
4.  $\Pr_{(E)}(\sqrt{I}_{|\frac{1}{2}\rangle}^{(3)}\rho) = \frac{1}{2} - \frac{r_4}{\sqrt{3}}$

5.  $\Pr_{(E)}(\sqrt{I}_{|1\rangle}^{(3)}\rho) = \frac{1}{6}(3 - \sqrt{3}r_1 - 3r_8)$
6.  $\Pr_{(E)}(\sqrt{\text{Not}}_{|0\rangle}^{(3)}\rho) = \frac{1}{12}(6 - 3\sqrt{3}r_3 - 2\sqrt{3}r_7 - 3r_8)$
7.  $\Pr_{(E)}(\sqrt{\text{Not}}_{|\frac{1}{2}\rangle}^{(3)}\rho) = \frac{1}{2} - \frac{r_5}{\sqrt{3}}$
8.  $\Pr_{(E)}(\sqrt{\text{Not}}_{|1\rangle}^{(3)}\rho) = \frac{1}{6}(3 - \sqrt{3}r_2 - 3r_8)$ .

*Proof* Simply notice that the linearly independent Eqs. (1)–(8) uniquely characterize the Bloch vector associated with  $\rho$ .

## 6 Conclusions

The aim of this paper was foundational. In particular, we tried to spell out how the concepts of uncertainty and truth-degree meet new degrees of freedom in the framework of quantum computational logics. A natural future development of the ideas in this paper would include:

- a study of binary qutrit-gates;
- a generalization of Proposition 1 to an arbitrary Hilbert space.

## References

Anwar H, Campbell ET, Browne DE (2012) Qutrit magic state distillation. [arXiv:1202.2326v2](https://arxiv.org/abs/1202.2326v2)

Bertini C, Leporini R (2007) Quantum computational finite-valued logics. *Int J Quantum Inf* 5:641–665

Broers B, van den Heuvel HBV, Noordam LD (1992) Efficient population transfer in a 3-level ladder system by frequency-swept ultrashort laser-pulses. *Phys Rev Lett* 69(14):2062–2065

Bush P, Grabowski M, Lahti PJ (1995) *Operational quantum physics*. Springer, Berlin

Cattaneo G, Dalla Chiara ML, Giuntini R, Leporini R (2004) An unsharp logic from quantum computation. *Int J Theor Phys* 43(7–8):1803–1817

Cattaneo G, Dalla Chiara ML, Giuntini R, Leporini R (2004) Quantum computational structures. *Math Slovaca* 54:87–108

Cattaneo G, Dalla Chiara ML, Giuntini R, Paoli F (2009) Quantum logic and nonclassical logics. In: Engesser K, Gabbay DM, Lehmann D (eds) *Handbook of quantum logic and quantum structures*. Elsevier, Amsterdam, pp 127–235

Chernoff H, Moses LE (2012) *Elementary decision theory*. Wiley, New York

Dalla Chiara ML, Giuntini R, Greechie R (2004) Reasoning in quantum theory. Kluwer, Dordrecht

Dalla Chiara ML, Giuntini R, Leporini R (2003) Quantum computational logics: a survey. In: Hendricks VF, Malinowski J (eds) *Trends in logic: 50 years of studia logica*. Kluwer, Dordrecht, pp 229–271

Dalla Chiara ML, Giuntini R, Leporini R (2005) Logics from quantum computation. *Int J Quantum Inf* 3(2):293–337

Dalla Chiara ML, Giuntini R, Freytes H, Ledda A, Sergioli G (2009) The algebraic structure of an approximately universal system of quantum computational gates. *Found Phys* 39(6):559–572

- Dalla Chiara ML, Giuntini R, Ledda A, Sergioli G (2013) The Toffoli–Hadamard gate system: an algebraic approach. *J Philos Log* 42(3):467–481
- Dalla Chiara ML, Giuntini R, Sergioli G (2014) Probability in quantum computation and in quantum computational logics. A survey. *Math Struct Comput Sci* 24(03):1–14
- Der Kiureghiana A, Ditlevsen O (2009) Aleatory or epistemic? Does it matter? *Struct Saf* 31(2):105–112
- Freytes H, Ledda A, Sergioli G, Giuntini R (2013) Probabilistic logics in quantum computation. In: Andersen H, Dieks D, Gonzalez WJ, Uebel T, Wheeler G (eds) *New challenges to philosophy of science. The philosophy of science in a European perspective*, vol 4. Springer, Dordrecht, pp 49–57
- Fine K (1975) Vagueness, truth and logic. *Synthese* 30(3–4):265–300
- Giuntini R, Freytes H, Ledda A, Paoli F (2009) A discriminator variety of Gödel algebras with operators arising in quantum computation. *Fuzzy Sets Syst* 160(8):1082–1098
- Giuntini R, Greulin H (1989) Toward a formal language for unsharp properties. *Found Phys* 19(7):931–945
- Giuntini R, Ledda A, Sergioli G, Paoli F (2011) Some generalisations of fuzzy structures arising in quantum computational logic. *Int J Gen Syst* 40(1):61–83
- Goguen JA (1969) The logic of inexact concepts. *Synthese* 19(3–4):325–373
- Goyal SK, Simon NB, Singh R, Simon S (2011) Geometry of the generalized Bloch sphere for qutrit. [arXiv:1111.4427](https://arxiv.org/abs/1111.4427)
- Gudder S (1998) Sharp and unsharp quantum effects. *Adv Appl Math* 20(2):169–187
- Hájek P (1998) *Metamathematics of fuzzy logic*. Kluwer, Dordrecht
- Keefe R, Smith P (1996) *Vagueness: a reader*. MIT press, Cambridge
- Lindley DV (2006) *Understanding uncertainty*. Wiley, New York
- Matthies HG (2007) Quantifying uncertainty: modern computational representation of probability and applications. In: Ibrahimbegovic A, Kozar I (eds) *Extreme man-made and natural hazards in dynamics of structures. NATO security through science series*, pp 105–135
- Muthukrishnan A, Stroud CR Jr (2000) Multi-valued logic gates for quantum computation. *Phys Rev A* 62(5):0523091–0523098
- Paté-Cornell ME (1999) Uncertainties in risk analysis: six levels of treatment. *Reliab Eng Syst Saf* 54(2–3):95–111
- Schlienz J, Mahler G (1995) Description of entanglement. *Phys Rev A* 52(6):4396–4404
- Schlosshauer M (2007) *Decoherence and the quantum-to-classical transition*, 1st edn. Springer, Berlin
- Tarasov VE (2002) Quantum computation by quantum operations on mixed states. [arXiv:quant-ph/0201033](https://arxiv.org/abs/quant-ph/0201033)
- Timpson CG, Brown HR (2005) Proper and improper separability. *Int J Quantum Inf* 3:679–690
- Williamson T (2002) *Vagueness*. Routledge, London
- Zurek WH (2007) Decoherence and the transition from quantum to classical—revisited. *Quantum Decoherence Prog Math Phys* 48:1–31