



On decidability and axiomatizability of some ordered structures

Ziba Assadi¹ · Saeed Salehi^{2,3}

Published online: 5 June 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

The ordered structures of natural, integer, rational and real numbers are studied here. It is known that the theories of these numbers in the language of order are decidable and finitely axiomatizable. Also, their theories in the language of order and addition are decidable and infinitely axiomatizable. For the language of order and multiplication, it is known that the theories of \mathbb{N} and \mathbb{Z} are not decidable (and so not axiomatizable by any computably enumerable set of sentences). By Tarski's theorem, the multiplicative ordered structure of \mathbb{R} is decidable also; here we prove this result directly and present an axiomatization. The structure of \mathbb{Q} in the language of order and multiplication seems to be missing in the literature; here we show the decidability of its theory by the technique of quantifier elimination, and after presenting an infinite axiomatization for this structure, we prove that it is not finitely axiomatizable.

Keywords Decidability · Undecidability · Completeness · Incompleteness · First-order theory · Quantifier elimination · Ordered structures

1 Introduction and preliminaries

Entscheidungsproblem, one of the fundamental problems of (mathematical) logic, asks for a single-input Boolean-output algorithm that takes a formula φ as input and outputs ‘yes’ if φ is logically valid and outputs ‘no’ otherwise. Now, we know that this problem is not (computably) solvable. One reason for this is the existence of an essentially undecidable and finitely axiomatizable theory, see, for example, Visser

(2017); for another proof see Boolos et al. (2007, Theorem 11.2). However, by Gödel's completeness theorem, the set of logically valid formulas is computably enumerable, i.e., there exists an input-free algorithm that (after running) lists all the valid formulas (and nothing else). For the structures, since their theories are complete, the story is different: the theory of a structure is either decidable or that structure is not axiomatizable [by any computably enumerable set of sentences; see, for example, Enderton (2001, Corollaries 25G and 26I) or Monk (1976, Theorem 15.2)]. For example, the additive theory of natural numbers $\langle \mathbb{N}; + \rangle$ was shown to be decidable by Presburger in 1929 (and by Skolem in 1930; see Smoryński 1991). The multiplicative theory of the natural numbers $\langle \mathbb{N}; \times \rangle$ was announced to be decidable by Skolem in 1930. Then it was expected that the theory of addition and multiplication of natural numbers would be decidable too, confirming Hilbert's program. But the world was shocked in 1931 by Gödel's incompleteness theorem which implies that the theory of $\langle \mathbb{N}; +, \times \rangle$ is undecidable (see Sect. 1.3.1). In this paper we study the theories of the sets \mathbb{N} , \mathbb{Z} , \mathbb{Q} and \mathbb{R} in the languages $\{<\}$, $\{<, +\}$ and $\{<, \times\}$; see the table below.

Communicated by A. Di Nola.

This is a part of the Ph.D. thesis of the first author written under the supervision of the second author who is partially supported by Grant N° 90030053 of the Institute for Research in Fundamental Sciences (IPM), Tehran, IRAN.

✉ Saeed Salehi
salehipour@tabrizu.ac.ir; saeed-salehi@ipm.ir
http://saeedsalehi.ir

Ziba Assadi
z_assadi.golzar@tabrizu.ac.ir

¹ Department of Mathematics, University of Tabriz, 29 Bahman Blvd., P.O. Box 51666–16471, Tabriz, Iran

² Research Institute for Fundamental Sciences (RIFS), University of Tabriz, 29 Bahman Blvd., P.O. Box 51666–16471, Tabriz, Iran

³ School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box 19395–5746, Niavaran, Tehran, Iran

	\mathbb{N}	\mathbb{Z}	\mathbb{Q}	\mathbb{R}
$\{<\}$	Theorem 3	Theorem 2	Theorem 1	Theorem 1
$\{<, +\}$	Remark 4	Theorem 5	Theorem 4	Theorem 4
$\{<, \times\}$	Section 1.3.1	Section 1.3.2	Theorem 7	Theorem 6
$\{+, \times\}$	Section 1.3.1	Section 1.3.2	Section 2.2	Section 2.1

Let us note that order is definable in the language $\{+, \times\}$ in these sets: in \mathbb{N} by $x < y \iff \exists z(z+z \neq z \wedge x+z=y)$, and in \mathbb{Z} by Lagrange's four-square theorem $x < y$ is equivalent with $\exists t, u, v, w(x \neq y \wedge x+t+t+u+u+v+v+w+w=y)$. The four-square theorem holds in \mathbb{Q} too: for any $p/q \in \mathbb{Q}^+$ we have $p/q > 0$ so $pq = a^2 + b^2 + c^2 + d^2$ for some integers a, b, c, d ; therefore, $p/q = pq/q^2 = (a/q)^2 + (b/q)^2 + (c/q)^2 + (d/q)^2$ holds. Thus, the same formula defines the order ($x < y$) in \mathbb{Q} as well. Finally, in \mathbb{R} the relation $x < y$ is equivalent with the formula $\exists z(z+z \neq z \wedge x+z \cdot z = y)$.

The decidability of $\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}$ in the languages $\{<\}$ and $\{<, +\}$ is already known. It is also known that the theories of \mathbb{N} and \mathbb{Z} in the language $\{<, \times\}$ are undecidable. The theory of \mathbb{R} in the language $\{<, \times\}$ is decidable too by Tarski's theorem (which states the decidability of $\langle \mathbb{R}; <, +, \times \rangle$). Here, we prove this directly by presenting an explicit axiomatization. Finally, the structure $\langle \mathbb{Q}; <, \times \rangle$ is studied in this paper (seemingly, for the first time). We show, by the method of quantifier elimination, that the theory of this structure is decidable. Here, the (super-)structure $\langle \mathbb{Q}; +, \times \rangle$ is not usable since it is undecidable [proved by Robinson (1949); see also Smoryński (1991, Theorem 8.30)]. On the other hand, its (sub-)structure $\langle \mathbb{Q}; \times \rangle$ is decidable [proved by Mostowski (1952); see also Salehi (2012a, 2018)]. So, the three structures $\langle \mathbb{Q}; +, \times \rangle$ and $\langle \mathbb{Q}; <, \times \rangle$ and $\langle \mathbb{Q}; \times \rangle$ are different from each other; the order relation $<$ is not definable in $\langle \mathbb{Q}; \times \rangle$ and the addition operation $+$ is not definable in $\langle \mathbb{Q}; <, \times \rangle$ (by our results; see Corollary 2). The additive structures of \mathbb{Z}, \mathbb{Q} and \mathbb{R} , and also the multiplicative structures of \mathbb{Q}^+ and \mathbb{R}^+ are abelian groups, and the theory of all abelian groups is decidable [proved by Szmielew (1949, 1955)]. Also, the additive and order structures of \mathbb{Q} and \mathbb{R} , and the multiplicative and order structures of \mathbb{Q}^+ and \mathbb{R}^+ are (regularly dense) ordered abelian groups and the theory of all regularly dense ordered abelian groups is proved to be decidable in Robinson and Zakon (1960). The additive and order structure of \mathbb{Z} does not belong to this category (as \mathbb{Z} is not dense); this structure is a Z -group (see Prestel 1986; Prestel and Delzell 2011). This paper is a continuation of the conference paper (Salehi 2012b).

1.1 The ordered structure of numbers

Definition 1 (*Ordered structure*) An *ordered structure* is a triple $\langle A; <, \mathcal{L} \rangle$ where A is a non-empty set and $<$ is a binary relation on A which satisfies the following axioms:

- (O₁) $\forall x, y(x < y \rightarrow y \not< x)$,
- (O₂) $\forall x, y, z(x < y < z \rightarrow x < z)$ and
- (O₃) $\forall x, y(x < y \vee x = y \vee y < x)$;

and \mathcal{L} is a language. \diamond

Here, \mathcal{L} could be empty, or any language, for example, $\{+\}$ or $\{\times\}$ or $\{+, \times\}$.

Definition 2 (*Various types of orders*) A linear order relation $<$ is called *dense* if it satisfies

$$(O_4) \forall x, y(x < y \rightarrow \exists z[x < z < y]).$$

An order relation $<$ is called *without endpoints* if it satisfies

$$(O_5) \forall x \exists y(x < y) \text{ and}$$

$$(O_6) \forall x \exists y(y < x).$$

A *discrete* order has the property that any element has an immediate successor (i.e., there is no other element in between them). If the successor of x is denoted by $s(x)$, then a discrete order satisfies

$$(O_7) \forall x, y(x < y \leftrightarrow s(x) < y \vee s(x) = y).$$

The successor of an integer x is $s(x) = x + 1$. \diamond

Remark 1 (The main lemma of quantifier elimination) It is known that a theory (or a structure) admits quantifier elimination if and only if every formula of the form $\exists x(\bigwedge_i \alpha_i)$ is equivalent with a quantifier-free formula, where each α_i is either an atomic formula or the negation of an atomic formula. This has been proved in, for example, Ender-ton (2001, Theorem 31F), Hinman (2005, Lemma 2.4.30), Kreisel and Krivine (1971, Theorem 1, Chapter 4), Marker (2002, Lemma 3.1.5) and Smoryński (1991, Lemma 4.1). In the presence of a linear order relation ($<$) by the equivalences $(s \neq t) \leftrightarrow (s < t \vee t < s)$ and $(s \not< t) \leftrightarrow (t \leq s)$, which follow from the axioms $\{O_1, O_2, O_3\}$ (of Definition 1), we do not need to consider the negated atomic formulas (when there is no relation symbol in the language other than $<, =$). \diamond

Convention Let \perp denote the (propositional constant of) contradiction, and \top the truth. By convention, $a \leq b$ abbreviates $a < b \vee a = b$. The symbols \times and \cdot are used interchangeably throughout the paper. For convenience, let us agree that $0^{-1} = 0$ as this does not contradict our intuition. Needless to say, x^n symbolizes $x \cdot x \cdot \dots \cdot x$ (n -times); also $x + x + \dots + x$ (n -times) is abbreviated as $n \cdot x$. \diamond

The following theorem has been proved in Marker (2002, Theorems 2.4.1 and 3.1.3). For making this paper as self-contained as possible, we present a syntactic (proof-theoretic) proof for it in "Appendix."

Theorem 1 (Axiomatizability of $\langle \mathbb{R}; < \rangle$ and $\langle \mathbb{Q}; < \rangle$) *The finite theory $\{O_1, O_2, O_3, O_4, O_5, O_6\}$ (of dense linear orders without endpoints—see Definitions 1 and 2) completely*

axiomatizes the theory of $\langle \mathbb{R}; < \rangle$ and $\langle \mathbb{Q}; < \rangle$, and so these structures are decidable. Moreover, (the theory of) those structures admit quantifier elimination.

In fact for any set A such that $\mathbb{Q} \subseteq A \subseteq \mathbb{R}$, the structure $\langle A; < \rangle$ can be completely axiomatized by the finite set of axioms $\{O_1, O_2, O_3, O_4, O_5, O_6\}$ in Definitions 1 and 2.

The theory of the structure $\langle \mathbb{Z}; < \rangle$ does not admit quantifier elimination: for example, the formula $\exists x(y < x < z)$ is not equivalent with any quantifier-free formula in the language $\{<\}$ (note that it is not equivalent with $y < z$). If we add the successor operation s to the language, then that formula will be equivalent with $s(y) < z$ and the process of quantifier elimination will go through; the proof of the following theorem appears in ‘‘Appendix.’’

Theorem 2 (Axiomatizability of $\langle \mathbb{Z}; < \rangle$) *The finite theory of discrete linear orders without endpoints, consisting of the axioms O_1, O_2, O_3, O_7 plus*

$$(O_8) \forall x \exists y (s(y) = x)$$

completely axiomatizes the order theory of the integer numbers, and so its theory is decidable. Moreover, the structure $\langle \mathbb{Z}; <, s \rangle$ admits quantifier elimination.

The structure $\langle \mathbb{N}; < \rangle$ can also be finitely axiomatized. The following theorem has been proved in Enderton (2001, Theorem 32A) so we do not present its proof in this paper.

Theorem 3 (Axiomatizability of $\langle \mathbb{N}; < \rangle$) *The finite theory consisting of the axioms $\{O_1, O_2, O_3, O_7\}$ (in Definitions 1 and 2) and also the following two axioms*

$$(O_8^0) \forall x \exists y (x \neq \mathbf{0} \rightarrow s(y) = x),$$

$$(O_9) \forall x (x \neq \mathbf{0}),$$

completely axiomatizes the order theory of the natural numbers, and so its theory is decidable. Moreover, the structure $\langle \mathbb{N}; <, s, \mathbf{0} \rangle$ admits quantifier elimination. \square

Let us note that the structure $\langle \mathbb{N}; <, s \rangle$ does not admit quantifier elimination, since, for example, $\exists x (s(x) = y)$ is not equivalent with any quantifier-free formula in the language $\{<, s\}$. However, this formula is equivalent with $\mathbf{0} < y$.

1.2 The additive ordered structures of numbers

Here we study the structures of the sets $\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}$ over the language $\{+, <\}$.

Definition 3 (Some group theory) A group is a structure $\langle G; *, e, \iota \rangle$ where $*$ is a binary operation on G , e is a constant (a special element of G) and ι is a unary operation on

G which satisfy the following axioms:

$$\forall x, y, z [x * (y * z) = (x * y) * z];$$

$$\forall x (x * e = x);$$

$$\forall x (x * \iota(x) = e).$$

It is called an *abelian* group when it also satisfies

$$\forall x, y (x * y = y * x).$$

A group is called *non-trivial* when

$$\exists x (x \neq e);$$

and it is called *divisible* when for each $n \in \mathbb{N}$ we have

$$\forall x \exists y [x = *^n(y)]$$

where $*^n(y) = y * \dots * y$ (n -times).

An *ordered group* is a group equipped with an order relation $<$ (which satisfies O_1, O_2, O_3) such that also the axiom

$$\forall x, y, z (x < y \rightarrow x * z < y * z \wedge z * x < z * y)$$

is satisfied in it. \diamond

The following has been proved in, for example, Marker (2002, Corollary 3.1.17); we also present a proof for it in ‘‘Appendix.’’

Theorem 4 (Axiomatizability of $\langle \mathbb{R}; <, + \rangle$ and $\langle \mathbb{Q}; <, + \rangle$) *The following infinite theory (of non-trivial ordered divisible abelian groups) completely axiomatizes the order and additive theory of the real and rational numbers, and so their theories are decidable. Moreover, the structures $\langle \mathbb{R}; <, +, -, \mathbf{0} \rangle$ and $\langle \mathbb{Q}; <, +, -, \mathbf{0} \rangle$ admit quantifier elimination.*

$$(O_1) \forall x, y (x < y \rightarrow y \neq x)$$

$$(O_2) \forall x, y, z (x < y < z \rightarrow x < z)$$

$$(O_3) \forall x, y (x < y \vee x = y \vee y < x)$$

$$(A_1) \forall x, y, z (x + (y + z) = (x + y) + z)$$

$$(A_2) \forall x (x + \mathbf{0} = x)$$

$$(A_3) \forall x (x + (-x) = \mathbf{0})$$

$$(A_4) \forall x, y (x + y = y + x)$$

$$(A_5) \forall x, y, z (x < y \rightarrow x + z < y + z)$$

$$(A_6) \exists y (y \neq \mathbf{0})$$

$$(A_7) \forall x \exists y (x = n \cdot y)$$

$$n \in \mathbb{N}^+$$

Remark 2 (Infinite axiomatizability) To see that the theories of $\langle \mathbb{R}; <, + \rangle$ and $\langle \mathbb{Q}; <, + \rangle$ are not finitely axiomatizable, it suffices to note that for a given natural number N , the set $\mathbb{Q}/N! = \{ \frac{m}{(N!)^k} \mid m \in \mathbb{Z}, k \in \mathbb{N} \}$ of rational numbers, where $N! = 1 \times 2 \times 3 \times \dots \times N$, is closed under addition and so satisfies the axioms $O_1, O_2, O_3, A_1, A_2, A_3, A_4, A_5, A_6$ and the finite number of the instances of the axiom A_7 (for $n = 1, \dots, N$) but does not satisfy the instance of A_7 for $n = p$ where p is a prime number larger than $N!$. \diamond

For eliminating the quantifiers of the formulas of the structure $\langle \mathbb{Z}; <, + \rangle$, we add the (binary) congruence relations $\{ \equiv_n \}_{n \geq 2}$ (modulo standard natural numbers) to the language; let us note that $a \equiv_n b$ is equivalent with $\exists x(a + n \cdot x = b)$. The following theorem has been proved, in various formats, in, for example, the books Boolos et al. (2007, Chapter 24), Enderton (2001, Theorem 32E), Hinman (2005, Corollary 2.5.18), Kreisel and Krivine (1971, Section III, Chapter 4), Marker (2002, Corollary 3.1.21), Monk (1976, Theorem 13.10) and Smoryński (1991, Section 4, Chapter III). In “Appendix,” we present a slightly different proof.

Theorem 5 (Axiomatizability of $\langle \mathbb{Z}; <, + \rangle$) *The infinite theory of non-trivial discretely ordered abelian groups with the division algorithm, that is $O_1, O_2, O_3, A_1, A_2, A_3, A_4, A_5$ and*

$$(O_7^o) \forall x, y (x < y \leftrightarrow x + \mathbf{1} \leq y)$$

$$(A_7^o) \forall x \exists y \left(\bigvee_{i < n} x = n \cdot y + \bar{i} \right) \quad n \in \mathbb{N}^+$$

where $\bar{i} = \mathbf{1} + \dots + \mathbf{1}$ (i – times)

completely axiomatizes the order and additive theory of the integer numbers, and so its theory is decidable. Moreover, the (theory of the) structure $\langle \mathbb{Z}; <, +, -, \mathbf{0}, \mathbf{1}, \{ \equiv_n \}_{n \geq 2} \rangle$ admits quantifier elimination.

Remark 3 (Infinite axiomatizability) The theory of the structure $\langle \mathbb{Z}; <, + \rangle$ cannot be axiomatized finitely, because $O_1, O_2, O_3, A_1, A_2, A_3, A_4, A_5, O_7^o$ and any finite number of the instances of A_7^o cannot prove all the instances of A_7^o . To see this take p to be a sufficiently large prime number and put $N = (p - 1)!$. Recall that $\mathbb{Q}/N = \{ m/N^k \mid m \in \mathbb{Z}, k \in \mathbb{N} \}$ is closed under addition and the operation $x \mapsto x/n$ for any $1 < n < p$. Let $\mathcal{A} = (\mathbb{Q}/N) \times \mathbb{Z}$ and define the structure $\mathfrak{A} = \langle \mathcal{A}; <_{\mathfrak{A}}, +_{\mathfrak{A}}, -_{\mathfrak{A}}, \mathbf{0}_{\mathfrak{A}}, \mathbf{1}_{\mathfrak{A}} \rangle$ by the following:

$$\begin{aligned} (<_{\mathfrak{A}}): & (a, \ell) <_{\mathfrak{A}} (b, m) \iff (a < b) \vee (a = b \wedge \ell < m); \\ (+_{\mathfrak{A}}): & (a, \ell) +_{\mathfrak{A}} (b, m) = (a + b, \ell + m); \\ (-_{\mathfrak{A}}): & -_{\mathfrak{A}}(a, \ell) = (-a, -\ell); \\ (\mathbf{0}_{\mathfrak{A}}): & \mathbf{0}_{\mathfrak{A}} = (0, 0); \\ (\mathbf{1}_{\mathfrak{A}}): & \mathbf{1}_{\mathfrak{A}} = (0, 1). \end{aligned}$$

It is straightforward to see that \mathfrak{A} satisfies the axioms $O_1, O_2, O_3, A_1, A_2, A_3, A_4, A_5$ and O_7^o ; but does not satisfy A_7^o for $n = p$ since $(1, 0) = p \cdot (a, \ell) + \bar{i}$ for any $a \in \mathbb{Q}/N, \ell \in \mathbb{Z}, i \in \mathbb{N}$ (with $i < p$) implies that $a = 1/p$ but $1/p \notin \mathbb{Q}/N$. However, \mathfrak{A} satisfies the finite number of the instances of A_7^o (for any $1 < n < p$): for any $(a, \ell) \in \mathcal{A}$ we have $a = m/N^k$ for some $m \in \mathbb{Z}, k \in \mathbb{N}$, and $\ell = nq + r$ for some q, r with $0 \leq r < n$; now, $(a, \ell) = n \cdot (m'/N^{k+1}, q) +_{\mathfrak{A}} (0, r)$ (where $m' = m \cdot (N/n) \in \mathbb{Z}$) and so $(a, \ell) = n \cdot (m'/N^{k+1}, q) +_{\mathfrak{A}} \bar{r}$ (where $\bar{r} = \mathbf{1}_{\mathfrak{A}} +_{\mathfrak{A}} \dots +_{\mathfrak{A}} \mathbf{1}_{\mathfrak{A}}$ for r times). \square

Remark 4 ($\langle \mathbb{N}; <, + \rangle$) Since \mathbb{N} is definable in the structure $\langle \mathbb{Z}; <, + \rangle$ by “ $x \in \mathbb{N}'' \iff \exists y(y + y = x \wedge y \leq x)$,” we do not study the theory of the structure $\langle \mathbb{N}; <, + \rangle$ separately (see Enderton 2001, Theorem 32E). In fact the decidability of $\langle \mathbb{Z}; <, + \rangle$ implies the decidability of $\langle \mathbb{N}; <, + \rangle$: relativization $\psi^{\mathbb{N}}$ of a $\{ <, + \}$ -formula ψ resulted from substituting any subformula of the form $\forall x \theta(x)$ by $\forall x [“x \in \mathbb{N}'' \rightarrow \theta(x)]$ and $\exists x \theta(x)$ by $\exists x [“x \in \mathbb{N}'' \wedge \theta(x)]$ has the following property: $\langle \mathbb{N}; <, + \rangle \models \psi \iff \langle \mathbb{Z}; <, + \rangle \models \psi^{\mathbb{N}}$. \diamond

1.3 The multiplicative ordered structures of numbers

Finally, we consider the theories of the number sets $\mathbb{N}, \mathbb{Z}, \mathbb{R}$ and \mathbb{Q} in the language $\{ <, \times \}$.

1.3.1 Natural numbers with order and multiplication

The theory of the structure $\langle \mathbb{N}; <, \times \rangle$ is not decidable (and so no computably enumerable set of sentences can axiomatize this structure). This is because:

- The addition operation is definable in $\langle \mathbb{N}; <, \times \rangle$, since
 - the successor operation s is definable from order:

$$y = s(x) \iff x < y \wedge \neg \exists z(x < z < y),$$

- and the addition operation is definable from the successor and multiplication operations:

$$z = x + y \iff [\neg \exists u(s(u) = z) \wedge x = y = z] \vee [\exists u(s(u) = z) \wedge s(z \cdot x) \cdot s(z \cdot y) = s(z \cdot z \cdot s(x \cdot y))].$$

This identity was first introduced by Robinson (1949); also see, for example, Boolos et al. (2007, Chapter 24) or Enderton (2001, Exercise 2 on page 281).

- Thus, the structure $\langle \mathbb{N}; <, \times \rangle$ can interpret the structure $\langle \mathbb{N}; +, \times \rangle$ whose theory is undecidable [see, for example, Boolos et al. (2007, Theorem 17.4), Enderton (2001, Corollary 35A), Hinman (2005, Theorem 4.1.7), Monk (1976, Chapter 15) or Smoryński (1991, Corollary 6.4 in Chapter III)].

1.3.2 Integer numbers with order and multiplication

The undecidability of the theory of the structure $\langle \mathbb{N}; +, \times \rangle$ also implies the undecidability of the theories of the structures $\langle \mathbb{Z}; +, \times \rangle$ and $\langle \mathbb{Z}; <, \times \rangle$ as follows:

- By Lagrange’s four-square theorem [see, for example, Monk (1976, Theorem 16.6)] \mathbb{N} is definable in $\langle \mathbb{Z}; +, \times \rangle$, and so $\langle \mathbb{Z}; +, \times \rangle$ has an undecidable theory [see, for example, Monk (1976, Theorem 16.7) or Smoryński (1991, Corollary 8.29 in Chapter III)].
- The following numbers and operations are definable in the structure $\langle \mathbb{Z}; <, \times \rangle$:
 - The number zero: $u = 0 \iff \forall x(x \cdot u = u)$.
 - The number one: $u = 1 \iff \forall x(x \cdot u = x)$.
 - The number -1 : $u = -1 \iff u \cdot u = 1 \wedge u \neq 1$.
 - The additive inverse: $y = -x \iff y = (-1) \cdot x$.
 - The successor: $y = s(x) \iff x < y \wedge \nexists z(x < z < y)$.
 - The addition: $z = x + y \iff [z = 0 \wedge y = -x] \vee [z \neq 0 \wedge s(z \cdot x) \cdot s(z \cdot y) = s(z \cdot z \cdot s(x \cdot y))]$.

There is another beautiful definition for $+$ in terms of s and \times in \mathbb{Z} on page 187 of Hinman (2005):

$$z = x + y \iff [z \cdot s(z) = z \wedge s(x \cdot y) = s(x) \cdot s(y)] \vee [z \cdot s(z) \neq z \wedge s(z \cdot x) \cdot s(z \cdot y) = s(z \cdot z \cdot s(x \cdot y))].$$

- Whence, the structure $\langle \mathbb{Z}; <, \times \rangle$ can interpret the undecidable structure $\langle \mathbb{Z}; +, \times \rangle$.

2 Reals and rationals with order and multiplication

2.1 Real numbers with order and multiplication

The structure $\langle \mathbb{R}; <, \times \rangle$ is decidable, since by a theorem of Tarski the structure $\langle \mathbb{R}; <, +, \times \rangle$ can be completely axiomatized by the theory of *real closed ordered fields*, and so has a decidable theory; see, for example, Kreisel and Krivine (1971, Theorem 7, Chapter 4), Marker (2002, Theorem 3.3.15) or Monk (1976, Theorem 21.36). Here, we prove the decidability of the theory of $\langle \mathbb{R}; <, \times \rangle$ directly (without using Tarski’s theorem) and provide an explicit axiomatization for it. Before that let us make a little note about the theory $\langle \mathbb{R}^+; <, \times \rangle$ (of the positive real numbers) which is isomorphic to $\langle \mathbb{R}; <, + \rangle$ by the mapping $x \mapsto \log(x)$. Thus, we have the following immediate corollary of Theorem 4:

Proposition 1 (Axiomatizability of $\langle \mathbb{R}^+; <, \times \rangle$) *The following infinite theory completely axiomatizes the order and multiplicative theory of non-trivial ordered divisible abelian*

groups) completely axiomatizes the order and multiplicative theory of the positive real numbers, and so its theory is decidable. Moreover, the structure $\langle \mathbb{R}^+; <, \times, \square^{-1}, \mathbf{1} \rangle$ admits quantifier elimination.

- $(O_1) \forall x, y(x < y \rightarrow y \not< x)$
- $(O_2) \forall x, y, z(x < y < z \rightarrow x < z)$
- $(O_3) \forall x, y(x < y \vee x = y \vee y < x)$
- $(M_1) \forall x, y, z(x \cdot (y \cdot z) = (x \cdot y) \cdot z)$
- $(M_2) \forall x(x \cdot \mathbf{1} = x)$
- $(M_3) \forall x(x \cdot x^{-1} = \mathbf{1})$
- $(M_4) \forall x, y(x \cdot y = y \cdot x)$
- $(M_5) \forall x, y, z(x < y \rightarrow x \cdot z < y \cdot z)$
- $(M_6) \exists y(y \neq \mathbf{1})$
- $(M_7) \forall x \exists y(x = y^n) \quad n \geq 2$

Proof For the infinite axiomatizability it suffices to note that for a sufficiently large N the set $\{2^{m \cdot (N!)^k} \mid m \in \mathbb{Z}, k \in \mathbb{N}\}$ of positive real numbers (cf. Remark 2) satisfies all the axioms $(O_1, O_2, O_3, M_1, M_2, M_3, M_4, M_5, M_6)$ and finitely many instances of the axiom M_7 (for $n \leq N$) but not all the instances of M_7 (for example, when $n = p$ is a prime larger than $N!$). \square

Theorem 6 (Axiomatizability of $\langle \mathbb{R}; <, \times \rangle$) *The following infinite theory completely axiomatizes the order and multiplicative theory of the real numbers, and so its theory is decidable. Moreover, the structure $\langle \mathbb{R}; <, \times, \square^{-1}, -\mathbf{1}, \mathbf{0}, \mathbf{1} \rangle$ admits quantifier elimination.*

- $(O_1) \forall x, y(x < y \rightarrow y \not< x)$
- $(O_2) \forall x, y, z(x < y < z \rightarrow x < z)$
- $(O_3) \forall x, y(x < y \vee x = y \vee y < x)$
- $(M_1) \forall x, y, z(x \cdot (y \cdot z) = (x \cdot y) \cdot z)$
- $(M_2^0) \forall x(x \cdot \mathbf{1} = x \wedge x \cdot \mathbf{0} = \mathbf{0} = \mathbf{0}^{-1})$
- $(M_3^0) \forall x(x \neq \mathbf{0} \rightarrow x \cdot x^{-1} = \mathbf{1})$
- $(M_4) \forall x, y(x \cdot y = y \cdot x)$
- $(M_5^0) \forall x, y, z(x < y \wedge \mathbf{0} < z \rightarrow x \cdot z < y \cdot z)$
- $(M_5^1) \forall x, y, z(x < y \wedge z < \mathbf{0} \rightarrow y \cdot z < x \cdot z)$
- $(M_6^0) \exists y(-\mathbf{1} < \mathbf{0} < \mathbf{1} < y)$
- $(M_7^0) \forall x \exists y(x = y^{2n+1})$
- $(M_8) \forall x(x^{2n} = \mathbf{1} \iff x = \mathbf{1} \vee x = -\mathbf{1})$
- $(M_9) \forall x(\mathbf{0} < x \iff \exists y[y \neq \mathbf{0} \wedge x = y^2])$

Proof We have $(x < \mathbf{0}) \leftrightarrow (\mathbf{0} < -x)$ by M_5^*, M_2^*, M_6^* and M_8 , where $-x = (-\mathbf{1}) \cdot x$. Whence, for any quantifier-free formula η we have $\exists x \eta(x) \equiv \exists x > \mathbf{0} \eta(x) \vee \eta(\mathbf{0}) \vee \exists y > \mathbf{0} \eta(-y)$. Also, if z is another variable in η then $\eta(x, z)$ is equivalent with $[\mathbf{0} < z \wedge \eta(x, z)] \vee \eta(x, \mathbf{0}) \vee [\mathbf{0} < -z \wedge \eta(x, z)]$. For the last disjunct, if we let $z' = -z$ then $\mathbf{0} < -z \wedge \eta(x, z)$ will be $\mathbf{0} < z' \wedge \eta(x, -z')$. Thus, by introducing the constants $\mathbf{0}$ and $-\mathbf{1}$ (and renaming the variables if necessary), we can assume that all the variables of a quantifier-free formula are positive. Now, the process of eliminating the quantifier of the formula $\exists x \eta(x)$, where η is the conjunction of some atomic formulas (cf. Remark 1) goes as follows: we first eliminate the constants $\mathbf{0}$ and $-\mathbf{1}$ and then reduce the desired conclusion to Proposition 1. For the first part, we simplify terms so that each term is either positive (all the variables are positive) or equals to $\mathbf{0}$ or is the negation of a positive term (is $-t$ for some positive term t). Then by replacing $\mathbf{0} = \mathbf{0}$ with \top and $\mathbf{0} < \mathbf{0}$ with \perp we can assume that $\mathbf{0}$ appears at most once in any atomic formula; also $-\mathbf{1}$ appears at most once since $-t = -s$ is equivalent with $t = s$ and $-t < -s$ with $s < t$. Now, we can eliminate the constant $-\mathbf{1}$ by replacing the atomic formulas $-t = s, t = -s$ and $t < -s$ by \perp and $-t < s$ by \top for positive or zero terms t, s (note that $-\mathbf{0} = \mathbf{0}$ by M_2^*). Also the constant $\mathbf{0}$ can be eliminated by replacing $\mathbf{0} < t$ with \top and $t < \mathbf{0}$ and $t = \mathbf{0}$ (also $\mathbf{0} = t$) with \perp for positive terms t . Thus, we get a formula whose all variables are positive, and so we are in the realm of \mathbb{R}^+ . Finally, for the second part we have the equivalence of thus resulted formula with a quantifier-free formula by Proposition 1 provided that the relativized form of the axioms $O_1, O_2, O_3, M_1, M_2, M_3, M_4, M_5, M_6$ and M_7 to \mathbb{R}^+ can be proved from the axioms $O_1, O_2, O_3, M_1, M_2^*, M_3^*, M_4, M_5^*, M_6^*, M_7^*, M_8$, and M_9 . We need to consider M_6 and M_7 only, when relativized to \mathbb{R}^+ , i.e., $\exists y(\mathbf{0} < y \wedge y \neq \mathbf{1})$ and $\forall x \exists y[\mathbf{0} < x \rightarrow \mathbf{0} < y \wedge x = y^n]$. The relativization of M_6 immediately follows from M_6^* . For the relativization of M_7 take any $a > \mathbf{0}$, and any $n \in \mathbb{N}$. Write $n = 2^k(2m + 1)$; by M_7^* there exists some c such that $c^{2m+1} = a$, and by M_2^* and M_5^* we should have $c > \mathbf{0}$. Now, by using M_9 for k times there must exist some b such that $b^{2^k} = c$ and we can have $b > \mathbf{0}$ (since otherwise we can take $-b$ instead of b). Now, we have $b^{2^k(2m+1)} = c^{2m+1} = a$ and so $a = b^n$. \square

That no finite set of axioms can completely axiomatize the theory of $(\mathbb{R}; <, \times)$ can be seen from the fact that the set $\{0\} \cup \{-2^{m \cdot (N!)^{-k}}, 2^{m \cdot (N!)^{-k}} \mid m \in \mathbb{Z}, k \in \mathbb{N}\}$ of real numbers, for some $N > 2$, satisfies all the axioms of Theorem 6 except M_7^* ; however it satisfies a finite number of its instances (when $2n + 1 \leq N$) but not all the instances (e.g., when $2n + 1$ is a prime greater than $N!$) of M_7^* (cf. the proof of Proposition 1 and Remark 2).

2.2 Rational numbers with order and multiplication

The technique of the proof of Theorem 6 enables us to consider first the multiplicative and order structure of the positive rational numbers $(\mathbb{Q}^+; <, \times)$. The formula $\exists x(y = x^n)$ (for $n > 1$) is not equivalent with any quantifier-free formula in $(\mathbb{Q}^+; <, \times)$; so let us introduce the following notation.

Definition 4 (\mathfrak{R}) Let $\mathfrak{R}_n(y)$ be the formula $\exists x(y = x^n)$, stating that “ y is the n th power of a number” (for $n > 1$). \diamond

Now we can introduce our candidate axiomatization for the theory of the structure $(\mathbb{Q}^+; <, \times)$.

Definition 5 (TQ) Let TQ be the theory axiomatized by the axioms $O_1, O_2, O_3, M_1, M_2, M_3, M_4, M_5$ and M_6 of Proposition 1 plus the following two axiom schemes:

$$(M_{10}) \forall x, z \exists y(x < z \rightarrow x < y^n < z), \text{ and}$$

$$(M_{11}) \forall \{x_j\}_{j < q} \exists y \forall z \bigwedge_{m_j \nmid n(j < q)} (y^n \cdot x_j \neq z^{m_j});$$

for each $n \geq 1$ (and $m_j > 1$). \diamond

Some explanations on the new axioms M_{10} and M_{11} are in order. The axiom M_{10} , interpreted in \mathbb{Q}^+ , states that \mathbb{Q}^+ is dense not only in itself but also in the radicals of its elements (or more generally in \mathbb{R}^+ : for any $x, z \in \mathbb{Q}^+$ there exists some $y \in \mathbb{Q}^+$ that satisfies $\sqrt[n]{x} < y < \sqrt[n]{z}$). The axiom M_{11} , interpreted in \mathbb{Q}^+ , is actually equivalent with the fact that for any sequences $x_1, \dots, x_q \in \mathbb{Q}^+$ and $m_1, \dots, m_q \in \mathbb{N}^+$ none of which divides n (in symbols $m_j \nmid n$), there exists some $y \in \mathbb{Q}^+$ such that $\bigwedge_j \neg \mathfrak{R}_{m_j}(y^n \cdot x_j)$. This axiom is not true in \mathbb{R}^+ (while M_{10} is true in it) and to see that why M_{11} is true in \mathbb{Q}^+ it suffices to note that for given x_1, \dots, x_q one can take y to be a prime number which does not appear in the unique factorization (of the numerators and denominators of the reduced forms) of any of x_j 's. In this case $y^n \cdot x_j$ can be an m_j 's power (of a rational number) only when m_j divides n . The condition $m_j \nmid n$ is necessary, since otherwise (if $m_j \mid n$ and) if x_j happens to satisfy $\mathfrak{R}_{m_j}(x_j)$ then no y can satisfy the relation $\neg \mathfrak{R}_{m_j}(y^n \cdot x_j)$.

We now show that TQ completely axiomatizes the theory of the structure $(\mathbb{Q}^+; <, \times, \square^{-1}, \mathbf{1}, \{\mathfrak{R}_n\}_{n > 1})$ and moreover this structure admits quantifier elimination; thus, the theory of the structure $(\mathbb{Q}^+; <, \times)$ is decidable. For that, we will need the following lemmas.

Lemma 1 For any $x \in \mathbb{Q}^+$ and any natural $n_1, n_2 > 1$,

$$\mathfrak{R}_{n_1}(x) \wedge \mathfrak{R}_{n_2}(x) \iff \mathfrak{R}_n(x)$$

where n is the least common multiplier of n_1 and n_2 .

Proof The \Leftarrow part is straightforward; for the \Rightarrow direction suppose that $x = y^{n_1} = z^{n_2}$. By Bézout's Identity there are

some $c_1, c_2 \in \mathbb{Z}$ such that $c_1n/n_1 + c_2n/n_2 = 1$; therefore, $x = x^{c_1n/n_1} \cdot x^{c_2n/n_2} = y^{c_1n} \cdot z^{c_2n} = (y^{c_1}z^{c_2})^n$. \square

Lemma 2 For natural numbers $\{n_i\}_{i < p}$ with $n_i > 1$ and positive rational numbers $\{t_i\}_{i < p}$ and x ,

$$\bigwedge_{i < p} \mathfrak{R}_{n_i}(x \cdot t_i) \iff \mathfrak{R}_n(x \cdot \beta) \wedge \bigwedge_{i \neq j} \mathfrak{R}_{d_{i,j}}(t_i \cdot t_j^{-1})$$

where n is the least common multiplier of n_i 's, $d_{i,j}$ is the greatest common divisor of n_i and n_j (for each $i \neq j$) and $\beta = \prod_{i < p} t_i^{c_i(n/n_i)}$ in which c_i 's satisfy $\sum_{i < p} c_i(n/n_i) = 1$.

Proof For t_i 's, n_i 's, c_i 's, $d_{i,j}$'s and n as given above, we show that the relation $\mathfrak{R}_{n_k}(t_k \cdot \beta^{-1})$ holds for each fixed $k < p$ when $\bigwedge_{i \neq j} \mathfrak{R}_{d_{i,j}}(t_i \cdot t_j^{-1})$ holds. Let $m_{k,i}$ be the least common multiplier of n_k and n_i (which is a divisor of n then). Let us note that $d_{k,i}/n_i = n_k/m_{k,i}$. Since $\mathfrak{R}_{d_{k,i}}(t_k \cdot t_i^{-1})$ there should exist some $w_{k,i}$'s (for $i \neq k$) such that $t_k \cdot t_i^{-1} = w_{k,i}^{d_{k,i}}$. Now, the relation $\mathfrak{R}_{n_k}(t_k \cdot \beta^{-1})$ follows from the following identities:

$$\begin{aligned} t_k \cdot \beta^{-1} &= t_k^{\sum_i c_i(n/n_i)} \cdot \prod_i t_i^{-c_i(n/n_i)} \\ &= \prod_{i \neq k} (t_k \cdot t_i^{-1})^{c_i(n/n_i)} \\ &= \prod_{i \neq k} (w_{k,i}^{d_{k,i}})^{c_i(n/n_i)} \\ &= \prod_{i \neq k} w_{k,i}^{c_i \cdot n_k(n/m_{k,i})} \\ &= (\prod_{i \neq k} w_{k,i}^{c_i(n/m_{k,i})})^{n_k}. \end{aligned}$$

- (\Rightarrow): The relations $\mathfrak{R}_{n_i}(x \cdot t_i)$ and $\mathfrak{R}_{n_j}(x \cdot t_j)$ immediately imply that $\mathfrak{R}_{d_{i,j}}(x \cdot t_i)$ and $\mathfrak{R}_{d_{i,j}}(x \cdot t_j)$ and so $\mathfrak{R}_{d_{i,j}}(t_i \cdot t_j^{-1})$. For showing $\mathfrak{R}_n(x \cdot \beta)$ it suffices, by Lemma 1, to show that $\mathfrak{R}_{n_i}(x \cdot \beta)$ holds for each $i < p$. This immediately follows from $\mathfrak{R}_{n_i}(t_i \cdot \beta^{-1})$ which was proved above, and the assumption $\mathfrak{R}_{n_i}(x \cdot t_i)$.
- (\Leftarrow): From the first part of the proof we have $\mathfrak{R}_{n_k}(t_k \cdot \beta^{-1})$ for each $k < p$; now by $\mathfrak{R}_n(x \cdot \beta)$ we have $\mathfrak{R}_{n_k}(x \cdot \beta)$ and so $\mathfrak{R}_{n_k}(x \cdot t_k)$ for each $k < p$. \square

Let us note that Lemmas 1 and 2 are provable in TQ. The idea of the proof of Lemma 2 is taken from (Ore 1952).

Lemma 3 The following sentences are provable in TQ for any $n > 1$:

$$\begin{aligned} &\forall u \exists y [\mathfrak{R}_n(y \cdot u)], \\ &\forall x, u \exists y [x < y \wedge \mathfrak{R}_n(y \cdot u)], \\ &\forall z, u \exists y [y < z \wedge \mathfrak{R}_n(y \cdot u)] \text{ and} \\ &\forall x, z, u \exists y [x < z \rightarrow x < y < z \wedge \mathfrak{R}_n(y \cdot u)]. \end{aligned}$$

Proof We show the last formula only. By M_{10} (of Definition 5) there exists some v such that $x \cdot u < v^n < z \cdot u$. Then for $y = v^n \cdot u^{-1}$ we will have $x < y < z$ and $\mathfrak{R}_n(y \cdot u)$. \square

Lemma 4 The following sentences are provable in TQ for any $\{m_j > 1\}_{j < q}$:

$$\begin{aligned} &\forall \{x_j\}_{j < q} \exists y [\bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(y \cdot x_j)], \\ &\forall \{x_j\}_{j < q}, u \exists y [u < y \wedge \bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(y \cdot x_j)], \\ &\forall \{x_j\}_{j < q}, v \exists y [y < v \wedge \bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(y \cdot x_j)] \text{ and} \\ &\forall \{x_j\}_{j < q}, u, v \exists y [u < v \rightarrow u < y < v \wedge \\ &\quad \bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(y \cdot x_j)]. \end{aligned}$$

Proof The first sentence is an immediate consequence of M_{11} (of Definition 5) for $n = 1$. We show the last sentence. There exists γ , by M_{11} , such that $\bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(\gamma \cdot x_j)$. Let $M = \prod_j m_j$; by M_{10} there exists δ such that $u \cdot \gamma^{-1} < \delta^M < v \cdot \gamma^{-1}$. Now for $y = \gamma \cdot \delta^M$ we have $u < y < v$ and $\bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(y \cdot x_j)$ since if (otherwise) we had $\mathfrak{R}_{m_j}(y \cdot x_j)$ then $\mathfrak{R}_{m_j}(\gamma \cdot \delta^M \cdot x_j)$ and so $\mathfrak{R}_{m_j}(\gamma \cdot x_j)$ would hold; a contradiction. \square

Lemma 5 In the theory TQ the following formulas

$$\begin{aligned} &\exists x [\mathfrak{R}_n(x \cdot t) \wedge \bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(x \cdot s_j)], \\ &\exists x [u < x \wedge \mathfrak{R}_n(x \cdot t) \wedge \bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(x \cdot s_j)] \text{ and} \\ &\exists x [x < v \wedge \mathfrak{R}_n(x \cdot t) \wedge \bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(x \cdot s_j)] \end{aligned}$$

are equivalent with

$$\bigwedge_{m_j | n (j < q)} \neg \mathfrak{R}_{m_j}(t^{-1} \cdot s_j);$$

and the formula

$$\exists x \left[u < x < v \wedge \mathfrak{R}_n(x \cdot t) \wedge \bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(x \cdot s_j) \right]$$

is equivalent with

$$\bigwedge_{m_j | n (j < q)} \neg \mathfrak{R}_{m_j}(t^{-1} \cdot s_j) \wedge u < v.$$

Proof If $m_j \mid n$ then $\mathfrak{R}_n(x \cdot t)$ implies $\mathfrak{R}_{m_j}(x \cdot t)$. Now, if $\mathfrak{R}_{m_j}(t^{-1} \cdot s_j)$ were true then $\mathfrak{R}_{m_j}(x \cdot s_j)$ would be true too; contradicting $\bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(x \cdot s_j)$. Suppose now that the relation $\bigwedge_{m_j \nmid n} \neg \mathfrak{R}_{m_j}(t^{-1} \cdot s_j)$ holds. By M_{11} there exists some γ such that $\bigwedge_{m_j \nmid n} \neg \mathfrak{R}_{m_j}(\gamma \cdot t^{-1} \cdot s_j)$. By M_{10} there exists some δ such that $u \cdot t \cdot \gamma^{-n} < \delta^{M \cdot n} < v \cdot t \cdot \gamma^{-n}$ (if $u < v$) where $M = \prod_{j < q} m_j$. For $x = \delta^{M \cdot n} \cdot \gamma^n \cdot t^{-1}$ we have $u < x < v$ and $\mathfrak{R}_n(x \cdot t)$. We show $\neg \mathfrak{R}_{m_j}(x \cdot s_j)$ for each $j < q$ by distinguishing two cases: if $m_j \mid n$ then $\neg \mathfrak{R}_{m_j}(t^{-1} \cdot s_j)$ implies $\neg \mathfrak{R}_{m_j}(\delta^{M \cdot n} \cdot \gamma^n \cdot t^{-1} \cdot s_j)$; if $m_j \nmid n$ then $\neg \mathfrak{R}_{m_j}(\gamma \cdot t^{-1} \cdot s_j)$ implies $\neg \mathfrak{R}_{m_j}(\delta^{M \cdot n} \cdot \gamma^n \cdot t^{-1} \cdot s_j)$. \square

Theorem 7 (Axiomatizability of $\langle \mathbb{Q}; <, \times \rangle$) The infinite theory TQ completely axiomatizes the theory of $\langle \mathbb{Q}^+; <, \times \rangle$, and

moreover the structure $\langle \mathbb{Q}^+; <, \times, \square^{-1}, \mathbf{1}, \{\mathfrak{R}_n\}_{n>1} \rangle$ admits quantifier elimination.

Also, the structure $\langle \mathbb{Q}; <, \times \rangle$ can be completely axiomatized by the theory that results from TQ by adding the axioms M_8 (in Theorem 6) and substituting its M_2, M_3, M_5, M_6 and M_{10} , respectively, with the axioms $M_2^o, M_3^o, M_5^o, M_5^*, M_6^o$ and $(M_{10}^o) \forall x, z \exists y (\mathbf{0} < x < z \rightarrow x < y^n < z)$.

Moreover, $\langle \mathbb{Q}; <, \times, \square^{-1}, -\mathbf{1}, \mathbf{0}, \mathbf{1}, \{\mathfrak{R}_n\}_{n>1} \rangle$ admits quantifier elimination.

Proof Let us prove the \mathbb{Q}^+ part only. We are to eliminate the quantifier of the formula

$$\exists x \left(\bigwedge_{i < p} \mathfrak{R}_{n_i}(x^{a_i} \cdot t_i) \wedge \bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(x^{b_j} \cdot s_j) \wedge \bigwedge_{k < f} u_k < x^{c_k} \wedge \bigwedge_{\ell < g} x^{d_\ell} < v_\ell \wedge \bigwedge_{i < h} x^{e_i} = w_i \right). \tag{1}$$

By $a^n < b^n \Leftrightarrow a < b$ and $\mathfrak{R}_{m-n}(a^n) \Leftrightarrow \mathfrak{R}_m(a)$ we can assume that all the a_i 's, b_j 's, c_k 's, d_ℓ 's and e_i 's are equal to each other, and moreover, equal to one (cf. the proof of Theorem 5). We can also assume that $h = 0$ and $f, g \leq 1$. By Lemma 2 we can also assume that $p \leq 1$. If $q = 0$ then Lemma 3 implies that the quantifier of formula (1) can be eliminated. So, we assume that $q > 0$. If $p = 0$ then the quantifier of (1) can be eliminated by Lemma 4. Finally, if $p = 1$ (and $q \neq 0 = h$ and $f, g \leq 1$) then Lemma 5 implies that formula (1) is equivalent with a quantifier-free formula. \square

Corollary 1 (Decidability of $\langle \mathbb{Q}; <, \times \rangle$) *The (first-order) theory of the structure $\langle \mathbb{Q}; <, \times \rangle$ (and also $\langle \mathbb{Q}^+; <, \times \rangle$) is decidable.*

Proof By Theorem 7 it suffices to note that the atomic formulas of the language $\{<, \times, \square^{-1}, -\mathbf{1}, \mathbf{0}, \mathbf{1}, \{\mathfrak{R}_n\}_{n>1}\}$ are decidable in \mathbb{Q} . For any $r \in \mathbb{Q}$ and any natural $n > 1$ the formula $\mathfrak{R}_n(r)$ holds if and only if every exponent of the unique factorization (of the numerators and denominators of the reduced form) of r is divisible by n . \square

Corollary 2 (Non-definability of addition) *The addition operation (+) is not definable in the structure $\langle \mathbb{Q}; <, \times \rangle$.*

Proof If it were, then the structure $\langle \mathbb{Q}; <, +, \times \rangle$ would be decidable by Theorem 7; but (Robinson 1949) proved that this structure is not decidable. \square

Remark 5 (Infinite axiomatizability) To see that the structure $\langle \mathbb{Q}^+; <, \times \rangle$ cannot be finitely axiomatized, we present an ordered multiplicative structure that satisfies any sufficiently large finite number of the axioms of TQ but does not satisfy all of its axioms. Let p be a sufficiently large prime number.

The set $\mathbb{Q}/p = \{m/p^k \mid m \in \mathbb{Z}, k \in \mathbb{N}\}$ is closed under addition and the operation $x \mapsto x/p$, and also $\mathbb{Z} \subset \mathbb{Q}/p \subset \mathbb{Q}$. Let $\rho_0, \rho_1, \rho_2, \dots$ denote the sequence of all prime numbers $(2, 3, 5, \dots)$. Let

$$(\mathbb{Q}/p)^* = \left\{ \prod_{i < \ell} \rho_i^{r_i} \mid \ell \in \mathbb{N}, r_i \in \mathbb{Q}/p \right\};$$

this is closed under multiplication and $x \mapsto x^{1/p}$, and also $\mathbb{Q}^+ \subset (\mathbb{Q}/p)^* \subset \mathbb{R}^+$. Thus, $(\mathbb{Q}/p)^*$ satisfies the axioms $O_1, O_2, O_3, M_1, M_2, M_3, M_4, M_5$ and M_6 of Proposition 1, and also the axiom M_{10} . However, it does not satisfy the axiom M_{11} for $n = q = x_0 = 1$ and $m_0 = p$ because $(\mathbb{Q}/p)^* \models \forall y \mathfrak{R}_p(y)$. We show that $(\mathbb{Q}/p)^*$ satisfies the instances of the axiom M_{11} when $1 < m_j < p$ (for each $j < q$ and arbitrary n, q). Thus, no finite number of the instances of M_{11} can prove all of its instances (with the rest of the axioms of TQ). Let x_j 's be given from $(\mathbb{Q}/p)^*$; write $x_j = \prod_{i < \ell_j} \rho_i^{r_{i,j}}$ where we can assume that $\ell_j \geq q$. Put $r_{j,j} = u_j/p^{v_j}$ where $u_j \in \mathbb{Z}$ and $v_j \in \mathbb{N}$ (for each $j < q$). Define t_j to be 1 when $m_j \mid u_j$ and be m_j when $m_j \nmid u_j$. Let $y = \prod_{i < q} \rho_i^{(t_i/p^{v_i+1})} \in (\mathbb{Q}/p)^*$. We show $\bigwedge_{j < q} \neg \mathfrak{R}_{m_j}(y^n \cdot x_j)$ under the assumption $\bigwedge_{j < q} m_j \nmid n$. Take a $k < q$, and assume (for the sake of contradiction) that $\mathfrak{R}_{m_k}(y^n \cdot x_k)$. Then $\mathfrak{R}_{m_k}(\rho_k^{nt_k/p^{v_k+1}} \cdot \rho_k^{u_k/p^{v_k}})$ holds, and so there exist a, b such that $\rho_k^{(nt_k+pu_k)/p^{v_k+1}} = \rho_k^{(m_k \cdot a)/p^b}$. Therefore, $m_k \mid nt_k + pu_k$. We reach to a contradiction by distinguishing two cases:

- (i) If $m_k \mid u_k$ then $t_k = 1$ and so $m_k \mid n + pu_k$ whence $m_k \mid n$, contradicting $\bigwedge_{j < q} m_j \nmid n$;
- (ii) If $m_k \nmid u_k$ then $t_k = m_k$ and so $m_k \mid nm_k + pu_k$ whence $m_k \mid pu_k$ which by $(m_k, p) = 1$ implies that $m_k \mid u_k$, contradicting the assumption (of $m_k \nmid u_k$). \diamond

3 Conclusions

In the following table the decidable structures are denoted by Δ and the undecidable ones by \mathbb{A} :

	\mathbb{N}	\mathbb{Z}	\mathbb{Q}	\mathbb{R}
$\{<\}$	Δ	Δ	Δ	Δ
$\{<, +\}$	Δ	Δ	Δ	Δ
$\{<, \times\}$	\mathbb{A}	\mathbb{A}	Δ	Δ
$\{+, \times\}$	\mathbb{A}	\mathbb{A}	\mathbb{A}	Δ

The decidability of the structure $\langle \mathbb{Q}; <, \times \rangle$ is a new result of this paper, along with the explicit axiomatization for the already known decidable structure $\langle \mathbb{R}; <, \times \rangle$. For the other decidable structures (other than $\langle \mathbb{N}; < \rangle$ and $\langle \mathbb{N}; <, + \rangle$), some

old and some new (syntactic) proofs are given for their decidability, with explicit axiomatizations (see ‘‘Appendix’’). It is interesting to note that the undecidability of $\langle \mathbb{N}; <, \times \rangle$ and $\langle \mathbb{Z}; <, \times \rangle$ is inherited from the undecidability of $\langle \mathbb{N}; +, \times \rangle$ and $\langle \mathbb{Z}; +, \times \rangle$ (and the definability of the addition operation $+$ in terms of order $<$ and multiplication \times in \mathbb{N} and \mathbb{Z}), and the decidability of $\langle \mathbb{R}; <, \times \rangle$ comes from the decidability of $\langle \mathbb{R}; +, \times \rangle$ (and the definability of order $<$ in terms of addition $+$ and multiplication \times in \mathbb{R}). Nonetheless, the undecidability of $\langle \mathbb{Q}; +, \times \rangle$ has nothing to do with the (decidable) structure $\langle \mathbb{Q}; <, \times \rangle$; indeed the addition operation $+$ is not definable in $\langle \mathbb{Q}; <, \times \rangle$ even though the order relation $<$ is definable in $\langle \mathbb{Q}; +, \times \rangle$.

Acknowledgements The authors gratefully thank the two anonymous referees of the journal for their valuable comments and suggestions.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

Appendix

Theorem 1. *The finite theory $\{O_1, O_2, O_3, O_4, O_5, O_6\}$ (of dense linear orders without endpoints—see Definitions 1 and 2) completely axiomatizes the theory of $\langle \mathbb{R}; < \rangle$ and $\langle \mathbb{Q}; < \rangle$, and so these structures are decidable. Moreover, (the theory of) those structures admit quantifier elimination.*

Proof All the atomic formulas are either of the form $u < v$ or $u = v$ for some variables u and v . If both of the variables are equal, then $u < u$ is equivalent with \perp by O_1 and $u = u$ is equivalent with \top . So, by Remark 1, it suffices to eliminate the quantifier of the formulas of the form

$$\exists x \left(\bigwedge_{i < \ell} y_i < x \wedge \bigwedge_{j < m} x < z_j \wedge \bigwedge_{k < n} x = u_k \right) \tag{2}$$

where y_i ’s, z_j ’s and u_k ’s are variables. Now, if $n \neq 0$ then formula (2) is equivalent with the quantifier-free formula

$$\bigwedge_{i < \ell} y_i < u_0 \wedge \bigwedge_{j < m} u_0 < z_j \wedge \bigwedge_{k < n} u_0 = u_k.$$

So, let us suppose that $n = 0$. Then if $m = 0$ or $\ell = 0$ formula (2) is equivalent with the quantifier-free formula \top , by the axioms O_5 and O_6 (with O_2 and O_3) respectively, and if $\ell, m \neq 0$ it is equivalent with the quantifier-free formula $\bigwedge_{i < \ell, j < m} y_i < z_j$ by the axiom O_4 (with O_2 and O_3). \square

Theorem 2 *The finite theory of discrete linear orders without endpoints, consisting of the axioms O_1, O_2, O_3, O_7 plus*

$$(O_8) \forall x \exists y (\mathfrak{s}(y) = x)$$

completely axiomatizes the order theory of the integer numbers, and so its theory is decidable. Moreover, the structure $\langle \mathbb{Z}; <, \mathfrak{s} \rangle$ admits quantifier elimination.

Proof We note that all the terms in the language $\{<, \mathfrak{s}\}$ are of the form $\mathfrak{s}^n(y)$ for some variable y and $n \in \mathbb{N}$. So, all the atomic formulas are either of the form $\mathfrak{s}^n(u) = \mathfrak{s}^m(v)$ or $\mathfrak{s}^n(u) < \mathfrak{s}^m(v)$ for some variables u, v . If a variable x appears in the both sides of an atomic formula, then we have either $\mathfrak{s}^n(x) = \mathfrak{s}^m(x)$ or $\mathfrak{s}^n(x) < \mathfrak{s}^m(x)$. The formula $\mathfrak{s}^n(x) = \mathfrak{s}^m(x)$ is equivalent with \top when $n = m$ and with \perp otherwise; also $\mathfrak{s}^n(x) < \mathfrak{s}^m(x)$ is equivalent with \top when $n < m$ and with \perp otherwise. So, it suffices to consider the atomic formulas of the form $t < \mathfrak{s}^n(x)$ or $\mathfrak{s}^n(x) < t$ or $\mathfrak{s}^n(x) = t$ for some x -free term t and $n \in \mathbb{N}^+$. Now, by Remark 1, we eliminate the quantifier of the formulas

$$\exists x \left(\bigwedge_{i < \ell} t_i < \mathfrak{s}^{p_i}(x) \wedge \bigwedge_{j < m} \mathfrak{s}^{q_j}(x) < s_j \wedge \bigwedge_{k < n} \mathfrak{s}^{r_k}(x) = u_k \right). \tag{3}$$

The axioms prove the equivalences $[a < b] \leftrightarrow [\mathfrak{s}(a) < \mathfrak{s}(b)]$ and $[a = b] \leftrightarrow [\mathfrak{s}(a) = \mathfrak{s}(b)]$; so we can assume that p_i ’s and q_j ’s and r_k ’s in formula (3) are equal to each other, say to α . Then by O_8 formula (3) is equivalent with

$$\exists y \left(\bigwedge_{i < \ell} t'_i < y \wedge \bigwedge_{j < m} y < s'_j \wedge \bigwedge_{k < n} y = u'_k \right) \tag{4}$$

for some (possibly new) terms t'_i, s'_j, u'_k (and $y = \mathfrak{s}^\alpha(x)$). Now, if $n \neq 0$ then formula (4) is equivalent with the quantifier-free formula

$$\bigwedge_{i < \ell} t'_i < u'_0 \wedge \bigwedge_{j < m} u'_0 < s'_j \wedge \bigwedge_{k < n} u'_0 = u'_k.$$

Let us then assume that $n = 0$. The formula

$$\exists x \left(\bigwedge_{i < \ell} t_i < x \wedge \bigwedge_{j < m} x < s_j \right) \tag{5}$$

is equivalent, by the axiom O_7 (in Definition 2), with the quantifier-free formula $\bigwedge_{i < \ell, j < m} \mathfrak{s}(t_i) < s_j$. \square

Theorem 4 *The following infinite theory (of non-trivial ordered divisible abelian groups) completely axiomatizes the*

order and additive theory of the real and rational numbers, and so their theories are decidable. Moreover, the structures $\langle \mathbb{R}; <, +, -, \mathbf{0} \rangle$ and $\langle \mathbb{Q}; <, +, -, \mathbf{0} \rangle$ admit quantifier elimination.

- (O₁) $\forall x, y(x < y \rightarrow y \not< x)$
- (O₂) $\forall x, y, z(x < y < z \rightarrow x < z)$
- (O₃) $\forall x, y(x < y \vee x = y \vee y < x)$
- (A₁) $\forall x, y, z(x + (y + z) = (x + y) + z)$
- (A₂) $\forall x(x + \mathbf{0} = x)$
- (A₃) $\forall x(x + (-x) = \mathbf{0})$
- (A₄) $\forall x, y(x + y = y + x)$
- (A₅) $\forall x, y, z(x < y \rightarrow x + z < y + z)$
- (A₆) $\exists y(y \neq \mathbf{0})$
- (A₇) $\forall x \exists y(x = n \cdot y) \quad n \in \mathbb{N}^+$

Proof Firstly, let us note that O₄, O₅ and O₆ can be proved from the presented axioms: if $a < b$ then by A₇ there exists some c such that $c + c = a + b$; one can easily show that $a < c < b$ holds. Thus, O₄ is proved; for O₅ note that for any $\mathbf{0} < a$ we have $a < a + a$ by A₅. A dual argument can prove the axiom O₆. Also, the equivalences

- (i) $[a < b] \leftrightarrow [n \cdot a < n \cdot b]$ and
- (ii) $[a = b] \leftrightarrow [n \cdot a = n \cdot b]$

can be proved from the axioms: (i) follows from A₅ (with O₁, O₂, O₃) and (ii) follows from $\forall x(n \cdot x = \mathbf{0} \rightarrow x = \mathbf{0})$ which is derived from A₅ (with O₁, O₂, O₃).

Secondly, every term containing x is equal to $n \cdot x + t$ for some x -free term t and $n \in \mathbb{Z} - \{0\}$. So, every atomic formula containing x is equivalent with $n \cdot x \square t$ where $\square \in \{=, <, >\}$. Whence, by Remark 1, it suffices to prove the equivalence of the formula

$$\exists x \left(\bigwedge_{i < \ell} t_i < p_i \cdot x \wedge \bigwedge_{j < m} q_j \cdot x < s_j \wedge \bigwedge_{k < n} r_k \cdot x = u_k \right) \tag{6}$$

with a quantifier-free formula. By the equivalences (i) and (ii) above we can assume that p_i 's and q_j 's and r_k 's in formula (6) are equal to each other, say to α . Then by A₇ formula (6) is equivalent with

$$\exists y \left(\bigwedge_{i < \ell} t'_i < y \wedge \bigwedge_{j < m} y < s'_j \wedge \bigwedge_{k < n} y = u'_k \right) \tag{7}$$

for some (possibly new) terms t'_i, s'_j, u'_k (and $y = \alpha \cdot x$). Now, the quantifier of this formula can be eliminated just like the way that the quantifier of formula (2) was eliminated in the proof of Theorem 1. \square

About the congruence relations, a useful fact is the following generalized Chinese remainder theorem; below we present a proof of this theorem from (Fraenkel 1963).

Proposition 2 (Generalized Chinese remainder) *For integers $n_0, n_1, \dots, n_k \geq 2$ and t_0, t_1, \dots, t_k there exists some x such that $x \equiv_{n_i} t_i$ for $i = 0, \dots, k$ if and only if $t_i \equiv_{d_{i,j}} t_j$ holds for each $0 \leq i < j \leq k$, where $d_{i,j}$ is the greatest common divisor of n_i and n_j .*

Proof The ‘only if’ part is easy. We prove the ‘if’ part by induction on k . For $k = 0$ there is nothing to prove, and for $k = 1$ we note that by Bézout’s Identity there are a_0, a_1 such that $a_0 n_0 + a_1 n_1 = d_{0,1}$. Also, by the assumption there exists some c such that $t_0 - t_1 = c d_{0,1}$. Now, if we take x to be $a_0(n_0/d_{0,1})t_1 + a_1(n_1/d_{0,1})t_0$ then we have $x = t_0 - a_0 n_0 c$ and $x = t_1 + a_1 n_1 c$ so $x \equiv_{n_0} t_0$ and $x \equiv_{n_1} t_1$ hold. For the induction step ($k + 1$) suppose that $x \equiv_{n_i} t_i$ holds for $i = 0, \dots, k$ (and that $t_i \equiv_{d_{i,j}} t_j$ holds for each $0 \leq i < j \leq k + 1$). Let n be the least common multiplier of n_0, \dots, n_k ; then the greatest common divisor of n and n_{k+1} is the least common multiplier of $d_{0,k+1}, \dots, d_{k,k+1}$. Now $x \equiv_{d_{i,k+1}} t_i$ holds for $0 \leq i \leq k$ and so by the assumption $t_i \equiv_{d_{i,k+1}} t_{k+1}$ we have $x \equiv_{d_{i,k+1}} t_{k+1}$ (for $i = 0, \dots, k$). Therefore, $x \equiv_m t_{k+1}$ and so $x - t_{k+1} = mc$ for some c . By Bézout’s Identity there are a, b such that $an + bn_{k+1} = m$. Now, for $y = x - anc$ we have $y = t_{k+1} + bn_{k+1}c \equiv_{n_{k+1}} t_{k+1}$ and also $y \equiv_{n_i} x \equiv_{n_i} t_i$ holds for each $0 \leq i \leq k$. This proves the desired conclusion. \square

Theorem 5 *The infinite theory of non-trivial discretely ordered abelian groups with the division algorithm, that is O₁, O₂, O₃, A₁, A₂, A₃, A₄, A₅ and*

$$\begin{aligned} & (O_7^\circ) \forall x, y(x < y \leftrightarrow x + \mathbf{1} \leq y) \\ & (A_7^\circ) \forall x \exists y \left(\bigvee_{i < n} x = n \cdot y + \bar{i} \right) \quad n \in \mathbb{N}^+ \end{aligned}$$

where $\bar{i} = \mathbf{1} + \dots + \mathbf{1}$ (i - times)

completely axiomatizes the order and additive theory of the integer numbers, and so its theory is decidable. Moreover, the (theory of the) structure $\langle \mathbb{Z}; <, +, -, \mathbf{0}, \mathbf{1}, \{\equiv_n\}_{n \geq 2} \rangle$ admits quantifier elimination.

Proof The axiom A₇^o can be easily seen to be equivalent with the formula $\forall x \bigvee_{i < n} (x \equiv_n \bar{i} \wedge \bigwedge_{i \neq j < n} x \not\equiv_n \bar{j})$, and so the negation signs behind the congruences can be eliminated by $(a \not\equiv_n b) \leftrightarrow \bigvee_{0 < i < n} (a \equiv_n b + \bar{i})$. Whence, by Remark 1, it suffices to show the equivalence of

$$\exists x \left(\bigwedge_{i < m} a_i \cdot x \equiv_{n_i} t_i \wedge \bigwedge_{j < p} u_j < b_j \cdot x \right. \\ \left. \wedge \bigwedge_{k < q} c_k \cdot x < v_k \wedge \bigwedge_{\ell < r} d_\ell \cdot x = w_\ell \right) \tag{8}$$

with some quantifier-free formula, where a_i 's, b_j 's, c_k 's and d_ℓ 's are natural numbers and t_i 's, u_j 's, v_k 's and w_ℓ 's are x -free terms. By the equivalences

- (i) $[a < b] \leftrightarrow [n \cdot a < n \cdot b]$,
- (ii) $[a = b] \leftrightarrow [n \cdot a = n \cdot b]$ and
- (iii) $[a \equiv_m b] \leftrightarrow [n \cdot a \equiv_{nm} n \cdot b]$

which are provable from the axioms, we can assume that a_i 's, b_j 's, c_k 's and d_ℓ 's in formula (8) are equal to each other, say to α . Now, (8) is equivalent with

$$\exists y \left(y \equiv_\alpha \mathbf{0} \wedge \bigwedge_{i < m} y \equiv_{n_i} t'_i \wedge \bigwedge_{j < p} u'_j < y \right. \\ \left. \wedge \bigwedge_{k < q} y < v'_k \wedge \bigwedge_{\ell < r} y = w'_\ell \right) \tag{9}$$

for $y = \alpha \cdot x$ and some (possibly new) terms t'_i 's, u'_j 's, v'_k 's and w'_ℓ 's. If $r \neq 0$ then (9) is readily equivalent with the quantifier-free formula which results from substituting w'_0 with y . So, it suffices to eliminate the quantifier of

$$\exists x \left(\bigwedge_{i < m} x \equiv_{n_i} t_i \wedge \bigwedge_{j < p} u_j < x \wedge \bigwedge_{k < q} x < v_k \right). \tag{10}$$

By the equivalence of the formula $\exists x(\theta(x) \wedge u_0 < x \wedge u_1 < x)$ with the following formula

$$[\exists x(\theta(x) \wedge u_0 < x) \wedge u_1 \leq u_0] \vee [\exists x(\theta(x) \wedge u_1 < x) \wedge u_0 \leq u_1],$$

we can assume that $p \leq 1$ (and $q \leq 1$ by a dual argument). Also, the formula $\exists x(\theta(x) \wedge x \equiv_{n_0} t_0 \wedge x \equiv_{n_1} t_1)$ is equivalent with $\exists x(\theta(x) \wedge x \equiv_n t) \wedge t_0 \equiv_d t_1$ where d is the greatest common divisor of n_0 and n_1 , n is their least common multiplier, and $t = a_0(n_0/d)t_1 + a_1(n_1/d)t_0$ where a_0, a_1 satisfy Bézout's Identity $a_0n_0 + a_1n_1 = d$ (see the proof of Proposition 2). So, we can assume that $m \leq 1$ as well. Now, if $m = 0$ then formula (10) is equivalent with a quantifier-free formula by Theorem 2 (with $s(x) = x + \mathbf{1}$ just like the way formula (5) was equivalent with some quantifier-free formula). So, suppose that $m = 1$. In this case, if any of p or q is equal to 0 then (10) is equivalent with \top (since any congruence can have infinitely large or infinitely

small solutions). Finally, if $p = q = 1 = m$ then the formula

$$\exists x(x \equiv_n t \wedge u < x \wedge x < v)$$

is equivalent with $\exists y(r < n \cdot y \leq s)$ for $x = t + n \cdot y$, $r = u - t$ and $s = v - t - \mathbf{1}$. Now, $\exists y(r < n \cdot y \leq s)$ is equivalent with the quantifier-free formula

$$\bigvee_{i < n} (s \equiv_n \bar{i} \wedge r + \bar{i} < s)$$

since by the division algorithm there are some q and some $i < n$ such that $s = qn + i$. The existence of some y such that $r < ny \leq s$ is then equivalent with $r < nq (= s - i)$. \square

References

Boolos GS, Burgess JP, Jeffrey RC (2007) Computability and logic, 5th edn. Cambridge University Press, Cambridge (ISBN: 9780521701464)

Enderton HB (2001) A mathematical introduction to logic, 2nd edn. Academic Press, New York (ISBN: 9780122384523)

Fraenkel AS (1963) New proof of the generalized Chinese remainder theorem. Proc Am Math Soc 14(5):790–791. <https://doi.org/10.1090/S0002-9939-1963-0154841-6>

Hinman PG (2005) Fundamentals of mathematical logic. CRC Press, Boca Raton (ISBN: 9781568812625)

Kreisel G, Krivine JL (1971) Elements of mathematical logic: model theory. North-Holland, Amsterdam (ISBN: 9780720422658)

Marker D (2002) Model theory: an introduction. Springer, Berlin (ISBN: 9781441931573)

Monk JD (1976) Mathematical logic. Springer, Berlin (ISBN: 9780387901701)

Mostowski A (1952) On direct products of theories. J Symb Log 17:1–31. <https://doi.org/10.2307/2267454>

Ore O (1952) The general Chinese remainder theorem. Am Math Mon 59(6):365–370. <https://doi.org/10.2307/2306804>

Prestel A (1986) Einführung in die mathematische Logik und Modelltheorie. Vieweg, Berlin (ISBN: 9783528072605)

Prestel A, Delzell CN (2011) Mathematical logic and model theory: a brief introduction. Springer, Berlin (ISBN: 9781447121756)

Robinson J (1949) Definability and decision problems in arithmetic. J Symb Log 14(2):98–114. <https://doi.org/10.2307/2266510>

Robinson A, Zakon E (1960) Elementary properties of ordered Abelian groups. Trans Am Math Soc 96(2):222–236. <https://doi.org/10.2307/1993461>

Salehi S (2012a) Axiomatizing mathematical theories: multiplication. In: Kamali-Nejad A (ed) Proceedings of frontiers in mathematical sciences. Sharif University of Technology, Tehran, pp 165–176. [arxiv:1612.06525](https://arxiv.org/abs/1612.06525)

Salehi S (2012b) Computation in logic and logic in computation. In: Sadeghi-Bigham B (ed) Proceedings of the third international conference on contemporary issues in computer and information sciences (CICIS 2012). Brown Walker Press, New York, pp 580–583. [arxiv:1612.06526](https://arxiv.org/abs/1612.06526)

Salehi S (2018) On axiomatizability of the multiplicative theory of numbers. Fundam Inf 159(3):279–296. <https://doi.org/10.3233/FI-2018-1665>

- Smoryński C (1991) Logical number theory I: an introduction. Springer, Berlin (ISBN: 9783540522362)
- Szmielew W (1949) Decision problem in group theory. In: Beth EW, Pos HJ, Hollak JHA (eds) Proceedings of the tenth international congress of philosophy, vol 2. North-Holland, Amsterdam, pp 763–766. <https://doi.org/10.5840/wcp1019492212>
- Szmielew W (1955) Elementary properties of Abelian groups. *Fundam Math* 41:203–271. <https://doi.org/10.4064/fm-41-2-203-271>
- Visser A (2017) On Q. *Soft Comput* 21(1):39–56. <https://doi.org/10.1007/s00500-016-2341-5>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.