

Gamma convergence of an energy functional related to the fractional Laplacian

María del Mar González

Received: 18 May 2008 / Accepted: 9 January 2009 / Published online: 30 January 2009
© Springer-Verlag 2009

Abstract We prove a Γ -convergence result for an energy functional related to some fractional powers of the Laplacian operator, $(-\Delta)^s$ for $1/2 < s < 1$, with two singular perturbations, that leads to a two-phase problem. The case $(-\Delta)^{1/2}$ was considered by Alberti–Bouchitté–Seppecher in relation to a model in capillarity with line tension effect. However, the proof in our setting requires some new ingredients such as the Caffarelli–Silvestre extension for the fractional Laplacian and new trace inequalities for weighted Sobolev spaces.

Mathematics Subject Classification (2000) 49J45 · 35J20 · 82B26 · 47G30

1 Introduction and statement of the theorem

Let Ω be a bounded domain in \mathbb{R}^3 with smooth C^2 boundary $\partial\Omega$ and let $h : \Omega \rightarrow \mathbb{R}$ be the distance function to the boundary. Fix a real number $-1 < a < 0$. Let $\alpha, \beta, \alpha', \beta' \in \mathbb{R}$ such that $\alpha < \beta, \alpha' < \beta'$ and consider two double-well potentials $W, V : \mathbb{R} \rightarrow [0, \infty)$ such that W only vanishes at α, β , and V only vanishes at α', β' . For a function u defined in Ω , denote its trace on $\partial\Omega$ by Tu . Given $\epsilon > 0$, we study the following energy functional

$$F_\epsilon^a[u] := \epsilon^{1-a} \int_{\Omega} |\nabla u|^2 h^a + \frac{1}{\epsilon^{1-a}} \int_{\Omega} W(u) h^{-a} + \lambda_\epsilon \int_{\partial\Omega} V(Tu). \quad (1.1)$$

The aim of the present paper is to understand the Γ -convergence of this functional when $\epsilon \rightarrow 0$ and $\lambda_\epsilon \rightarrow \infty$.

This article was supported by Spain Government project MTM2005-07660-C02-01.

M. d. M. González (✉)
ETSEIB, Departament de Matemàtica Aplicada I, Universitat Politècnica de Catalunya,
Av. Diagonal 647, 08028 Barcelona, Spain
e-mail: mar.gonzalez@upc.edu

Note that (1.1) generalizes the two-phase model of Alberti–Bouchitté–Seppecher considered in [6] in relation to capillarity energy with line tension. They studied the Γ -convergence of

$$F_\epsilon[u] := \epsilon \int_\Omega |\nabla u|^2 + \frac{1}{\epsilon} \int_\Omega W(u) + \lambda_\epsilon^0 \int_{\partial\Omega} V(Tu), \tag{1.2}$$

where $\lambda_\epsilon^0 \rightarrow \infty$ is a sequence with some specific behavior as $\epsilon \rightarrow 0$.

Historically, this type of models appeared when studying phase transitions. Consider a container $\Omega \subset \mathbb{R}^3$ which is filled with two immiscible and incompressible fluids, or two different phases of the same fluid; equilibrium is achieved when we minimize

$$E[u] = \sigma \mathcal{H}^2(S_u) \tag{1.3}$$

among all the admissible configurations $u \in BV(\Omega, \{\alpha, \beta\})$ with fixed volume $\int_\Omega u = M_0$. Here \mathcal{H}^2 is the two-dimensional Hausdorff measure of S_u , the singular set of u , and σ is a constant parameter. The classical work by Modica [17] established that the following variational model for $u : \Omega \rightarrow [\alpha, \beta]$, $\int_\Omega u = M_0$, given by

$$E_\epsilon[u] = \epsilon \int_\Omega |\nabla u|^2 + \frac{1}{\epsilon} \int_\Omega W(u), \tag{1.4}$$

Γ -converges to E , for $\sigma := 2 \int_\alpha^\beta \sqrt{W(t)} dt$.

Here we consider the generalization given by (1.1), where the weight h^a is singular at the boundary $\partial\Omega$ since $-1 < a < 0$. With this modification, F_ϵ^a becomes the energy functional related to some fractional powers of the Laplacian, plus two double-well potential terms. Indeed, if $s = \frac{1-a}{2}$, then the Euler–Lagrange equation of the functional

$$J[u] = \int_{x \in \mathbb{R}^n, y \in \mathbb{R}^+} |\nabla u|^2 y^a dx dy \tag{1.5}$$

is just $(-\Delta)^s(Tu) = 0$ in \mathbb{R}^n , as it has been shown in the recent work by Caffarelli–Silvestre [9].

On the other hand, the boundary potential term in (1.1) constitutes a very interesting modification from (1.4), and adds new terms in the Γ -limit. The first result involving boundary integrals was obtained by Modica [18], when $\lambda_\epsilon = 1$. Other works can be found in Cabré and Solá-Morales [8], where they look at layer solutions for boundary reactions of the half-Laplacian ($s = 1/2$). A refinement of the Γ -convergence result for $s = 1/2$ is being studied by Cabré and Cónsul [7]. In addition, in the current work [20] by Monneau and the present author, we look at an homogenization problem for a reaction-diffusion equation with half-Laplacian, where we try to understand the interaction energy that appears in the problem.

The main theorem in the present paper states that the Γ -convergence of the sequence F_ϵ^a , for some suitable scaling λ_ϵ , has a similar behavior to the case $s = 1/2$ of Alberti–Bouchitté–Seppecher considered in [6]. However, the proof needs some new results on the fractional Laplacian: Sect. 4 contains a result on singular perturbations of the norm $H^s(\partial\Omega)$, that is indeed the norm of traces of functions in a weighted Sobolev space $W^{1,2}(\Omega, w_a)$ for a suitable weight w_a . The second main ingredient, in Sect. 5, deals with this trace embedding, and gives a more precise control of the Sobolev constant. In particular, in Theorem 19 we understand the relation between the functional (1.5) in \mathbb{R}_+^n and the non-local energy of the trace Tu in $H^s(\mathbb{R})$ given by

$$\int_{\mathbb{R} \times \mathbb{R}} \frac{|Tu(x') - Tu(x)|^2}{|x' - x|^{1+2s}} dx' dx.$$

The proof uses the interesting characterization of the fractional Laplacian in \mathbb{R} as an extension problem to $\mathbb{R} \times \mathbb{R}_+$, given by Caffarelli–Silvestre in [9].

Let us fix some notation. Set $-1 < a < 0$. Let $I := \{\alpha, \beta\}$, $I' := \{\alpha', \beta'\}$. Assume that there exists m such that $-m \leq \alpha, \alpha', \beta, \beta' \leq m$. Assume that both W and V are positive, increasing in $[m, +\infty)$ and decreasing in $(-\infty, -m]$, with growth at least linear at infinity. As we have mentioned, W only vanishes at the two wells $\alpha < \beta$, and V only at $\alpha' < \beta'$.

Consider the background space $X := L^1(\Omega) \times L^1(\partial\Omega)$. Fix $h : \Omega \rightarrow [0, +\infty)$ be the distance to the boundary of Ω . Fix the singular weight in Ω given by $w_a := h^a$, and consider the weighted Sobolev space $W^{1,2}(\Omega, w_a)$ with norm

$$\|u\|_{W^{1,2}(\Omega, w_a)}^2 := \int_{\Omega} |u|^2 h^a + \int_{\Omega} |\nabla u|^2 h^a.$$

Note that the trace on $\partial\Omega$ of a function $u \in W^{1,2}(\Omega, w_a)$ is well defined (see Theorem 3); denote it by Tu .

We define the functional F_ϵ^a as

$$F_\epsilon^a[u] := \begin{cases} \epsilon^{1-a} \int_{\Omega} |\nabla u|^2 h^a + \frac{1}{\epsilon^{1-a}} \int_{\Omega} W(u) h^{-a} + \lambda_\epsilon \int_{\partial\Omega} V(Tu), & u \in W^{1,2}(\Omega, w_a) \cap X, \\ +\infty, & \text{elsewhere in } X. \end{cases} \tag{1.6}$$

On the other hand, given $u \in BV(\Omega, I)$ and $v \in BV(\partial\Omega, I')$, set

$$\Phi(u, v) := \sigma \mathcal{H}^2(S_u) + \int_{\partial\Omega} |\mathcal{W}(Tu) - \mathcal{W}(v)| + \kappa_s \mathcal{H}^1(S_v), \tag{1.7}$$

and

$$F^a(u) := \begin{cases} \inf\{\Phi(u, v) : v \in BV(\partial\Omega, I')\}, & \text{if } u \in BV(\Omega, I), \\ +\infty, & \text{elsewhere in } X. \end{cases} \tag{1.8}$$

Here \mathcal{H}^i denotes the i -dimensional Hausdorff measure, that in this case is well defined because of the hypothesis on the bounded variation of u, v . Also, S_u is the set of all points where u is essentially discontinuous, and the same for S_v . The main result of the present paper states that F^a is the Γ -limit of the functionals F_ϵ^a , for suitable positive constants σ, κ_s and a function \mathcal{W} .

Finally, we set

$$\Lambda_\epsilon := \epsilon^{\frac{1-a}{2}}. \tag{1.9}$$

It will become clear in Sects. 4 and 6.3 that this is the natural scaling for the problem.

We have that, under the above conditions:

Theorem 1 Fix $-1 < a < 0$, $s = \frac{1-a}{2}$, and assume that

$$\lambda_\epsilon = \frac{1}{\Lambda_\epsilon}. \tag{1.10}$$

Then there exist constants $\sigma, \kappa_s > 0$ such that the functionals F_ϵ^a defined in (1.6) Γ -converge to the functional F^a given in (1.8) and any sequence (u_ϵ) with bounded energy is precompact

in X . Moreover, $\sigma := \mathcal{W}(\beta) - \mathcal{W}(\alpha)$ where \mathcal{W} is a primitive of $2\sqrt{W}$, and κ_s is a constant depending only on s , V whose exact value is given in (4.2).

Let us motivate the previous theorem. The potential in the interior, W , forces the minimizer u_ϵ to take values near the two wells α, β , while the gradient term in the functional penalizes the jump of the function; thus we create two bulk phases in the interior of the container Ω , namely, $\{u = \alpha\}, \{u = \beta\}$, with interphase S_u . When $\epsilon \rightarrow 0$, $u_\epsilon \rightarrow u$ and $Tu_\epsilon \rightarrow v$. On the other hand, the second double-well potential V forces the trace to take values near α', β' , and thus it creates two boundary phases $\{v = \alpha'\}, \{v = \beta'\}$, separated by $S_v \subset \partial\Omega$. However, we usually have $Tu \neq v$, and thus additional terms appear in the limit functional (1.7). Note that, although many of the arguments would work for a domain Ω contained in \mathbb{R}^n for any $n \geq 3$, we restrict ourselves to dimension three so that the energy on $\partial\Omega$ occurring from the boundary phases concentrates over a one-dimensional set S_v of $\partial\Omega$.

We remark here that the proof of the Gamma-convergence result in general follows some well established steps (see Alberti et al. [6] or Palatucci [22]). However, in our case the shape of the functional requires a deep understanding of the singular factor in the energy and its relation to the fractional Laplacian operator - this is precisely the main new idea of the present article.

The result is true if either W or V are identically zero. Indeed, if V is zero, we are in the situation of Sect. 3, while if W is zero, then we can ignore the first two terms in the Γ -limit (1.7).

Palatucci (cf. [22], or his Ph.D. thesis [23]), has considered the Γ -convergence of the super-quadratic functional

$$P_\epsilon^p[u] := \epsilon^{p-2} \int_{\Omega} |\nabla u|^p + \frac{1}{\epsilon^{\frac{p-2}{p-1}}} \int_{\Omega} W(u) + \frac{1}{\epsilon} \int_{\partial\Omega} V(Tu)$$

when $p > 2$. The main difference with ours is the lack of trace inequalities as in Theorem 19 for the case $p > 2$.

It has come to our attention that the generalization of the super-quadratic functional P_ϵ^p including a singular weight at the boundary $\partial\Omega$ is being completed by Palatucci and Sire [24]. It uses some of the results of the present article.

Several open questions arise: first, when $0 < s < 1/2$ we do not know yet how to formulate a Gamma-convergence result due to the lack of layer solutions for the functional (4.1). It may even happen that some other new non-local quantities appear. Also, not much is known in the anisotropic case.

The outline of the paper is the following: in Sect. 2 we give some general background on Γ -convergence and on the fractional Laplacian. In Sect. 3 we study the problem in the interior ignoring the boundary interaction. The following two sections contain the main ingredients: in Sect. 4 we look at a singular perturbation result for the norm $H^s(E)$ where E is an interval in \mathbb{R} , while the next section deals with some new trace embeddings for weighted Sobolev embeddings. Section 6 contains several technical results that are needed in the main proof, including a dimension reduction argument. Finally, we prove Theorem 1 in the last section.

2 Some background

In order to make this paper self-contained, we present to the reader some standard background. We will denote by $\epsilon \rightarrow 0$ any countable sequence converging to zero. The integrals in a domain Ω are taken with respect the standard Lebesgue measure, while the integrals on $\partial\Omega$ are with respect to the standard Hausdorff measure on the boundary. First, let us give the definition of Γ -convergence.

Definition 2 Let X be a metric space, and for $\epsilon > 0$, consider the functional $F_\epsilon : X \rightarrow [0, +\infty]$. We say that the sequence F_ϵ Gamma-converges to F on X as $\epsilon \rightarrow 0$ if the following conditions hold:

- i. Lower bound inequality: for every $u \in X$ and every sequence (u_ϵ) such that $u_\epsilon \rightarrow u$ in X , there holds

$$\liminf_{\epsilon \rightarrow 0} F_\epsilon[u_\epsilon] \geq F[u].$$

- ii. Upper bound inequality: for every $u \in X$ there exists (u_ϵ) such that $u_\epsilon \rightarrow u$ in X and

$$\lim_{\epsilon \rightarrow 0} F_\epsilon[u_\epsilon] = F[u].$$

Together with conditions *i.* and *ii.* a compactness condition is usually proved:

- iii. Given a sequence (u_ϵ) when $\epsilon \rightarrow 0$ such that $F_\epsilon[u_\epsilon]$ is bounded, then (u_ϵ) is pre-compact in X .

Consider a sequence of functionals F_ϵ that Γ -converges to F . If u_ϵ is a minimizer for F_ϵ , then conditions *i* and *ii.* imply that any limit point of the sequence (u_ϵ) is a minimizer for F . Condition *iii.* assures that this limit point exists in X .

The definition and properties of bounded variation functions can be found in [10] or [15]. Let Ω be an open subset of \mathbb{R}^n . Note that if $\partial\Omega$ is Lipschitz, the trace of a bounded variation function on $\partial\Omega$ is well defined and it belongs to $L^1(\partial\Omega)$. Given $f \in BV(\Omega, I)$, we define S_u to be the set of all points where u is essentially discontinuous, that is, it has no approximate limit, and it agrees with the measure theoretic boundary of the set $\{u = \alpha\}$ in Ω .

Now we give the relation between weighted Sobolev spaces and their traces. Let $n > 0$, $k \geq 0$ be integers, and a, p real numbers, $1 < p < \infty$. Let Ω be a non-empty, open, bounded subset of \mathbb{R}^n . Let M be a closed subset of $\partial\Omega$ and let $d_M(x)$ be the distance function, $d_M(x) := \text{dist}(x, M)$. For simplicity we shall write $d(x)$ instead of $d_M(x)$. For an integer m , $1 \leq m \leq n$, we set $Q_m = (0, 1)^m$.

We shall write $(\Omega, M) \in B(k, n)$ for $1 \leq k \leq n - 1$, $n \geq 2$ if and only if there exists a bi-Lipschitz mapping $B : Q_n \rightarrow \Omega$ such that $B(Q_k) = M$.

By $C^\infty(\bar{\Omega})$ we denote the set of real functions u defined on $\bar{\Omega}$ such that the derivatives $D^\alpha u$ can be continuously extended to $\bar{\Omega}$ for all multi-indexes α . Consider the weight $w = d^a$. Define the weighted Sobolev space $W^{1,p}(\Omega, w)$ as the closure of $C^\infty(\bar{\Omega})$ with respect to the norm

$$\|u\|_{W^{1,p}(\Omega,w)}^p := \int_{\Omega} |u|^p w \, dx + \int_{\Omega} |Du|^p w \, dx.$$

These norms have been well studied for a certain class of weights, called A_p weights (see [12,26]). We will also need some other norms for functions v defined on an interval $E \subset \mathbb{R}$ given by

$$\|v\|_{H^s(E)}^2 = \|v\|_{L^2(E)}^2 + \int_E \frac{|v(x') - v(x)|^2}{|x' - x|^{1+2s}} \, dx' dx$$

for $0 < s < 1$.

Although it can be stated more generally, we are just interested in the case $k = n - 1$:

Theorem 3 (theorem 2.8 in [21]) *Let $n \geq 2$, $-1 < a < p - 1$ and $(\Omega, M) \in B(n - 1, n)$. Then there exists a unique bounded linear operator*

$$T : W^{1,p}(\Omega, w) \rightarrow W^{1-\frac{1+a}{p},p}(M) \tag{2.1}$$

such that $Tu = u|_M$ for all $u \in C^\infty(\bar{Q}_n)$.

In the special case that $p = 2$, $s = \frac{1-a}{2}$, the trace operator (2.1) reads precisely

$$T : W^{1,2}(\Omega, w) \rightarrow H^s(M).$$

The classical reference for the fractional Laplacian $(-\Delta)^s$ is the book by Landkof [16], although a good reference is the Ph.D. thesis by Luis Silvestre [25]. Given $s \in (0, 1)$ we define the fractional Laplacian of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, as a pseudo-differential operator by

$$\widehat{(-\Delta)^s f}(\xi) = |\xi|^{2s} \hat{f}(\xi),$$

i.e., its principal symbol is $|\xi|^{2s}$. It can also be written as the singular integral

$$(-\Delta)^s f(x) = C_{n,s} \int_{\mathbb{R}^n} \frac{f(x) - f(\xi) - \nabla f(x) \cdot (x - \xi) \chi_{(|x-\xi|<1)}}{|x - \xi|^{n+2s}} d\xi.$$

Caffarelli–Silvestre have developed in [9] an equivalent definition using an extension problem, that is crucial in the present work. For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, we construct the extension $u : \mathbb{R}^n \times [0, +\infty) \rightarrow \mathbb{R}$, $u = u(x, y)$, as the solution of the equation

$$\begin{cases} \Delta_x u + \frac{a}{y} \partial_y u + \partial_{yy} u = 0 & \text{for } x \in \mathbb{R}^n, y \in [0, +\infty), \\ u(x, 0) = f(x), \end{cases} \tag{2.2}$$

for $s = \frac{1-a}{2}$. This type of degenerate elliptic equations have been studied in [11]. Then the fractional Laplacian of f can be recovered as

$$(-\Delta)^s f = c_{n,s} \lim_{y \rightarrow 0} y^a \partial_y u,$$

i.e., we are looking at a non-local Dirichlet-to-Neumann operator. Note that Eq. (2.2) can be written in divergence form as

$$\operatorname{div}(y^a \nabla u) = 0,$$

which is the Euler–Lagrange equation for the functional

$$J[u] = \int_{x \in \mathbb{R}^n, y > 0} |\nabla u|^2 y^a dx dy.$$

To finish, just mention that the Poisson kernel for the fractional Laplacian $(-\Delta)^s$ is given by

$$P(x, y) = c_{n,s} \frac{y^{1-a}}{(|x|^2 + |y|^2)^{\frac{n+1-a}{2}}}, \tag{2.3}$$

and thus $u = P *_x f$.

3 Γ -convergence in the interior

The theory of phase transitions (see Modica [17], Modica and Mortola [19] for the classical references, or Alberti [3], for a very well written survey), studies the interface between two fluids in a container Ω neglecting the interaction with the boundary. In particular, it is proven that the functionals E_ϵ defined in (1.4), Γ -converge to E given in (1.3). In this section we consider the generalization to E_ϵ^a given in (3.1), that involves a singular weight $w_a := h^a$. However, since the interaction with the boundary is neglected and the weight is regular in the interior, the behavior of E_ϵ^a is going to be very similar to E_ϵ . Indeed, we can easily modify the argument of Modica to prove:

Proposition 4 *Let A be a domain contained in Ω , $h := \text{dist}(\cdot, \partial\Omega)$, $w_a := h^a$. Assume, in addition, that $\text{dist}(A, \partial\Omega) \geq r$ for some $r > 0$ fixed. Set*

$$E_\epsilon^a[u, A] := \epsilon^{1-a} \int_A |\nabla u|^2 h^a + \frac{1}{\epsilon^{1-a}} \int_A W(u) h^{-a} \tag{3.1}$$

and

$$E^a[u, A] := \sigma \mathcal{H}^2(S_u \cap A) \tag{3.2}$$

for $\sigma := 2 \int_\alpha^\beta \sqrt{W}$. Then

- i. $\forall u \in BV(A, I), \forall (u_\epsilon) \subset W^{1,2}(A, w_a)$ such that $u_\epsilon \rightarrow u$ in $L^1(A)$, we have

$$\liminf_{\epsilon \rightarrow 0} E_\epsilon^a[u_\epsilon, A] \geq \sigma E^a[u, A].$$

- ii. $\forall u \in BV(A, I)$, there exists $(u_\epsilon) \subset W^{1,2}(A, w_a)$ such that $u_\epsilon \rightarrow u$ in $L^1(A)$ and

$$\limsup_{\epsilon \rightarrow 0} E_\epsilon^a[u_\epsilon, A] \leq \sigma E^a[u, A].$$

Moreover, when S_u is a closed Lipschitz surface in A , the functions u_ϵ may be required to be Lipschitz continuous with constant $\frac{C}{\epsilon^{1-a} r^a}$, and to converge uniformly to u in every set with positive distance from S_u and away from the boundary.

- iii. Any sequence $(u_\epsilon) \subset W^{1,2}(A, w_a)$ with uniformly bounded energies $E_\epsilon^a[u_\epsilon, A]$ is pre-compact in X and every cluster point belongs to $BV(A, I)$.

Proof The proof is essentially the one of Modica and can be found in Alberti [3]. By a well known truncation argument [4, lemma 1.14], we can assume that $u : A \rightarrow [\alpha, \beta]$. Now, use the inequality $x_1^2 + x_2^2 \geq 2x_1x_2$ with $x_1 = \epsilon^{(1-a)/2} |\nabla u| h^{a/2}$ and $x_2 = \epsilon^{-(1-a)/2} W^{1/2}(u) h^{-a/2}$, then

$$E_\epsilon^a[u, A] \geq 2 \int_A \sqrt{W(u)} |\nabla u| = \int_A |\nabla(\mathcal{W}(u))|, \tag{3.3}$$

where $\mathcal{W} : [\alpha, \beta] \rightarrow \mathbb{R}$ is a primitive of $2\sqrt{W}$. This gives *iii.* and *i.* using standard arguments, and can be found exactly in [3, paragraph 4.5].

For *ii.* we need to take care of the weight w_a in the construction. Let $u \in BV(A, I)$. Without loss of generality, we can assume that its singular set S_u is a Lipschitz surface in A , even a polyhedral surface of dimension 2 (see [15, theorem 1.24]). We would like to construct a sequence of functions u_ϵ that converges to u in $L^1(A)$. First, it is possible to give

coordinates $(e(x), \omega)$ in Ω such that ω parameterizes S_u , and $e(x) \in \mathbb{R}$ is just the signed distance to S_u (positive where $u = \beta$ and negative where $u = \alpha$). Next, solve the ODE

$$\theta' = \sqrt{W(\theta)} \tag{3.4}$$

with initial condition $\theta(0) = \frac{\alpha+\beta}{2}$. This θ is the well known optimum profile for the case $a = 0$. In order to take into account the weight w_a we set, for every fixed ω , (note that $(0, \omega)$ is any point in S_u),

$$\phi_\omega(t) := \phi(t, \omega) = \theta\left(\frac{t}{h^a(0, \omega)}\right). \tag{3.5}$$

Now, for each $\epsilon > 0$, let $t = e(x)/\epsilon^{1-a}$ and

$$u_\epsilon(x) := \phi_\omega\left(\frac{e(x)}{\epsilon^{1-a}}\right). \tag{3.6}$$

When $\epsilon \rightarrow 0$ we can compute that

$$|\nabla u_\epsilon|^2(x) = \frac{1}{\epsilon^{2(1-a)}} [\phi'_\omega(t)^2 + R(\omega, t)o(1)].$$

Then we can use the coarea formula to calculate the energy of this function:

$$\begin{aligned} E_\epsilon^a[u_\epsilon, A] &= \epsilon^{1-a} \int_A |\nabla u_\epsilon|^2 h^a + \frac{1}{\epsilon^{1-a}} \int_A W(u_\epsilon) h^{-a} \\ &= \frac{1}{\epsilon^{1-a}} \int_A [(\phi'_\omega(t))^2 h^a dx + W(\phi_\omega(t)) h^{-a} + o(1)] dx \\ &= \int_{-\infty}^{+\infty} \int_{\Sigma_{\epsilon^{1-a}t}} [(\phi'_\omega(t))^2 h^a + W(\phi_\omega(t)) h^{-a} + o(1)] d\omega dt. \end{aligned}$$

When $\epsilon \rightarrow 0$, the level set $\Sigma_{\epsilon^{1-a}t}$ converges to $S_u \cap A$, and if x is written in the new coordinates $(e(x), \omega)$, then $h(t, \omega)$ converges to $\text{dist}((0, \omega), \partial\Omega) = h(0, \omega)$. Taking the limit we have that

$$\limsup_{\epsilon \rightarrow 0} E_\epsilon^a[u_\epsilon, A] = \int_{-\infty}^{+\infty} \int_{S_u \cap A} [(\phi'_\omega(t))^2 h^a(0, \omega) + W(\phi_\omega(t)) h^{-a}(0, \omega)] d\omega dt.$$

But because of (3.4) and (3.5), both terms in the above integration are equal. Then the inequality $x_1^2 + x_2^2 \geq 2x_1x_2$ for two positive numbers $x_1 = x_2$ becomes an equality and thus

$$\begin{aligned} \limsup_{\epsilon \rightarrow 0} E_\epsilon^a[u_\epsilon, A] &= \int_{S_u \cap A} \int_{-\infty}^{+\infty} 2\sqrt{W(\phi_\omega(t))} \phi'_\omega(t) dt d\omega \\ &= \int_{S_u \cap A} \int_\alpha^\beta 2\sqrt{W(r)} dr d\omega = \sigma \mathcal{H}^2(S_u \cap A) \end{aligned}$$

as we wished. The Lipschitz constant of u_ϵ is computed from (3.5) and (3.6). □

4 Perturbation of the norm H^s

Let $1/2 < s < 1$. In this section we consider a singular perturbation of the norm $H^s(E)$ when E is a bounded interval in \mathbb{R} . As usual, denote $s = \frac{1-a}{2}$, so $-1 < a < 0$, and $I' := \{\alpha', \beta'\}$. Consider a double-well potential $V : \mathbb{R} \rightarrow [0, \infty)$ vanishing only at I' , with the same hypothesis as in the introduction.

More precisely, we will study the Γ -convergence of the functional

$$G_\epsilon^a[v, E] := \frac{\epsilon^{1-a}}{D_s} \int_{E^2} \frac{|v(x') - v(x)|^2}{|x' - x|^{1+2s}} dx'dx + \lambda_\epsilon \int_E V(v)dx, \tag{4.1}$$

when $\epsilon \rightarrow 0$. Although D_s could be any positive constant in this section, we will fix its value as given in (5.20).

The functional G_ϵ^a presents the following scaling property that justifies the election of λ_ϵ in (1.10). Indeed, if we set $u^\epsilon(x) := u(\Lambda_\epsilon x)$, $E_\epsilon := \{x : \Lambda_\epsilon x \in E\}$, then we immediately see that

$$G_\epsilon^a[u, E] = G_1^a[u^\epsilon, E_\epsilon].$$

It is interesting to observe the deterioration of λ_ϵ when $a \rightarrow 0$. In particular, the functional for $s = 1/2, a = 0$ studied in [5] reads

$$G_\epsilon[v, E] := \frac{\epsilon}{2\pi} \int_{E^2} \frac{|v(x') - v(x)|^2}{|x' - x|^2} dx'dx + \lambda_\epsilon^0 \int_E V(v)dx,$$

for $\lim_{\epsilon \rightarrow 0} \epsilon \log \lambda_\epsilon^0 = K, 0 < K < \infty$.

It is natural then to consider profiles on the whole real line that minimize the energy, i.e., we consider the following optimal profile problem

$$\kappa_s := \inf \left\{ \frac{1}{D_s} \int_{\mathbb{R}^2} \frac{|v(x) - v(x')|^2}{|x - x'|^{1+2s}} dx dx' + \int_{\mathbb{R}} V(v) dx \right\}, \tag{4.2}$$

where the infimum is taken among all the functions in the set

$$\left\{ v \in H^s(\mathbb{R}) : \lim_{x \rightarrow -\infty} v(x) = \alpha', \quad \lim_{x \rightarrow +\infty} v(x) = \beta' \right\}.$$

Then we can prove:

Proposition 5 *Let E be an interval in \mathbb{R} . The functional defined on $L^1(E)$ given by*

$$\bar{G}_\epsilon^a[v, E] := \begin{cases} G_\epsilon^a[v, E], & \text{if } v \in H^s(E), \\ +\infty, & \text{otherwise,} \end{cases}$$

Γ -converges in $L^1(E)$ to

$$\bar{G}_0^a[v, E] := \begin{cases} \kappa_s \mathcal{H}^0(S_v) & \text{if } v \in BV(E, I'), \\ +\infty, & \text{otherwise,} \end{cases}$$

where $\mathcal{H}^0(S_v)$ represents number of points in the singular set of $v \in BV(E, I')$. The constant $\kappa_s > 0$ is given by the optimal profile problem (4.2). Moreover, every sequence $(v_\epsilon) \subset L^1(E)$ with uniformly bounded energies $G_\epsilon^a[v_\epsilon, E]$ is precompact in $L^1(E)$ and every cluster point belongs to $BV(E, I')$.

The proof of this proposition is similar to the work of Garroni and Palatucci [13, theorem 2.1]. Indeed, they considered the functional

$$\tilde{G}_{\tilde{\epsilon}}^p[v, E] := \tilde{\epsilon}^{p-2} \int_{E^2} \frac{|v(x) - v(x')|^p}{|x - x'|^p} dx dx' + \frac{1}{\tilde{\epsilon}} \int_E V(v) dx \tag{4.3}$$

for $p > 2$. Our case is analogous if we take $p = 2s + 1$ and $\tilde{\epsilon} = \Lambda_\epsilon$ since the exponent of the term $|v(x) - v(x')|$ does not play any special role in their proof.

We remark that the case $s = 1/2$ was considered in [5]. The main difference with respect to the case $1/2 < s < 1$, is that here the optimal profile is characterized by the equipartition of the energy between the two terms in the functional. Instead, the logarithmic scaling for $s = 1/2$ produces no equipartition of the energy: the limit comes only from the non-local part and does not depend on V , i.e., any profile is optimal as far as the transition occurs on a layer of order Λ_ϵ . This does not happen in our case; both terms of the energy (4.1) are equally important.

A related result on Gamma-convergence and optimal profiles was obtained by Alberti–Bellettini [4] for anisotropic singularities that are integrable, unlike our $\frac{1}{|x-x'|^{1+2s}}$.

The proof of Proposition 5 is given in the following, and it is divided into three parts: first we show compactness, then we give some preliminary results on the optimal profile problem (4.2), and in the last part we show the upper and lower bounds of the Γ -convergence.

4.1 Compactness

We start with a (non-optimal) bound from below:

Lemma 6 *Let δ be given such that $0 < \delta < (\beta' - \delta')/2$. For every interval $J \subset E$, $\epsilon > 0$, and $(v_\epsilon) \subset L^1(E)$, let A_ϵ and B_ϵ be the sets of all points $x \in J$ such that $v_\epsilon(x) \leq \alpha' + \delta$ and $v_\epsilon(x) \geq \beta' - \delta$, respectively. Set*

$$a_\epsilon := \frac{|A_\epsilon \cap J|}{|J|}, \quad b_\epsilon := \frac{|B_\epsilon \cap J|}{|J|}. \tag{4.4}$$

Then

$$G_\epsilon^a[v_\epsilon, J] \geq \epsilon^{1-a} C_s \frac{(\beta' - \alpha' - 2\delta)^2}{|J|^{2s-1}} \left\{ 1 - \frac{1}{(1 - a_\epsilon)^{2s-1}} - \frac{1}{(1 - b_\epsilon)^{2s-1}} \right\} + C_\delta, \tag{4.5}$$

where the constant C_δ , does not depend on ϵ , and C_s only depends on s .

Proof First note that the non-local part of the energy decreases under monotone rearrangements (see Lemma 7). Set $J = (a_0, b_0)$. Then, if we define v^* to be the non-decreasing rearrangement of v , we have that

$$\begin{aligned} \int_{J^2} \frac{|v(x') - v(x)|^2}{|x' - x|^{1+2s}} dx' dx &\geq \int_{J^2} \frac{|v^*(x') - v^*(x)|^2}{|x' - x|^{1+2s}} dx' dx \\ &\geq 2(\beta' - \alpha' - 2\delta)^2 \int_{a_0}^{a_0+a_\epsilon|J|} \int_{b_0-b_\epsilon|J|}^{b_0} \frac{1}{|x - x'|^{1+2s}} dx' dx \\ &= \frac{2(\beta' - \alpha' - 2\delta)^2}{2s(2s - 1) |J|^{2s-1}} \left[1 - \frac{1}{(1 - a_\epsilon)^{2s-1}} - \frac{1}{(1 - b_\epsilon)^{2s-1}} + \frac{1}{(1 - a_\epsilon - b_\epsilon)^{2s-1}} \right]. \end{aligned}$$

On the other hand, let $m_\delta := \inf \{V(t) : \alpha' + \delta \leq t \leq \beta' - \delta\}$, we obtain

$$\int_J V(v)dx \geq m_\delta |J| (1 - a_\epsilon - b_\epsilon),$$

so then

$$G_\epsilon^a[v_\epsilon, J] \geq \epsilon^{1-a} \frac{2(\beta' - \alpha' - 2\delta)^2}{2s(2s - 1)D_s |J|^{2s-1}} + \lambda_\epsilon m_\delta |J| (1 - a_\epsilon - b_\epsilon).$$

Minimizing with respect to $|J| (1 - a_\epsilon - b_\epsilon)$, and taking into account that $s = \frac{1-a}{2}$ we get

$$G_\epsilon^a[v_\epsilon, J] \geq \epsilon^{1-a} \frac{2(\beta' - \alpha' - 2\delta)^2}{2s(2s - 1)D_s |J|^{2s-1}} \left[1 - \frac{1}{(1 - a_\epsilon)^{2s-1}} - \frac{1}{(1 - b_\epsilon)^{2s-1}} \right] + \frac{2^{\frac{1}{2s}} (2s)^{\frac{2s-1}{2s}} (\beta' - \alpha' - 2\delta)^{\frac{2}{2s}} m_\delta^{\frac{2s-1}{2s}}}{2s - 1}$$

for every $0 < \delta < (\beta' - \alpha')/2$, and the lemma is proved. □

As we have mentioned, the main ingredient in the proof above is the following rearrangement result. In particular, it tells us that the infimum of the functional must be attained at a non-decreasing function.

Lemma 7 [14] *Let*

$$I_{\Psi,p}(v) = \int_0^1 \int_0^1 \Psi \left(\frac{v(x) - v(x')}{p(x - x')} \right) dx dx',$$

where Ψ and p are restricted as follows:

1. $\Psi(t)$ is defined and continuous on \mathbb{R} and $\Psi(t) = \Psi(-t)$ is strictly increasing as $|t| \rightarrow \infty$.
2. $p(t)$ is defined an continuous on $(-1, 1)$ and $p(u) = p(-u)$ is strictly decreasing as $|t| \rightarrow 0$.
3. $\Psi(e^x)$ is convex.

We also define the non-decreasing rearrangement of f as

$$f^*(x) = \inf \{ \lambda : m\{t : f(t) \geq \lambda\} \leq x \}.$$

Then

$$I_{\Psi,p}(f^*) \leq I_{\Psi,p}(f).$$

Now we are ready to prove compactness:

Proposition 8 *Let (v_ϵ) be a sequence in $H^s(E)$ with equibounded energy $G_\epsilon^a[v_\epsilon, E] \leq C$. Then (v_ϵ) is precompact in $L^1(E)$ and every cluster point belongs to $BV(E, I')$.*

Proof Once we have the estimate from Lemma 6, it is a standard argument that we rewrite here for completeness (see theorem 4.4, part (i), in [6]), and to show how the estimate (4.5) is used. First, the condition $G_\epsilon^a[v_\epsilon, E] \leq C$ implies that

$$\int_E V(v_\epsilon)dx \leq C\lambda_\epsilon^{-1}, \tag{4.6}$$

and we obtain that $V(v_\epsilon) \rightarrow 0$ in $L^1(E)$ when $\epsilon \rightarrow 0$. Thanks to the growth assumption on V , (v_ϵ) is weakly relatively compact in $L^1(E)$, and some subsequence, still denoted by (v_ϵ) , converges weakly in $L^1(E)$ to some v .

In order to prove that this convergence is strong in $L^1(E)$ and that $v \in BV(E, I')$, we need to use the properties of Young measures (see the notes [27]). Let ν_x be the Young measure associated with (v_ϵ) . Since V is a non-negative continuous function in \mathbb{R} , then

$$\int_E \int_{\mathbb{R}} V(t) d\nu_x(t) \leq \liminf_{\epsilon \rightarrow 0} \int_E V(v_\epsilon) dx.$$

Hence, by (4.6) we have that

$$\int_{\mathbb{R}} V(t) d\nu_x(t) = 0, \quad \text{a.e. } x \in E.$$

Since $V(t) = 0$ if and only if $t = \alpha'$ or $t = \beta'$, the probability measure ν_x is supported on $I' = \{\alpha', \beta'\}$ for a.e. x . In other words, there exists a function $\theta : E \rightarrow [0, 1]$ such that

$$\nu_x(dt) = \theta(x)\delta_{\alpha'}(dt) + (1 - \theta(x))\delta_{\beta'}(dt), \quad x \in E,$$

and

$$v(x) = \theta(x)\alpha' + (1 - \theta(x))\beta', \quad x \in E.$$

It remains to prove that θ belongs to $BV(E, \{0, 1\})$. Let us consider the set S of the points where the approximate limits of θ is neither 0 nor 1. For every $N \leq \mathcal{H}^0(S)$, we can find N disjoint intervals $\{J_j\}_{j=1, \dots, N}$ such that $J_j \cap S \neq \emptyset$ and such that the quantities a_ϵ^j and b_ϵ^j defined by (4.4) replacing J by J_j satisfy

$$a_\epsilon^j \rightarrow a_j \in (0, 1) \quad \text{and} \quad b_\epsilon^j \rightarrow b_j \in (0, 1) \quad \text{as } \epsilon \rightarrow 0.$$

Then we can apply Lemma 6 in the interval J_j and, taking the limit as $\epsilon \rightarrow 0$ in the inequality (4.5) we obtain that

$$\liminf_{\epsilon \rightarrow 0} G_\epsilon^a[v_\epsilon, J_j] \geq C_\delta.$$

Finally, we use the sub-additivity of G_ϵ^a , and we get

$$\liminf_{\epsilon \rightarrow 0} G_\epsilon^a[v_\epsilon, E] \geq \sum_{j=1}^N \liminf_{\epsilon \rightarrow 0} G_\epsilon^a[v_\epsilon, J_j] \geq NC_\delta.$$

Since (v_ϵ) has equi-bounded energy, this implies that S is a finite set. Hence, $\theta \in BV(E, \{0, 1\})$ and the proof of the compactness for G_ϵ^a is complete. \square

4.2 The optimal profile

We remind the reader that we have set the following optimal profile problem

$$\kappa_S := \inf \left\{ G_1^a[v, \mathbb{R}] : v \in H^s(\mathbb{R}), \lim_{x \rightarrow -\infty} v(x) = \alpha', \lim_{x \rightarrow +\infty} v(x) = \beta' \right\}. \quad (4.7)$$

We would like to show that this infimum is attained.

Lemma 9 *Let $0 < \delta < \frac{1}{2}(\beta' - \alpha')$ and set*

$$m_\delta := \min\{V(t) : \alpha' + \delta \leq t \leq \beta' - \delta\}.$$

Then, for any $v \in H^s(\mathbb{R})$ with $\lim_{x \rightarrow -\infty} v(x) = \alpha'$ and $\lim_{x \rightarrow +\infty} v(x) = \beta'$, it holds that

$$G_1^a[v, \mathbb{R}] \geq \frac{(2s)^{\frac{2s-1}{2s}}}{2s-1}(\beta' - \alpha' - 2\delta)^{\frac{2}{2s}} m_\delta^{\frac{2s-1}{2s}} > 0.$$

Proof It is analogous to proposition 3.1. in [13] for the super-quadratic case, and follows the lines of Lemma 6. Fix $\delta > 0$ and fix $v \in H_{loc}^s(\mathbb{R})$ such that $\lim_{x \rightarrow -\infty} v(x) = \alpha'$, $\lim_{x \rightarrow +\infty} v(x) = \beta'$ and $G_1^a[v, \mathbb{R}] < \infty$. Define

$$I_{\alpha'} := \{x \in \mathbb{R} : v(x) \leq \alpha' + \delta\} \quad \text{and} \quad I_{\beta'} := \{x \in \mathbb{R} : v(x) \geq \beta' - \delta\},$$

and also $J_\delta := \mathbb{R} \setminus (I_{\alpha'} \cup I_{\beta'})$. Notice that $I_{\alpha'}$, $I_{\beta'}$ and J_δ are non-empty, and that J_δ is bounded, for every fixed $\delta \in (0, (\beta' - \alpha')/2)$. Consider the truncated function

$$v_\delta(x) := (v(x) \vee (\alpha' + \delta)) \wedge (\beta' - \delta) \quad \text{for every } x \in \mathbb{R}.$$

It is easy to see that the non-local energy decreases under truncation and then it follows that

$$\begin{aligned} G_1^a[v, \mathbb{R}] &\geq \int_{\mathbb{R} \times \mathbb{R}} \frac{|v_\delta(x) - v_\delta(x')|^2}{|x - x'|^{1+2s}} dx dx' + \int_{\mathbb{R}} V(v) dx \\ &\geq \int_{\mathbb{R} \times \mathbb{R}} \frac{|v_\delta(x) - v_\delta(x')|^2}{|x - x'|^{1+2s}} dx dx' + m_\delta |J_\delta|. \end{aligned} \tag{4.8}$$

We set

$$x_{\alpha'} := \min\{x : v(x) > \alpha' + \delta\}, \quad x_{\beta'} := \max\{x : v(x) < \beta' - \delta\}.$$

Since $v_\delta(x) = \alpha' + \delta$ for every $x < x_{\alpha'}$ and $v_\delta(x) = \beta' - \delta$ for every $x > x_{\beta'}$, for any interval $J \supset [x_{\alpha'}, x_{\beta'}]$ the non-decreasing rearrangement v_δ^* of v_δ in J does not depend on J . Because the rearrangement decreases the non-local energy (Lemma 7), we have that

$$\begin{aligned} \int_{\mathbb{R} \times \mathbb{R}} \frac{|v_\delta(x) - v_\delta(x')|^2}{|x - x'|^{1+2s}} dx dx' &\geq \int_{\mathbb{R} \times \mathbb{R}} \frac{|v_\delta^*(x) - v_\delta^*(x')|^2}{|x - x'|^{1+2s}} dx dx' \\ &\geq \int_{-\infty}^{x_{\alpha'}^*} \int_{x_{\beta'}^*}^{+\infty} \frac{|v_\delta^*(x) - v_\delta^*(x')|^2}{|x - x'|^{1+2s}} dx dx', \end{aligned}$$

where $x_{\alpha'}^* := \sup\{x : v_\delta^*(x) = \alpha' + \delta\}$ and $x_{\beta'}^* := \inf\{x : v_\delta^*(x) = \beta' - \delta\}$. So from (4.8) we deduce that

$$\begin{aligned} G_1^a[v, \mathbb{R}] &\geq (\beta' - \alpha' - 2\delta)^2 \int_{-\infty}^{x_{\alpha'}^*} \int_{x_{\beta'}^*}^{+\infty} \frac{1}{|x - x'|^{1+2s}} dx dx' + m_\delta |J_\delta| \\ &= \frac{(\beta' - \alpha' - 2\delta)^2}{2s(2s - 1) |J_\delta|^{2s-1}} + m_\delta |J_\delta|. \end{aligned}$$

Minimizing with respect to $|J_\delta|$ we obtain

$$G_1^a[v, \mathbb{R}] \geq \frac{(2s)^{\frac{2s-1}{2s}}}{2s-1} (\beta' - \alpha' - 2\delta)^{\frac{2}{2s}} m_\delta^{\frac{2s-1}{2s}} > 0,$$

and the lemma is proved. □

Corollary 10 *The constant κ_s is strictly positive.*

We also need an auxiliary optimal profile problem: for every $T > 0$, we minimize

$$\kappa_s^T := \inf \{ G_1^a[v, \mathbb{R}] : v \in H^s(\mathbb{R}), v(x) = \alpha' \forall x \leq T, v(x) = \beta' \forall x \geq T \}.$$

Proposition 11 *The minimum is achieved by a function $\phi^T \in H^s(\mathbb{R})$ which is non-decreasing and satisfies $\alpha' \leq \phi^T \leq \beta'$. Moreover the sequence κ_s^T is non-increasing in T and $\lim_{T \rightarrow +\infty} \kappa_s^T = \kappa_s$.*

Proof This is essentially proposition 3.2 in [13]. □

Then we have that

Proposition 12 *The infimum for the optimum profile problem (4.7) is achieved by a non-decreasing function ϕ satisfying $\alpha' \leq \phi \leq \beta'$.*

Proof It is a straightforward modification of proposition 3.3 in [13]. Let $T > 0$ and let ϕ^T be a non-decreasing minimizer for κ_s^T . Since the functions ϕ^T are monotone and bounded, by Helly’s theorem, there exist a subsequence ϕ^{T_k} of ϕ^T and a non-decreasing function ϕ , bounded by α' and β' , such that ϕ^{T_k} converges pointwise in \mathbb{R} to ϕ . By Fatou’s lemma and Proposition 11 we also have

$$\int_{\mathbb{R} \times \mathbb{R}} \frac{|\phi(x) - \phi(x')|^2}{|x - x'|^{1+2s}} dx dx' + \int_{\mathbb{R}} V(\phi) dx \leq \lim_{k \rightarrow \infty} \kappa_s^{T_k} = \kappa_s.$$

This ϕ is the minimizer we seek. □

4.3 Lower and upper bound inequalities

The proof of the upper and lower bound inequalities of Proposition 5 use the optimal profile obtained in Proposition 12. The lower bound inequality is a consequence of the following:

Proposition 13 *Let J be an open interval of \mathbb{R} . Let (v_ϵ) be a non-decreasing sequence in $H^s(J)$ and assume that there exist $\bar{a}, \bar{b} \in J, \bar{a} < \bar{b}$, such that for every $\delta > 0$, there exists ϵ_δ such that*

$$v_\epsilon(\bar{a}) \leq \alpha' + \delta \text{ and } v_\epsilon(\bar{b}) \geq \beta' - \delta, \quad \forall \epsilon \leq \epsilon_\delta.$$

Then

$$\liminf_{\epsilon \rightarrow 0} G_\epsilon^a[v_\epsilon, J] \geq \kappa_s.$$

Proof It is exactly the same as proposition 5.1 in [13] with the modifications indicated after (4.3) and we do not find it necessary to include it here. □

Clearly, an analog proposition holds in the case of u_ϵ non-increasing satisfying the hypothesis with $\bar{a} > \bar{b}$. Next, thanks to the compactness result for G_ϵ^a from Proposition 8, we may assume that the sequence (v_ϵ) converges in $L^1(E)$ to some $v \in BV(E, I')$. Hence, the jump set S_v is finite and we can find $N := \mathcal{H}^0(S_u)$ disjoint subintervals $\{J_j\}_{j=1, \dots, N}$ such that $S_v \cap J_j \neq \emptyset$ for every $j = 1, \dots, N$.

Now, let us consider the monotone rearrangement $v_{\epsilon, j}^*$ of v_ϵ in J_j . This rearrangement is non-decreasing if v is non-decreasing in J_j , and non-increasing otherwise. With this choice, clearly $v_{\epsilon, j}^*$ converges to v in $L^1(J_j)$ and thus it satisfies the assumptions of Proposition 13 with J replaced by J_j . Then, for every $j = 1, \dots, N$, we may conclude that

$$\liminf_{\epsilon \rightarrow 0} G_\epsilon^a[v_\epsilon, J_j] \geq \liminf_{\epsilon \rightarrow 0} G_\epsilon^a[v_{\epsilon, j}^*, J_j] \geq \kappa_s.$$

Finally, using the subadditivity of $G_\epsilon^a[v_\epsilon, \cdot]$, we get

$$\liminf_{\epsilon \rightarrow 0} G_\epsilon^a[v_\epsilon, J] \geq \sum_{j=1}^N G_\epsilon^a[v_\epsilon, J_j] \geq N\kappa_s = \kappa_s \mathcal{H}^0(S_v).$$

The lower bound is shown.

We prove first the upper bound inequality for a v of the form

$$v(x) = \begin{cases} \alpha', & \text{if } x \leq x_0, \\ \beta', & \text{if } x > x_0. \end{cases}$$

Let $T > 0$ be fixed and let $\phi^T \in H_{\text{loc}}^s(\mathbb{R})$ be the minimizer for κ_s^T found in Proposition 11. It satisfies $\phi^T(x) = \alpha'$ for all $x \leq -T$, $\phi^T(x) = \beta'$ for all $x \geq T$ and $G_1^a[\phi^T, \mathbb{R}] = \kappa_s^T$.

Let us define, for every $\epsilon > 0$, $v_\epsilon(x) := \phi^T\left(\frac{x-x_0}{\Lambda\epsilon}\right)$, for every $x \in E$. We have that $v_\epsilon \rightarrow v$ in $L^1(E)$ and

$$G_\epsilon^a[v_\epsilon, E] = G_1^a[\phi^T, (E - x_0)/\Lambda\epsilon] \leq G_1^a[\phi^T, \mathbb{R}] = \kappa_s^T.$$

By Proposition 11 again, we get

$$\lim_{T \rightarrow +\infty} \limsup_{\epsilon \rightarrow 0} G_\epsilon^a[v_\epsilon, E] \leq \kappa_s.$$

Then, by a diagonalization argument, we can construct a sequence \tilde{v}_ϵ converging to v in $L^1(E)$, which satisfies,

$$\limsup_{\epsilon \rightarrow 0} G_\epsilon^a[\tilde{v}_\epsilon, E] \leq \kappa_s.$$

The optimal sequence for an arbitrary $v \in BV(E, I')$ can be easily obtained gluing the sequences constructed above for each single jump of v and taking into account that the long range interactions decay as $\epsilon \rightarrow 0$.

5 Some new trace inequalities

It is well known that traces of functions in some Sobolev spaces are represented by functions in Besov spaces (see the book [1, chapters 4 and 7]). Here we would like to consider extensions of this result to weighted Sobolev spaces, for the case of domains in \mathbb{R}^2 . Let $-1 < a < 0$, and $\Omega \subset \mathbb{R}_+^2$ a bounded domain with Lipschitz boundary. Theorem 3 gives that the class of traces on M , a piece of $\partial\Omega$, of the space $W^{1,2}(\Omega, w_a)$ is just the space $H^s(M)$. However,

Nekvinda’s proof does not give an exact value of the constant in the embedding. We seek to give a precise inequality between functions in $W^{1,2}(\Omega, w_a)$ and their traces.

This section is divided into four parts. First we summarize some lemmas that will be needed in the proof. In the second part, we prove the embedding inequality when Ω is the half-plane \mathbb{R}_+^2 (Lemma 17). Next, we localize the inequality for a domain $\Omega = (-1, 1) \times (0, 1)$; this is the main result of the section and it is contained in Theorem 19. Last, we show that the inequality is optimal in some sense.

5.1 Technical lemmas

In order to understand the energy term related to the fractional Laplacian, first we need to study Eq. (2.2). We prove a basic auxiliary result:

Lemma 14 *Consider the following ODE defined for $y \in \mathbb{R}^+$:*

$$-\varphi(y) + \frac{a}{y}\varphi_y(y) + \varphi_{yy}(y) = 0.$$

It has two linearly independent solutions. In particular,

$$\varphi(y) = y^s [c_1 I_s(y) + c_2 K_s(y)] \tag{5.1}$$

for constants $c_1, c_2 \in \mathbb{R}$. Here I_s, K_s are the modified Bessel functions; their asymptotic behavior is given precisely in (5.3) and (5.4).

Proof We actually have explicit formulas for the solution. Indeed, the change $\varphi(y) = y^s \psi(y)$ gives that ψ must be a solution of the modified Bessel equation

$$y\psi'' + y\psi' - (y^2 + s^2)\psi = 0. \tag{5.2}$$

The books [2, 28] are classical references for this equation. There are two independent solutions of (5.2): I_s , the modified Bessel function of the first kind, and K_s , the modified Bessel function of second kind (see 9.6.1. in [2]). In particular, they have the following asymptotic behavior (cf. 9.6.7 and 9.7.1 in [2]):

$$I_s(y) \sim \frac{1}{\Gamma(s+1)} \left(\frac{y}{2}\right)^s, \quad K_s(y) \sim \frac{\Gamma(s)}{2} \left(\frac{2}{y}\right)^s \quad \text{when } y \rightarrow 0^+. \tag{5.3}$$

And when $y \rightarrow +\infty$,

$$I_s(y) \sim \frac{1}{\sqrt{2\pi y}} e^y, \quad K_s(y) \sim \sqrt{\frac{\pi}{2y}} e^{-y}. \tag{5.4}$$

□

We show now a result from [9] that characterizes the minimizers of the energy in \mathbb{R}_+^2 :

Lemma 15 *Given $v \in H^s(\mathbb{R})$ for $1/2 < s < 1$, there exists a minimizer u_0 of the following variational problem*

$$J(u) = \int_{\mathbb{R}_+^2} |\nabla u|^2 w_a \tag{5.5}$$

in the space $Y := \{u \in L^2_{\text{loc}}(\mathbb{R}_+^2, w_a) : \nabla u \in L^2(\mathbb{R}_+^2, w_a)\}$, subject to the constraint $u(x, 0) = v(x)$ for all $x \in \mathbb{R}$, and the weight is taken as $w_a := y^a$. Moreover,

$$u_0 = P *_x v,$$

where P is the Poisson kernel defined in (2.3) and

$$\int_{\mathbb{R}_+^2} |\nabla u_0|^2 w_a \, dx dy = e_s \int_{\mathbb{R}} |\hat{v}(\xi)|^2 |\xi|^{2s} \, d\xi$$

for some positive constant e_s , depending only on s , whose precise value is given in (5.8).

Proof The Euler–Lagrange equation of J is

$$\begin{cases} \operatorname{div}(y^a \nabla u) = 0 \text{ in } \mathbb{R}_+^2 \\ u = v \text{ on } \mathbb{R} \times \{0\}. \end{cases} \tag{5.6}$$

The main idea is to reduce (5.6) to an ODE by taking Fourier transform in x . We obtain

$$-|\xi|^2 \hat{u}(\xi, y) + \frac{a}{y} \hat{u}_y(\xi, y) + \hat{u}_{yy}(\xi, y) = 0,$$

that is ODE for each fixed value of ξ .

On the other hand, consider the one-dimensional functional for each $\varphi : [0, +\infty) \rightarrow \mathbb{R}$ given by

$$\bar{J}[\varphi] = \int_{\mathbb{R}_+} \{(\varphi')^2(t) + \varphi^2(t)\} t^a \, dt$$

subject to the conditions $\varphi(0) = 1$ and $\lim_{t \rightarrow +\infty} \varphi(t) = 0$. Its Euler–Lagrange equation is given by

$$-\varphi(y) + \frac{a}{y} \varphi_y(y) + \varphi_{yy}(y) = 0,$$

that has been well understood in Lemma 14. Indeed, if φ_0 is the minimizer of \bar{J} , its precise formula is given by (5.1) with $c_1 = 0, c_2 = cst(s)$.

Thus we conclude that the minimizer we seek in the present lemma satisfies $\hat{u}_0(\xi, y) = \hat{v}(\xi)\varphi_0(|\xi|y)$. Next, just compute

$$\begin{aligned} \int_{\mathbb{R}_+^2} |\nabla u_0|^2 y^a &= \int_{\mathbb{R}} \int_0^\infty (|\xi|^2 |\hat{u}_0|^2 + |\partial_y \hat{u}_0|^2) y^a \, dy d\xi \\ &= \int_{\mathbb{R}} \int_0^\infty |\hat{v}(\xi)|^2 |\xi|^2 (|\varphi_0(|\xi|y)|^2 + |\varphi_0'(|\xi|y)|^2) y^a \, dy d\xi \\ &= \int_{\mathbb{R}} |\hat{v}(\xi)|^2 |\xi|^{1-a} \int_0^\infty (|\varphi_0(t)|^2 + |\varphi_0'(t)|^2) t^a \, dt d\xi \\ &= \bar{J}[\varphi_0] \int_{\mathbb{R}} |\hat{v}(\xi)|^2 |\xi|^{1-a} \, d\xi. \end{aligned} \tag{5.7}$$

The lemma is proved, for a constant

$$e_s := \bar{J}[\varphi_0]. \tag{5.8}$$

□

The following result allows to compare the energies in \mathbb{R}_+^2 and in $\mathbb{R} \times (0, M)$ for big M .

Lemma 16 *Given $v \in H^s(\mathbb{R})$, consider the functionals*

$$J[u] = \int_{\mathbb{R}_+^2} |\nabla u|^2 w_a \, dx dy \quad \text{and} \quad J_M[u] = \int_{\mathbb{R} \times (0, M)} |\nabla u|^2 w_a \, dx dy$$

subject to the constraint $Tu = v$ in \mathbb{R} . Let u_0 be the minimizer of J and u_M be the minimizer of J_M , in their corresponding weighted function spaces. Given $\epsilon > 0$, there exists $M > 0$ depending only on ϵ and not on v such that

$$\int_{\mathbb{R} \times (0, M)} |\nabla u_M|^2 w_a \, dx dy \geq (1 - \epsilon) \int_{\mathbb{R}_+^2} |\nabla u_0|^2 w_a \, dx dy.$$

Proof We use the ideas of Lemma 15 in order to reduce the two dimensional problem to an ordinary differential equation through Fourier transform. The Euler–Lagrange equation of J_M is

$$\begin{cases} \operatorname{div}(y^a \nabla u) = 0 \text{ in } \mathbb{R}_+^2, \\ \partial_\nu u = 0 \text{ on } \mathbb{R} \times \{M\}, \\ u = v \text{ on } \mathbb{R} \times \{0\}, \end{cases} \tag{5.9}$$

while the one for J is given by (5.6). Taking Fourier transform in x , we immediately realize that both minimizers u_0, u_M must satisfy the ODE (for each value of ξ)

$$-|\xi|^2 \hat{u}(\xi, y) + \frac{a}{y} \hat{u}_y(\xi, y) + \hat{u}_{yy}(\xi, y) = 0.$$

Now consider the two one-dimensional functionals for each $\varphi : [0, +\infty) \rightarrow \mathbb{R}$,

$$\bar{J}[\varphi] = \int_{\mathbb{R}_+} (|\varphi'|^2 + \varphi^2) y^a \, dy \quad \text{and} \quad \bar{J}_M[\varphi] = \int_{(0, M)} (|\varphi'|^2 + \varphi^2) y^a \, dy$$

subject to the condition $\varphi(0) = 1$. Let φ_0 and φ_M be the minimizers of \bar{J} and \bar{J}_M , respectively. Then $\hat{u}_0(\xi, y) = \hat{v}(\xi)\varphi_0(|\xi|y)$ and $\hat{u}_M(\xi, y) = \hat{v}(\xi)\varphi_M(|\xi|y)$ because both φ_0 and φ_M satisfy the Euler equation

$$-\varphi(y) + \frac{a}{y} \varphi_y(y) + \varphi_{yy}(y) = 0. \tag{5.10}$$

with boundary conditions

$$\begin{aligned} \varphi_0(0) &= 1, & \varphi_0(y) &\rightarrow 0 \quad \text{when } y \rightarrow +\infty, \\ \varphi_M(0) &= 1, & \varphi'(M) &= 0. \end{aligned} \tag{5.11}$$

Thus Lemma (14) gives that

$$\varphi_0(y) = y^s [c_1 I_s(y) + c_2 K_s(y)], \quad \varphi_M(y) = y^s [c_1^M I_s(y) + c_2^M K_s(y)].$$

If we impose the boundary conditions (5.11) at $y = 0$, we obtain the value of $c_2 = c_2^M = \operatorname{cst}(s)$. Now, note that $\partial_y \varphi_M(M) = 0$ while φ decays at infinity, this fixes c_1, c_1^M . In particular, $c_1^M \rightarrow c_1 = 0$ exponentially. We have then that φ_M converges to φ_0 and φ'_M to φ'_0 when $M \rightarrow \infty$, and thus, also

$$\bar{J}_M[\varphi_M] \rightarrow \bar{J}[\varphi_0]. \tag{5.12}$$

Then, the computation in (5.7) gives that

$$\int_{\mathbb{R}_+^2} |\nabla u_0|^2 w_a = \bar{J}[\varphi_0] \int_{\mathbb{R}} |\hat{v}(\xi)|^2 |\xi|^{1-a} d\xi \tag{5.13}$$

while

$$\int_{\mathbb{R} \times (0, M)} |\nabla u_M|^2 w_a = \bar{J}_M[\varphi_M] \int_{\mathbb{R}} |\hat{v}(\xi)|^2 |\xi|^{1-a} d\xi. \tag{5.14}$$

The lemma is proved by comparing (5.14) and (5.13), because of (5.12). □

5.2 Inequality for a half-plane

Denote by \mathbb{R}_+^2 the half-plane $\mathbb{R} \times (0, +\infty)$, and consider the weight in \mathbb{R}_+^2 given by $w_a(x, y) := y^a$ for $x \in \mathbb{R}, y > 0$. When $a = 0$, Fourier transform methods quickly give that a function u defined on \mathbb{R}_+^2 has a well defined trace on $\mathbb{R} \times \{0\}$ and moreover (see [6, lemma 6.2])

$$\int_{\mathbb{R}^2} \left| \frac{v(x) - v(x')}{x - x'} \right|^2 dx dx' \leq 2\pi \int_{\mathbb{R}_+^2} |\nabla u|^2 dx dy. \tag{5.15}$$

This constant 2π is sharp.

Trace inequalities for the general case $-1 < a < 0$ require the characterization of the fractional Laplacian by Caffarelli and Silvestre [9] as a Dirichlet-to-Neumann operator. The first result of this subsection deals with the generalization of (5.15).

Lemma 17 *Let u be a function in $L^2_{loc}(\mathbb{R}_+^2, w_a)$ with derivative in $L^2(\mathbb{R}_+^2, w_a)$. Then the trace of u on $\mathbb{R} \times \{0\}$, call it v , is a well defined function $v \in L^2_{loc}(\mathbb{R})$. Moreover,*

$$\int_{\mathbb{R}^2} \frac{|v(x) - v(x')|^2}{|x - x'|^{1+2s}} dx dx' \leq D_s \int_{\mathbb{R}_+^2} |\nabla u|^2 w_a dx dy, \tag{5.16}$$

where D_s is a constant depending only on s but not on u , and it is given precisely in (5.20).

Proof First we assume that u is smooth. The result will then follow by approximation. Next, if we apply Plancherel theorem for the Fourier transform we obtain

$$\begin{aligned} \int_{\mathbb{R}^2} \frac{|v(x) - v(x')|^2}{|x - x'|^{1+2s}} dx dx' &= \int_{\mathbb{R}} \left[\int_{\mathbb{R}} |v(x+h) - v(x)|^2 dx \right] \frac{1}{h^{1+2s}} dh \\ &= \int_{\mathbb{R}} \left[\int_{\mathbb{R}} |\hat{v}(\xi)(e^{2\pi i h \xi} - 1)|^2 d\xi \right] \frac{1}{h^{1+2s}} dh \\ &= \int_{\mathbb{R}} \left[\int_{\mathbb{R}} \frac{2 - 2 \cos(2\pi h \xi)}{h^{1+2s}} dh \right] |\hat{v}(\xi)|^2 d\xi \\ &= d_s (2\pi)^{2s} \int_{\mathbb{R}} |\hat{v}(\xi)|^2 |\xi|^{2s} d\xi, \end{aligned} \tag{5.17}$$

for

$$d_s = \int_{\mathbb{R}} \frac{2 - 2 \cos z}{z^{1+2s}} dz. \tag{5.18}$$

On the other hand, let u_0 be the minimizer of the functional

$$J(u) = \int_{\mathbb{R}_+^2} |\nabla u|^2 w_a$$

found in Lemma 15. Then

$$\int_{\mathbb{R}_+^2} |\nabla u_0|^2 w_a dx dy = e_s \int_{\mathbb{R}} |\hat{v}(\xi)|^2 |\xi|^{2s} d\xi. \tag{5.19}$$

We set

$$D_s := d_s(2\pi)^{2s} / e_s, \tag{5.20}$$

where e_s is given in (5.8) and d_s in (5.18).

The proof of (5.16) is completed from (5.17), (5.19), and using the fact that u_0 is a minimizer of J . □

Corollary 18 *Inequality (5.16) is achieved when $u = P *_x v$.*

Proof Indeed, the minimizer u_0 is given precisely by $u_0 = P *_x v$. □

5.3 Inequality of a bounded domain

Now we would like to localize inequality (5.16) in a bounded domain $\Omega \subset \mathbb{R}_+^2$ such that part of its boundary, say $M \subset \partial\Omega$ lies on $\mathbb{R} \times \{0\}$. Again, we fix the weight $w_a := y^a$. Nekvinda’s work, summarized in Theorem 3, assures that the trace of a function $u \in W^{1,2}(\Omega, w_a)$ belongs to the Besov space $H^s(M)$, but it does not give the explicit value of the constant in the embedding.

Here we claim that this constant can be taken as the constant in \mathbb{R}_+^2 , the D_s found in the previous subsection:

Theorem 19 *Let $\Omega := (-1, 1) \times (0, 1) \subset \mathbb{R}_+^2$ and $u \in W^{1,2}(\Omega, w_a)$. Then the trace of u on $(-1, 1) \times \{0\}$, call it v , is a well defined function $v \in H^s(-1, 1)$ and, for the same constant D_s as in the Lemma 17, we have*

$$\int_{(-1,1)^2} \frac{|v(x) - v(x')|^2}{|x - x'|^{1+2s}} dx dx' \leq D_s \int_{\Omega} |\nabla u|^2 w_a dx dy. \tag{5.21}$$

Proof First note that Theorem 3 quickly gives that $v \in H^s(-1, 1)$. Moreover, it is enough to prove inequality (5.21) for $u \in C^\infty(\bar{\Omega})$.

Fixed $\epsilon > 0$, consider the constant $M > 0$ given in Lemma 16, that is independent of u, v . We will show first that

$$\int_{\Omega_M} |\nabla u|^2 w_a dx dy \geq D_s^{-1}(1 - \epsilon) \int_{(-M,M)^2} \frac{|v(x) - v(x')|^2}{|x - x'|^{1+2s}} dx dx', \tag{5.22}$$

i.e., that the inequality holds in a domain $\Omega_M := (-M, M) \times (0, M)$ up to epsilon. Because the inequality (5.22) is invariant under the rescaling $(\tilde{x}, \tilde{y}) := \frac{1}{M}(x, y)$, it also holds for $(-1, 1) \times (0, 1)$, up to ϵ . This would finish the proof of (5.21).

We compare the energy of u to the energy of u_1 , the minimizer of the functional

$$J_{\Omega_M}[u] = \int_{\Omega_M} |\nabla u|^2 w_a \, dx dy,$$

subject to the restriction $Tu_1 = v$. We now try to extend u_1 defined on Ω_M to the whole \mathbb{R}_+^2 , in order to use Lemma 17. We do it in several steps.

First, because u_1 satisfies a zero Neumann condition on the boundary $\{-M, M\} \times (0, M)$, it is possible to reflect it evenly N times to obtain a solution on $\Omega_{NM} = A_{NM} \times (0, M)$ where A_{NM} is an interval of length NM . We still denote that extension by u_1 . Call v_1 to be the trace of u_1 on $y = 0$. Now we would like to extend u_1 to the whole $\mathbb{R} \times (0, M)$. For this, first extend v_1 continuously to \mathbb{R} such that the extension, call it v_2 , belongs to $H^s(\mathbb{R})$, and solve the Caffarelli–Silvestre extension problem

$$\begin{cases} \operatorname{div}(y^a \nabla u_2) = 0 \text{ in } \mathbb{R}_+^2 \\ u_2 = v_2 \text{ on } \mathbb{R} \times \{0\}. \end{cases}$$

Let $B := [b_1 - 1, b_1] \cup [b_2, b_2 + 1]$ where $b_1 = \inf(A_{NM})$, $b_2 = \sup(A_{NM})$, and let $\eta : \mathbb{R} \rightarrow \mathbb{R}^+$ be a smooth cutoff such that $\eta = 1$ in A_{NM} and $\eta = 0$ outside $A_{NM} \cup B$. Consider the function obtained by the gluing

$$u_3 := \eta u_1 + (1 - \eta)u_2,$$

it is defined on $\mathbb{R} \times (0, M)$.

Next, let $u_M : \mathbb{R} \times (0, M) \rightarrow \mathbb{R}$ be the minimizer of the functional

$$J_M[u] = \int_{\mathbb{R} \times (0, M)} |\nabla u|^2 w_a \, dx dy$$

and $u_0 : \mathbb{R}_+^2 \rightarrow \mathbb{R}$ be the minimizer of

$$J[u] = \int_{\mathbb{R}_+^2} |\nabla u|^2 w_a \, dx dy,$$

both subject to the condition $Tu = v_2$ on $y = 0$. In particular, we can use the Poisson kernel (2.3) to write $u_0 = P *_x v_2$. Lemma 16 implies that

$$\int_{\mathbb{R} \times (0, M)} |\nabla u_3|^2 w_a \geq \int_{\mathbb{R} \times (0, M)} |\nabla u_M|^2 w_a \geq (1 - \epsilon) \int_{\mathbb{R}_+^2} |\nabla u_0|^2 w_a. \tag{5.23}$$

Applying Corollary 18 to $u_0 = P *_x v_2$, we obtain that

$$\begin{aligned} \int_{\mathbb{R}_+^2} |\nabla u_0|^2 w_a &= D_s^{-1} \int_{\mathbb{R} \times \mathbb{R}} \frac{|v_2(x) - v_2(x')|^2}{|x - x'|^{1+2s}} \, dx dx' \\ &\geq N D_s^{-1} \int_{(-M, M)^2} \frac{|v(x) - v(x')|^2}{|x - x'|^{1+2s}} \, dx dx', \end{aligned} \tag{5.24}$$

where the last inequality holds just by dropping part of the domain of integration. On the other hand, because of our gluing construction,

$$\begin{aligned} \int_{\mathbb{R} \times (0, M)} |\nabla u_3|^2 w_a &= N \int_{\Omega_M} |\nabla u_1|^2 w_a + \int_{B \times (0, M)} |\nabla u_3|^2 w_a \\ &+ \int_{[\mathbb{R} \setminus (A_{NM} \cup B)] \times (0, M)} |\nabla u_2|^2 w_a \\ &=: N \int_{\Omega_M} |\nabla u_1|^2 w_a + I_3 + I_2. \end{aligned} \tag{5.25}$$

We would like to show that the last two terms I_2, I_3 above are bounded independently of N . First of all, we can use the Poisson kernel (2.3) to write explicit formulas for $u_2 = P *_x v_2$, i.e.,

$$u_2(x, y) = c_{2,s} \int_{\mathbb{R}} \frac{y^{1-a} v_2(\xi)}{(|x - \xi|^2 + |y|^2)^{\frac{2-a}{2}}} d\xi = c_{2,s} \int_{\mathbb{R}} \frac{v_2(x - zy)}{(z^2 + 1)^{\frac{2-a}{2}}} dz$$

after the change $z = \frac{x-\xi}{y}$. Because the original v was differentiable, we have that v_2 has bounded derivative (it is smooth except perhaps at the reflection points). Then

$$\partial_x u_2(x, y) = C \int_{\mathbb{R}} \frac{v'_2(x - zy)}{(z^2 + 1)^{\frac{2-a}{2}}} dz$$

and

$$\partial_y u_2(x, y) = C \int_{\mathbb{R}} \frac{(-z)v'_2(x - zy)}{(z^2 + 1)^{\frac{2-a}{2}}} dz.$$

In the following, C will be a positive constant that may change from line to line. We consider the following auxiliary term I and can compute, using Hölder, that

$$\begin{aligned} I &:= \int_{x \in (b_2, +\infty), y \in (0, M)} |\nabla u_2|^2 y^a dx dy \\ &\leq C \int_{x \in (b_2, +\infty), y \in (0, M)} \int_{z \in \mathbb{R}} y^a \frac{[v'_2(x - zy)]^2 \max\{1, |z|^2\}}{(z^2 + 1)^{2-a}} dz dx dy. \end{aligned} \tag{5.26}$$

We could have extended v_2 to constant (or very decaying to constant) on the interval $[b_2, +\infty)$. Thus it is enough to consider the change $\theta = x - zy - b_2$ so that

$$\begin{aligned} I &\leq C \int_{z \in \mathbb{R}} \frac{\max\{1, |z|^2\}}{(z^2 + 1)^{2-a}} \int_{y \in (0, M)} y^a \int_{\theta \in (-zy, 0)} v'_2(\theta + b_2)^2 d\theta dy dz \\ &\leq C \int_{z \in \mathbb{R}} \frac{|z| \max\{1, |z|^2\}}{(z^2 + 1)^{2-a}} dx \int_{y \in (0, M)} y^{a+1} dy. \end{aligned} \tag{5.27}$$

This last integral is bounded independently of N because $-1 < a < 0$. Similar arguments give that the terms I_2, I_3 from (5.25) are $o(N)$.

Then, combining (5.23), (5.24) and (5.25) we obtain that

$$N \int_{\Omega_M} |\nabla u_1|^2 w_a + o(N) \geq (1 - \epsilon) D_s^{-1} N \int_{(-M, M)^2} \frac{|v(x) - v(x')|^2}{|x - x'|^{1+2s}} dx dx'.$$

Divide the previous inequality by N and let $N \rightarrow \infty$, so that

$$\int_{\Omega_M} |\nabla u_1|^2 w_a \geq (1 - \epsilon) D_s^{-1} \int_{(-M, M)^2} \frac{|v(x) - v(x')|^2}{|x - x'|^{1+2s}} dx dx'. \tag{5.28}$$

To finish just note that

$$\int_{\Omega_M} |\nabla u|^2 w_a dx dy \geq \int_{\Omega_M} |\nabla u_1|^2 w_a dx dy \tag{5.29}$$

because u_1 was a minimizer of J_{Ω_M} . The proposition follows from (5.28) and (5.29), after rescaling back. □

5.4 Optimality

In the remaining of the section we would like to show that the constant D_s in (5.21) is optimal in some sense.

Proposition 20 *For each $\epsilon > 0$, there exists a function u_ϵ defined on $(-1, 1) \times (0, 1)$ such that*

$$\epsilon^{1-a} \int_{(-1, 1) \times (0, 1)} |\nabla u_\epsilon|^2 w_a = D_s^{-1} \epsilon^{1-a} \int_{(-1, 1)^2} \frac{|Tu_\epsilon(x) - Tu_\epsilon(x')|^2}{|x - x'|^{1+2s}} dx dx' + R_\epsilon \tag{5.30}$$

with $\lim_{\epsilon \rightarrow 0} |R_\epsilon| = 0$, and this term is of lower order when $\epsilon \rightarrow 0$.

Proof Fix a domain $\Omega = (-1, 1) \times (0, 1)$, and for each $\epsilon > 0$ consider the scaling $\Lambda_\epsilon > 0$ given by (1.9). Set $v_\epsilon : (-1, 1) \rightarrow \mathbb{R}$ to be the function that satisfies $v_\epsilon(x) = 0$ if $x \in (-1, -\Lambda_\epsilon/2)$, $v_\epsilon(x) = 1$ if $x \in [\Lambda_\epsilon/2, 1)$, and linear in the interval $[-\Lambda_\epsilon/2, \Lambda_\epsilon/2]$. Extend it to \mathbb{R} , by making it constant on $(-\infty, -1)$ and $(1, +\infty)$, and denote this extension by $\tilde{v} := \tilde{v}_\epsilon$. Now construct a function $u := u_\epsilon$ defined on \mathbb{R}_+^2 with trace \tilde{v} as $u = P *_x \tilde{v}$. Corollary 18 tells that for u constructed this way,

$$\int_{\mathbb{R}_+^2} |\nabla u|^2 w_a dx dy = D_s^{-1} \int_{\mathbb{R} \times \mathbb{R}} \frac{|\tilde{v}(x) - \tilde{v}(x')|^2}{|x - x'|^{1+2s}} dx dx'. \tag{5.31}$$

We try now to restrict the domain of integration to Ω by estimating the remaining terms. Let $\Omega_\epsilon := ((-1, -\Lambda_\epsilon/2) \cup [\Lambda_\epsilon/2, 1)) \times (\Lambda_\epsilon/2, 1)$. A straightforward computation from the Poisson formula

$$u(x, y) = C \int_{\mathbb{R}} \frac{y^{1-a} \tilde{v}(\xi)}{(|x - \xi|^2 + |y|^2)^{(2-a)/2}} d\xi$$

gives that

$$\int_{\Omega_\epsilon} |\nabla u|^2 w_a \, dx dy = \frac{C}{\Lambda_\epsilon^{2s-1}} + o\left(\frac{1}{\Lambda_\epsilon^{2s-1}}\right),$$

while

$$\int_{\mathbb{R}_+^2} |\nabla u|^2 w_a \, dx dy - \int_{\Omega_\epsilon} |\nabla u|^2 w_a \, dx dy = o\left(\frac{1}{\Lambda_\epsilon^{2s-1}}\right).$$

On the other hand, let $A_\epsilon = (-1, \Lambda_\epsilon/2]$, $B_\epsilon = [\Lambda_\epsilon/2, 1)$. We also have

$$\int_{A_\epsilon \times B_\epsilon} \frac{|\tilde{v}(x) - \tilde{v}(x')|^2}{|x - x'|^{1+2s}} \, dx dx' = \frac{C'}{\Lambda_\epsilon^{2s-1}} + o\left(\frac{1}{\Lambda_\epsilon^{2s-1}}\right)$$

while

$$\int_{\mathbb{R}^2 \setminus ((A_\epsilon \times B_\epsilon) \cup (B_\epsilon \times A_\epsilon))} \frac{|\tilde{v}(x) - \tilde{v}(x')|^2}{|x - x'|^{1+2s}} \, dx dx' = o\left(\frac{1}{\Lambda_\epsilon^{2s-1}}\right)$$

Because of equality (5.31), the previous estimates, and the fact that

$$\epsilon^{1-a} \frac{1}{\Lambda_\epsilon^{2s-1}} = 1$$

we conclude that the u_ϵ, v_ϵ we have constructed satisfy (5.30). □

Remark 1 We have constructed a piecewise linear function. However, the above proposition is still true as long as the transition occurs on a layer of length Λ_ϵ .

6 Preliminary results

We are going to prove the main Theorem 1 by localizing the functional into several regions. Section 3 takes care of the behavior of the functional in the interior of the domain Ω . Thus, it remains to study the interaction the interaction with the wall of the container $\partial\Omega$. In the present section we give some preliminary results in that regard.

First, we look at a small neighborhood of $x \in \partial\Omega$, and reduce the problem to the study of a small neighborhood $B_r \times (0, r) \subset \mathbb{R}_+^3$. Then we reduce the dimension from three to two through a slicing argument, so that it is enough to consider a subdomain $\Omega = (-r, r) \times (0, r) \subset \mathbb{R}_+^2$, and thus, the results of Sects. 4 and 5 can be applied.

Let us remind the reader that the Gamma-limit is going to be expressed in terms of the functional (1.7), whose exact expression is

$$\Phi(u, v) := \sigma \mathcal{H}^2(S_u) + \int_{\partial\Omega} |\mathcal{W}(Tu) - \mathcal{W}(v)| + \kappa_s \mathcal{H}^1(S_v).$$

The first term in Φ comes from the behavior in the interior (Sect. 3). In Proposition 26 we will consider the “wall effect”, that explains the presence of the second term; while the remaining of the section is devoted to the “boundary effect”, that deals with the third term.

We define the localization of the functional F_ϵ^a as follows. For every open set $A \subset \Omega$ and any $A' \subset \partial A$, we set

$$F_\epsilon^a[u, A, A'] := \epsilon^{1-a} \int_A |\nabla u|^2 w_a + \frac{1}{\epsilon^{1-a}} \int_A W(u) w_a^{-1} + \lambda_\epsilon \int_{A'} V(Tu). \tag{6.1}$$

Note that we easily recover the original functional as $F_\epsilon^a[u] = F_\epsilon^a[u, \Omega, \partial\Omega]$.

6.1 Reduction to the flat case

Fixed $x \in \partial\Omega$, we will consider small ‘‘cubical’’ neighborhoods $Q_r(x)$ near the boundary, of size r , such that $\Omega \cap Q_r(x)$ is equivalent to the cylinder $D_r := B_r \times (0, r)$. Here B_r is the two-dimensional ball of radius r centered an the origin. Let E_r be the boundary part given by $E_r := B_r \times \{0\}$. In order to evaluate the error in the deformation, we need to introduce the notion of isometry defect.

Given two domains $A_1, A_2 \subset \mathbb{R}^3$ and a bi-Lipschitz homeomorphism $\Psi : \bar{A}_1 \rightarrow \bar{A}_2$, the isometry defect $\delta(\Psi)$ of Ψ is the smallest constant δ such that

$$\text{dist}(D\Psi(x), O(3)) \leq \delta \quad \text{for a.e. } x \in A_1. \tag{6.2}$$

Here $O(3)$ is the set of linear isometries on \mathbb{R}^3 , and $D\Psi(x)$ is regarded as a linear mapping of \mathbb{R}^3 into \mathbb{R}^3 . Let I be the identity map on \mathbb{R}^3 . The distance between linear mappings is induced by the norm $\|\cdot\|$, which for every linear map T is defined as

$$\|T\| = \sup_{|v| \leq 1} |Tv|.$$

The following proposition shows that the localized energy $F_\epsilon^a[u, \Omega \cap Q_r(x), \partial\Omega \cap Q_r(x)]$ can be replaced by the energy $F_\epsilon^a[u, D_r, E_r]$.

Proposition 21 *Let Ω be a domain in \mathbb{R}^3 with C^2 boundary $\partial\Omega$ in C^2 . Then for every $x \in \partial\Omega$ and every positive r smaller than a certain critical value $r_x > 0$, there exists a diffeomorphism $\Psi_r : \bar{D}_r \rightarrow \overline{\Omega \cap Q_r(x)}$ such that*

1. Ψ_r takes D_r onto $\Omega \cap Q_r(x)$ and E_r onto $\partial\Omega \cap Q_r(x)$.
2. Ψ_r is of class C^1 on D_r and $\|D\Psi_r - Id\| \leq \delta_r$ everywhere on D_r , where $\delta_r \rightarrow 0$ as $r \rightarrow 0$.

In particular, the isometry defect of Ψ_r vanishes as $r \rightarrow 0$. Moreover,

$$F_\epsilon^a[u, Q_r(x) \cap \Omega, Q_r(x) \cap \partial\Omega] \geq (1 - \delta(\Psi_r))^5 F_\epsilon^a[u \circ \Psi_r, D_r, E_r].$$

Proof It is essentially contained in propositions 4.9 and 4.10 from [6], although in our case we need to make some modifications due to the presence of a weight.

Because of the smoothness assumptions on $\partial\Omega$, we can parameterize a small neighborhood of $x \in \partial\Omega$ with coordinates (t, ρ) , where $\rho = \text{dist}(\cdot, \partial\Omega)$, t is the coordinate parameterizing each level set of ρ and such that $\rho \in (0, r)$, and $t \in B_r$. The change of coordinates map, call it Ψ_r is a diffeomorphism with $D\Psi_r(x) = I$. It is clear than

$$\delta := \delta(\Psi_r) \rightarrow 0, \quad \text{when } r \rightarrow 0. \tag{6.3}$$

Moreover,

$$|D(u \circ \Psi_r)| \leq (1 + \delta) |(Du) \circ \Psi_r|. \tag{6.4}$$

Let J be the Jacobian determinant of Ψ_r on D_r and J' be the one of $\Psi_r|_{E_r}$. They satisfy $|J|, |J'| \leq (1 + \delta)^3$ a.e.. Using the change of variable formula and (6.4) we see that

$$\begin{aligned} F_\epsilon^a[u \circ \Psi_r, D_r, E_r] &= \epsilon^{1-a} \int_{D_r} |\nabla(u \circ \Psi_r)|^2 \rho^a J d\rho dt \\ &\quad + \frac{1}{\epsilon^{1-a}} \int_{D_r} W(u \circ \Psi_r) \rho^{-a} J d\rho dt + \lambda_\epsilon \int_{E_r} V(T(u \circ \Psi_r)) J' dt \\ &\leq (1 + \delta(\Psi_r))^5 F_\epsilon^a[u, Q_r(x) \cap \Omega, Q_r(x) \cap \partial\Omega], \end{aligned}$$

as we wished, because $\frac{1}{1+\delta} \geq 1 - \delta$. □

Remark 2 The regularity of $\partial\Omega$ could be reduced to $C^{1,\alpha}$ and still have (6.3).

6.2 Dimension reduction

The next step is to reduce the problem from three to two dimensions through slicing. Before we give the main result, we state some classical slicing results in \mathbb{R}^n :

We fix $m > 0$ and assume that every function in this subsection takes values in $[-m, m]$. Let A be a bounded open subset of \mathbb{R}^n , e is a unit vector in \mathbb{R}^n and u a function on A . We denote by M the orthogonal complement of e , and by A_e the projection of A onto M . For every $z \in M$, we set $A_e^z := \{t \in \mathbb{R} : z + te \in A\}$; and u_e^z to be the trace of u on A_e^z , that is $u_e^z := u(z + te)$.

Proposition 22 (section 5.10, p. 216, in [10]) *Let $B \subset A$ be a given Borel set. If B has finite perimeter in A , then B_e^z has finite perimeter in A_e^z and $\partial(B_e^z \cap A_e^z) = (\partial B \cap A)_e^z$ for a.e. $z \in A_e$, and*

$$\int_{A_e} \mathcal{H}^0(\partial B_e^z \cap A_e^z) dz = \int_{\partial B \cap A} \langle \nu_B, e \rangle. \tag{6.5}$$

Conversely, B has finite perimeter in A if there exist n linearly independent unit vectors e such that the integral of $\mathcal{H}^0(\partial B_e^z \cap A_e^z)$ over all $z \in A_e$ is finite.

From here we can establish a connection between the compactness of a family of functions in $L^1(\mathbb{R}^3)$ and the compactness of the traces of these functions. For every family \mathcal{F} of functions of A , we set $\mathcal{F}_e^z := \{u_e^z : u \in \mathcal{F}\}$, so that \mathcal{F}_e^z is a family of functions on A_e^z . We say that a family \mathcal{F}' is δ -dense in \mathcal{F} if \mathcal{F} lies in a δ -neighborhood of \mathcal{F}' with respect to the $L^1(A)$ topology. Then

Theorem 23 (theorem 6.6 in [6]) *Let \mathcal{F} be a family of functions $v : A \rightarrow [-m, m]$ and assume that there exist n linearly independent unit vectors e which satisfy the following property:*

For every $\delta > 0$ there exists a family \mathcal{F}_δ δ -dense in \mathcal{F} such that $(\mathcal{F}_\delta)_e^z$ is pre-compact in $L^1(A_e^z)$ for \mathcal{H}^{n-1} a.e. $z \in A_e$.

Then \mathcal{F} is precompact in $L^1(A)$.

Now we can give the main proposition of the subsection, where we slice a cylinder $D_r = B_r \times (0, r) \subset \mathbb{R}_+^3$. We drop the subindex r in the notation. We set coordinates in D

as (x_1, x_2, y) where x_1, x_2 parameterize B_r and $y \in (0, r)$. Then $E = \{y = 0\} \cap \bar{D}$. Fix an arbitrary unit vector e in the plane $\{y = 0\}$, and let E_e the projection of the set E . We will slice D in the direction of y , perpendicularly to e . The slice corresponding for each $z \in E_e$ is denoted by D^z ; let E^z its projection onto the plane $\{y = 0\}$. For a nice picture, see figure 4 in [6].

Proposition 24 *Let $(u_\epsilon) \subset W^{1,2}(D, w_a)$ be a sequence with uniformly bounded energies $F_\epsilon^a[u_\epsilon, D, E]$. Then the traces of u_ϵ are pre-compact in $L^1(E)$ and every cluster point belongs to $BV(E, I')$. Moreover, if $Tu_\epsilon \rightarrow v$ in $L^1(E)$, then*

$$\liminf_{\epsilon \rightarrow 0} F_\epsilon^a[u_\epsilon, D, E] \geq \kappa_s \left| \int_{E \cap S_v} v_v \right| d\mathcal{H}^1, \tag{6.6}$$

where the constant κ_s is defined in (4.2).

Proof We slice the region D using Fubini’s theorem, so then

$$\begin{aligned} F_\epsilon^a[u_\epsilon, D, E] &\geq \epsilon^{1-a} \int_D |\nabla u_\epsilon|^2 w_a + \lambda_\epsilon \int_E V(Tu) \\ &\geq \int_{E_e} \left(\epsilon^{1-a} \int_{D^z} |\nabla u_\epsilon^z|^2 y^a + \lambda_\epsilon \int_{E^z} V(Tu_\epsilon^z) \right) dz. \end{aligned} \tag{6.7}$$

Next, we use the trace inequalities of Sect. 5 in each domain D^z, E^z (note that the inequalities are invariant by rescaling). Indeed, because of (5.21), we have the estimate

$$\begin{aligned} F_\epsilon^a[u_\epsilon, D, E] &\geq \int_{E_e} \left(\epsilon^{1-a} D_s^{-1} \int_{E^z \times E^z} \frac{|Tu_\epsilon^z(x) - Tu_\epsilon^z(x')|^2}{|x - x'|^{1+2s}} + \lambda_\epsilon \int_{E^z} V(Tu_\epsilon^z) \right) dz \\ &= \int_{E_e} G_\epsilon^a[Tu_\epsilon^z, E^z] dz. \end{aligned} \tag{6.8}$$

This last functional G_ϵ^a has been well studied in Sect. 4 when E^z is an interval in \mathbb{R} .

The rest of the proof follows exactly as proposition 4.7 in [6]. For commodity of the reader, we give the main ideas. We check first that (Tu_ϵ) is pre-compact in $L^1(E)$. Thanks to Theorem 23, it suffices to show that the family $\mathcal{F} := (Tu_\epsilon)$ satisfy the following property: for every $\delta > 0$, there exists a family \mathcal{F}_δ δ -dense in \mathcal{F} such that $(\mathcal{F}_\delta)_\epsilon^z$ is pre-compact in $L^1(E)$ for \mathcal{H}^2 -a.e. $z \in E_e$. By assumption $F_\epsilon^a[u_\epsilon, D, E] \leq C$, so that (6.8) implies also that

$$\int_{E_e} G_\epsilon^a[u_\epsilon^z, E^z] \leq C. \tag{6.9}$$

Fix $\delta > 0$. For every $\delta > 0$, define $v_\epsilon : E \rightarrow [-m, m]$ such that

$$v_\epsilon^z := \begin{cases} Tu_\epsilon^z, & \text{if } z \in E_e \text{ and } G_\epsilon^a[u_\epsilon^z, E^z] \leq 2mrC/\delta \\ \alpha', & \text{otherwise.} \end{cases} \tag{6.10}$$

By (6.8), (6.9), (6.10), we have $v_\epsilon^z = Tu_\epsilon^z$ for all $z \in E_e$ apart from a subset of measure smaller than $\delta/(2mr)$. Hence $v_\epsilon = Tu_\epsilon$ in E up to a set of measure smaller than δ/m . So,

from $|Tu_\epsilon| \leq m$, we deduce that $\|v_\epsilon - Tu_\epsilon\|_{L^1(E)} \leq \delta$. Therefore, the family $\mathcal{F}_\delta := \{v_\epsilon\}$ is δ -dense in \mathcal{F} .

Next, by (6.10), we have that $G_\epsilon^a[v_\epsilon^z, E^z] \leq 2mrC/\delta$ for every $z \in E_\epsilon$ and every ϵ . Hence, Theorem 5 implies that the sequence (v_ϵ^z) is pre-compact in $L^1(E^z)$. Thus \mathcal{F} satisfies the hypothesis of Theorem 23 for every e , and the sequence (Tu_ϵ) is pre-compact in $L^1(E)$.

It remains to prove that if $Tu_\epsilon \rightarrow E$ in $L^1(E)$, then v belongs to $BV(E, I')$, and inequality (6.6) holds. We have (up to a subsequence), that $Tu_\epsilon^z \rightarrow v^z$ in $L^1(E)$ for a.e. $z \in E_\epsilon$ (remark 6.7 in [6]). Then, Proposition 5 yields $v^z \in BV(E^z, I')$ and

$$\liminf_{\epsilon \rightarrow 0} F_\epsilon^a[u_\epsilon, D, E] \geq \int_{E_e} \kappa_s \mathcal{H}^0(Sv^z) dz. \tag{6.11}$$

The right-hand side of the formula above is finite, so then Proposition 22 implies that v belongs to $BV(E, I')$, and that S_{v^z} agrees with $S_v \cap E^z$ for a.e. $z \in E_\epsilon$. Then, by (6.5) we may rewrite (6.11) as

$$\liminf_{\epsilon \rightarrow 0} F_\epsilon^a[u_\epsilon, D, E] \geq \kappa_s \int_{E \cap S_v} \langle v_v, e \rangle dz.$$

Finally, (6.6) follows by choosing a suitable unit vector e in the expression above. □

6.3 The boundary effect

In the previous arguments we have reduced the dimension from three to two, so it is enough to understand the following functional on \mathbb{R}_+^2 . Set $D = (-1, 1) \times (0, 1)$, $E = (-1, 1)$, we define

$$H_\epsilon^a[w, D, E] := \epsilon^{1-a} \int_D |\nabla w|^2 y^a dx dy + \lambda_\epsilon \int_E V(Tw).$$

Trace inequalities allow to quickly relate this functional to the optimal profile obtained in Proposition 12. This link is precisely the missing ingredient in [22] for the superquadratic case.

Let $\phi : \mathbb{R} \rightarrow [\alpha', \beta']$ be such optimal profile. It achieves the infimum of

$$\kappa_s := \inf \left\{ G_1^a[v, \mathbb{R}] : v \in H^s(\mathbb{R}), \lim_{x \rightarrow -\infty} v(x) = \alpha', \lim_{x \rightarrow +\infty} v(x) = \beta' \right\}.$$

Now let $w_1 := P *_x \phi$. We rescale

$$w_\epsilon(x, y) := w_1 \left(\frac{x}{\Lambda_\epsilon}, \frac{y}{\Lambda_\epsilon} \right), \tag{6.12}$$

and

$$\phi_\epsilon(x) := \phi \left(\frac{x}{\Lambda_\epsilon} \right).$$

Note that $w_\epsilon = P *_x \phi_\epsilon$.

Lemma 25 *In the hypothesis above, we have that*

$$H_\epsilon^a[w_\epsilon, D, E] = \kappa_s + o(1), \text{ when } \epsilon \rightarrow 0.$$

Proof We note that, because of our rescaling,

$$H_\epsilon^a[w_\epsilon, D, E] = H_1^a[w_1, D/\Lambda_\epsilon, E/\Lambda_\epsilon].$$

But we can compute explicitly that

$$H_1^a[w_\epsilon, D/\Lambda_\epsilon, E/\Lambda_\epsilon] = H_1^a[w_1, \mathbb{R}_+^2, \mathbb{R}] - o(1).$$

On the other hand, because of the definition of w_1 , we have equality in Corollary 18, and thus

$$H_1^a[w_1, \mathbb{R}_+^2, \mathbb{R}] = \frac{1}{D_s} \int_{\mathbb{R}^2} \frac{|\phi(x) - \phi(x')|^2}{|x - x'|^{1+2s}} dx dx' + \int_{\mathbb{R}} V(\phi) dx = G_1^a[\phi, \mathbb{R}] = \kappa_s.$$

The lemma is proved. □

6.4 The wall effect

Here we deal with the second term in the limit functional (1.7):

Proposition 26 *Let $A \subset \Omega \subset \mathbb{R}^3$ be a domain with piecewise C^1 boundary, and $A' = \partial A \cap \partial\Omega$ with Lipschitz boundary. Let $u \in BV(A, I)$, $v \in BV(A', I')$ be given. Then*

- i. *For every sequence $(u_\epsilon) \subset W^{1,2}(A, w_a)$ such that $u_\epsilon \rightarrow u$ in $L^1(A)$ and $Tu_\epsilon \rightarrow v$ in $L^1(A')$,*

$$\liminf_{\epsilon \rightarrow 0} E_\epsilon^a[u_\epsilon, A] \geq \int_{A'} |\mathcal{W}(Tu) - \mathcal{W}(v)|.$$

- ii. *If v is constant on A' and u is constant in A with $u = \alpha$ or $u = \beta$, there exists a sequence (u_ϵ) such that $Tu_\epsilon = v$ in A' , (u_ϵ) converges to u , uniformly on every set with positive distance from A' and*

$$\limsup_{\epsilon \rightarrow 0} E_\epsilon^a[u_\epsilon, A] \leq \int_{A'} |\mathcal{W}(Tu) - \mathcal{W}(v)|.$$

Moreover, the function u_ϵ may be required to be $\frac{C}{\epsilon} \left(\frac{r}{\epsilon}\right)^{-a}$ -Lipschitz continuous in

$$A_r := \{x \in A : \text{dist}(x, \partial A) \leq r\}.$$

Proof The proof is a modification of propositions 1.2 and 1.4 in [18], and it is very well written for the case $p > 2$ in Palatucci’s Ph.D. thesis [23, proposition 4.3]. Here we indicate the steps required, and only give the proof for the ones that require any modification.

We may assume that $E_\epsilon^a[u_\epsilon, A] \leq C$. For every $\epsilon > 0$, let us denote

$$w_\epsilon(x) := (\mathcal{W} \circ u_\epsilon)(x), \quad \text{for } x \in A. \tag{6.13}$$

Step 1: We claim that $\int_A |Dw_\epsilon| \leq \text{constant}$. Indeed, by Young’s inequality we have

$$\int_A |Dw_\epsilon| = \int_A |\mathcal{W}'(u_\epsilon)| |Du_\epsilon| = 2 \int_A \sqrt{\mathcal{W}(u_\epsilon)} |Du_\epsilon| \leq E_\epsilon^a[u_\epsilon, A] \leq C.$$

Step 2: $w_\epsilon \rightarrow \mathcal{W} \circ u \in BV(A)$ in $L^1(A)$.

Step 3: The functional

$$G_0(z) := \int_A |Dz(x)| + \int_{\partial A} |Tz - \mathcal{W}(v)| d\mathcal{H}^2$$

is l.s.c. on $BV(A)$ with respect to the topology in $L^1(A)$.

Step 4: Proof of statement *i*. Applying lower semi-continuity of the functional G_0 to the sequence w_ϵ defined by (6.13) we obtain the following inequality

$$\begin{aligned} & \int_A |D(\mathcal{W} \circ u)(x)| + \int_{\partial A} |\mathcal{W}(Tu) - W(v)| d\mathcal{H}^2 \\ & \leq \liminf_{\epsilon \rightarrow 0} \left(\int_A |D(\mathcal{W} \circ u_\epsilon)(x)| + \int_{\partial A} |\mathcal{W}(Tu_\epsilon) - W(v)| d\mathcal{H}^2 \right) \\ & \leq \liminf_{\epsilon \rightarrow 0} \left(E_\epsilon^a[u_\epsilon, A] + \int_{\partial A} |\mathcal{W}(Tu_\epsilon) - W(v)| d\mathcal{H}^2 \right), \end{aligned} \tag{6.14}$$

by Young’s inequality. Next, since $Tu_\epsilon \rightarrow v$ in $L^1(A')$, we deduce that

$$\liminf_{\epsilon \rightarrow 0} \int_{\partial A} |\mathcal{W}(Tu_\epsilon) - \mathcal{W}(v)| d\mathcal{H}^2 = 0. \tag{6.15}$$

From (6.14) and (6.15) we obtain the lower bound inequality *i*.

Step 5: Proof of the upper bound *ii*. The weight w_a needs to be taken into account. Without loss of generality, consider the case $u = \beta$ and $v = \gamma$ with $\alpha < \gamma < \beta$; the other cases are similar. Let $\theta : [0, +\infty) \rightarrow [\gamma, \beta]$ be the solution of the ODE written as

$$\begin{cases} \theta' = \sqrt{W(\theta)} \\ \theta(0) = \gamma. \end{cases} \tag{6.16}$$

Let $d(x) = \text{dist}(x, A') = \text{dist}(x, \partial\Omega)$. We set $\phi(t) := \theta(\omega)$ for $\omega := \frac{t^{1-a}}{1-a}$, and $u_\epsilon(x) := \phi\left(\frac{d(x)}{\epsilon}\right)$. Then

$$E_\epsilon^a[u_\epsilon, A] = \epsilon^{1-a} \int_A |\nabla u_\epsilon|^2 h^a + \frac{1}{\epsilon^{1-a}} \int_A W(u_\epsilon) h^{-a},$$

that can be written by the coarea formula as

$$E_\epsilon^a[u_\epsilon, A] = \int_{\mathbb{R}^+} \int_{\Sigma_{\epsilon t}} [\phi'(t)^2 t^a + W(\phi(t)) t^{-a}] d\sigma dt,$$

and after the change $\omega = \frac{t^{1-a}}{1-a}$ we get

$$E_\epsilon^a[u_\epsilon, A] = \int_{\mathbb{R}^+} \int_{\Sigma_{\epsilon((1-a)\omega)^{\frac{1}{1-a}}}} [\theta'(\omega)^2 + W(\theta(\omega))] d\sigma d\omega,$$

where Σ_s is the set of points in A at a distance exactly s from A' . When $\epsilon \rightarrow 0$, we know that $\Sigma_{O(\epsilon)} \rightarrow A'$, and thus we have that

$$E_\epsilon^a[u_\epsilon, A] \rightarrow \int_{\mathbb{R}^+} \int_{A'} [\theta'(\omega)^2 + W(\theta(\omega))] d\sigma d\omega.$$

But because θ satisfies the ODE (6.16), then both terms in the above expression are equal, so the inequality $x_1^2 + x_2^2 \geq 2x_1x_2$ becomes an equality and we can conclude that

$$E_\epsilon^a[u_\epsilon, A] \rightarrow \int_{A'} \int_{\mathbb{R}^+} 2\sqrt{W(\theta)}\theta' d\omega d\sigma = \int_{A'} [\mathcal{W}(\beta) - \mathcal{W}(\gamma)] d\sigma,$$

where we take \mathcal{W} to be a primitive of $2\sqrt{W}$. To finish the proof of the proposition, just note that

$$\sup |\nabla u_\epsilon| \leq \frac{Cr^{-a}}{\epsilon^{1-a}}.$$

□

7 Proof of Theorem 1

Once we have the main ingredients from the previous sections, we can give the proof of the main theorem.

7.1 Compactness

Let (u_ϵ) be a sequence in $W^{1,2}(\Omega, w_a)$ such that $F_\epsilon^a[u_\epsilon]$ is bounded. Using the localization defined (6.1) and the functional E_ϵ^a from (3.1) we know that

$$F_\epsilon^a[u_\epsilon] \geq F_\epsilon^a[u_\epsilon, \Omega, \emptyset] = E_\epsilon^a[u_\epsilon, \Omega].$$

By statement iii. of Proposition 4 we conclude that (u_ϵ) is precompact in $L^1(\Omega)$ and there exists $u \in BV(\Omega, I)$ such that $u_\epsilon \rightarrow u$ in $L^1(\Omega)$.

It remains to prove that (Tu_ϵ) is pre-compact in $L^1(\partial\Omega)$ and every cluster point belongs to $BV(\partial\Omega, I')$. Thanks to Proposition 21, we can cover $\partial\Omega$ with finitely many ‘‘cubes’’ $(Q_j)_{j \in J}$, centered on $\partial\Omega$, of radius r_j , such that for every $j \in J$, there exists a bi-Lipschitz map Ψ_j with isometry defect $\delta(\Psi_j) < 1$, which satisfies $\Psi_j(D_{r_j} \cap Q_j) = \Omega \cap Q_j$ and $\Psi_j(E_{r_j} \cap Q_j) = \partial\Omega \cap B_j$.

We show that (Tu_ϵ) is pre-compact in $L^1(\partial\Omega \cap Q_j)$ for every $j \in J$. For every fixed j , let us set $u_\epsilon^j := u_\epsilon \circ \Psi_j$. We have that

$$F_\epsilon^a[u_\epsilon, \Omega \cap Q_j, \partial\Omega \cap Q_j] \geq (1 - \delta(\Psi_j))^5 F_\epsilon^a[u_\epsilon^j, D_{r_j} \cap B_j, E_{r_j} \cap B_j],$$

so we conclude that $F_\epsilon^a[u_\epsilon^j, D_{r_j} \cap B_j, E_{r_j} \cap B_j]$ is uniformly bounded in epsilon. Hence, the compactness of the traces Tu_ϵ^j in $L^1(E_{r_j})$ follows from Proposition 24. Finally, using the invertibility of Ψ_j , we have that (Tu_ϵ) is pre-compact in $L^1(\partial\Omega)$ and that its cluster points are in $BV(\partial\Omega, I')$.

7.2 Lower bound inequality

Now we continue with the proof of the theorem. Parts i. and iii. follow by putting together the results in the interior (Sect. 3) and the boundary (Sects. 4, 6).

Let (u_ϵ) be a sequence in $W^{1,2}(\Omega, w_a)$ satisfying $u \in BV(\Omega, I)$, $v \in BV(\partial\Omega, I')$, and such that $u_\epsilon \rightarrow u$ in $L^1(\Omega)$, $Tu_\epsilon \rightarrow v$ in $L^1(\partial\Omega)$. We have to show that

$$\liminf_{\epsilon \rightarrow 0} F_\epsilon^a[u_\epsilon] \geq \Phi(u, v), \tag{7.1}$$

where Φ is given by (1.7). Assume, without loss of generality, that $\liminf_{\epsilon \rightarrow 0} F_\epsilon^a[u_\epsilon] < +\infty$.

For every $\epsilon > 0$, let μ_ϵ be the energy distribution associated with F_ϵ^a with configuration u_ϵ , i.e., μ_ϵ is the positive measure given by

$$\mu_\epsilon(B) := \epsilon^{1-a} \int_{\Omega \cap B} |\nabla u|^2 h^a + \frac{1}{\epsilon^{1-a}} \int_{\Omega \cap B} W(u) h^{-a} + \lambda_\epsilon \int_{\partial\Omega \cap B} V(Tu),$$

for every $B \subset \mathbb{R}^3$. Similarly, we define

$$\begin{aligned} \mu^1(B) &:= \sigma \mathcal{H}^2(S_u \cap B), \\ \mu^2(B) &:= \int_{\partial\Omega \cap B} |\mathcal{W}(Tu) - \mathcal{W}(v)| d\mathcal{H}^2, \\ \mu^3(B) &:= \kappa_s \mathcal{H}^1(S_v \cap B). \end{aligned}$$

The total variation $\|\mu_\epsilon\|$ of the measure μ_ϵ is equal to $F_\epsilon^a[u_\epsilon]$, and $\|\mu^1\| + \|\mu^2\| + \|\mu^3\|$ is equal to $\Phi(u, v)$. Note that $\|\mu_\epsilon\|$ is bounded, so we can assume that μ_ϵ converges in the sense of measure to some finite measure μ . Then, by the lower semicontinuity of the total variation we have

$$\liminf_{\epsilon \rightarrow 0} F_\epsilon^a[u_\epsilon] = \liminf_{\epsilon \rightarrow 0} \|\mu_\epsilon\| \geq \|\mu\|.$$

Since the measures μ^i are mutually singular, we obtain the lower bound inequality (7.1) if we prove that

$$\mu \geq \mu^i \quad \text{for } i = 1, 2, 3.$$

It is enough to show that $\mu(B) \geq \mu^i(B)$ for all sets $B \subset \mathbb{R}^3$ such that $B \cap \Omega$ is a Lipschitz domain and $\mu(\partial B) = 0$.

First, because of Proposition 4 we have that

$$\mu(B) = \lim_{\epsilon \rightarrow 0} \mu_\epsilon(B) \geq \liminf_{\epsilon \rightarrow 0} F_\epsilon^a[u_\epsilon, \Omega \cap B, \emptyset] \geq \sigma \mathcal{H}^2(S_u \cap B) = \mu^1(B).$$

Similarly, we can prove that $\mu \geq \mu^2$. More precisely,

$$\mu(B) = \lim_{\epsilon \rightarrow 0} \mu_\epsilon(B) \geq \liminf_{\epsilon \rightarrow 0} F_\epsilon^a[u_\epsilon, \Omega \cap B, \emptyset] \geq \int_{\partial\Omega \cap B} |\mathcal{W}(T(u)) - \mathcal{W}(v)| d\mathcal{H}^2 = \mu^2(B),$$

where we have used Proposition 26 with $A = B \cap \Omega$ and $A' = B \cap \partial\Omega$.

The inequality $\mu \geq \mu^3$ requires a different argument. Notice that μ^3 is the restriction of \mathcal{H}^1 to the set S_v , multiplied by the factor κ_s . Thus, if we prove that

$$\liminf_{r \rightarrow 0} \frac{\mu(Q_r(x))}{2r} \geq \kappa_s, \quad \mathcal{H}^1 \text{ a.e. } x \in S_v, \tag{7.2}$$

for $Q_r(x)$ as in Proposition 21, we obtain the required inequality. Note that, in any case, μ is supported on $\bar{\Omega}$.

Let us fix $x \in S_v$ such that there exists $\lim_{r \rightarrow 0} \frac{\mu(Q_r(x))}{2r}$, and S_v has one-dimensional density equal to 1. We denote by ν_v the unit normal at x . For r small enough, we choose a

map Ψ_r as in Proposition 21. Set $\bar{u}_\epsilon := u_\epsilon \circ \Psi_r$ and $\bar{v} := v \circ \Psi_r$. Hence, $T\bar{u}_\epsilon \rightarrow \bar{v}$ in $L^1(E_r)$ and $\bar{v} \in BV(E_r, I')$. Moreover,

$$\begin{aligned} \mu(Q_r(x)) &= \lim_{\epsilon \rightarrow 0} \mu_\epsilon(Q_r(x)) \\ &= \lim_{\epsilon \rightarrow 0} F_\epsilon^a[u_\epsilon, \Omega \cap Q_r(x), \partial\Omega \cap Q_r(x)] \\ &\geq \liminf_{\epsilon \rightarrow 0} (1 - \delta(\Psi_r))^5 F_\epsilon^a[\bar{u}_\epsilon, D_r, E_r]. \end{aligned} \tag{7.3}$$

On the other hand, by Proposition 24, we have that

$$\liminf_{\epsilon \rightarrow 0} F_\epsilon^a[\bar{u}_\epsilon, D_r, E_r] \geq \kappa_s \left| \int_{S_{\bar{v}} \cap E_r} v_v d\mathcal{H}^1 \right|.$$

Finally, notice that $\delta(\Psi_r) \rightarrow 0$ as $r \rightarrow 0$, and that

$$\left| \int_{S_{\bar{v}} \cap E_r} v_v d\mathcal{H}^1 \right| = 2r + o(r).$$

Thus we obtain that

$$\frac{\mu(Q_r(x))}{2r} \geq \kappa_s \left(1 + \frac{o(r)}{2r} \right), \text{ as } r \rightarrow 0,$$

that implies $\mu \geq \mu^3$. The proof of the lower bound inequality is completed.

7.3 Upper bound inequality

For the proof of *ii.*, we use a standard construction piece by piece. We will require an extension lemma

Lemma 27 *Let A be a domain in \mathbb{R}^3 , that is contained in the strip $\{r < \text{dist}(\cdot, \partial\Omega) < 2r\}$, and let $A' \subset \partial A$, $v : A' \rightarrow [-m, m]$ a Lipschitz function. Then, for every $\epsilon > 0$, there exists an extension $u : \bar{A} \rightarrow [-m, m]$ such that*

$$\text{Lip}(u) \leq \frac{1}{\epsilon^{1-a} r^a} + \text{Lip}(v)$$

and

$$E_\epsilon^a[u, A] \lesssim \left[(\epsilon^{1-a} r^a \text{Lip}(v) + 1)^2 + C_m \right] (\mathcal{H}^2(\partial A) + o(1)) z, \text{ as } \epsilon \rightarrow 0.$$

where

$$C_m := \max_{t \in [-m, m]} W(t), \quad z = \min\{\|v - \alpha\|_{L^\infty}, \|v - \beta\|_{L^\infty}\}.$$

Proof It follows the ideas of lemma 4.11 in [6], but we need to take care of the weight h . First, we assume without loss of generality, that $A' = \partial A$. In fact, we can extend v to ∂A without increasing its Lipschitz constant. We additionally suppose that $z = \|v - \alpha\|_\infty$; the other case $z = \|v - \beta\|_\infty$ is similar.

Let $\delta = \epsilon^{1-a} r^a$, and set

$$u(x) := \begin{cases} v(x), & \text{on } \partial A, \\ \alpha, & \text{on } A \setminus A_{z\delta}, \end{cases}$$

where A_t is the set of all x in A such that $0 < \text{dist}(x, \partial A) < t$. Then, u is $(\frac{1}{\delta} + \text{Lip}(v))$ -Lipschitz continuous on $\bar{A} \setminus A_{z\delta}$. Finally, u can be extended to \bar{A} , without increasing its Lipschitz constant. We have

$$\begin{aligned} E_\epsilon^a[u, A] &= \epsilon^{1-a} \int_{A_{z\delta}} |\nabla u|^2 h^a + \frac{1}{\epsilon^{1-a}} \int_{A_{z\delta}} W(u) h^{-a} \\ &\lesssim |A_{z\delta}| \left[\epsilon^{1-a} \left(\frac{1}{\delta} + \text{Lip}(v) \right)^2 r^a + \frac{1}{\epsilon^{1-a}} C_m r^{-a} \right] \\ &\leq [(\mathcal{H}^2(\partial A) + o(1)) (\delta \text{Lip}(v) + 1)^2 + C_m] z, \quad \text{as } \epsilon \rightarrow 0, \end{aligned}$$

where we have used that $|A_t| = t (\mathcal{H}^2(\partial A) + o(1))$ as $t \rightarrow 0$. □

Now we are ready for the proof of the upper bound in Theorem 1. Fix $u \in BV(\Omega, I)$ and $v \in BV(\partial\Omega, I')$. It is enough to assume that the singular sets of u and v , S_u and S_v respectively, are closed manifolds of class \mathcal{C}^2 without boundary. This is so because every pair $(u, v) \in BV(\Omega, I) \times BV(\partial\Omega, I')$ can be approximated in $L^1(\Omega) \times L^1(\partial\Omega)$ by pairs that fulfils those regularity assumptions (see theorem 1.24 of [15]). We assume that u and v , up to modifications on negligible sets, are constant in each connected component of $\Omega \setminus S_u$ and $\partial\Omega \setminus S_v$ respectively.

The idea is to construct a partition of Ω into four subsets, and to use the preliminary convergence results of the previous sections to obtain the upper bound inequality.

For every $x \in \Omega$, set $d(x) = \text{dist}(x, \partial\Omega)$ and $d' : \partial\Omega \rightarrow \mathbb{R}$ is the oriented distance from S_v defined by

$$d'(x) = \begin{cases} \text{dist}(x, S_v) & \text{if } x \in \{v = \beta'\}, \\ -\text{dist}(x, S_v) & \text{if } x \in \{v = \alpha'\}. \end{cases}$$

For every $r > 0$, set

$$\Gamma_r := \{x \in \Omega : \text{dist}(x, \partial\Omega) = r\}.$$

Fix $r > 0$ such that Γ_r and Γ_{2r} are Lipschitz surfaces and $S_u \cap \Gamma_r$ is a Lipschitz curve. With this in mind, we construct a partition of Ω . Let

$$\begin{aligned} B_1 &:= \{x \in \Omega : \text{dist}(x, S_v \cup (S_u \cap \Gamma_r)) < 3r\}, \\ A_1 &:= \{x \in \Omega \setminus \bar{B}_1 : d(x) < r\}, \\ B_2 &:= \{x \in \Omega \setminus \bar{B}_1 : r < d(x) < 2r\}, \\ A_2 &:= \{x \in \Omega \setminus \bar{B}_1 : 2r < d(x)\}. \end{aligned}$$

We will construct a Lipschitz function $u_\epsilon := u_{r,\epsilon}$ for every $\epsilon < r$, piece by piece, with controlled Lipschitz constant.

Step 1: In the set A_2 , we take u_ϵ as in part *ii.* of Proposition 4. We extend it to ∂A_2 by continuity. Hence u_ϵ is $\frac{C}{\epsilon^{1-a}}$ -Lipschitz on \bar{A}_2 , u_ϵ converges pointwise to u in A_2 , uniformly on $\partial A_2 \cap \partial B_2$, and

$$F_\epsilon^a[u_\epsilon, A_2, \emptyset] = E_\epsilon^a[u_\epsilon, A_2] \leq \sigma \mathcal{H}^2(S_u \cap A_2) + o(1) \leq \sigma \mathcal{H}^2(S_u) - \sigma \mathcal{H}^2(S_u \setminus A_2) + o(1)$$

as $\epsilon \rightarrow 0$.

Step 2: Now we consider the set A_1 . The function u is constant (equal to α or β) in every connected component A of A_1 on $\partial A \cap \partial\Omega$, and the function v is constant (equal to α' or β') on $\partial A \cap \partial\Omega$. Then we can use Proposition 26 to get a function u_ϵ such that $Tu_\epsilon = v$ on

$\partial A \cap \partial \Omega$ and u_ϵ converges to u pointwise in A_1 and uniformly on every subset with positive distance from $\partial A \cap \partial \Omega$. By the same proposition, we also have that u_ϵ is $\frac{C_W}{\epsilon^{1-a}r^a}$ -Lipschitz continuous on \bar{A}_1 , and we can extend it to ∂A_1 by continuity. Since the distance of two connected components of A_1 is larger than r and $\frac{1}{\epsilon^{1-a}r^a} > \frac{1}{r}$, choosing $C > \max\{2m, C_W\}$, it follows that u_ϵ is $\frac{C}{\epsilon^{1-a}r^a}$ -Lipschitz continuous on \bar{A}_1 and agrees with v on $\partial A_1 \cap \partial \Omega$. Moreover, the function u_ϵ satisfies

$$F_\epsilon^a[u_\epsilon, A_1, \partial A_1 \cap \partial \Omega] = E_\epsilon^a[u_\epsilon, A_1] \leq \int_{\partial A_1 \cap \partial \Omega} |\mathcal{W}(Tu_\epsilon) - \mathcal{W}(v)| + o(1), \quad \text{as } \epsilon \rightarrow 0.$$

Step 3: Note that in the previous steps we have constructed an optimal sequence in $\bar{A}_1 \cup \bar{A}_2$ that is $\frac{C}{\epsilon^{1-a}r^a}$ -Lipschitz continuous, in particular, it is defined and Lipschitz on $((\partial A_1 \cup \partial A_2) \cap \partial B)$, for every connected component B of B_2 .

By virtue of Lemma 27, we can extend u_ϵ to every B , obtaining a $\frac{C+1}{\epsilon^{1-a}r^a}$ -Lipschitz continuous function that satisfies

$$F_\epsilon^a[u_\epsilon, B_2, \emptyset] = E_\epsilon^a[u_\epsilon, B_2] \leq z_\epsilon ((C + 2)^2 + C_m) (\mathcal{H}^2(\partial B_2) + o(1)) = o(1)$$

as $\epsilon \rightarrow 0$, where we have used that $z_\epsilon := \inf_{(\partial A_1 \cup \partial A_2) \cap \partial B_2} |u_\epsilon - u| = o(1)$, since u_ϵ is constant on each connected components of B_2 .

Step 4: To construct the function in the piece B_1 is the most delicate step. First, we need some preliminaries: construct a function on the whole \mathbb{R}_+^2 with suitable behavior.

Consider the rescalings

$$\Lambda_\epsilon \ll \sigma_\epsilon \ll \rho_\epsilon \ll \epsilon^{1-a}$$

for some $\sigma_\epsilon = \epsilon^p, \rho_\epsilon = \epsilon^q$. Let \bar{w}_1 be the function on \mathbb{R}_+^2 defined in Proposition 20 and its rescaling $\bar{w}_\epsilon(x, y) := \bar{w}_1\left(\frac{x}{\Lambda_\epsilon}, \frac{y}{\Lambda_\epsilon}\right)$. We also consider the function w_1 from Lemma 25 and its rescaling $w_\epsilon(x, y) := w_1\left(\frac{x}{\Lambda_\epsilon}, \frac{y}{\Lambda_\epsilon}\right)$. We glue them, so that we obtain a function defined in the whole \mathbb{R}_+^2 , as

$$\tilde{w}_1 := \begin{cases} w_1, & \text{if } (x, y) \in D_{\sigma_\epsilon} \\ \bar{w} & \text{if } (x, y) \in \mathbb{R}_+^2 \setminus D_{\rho_\epsilon} \end{cases}$$

and smooth in between, with its corresponding rescaling

$$\tilde{w}_\epsilon(x, y) := \tilde{w}_1\left(\frac{x}{\Lambda_\epsilon}, \frac{y}{\Lambda_\epsilon}\right). \tag{7.4}$$

Because $\rho_\epsilon \gg \sigma_\epsilon \gg \Lambda_\epsilon$, we can apply Lemma 25 to obtain

$$H_\epsilon^a[\tilde{w}_\epsilon, D_{\rho_\epsilon}, E_{\rho_\epsilon}] = \kappa_s - o(1)$$

when $\epsilon \rightarrow 0$.

Now we pass from two to three dimensions. In particular, we set \tilde{u}_ϵ on $S_v \times \mathbb{R}_+^2$ to be

$$\tilde{u}_\epsilon(x, y, z) := \tilde{w}_\epsilon(x, y) \quad \text{for every } z \in S_v, (x, y) \in \mathbb{R}_+^2. \tag{7.5}$$

where \tilde{w}_ϵ is defined in (7.4). In addition, for any function \tilde{u} defined on $S_v \times D$, we define the following functional.

$$\begin{aligned} \tilde{F}_\epsilon^a[\tilde{u}, S_v \times D, S_v \times E] &:= \epsilon^{1-a} \int_{S_v \times D} |\nabla \tilde{u}|^2 y^a \, dx dy dz \\ &\quad + \frac{1}{\epsilon^{1-a}} \int_{S_v \times D} W(\tilde{u}) y^{-a} \, dx dy dz + \lambda_\epsilon \int_{S_v \times E} V(T\tilde{u}) \, dx dz. \end{aligned}$$

For the \tilde{u}_ϵ we have constructed in (7.5), Fubini’s theorem implies that

$$\begin{aligned} &\tilde{F}_\epsilon^a[\tilde{u}_\epsilon, S_v \times D_{\rho_\epsilon}, S_v \times E_{\rho_\epsilon}] \\ &= \mathcal{H}^1(S_v) \left(H_\epsilon^a[\tilde{w}_\epsilon, D_{\rho_\epsilon}, E_{\rho_\epsilon}] + \frac{1}{\epsilon^{1-a}} \int_{D_{\rho_\epsilon}} W(\tilde{w}_\epsilon) y^{-a} \, dx dy \right) \\ &\leq \mathcal{H}^1(S_v) \left(H_\epsilon^a[\tilde{w}_\epsilon, D_{\rho_\epsilon}, E_{\rho_\epsilon}] + \frac{C}{\epsilon^{1-a}} \int_{x \in (-\rho_\epsilon, \rho_\epsilon), y \in (0, \rho_\epsilon)} y^{-a} \, dx dy \right) \\ &= \mathcal{H}^1(S_v) \left(H_\epsilon^a[\tilde{w}_\epsilon, D_{\rho_\epsilon}, E_{\rho_\epsilon}] + \frac{C'}{\epsilon^{1-a}} \rho_\epsilon^{2-a} \right). \end{aligned} \tag{7.6}$$

We choose $1 - a < p < q < \frac{1-a}{-a}$ and $q > \frac{1-a}{2-a}$. Then from (7.6) we obtain that

$$\tilde{F}_\epsilon^a[\tilde{u}_\epsilon, S_v \times D_{\rho_\epsilon}, S_v \times E_{\rho_\epsilon}] \leq \mathcal{H}^1(S_v) [\kappa_s + o(1)] \tag{7.7}$$

as $\epsilon \rightarrow 0$.

Now we transplant the function \tilde{u}_ϵ obtained to our remaining piece B_1 . Since S_v is a boundary in $\partial\Omega$, we can construct a diffeomorphism between the intersection of a tubular neighborhood U of S_v and Ω and the product of S_v with a half-disk. More precisely, for every $r > 0$, we set

$$\mathcal{S}_r := \{x \in \Omega : 0 < \text{dist}(x, S_v) < r\}.$$

For every $x \in \bar{\Omega}$, define

$$\Psi(x) := (x'', d'(x'), \text{dist}(x, \partial\Omega)),$$

where x' is a projection of x on $\partial\Omega$ and x'' is a projection of x' on S_v . The function Ψ is well-defined and is a diffeomorphism of class \mathcal{C}^2 on $\bar{\Omega} \cap U$, and satisfies the following properties:

- $\Psi(\Omega \cap U) \subset S_v \times \mathbb{R}_+^2$,
- $\Psi(\partial\Omega \cap U) \subset S_v \times \mathbb{R} \times \{0\}$,
- $\Psi(x) = x$ for every $x \in \partial\Omega$.
- $D\Psi(x)$ is an isometry,
- $\lim_{r \rightarrow 0} \delta_r = 0$, where δ_r is the isometry defect of the restriction of Ψ to \mathcal{S}_r .

We construct u_ϵ on $S := \bar{\mathcal{S}}_{\rho_\epsilon/2}$. With small modifications, we can assume that S is a “cubical” neighborhood such that for ϵ small, the function Ψ maps S into $S_v \times D_{\rho_\epsilon}$ and $\partial S \cap \partial\Omega$ into

$S_v \times E_{\rho_\epsilon}$, Then we define $u_\epsilon := \tilde{u}_\epsilon \circ \Psi$, where \tilde{u}_ϵ is defined in (7.5). Thus Proposition 21 and (7.7) give that

$$\begin{aligned} F_\epsilon^a[u_\epsilon, S, \partial S \cap \partial \Omega] &\leq (1 - \delta_\epsilon)^{-5} F_\epsilon^a[\tilde{u}_\epsilon, S_v \times D_{\rho_\epsilon}, S_v \times E_{\rho_\epsilon}] \\ &\leq \mathcal{H}^1(S_v) (\kappa_S + o(1)) \end{aligned}$$

as $\epsilon \rightarrow 0$, because $\delta_\epsilon := \delta(\Psi|_{S_{\rho_\epsilon}})$ tends to zero as $\epsilon \rightarrow 0$.

Notice that for ϵ small enough, Ψ is 2-Lipschitz continuous. Using again Lemma 27, we can extend u_ϵ by setting $u_\epsilon = v$ on the remaining part of $\partial B_1 \cap \partial \Omega$. We have that u_ϵ is equal to v on $\partial \Omega \setminus S$. Thus, we can extend u_ϵ on the whole $B_1 \setminus S$ to a $\frac{2C+1}{\epsilon^{1-a}r^a}$ -Lipschitz continuous function, which satisfies

$$\begin{aligned} F_\epsilon^a[u_\epsilon, B_1 \setminus S, \partial(B_1 \setminus S) \cap \partial \Omega] &= H_\epsilon^a[u_\epsilon, B_1 \setminus S] \\ &\leq ((2C + 2)^2 + C_m) (\mathcal{H}^2(\partial B_1) + o(1)) 2m \end{aligned}$$

as $\epsilon \rightarrow 0$, where we have used $\|u_\epsilon - \alpha\|_\infty \wedge \|u_\epsilon - \beta\|_\infty \leq 2m$.

Step 5: We recall that for every $r > 0$ and every $\epsilon < r$ we have constructed a function u_ϵ defined on the whole Ω such that

$$\limsup_{\epsilon \rightarrow 0} \|u_\epsilon - u\|_{L^1(\Omega)} \leq 2m (|B_1| + |B_2|)$$

and

$$\limsup_{\epsilon \rightarrow 0} \|Tu_\epsilon - v\|_{L^1(\partial \Omega)} = 0.$$

Since $|B_1|$ and $|B_2|$ have order r^2 and r respectively, we get that $u_\epsilon \rightarrow u$ in $L^1(\Omega)$, first taking $\epsilon \rightarrow 0$ and then $r \rightarrow 0$.

Combining all the results above, we obtain

$$\begin{aligned} \limsup_{\epsilon \rightarrow 0} F_\epsilon^a[u_\epsilon] &\leq \sigma \mathcal{H}^2(S_u) + \int_{\partial \Omega} |\mathcal{W}(Tu(x)) - \mathcal{W}(v(x))| d\mathcal{H}^2 + \kappa_S \mathcal{H}^1(S_v) \\ &\quad - \sigma \mathcal{H}^2(S_u \setminus A_2) + ((2C + 2)^2 + C_m) (\mathcal{H}^2(\partial B_1) + o(1)) 2m. \end{aligned}$$

Since $\mathcal{H}^2(\partial B_1)$ has order r , taking $r \rightarrow 0$ above we deduce the upper bound inequality. Finally, applying a suitable diagonalization argument, to the sequence $u_\epsilon := u_{\epsilon,r}$ we obtain the desired sequence u_ϵ . Proof of *ii.* is completed.

Acknowledgments I would like to thank Prof. Luis Caffarelli for suggesting the study of fractional order operators and his support at UT Austin.

References

1. Adams, R.A.: Sobolev spaces, Pure and Applied Mathematics, vol. 65. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York (1975)
2. Abramowitz, M., Stegun, I.A.: Handbook of mathematical functions with formulas, graphs, and mathematical tables. National Bureau of Standards Applied Mathematics Series, vol. 55. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC (1964)
3. Alberti, G.: Variational models for phase transitions, an approach via Γ -convergence. In: Calculus of variations and partial differential equations (Pisa, 1996), pp. 95–114. Springer, Berlin (2000)
4. Alberti, G., Bellettini, G.: A non-local anisotropic model for phase transitions: asymptotic behaviour of rescaled energies. Eur. J. Appl. Math. **9**(3), 261–284 (1998)

5. Alberti, G., Bouchitté, G., Seppecher, P.: Un résultat de perturbations singulières avec la norme $H^{1/2}$. *C. R. Acad. Sci. Paris Sér. I Math.* **319**(4), 333–338 (1994)
6. Alberti, G., Bouchitté, G., Seppecher, P.: Phase transition with the line-tension effect. *Arch. Rational Mech. Anal.* **144**(1), 1–46 (1998)
7. Cabré, X., Cónsul, N.: Minimizers for boundary reactions: renormalized energy, location of singularities, and applications (in preparation)
8. Cabré, X., Solà-Morales, J.: Layer solutions in a half-space for boundary reactions. *Comm. Pure Appl. Math.* **58**(12), 1678–1732 (2005)
9. Caffarelli, L., Silvestre, L.: An extension problem related to the fractional Laplacian. *Comm. Partial Differ. Equ.* **32**(7-9), 1245–1260 (2007)
10. Evans, L.C., Gariepy, R.F.: Measure theory and fine properties of functions. *Studies in Advanced Mathematics*. CRC Press, Boca Raton (1992)
11. Fabes, E.B., Kenig, C.E., Serapioni, R.P.: The local regularity of solutions of degenerate elliptic equations. *Comm. Partial Differ. Equ.* **7**(1), 77–116 (1982)
12. García-Cuerva, J., Rubio de Francia J.L.: Weighted norm inequalities and related topics. *North-Holland Mathematics Studies*, vol. 116. North-Holland, Amsterdam, *Notas de Matemática [Mathematical Notes]*, 104 (1985)
13. Garroni, A., Palatucci, G.: A singular perturbation result with a fractional norm. In: *Variational problems in materials science. Progr. Nonlinear Differential Equations Appl.*, vol. 68, pp. 111–126. Birkhäuser, Basel (2006)
14. Garsia, A.M., Rodemich, E.: Monotonicity of certain functionals under rearrangement. *Ann. Inst. Fourier (Grenoble)* **24**(2):vi, 67–116, (1974) *Colloque International sur les Processus Gaussiens et les Distributions Aléatoires (Colloque Internat. du CNRS, No. 222, Strasbourg, 1973)*
15. Giusti, E.: Minimal surfaces and functions of bounded variation. *Monographs in Mathematics*, vol. 80. Birkhäuser Verlag, Basel (1984)
16. Landkof, N.S.: Foundations of modern potential theory. Springer, New York. Translated from the Russian by A. P. Doohovskoy, *Die Grundlehren der mathematischen Wissenschaften, Band 180* (1972)
17. Modica, L.: The gradient theory of phase transitions and the minimal interface criterion. *Arch. Rational Mech. Anal.* **98**(2), 123–142 (1987)
18. Modica, L.: Gradient theory of phase transitions with boundary contact energy. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **4**(5), 487–512 (1987)
19. Modica, L., Mortola, S.: Un esempio di Γ^- -convergenza. *Boll. Un. Mat. Ital. B* (5) **14**(1), 285–299 (1977)
20. del Mar González, M., Monneau, R.: Motion of particle systems as a limit of a reaction-diffusion equation with fractional laplacian (in preparation)
21. Nekvinda, A.: Characterization of traces of the weighted Sobolev space $W^{1,p}(\Omega, d_M^\epsilon)$ on M . *Czechoslovak Math. J.* **43**(118)(4), 695–711 (1993)
22. Palatucci, G.: Phase transitions with the line tension effect: the super-quadratic case. *Math. Models Methods Appl. Sciences*, accepted paper (2008)
23. Palatucci, G.: A class of phase transition problems with the line tension effect. Ph.D. thesis (2007)
24. Palatucci, G., Sire, Y.: Γ^- -convergence of some super quadratic functionals with singular weights. Preprint (2008)
25. Silvestre, L.: Regularity of the obstacle problem for a fractional power of the Laplace operator. Ph.D. thesis (2005)
26. Stein, E.M.: Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals. *Princeton Mathematical Series*, vol. 43. Princeton University Press, Princeton, NJ (1993) With the assistance of Timothy S. Murphy, *Monographs in Harmonic Analysis, III*
27. Valadier, M.: Young measures. In: *Methods of nonconvex analysis. Lecture Notes in Mathematics*, vol. 1446, pp. 152–188. Springer, Berlin (1990)
28. Watson, G.N.: A treatise on the theory of Bessel functions. *Cambridge Mathematical Library*. Cambridge University Press, Cambridge (1995). Reprint of the second (1944) edition