



Can we run to infinity? The diameter of the diffeomorphism group with respect to right-invariant Sobolev metrics

Martin Bauer¹ · Cy Maor²

Received: 3 October 2020 / Accepted: 1 January 2021 / Published online: 10 February 2021
© The Author(s), under exclusive licence to Springer-Verlag GmbH, DE part of Springer Nature 2021

Abstract

The group $\text{Diff}(\mathcal{M})$ of diffeomorphisms of a closed manifold \mathcal{M} is naturally equipped with various right-invariant Sobolev norms $W^{s,p}$. Recent work showed that for sufficiently weak norms, the geodesic distance collapses completely (namely, when $sp \leq \dim \mathcal{M}$ and $s < 1$). But when there is no collapse, what kind of metric space is obtained? In particular, does it have a finite or infinite diameter? This is the question we study in this paper. We show that the diameter is infinite for strong enough norms, when $(s - 1)p \geq \dim \mathcal{M}$, and that for spheres the diameter is finite when $(s - 1)p < 1$. In particular, this gives a full characterization of the diameter of $\text{Diff}(S^1)$. In addition, we show that for $\text{Diff}_c(\mathbb{R}^n)$, if the diameter is not zero, it is infinite.

Mathematics Subject Classification 58D05 · 58B20

Contents

1	Introduction and main results	2
1.1	Right-invariant Sobolev metrics on diffeomorphism groups: where they arise	3
1.2	Previous results on the geometry induced by right-invariant Sobolev metrics	4
1.3	Main ideas in the proofs	5
1.4	Some open questions	6
2	Right-invariant $W^{s,p}$ -norms on diffeomorphism groups	6
2.1	Fractional order Sobolev spaces on \mathbb{R}^n	7
2.2	Fractional order Sobolev norms on Riemannian manifolds	8
2.3	Right-invariant fractional order Sobolev metrics on diffeomorphism groups	8
3	The diameter of $\text{Diff}(S^1)$	9
3.1	The supercritical case $s > 1 + 1/p$	9
3.2	The critical case $s = 1 + 1/p$	10

Communicated by J. Jost.

✉ Cy Maor
cy.maor@mail.huji.ac.il
Martin Bauer
bauer@math.fsu.edu

¹ Department of Mathematics, Florida State University, Tallahassee, Florida, USA

² Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem, Israel

3.3 The subcritical case $s < 1 + 1/p$ 12
 3.4 The displacement energy 14
 4 The diameter of $\text{Diff}(M)$ for compact manifolds in higher dimensions 15
 4.1 The supercritical case $s > 1 + \frac{\dim \mathcal{M}}{p}$ 16
 4.2 The critical case $s = 1 + \frac{\dim \mathcal{M}}{p}$ 16
 4.3 The subcritical case $s < 1 + \frac{1}{p}$ 17
 4.4 The displacement energy 21
 5 The diameter of $\text{Diff}_c(\mathbb{R}^n)$ 24
 Appendix A: Proof of Lemma 3.9 25
 Appendix B: Sobolev norms of radial functions 29
 Appendix C: Diameter and displacement energy 31
 References 33

1 Introduction and main results

Right-invariant Sobolev metrics on diffeomorphism groups (or on subgroups thereof) arise naturally in several contexts—they play a central role in mathematical shape analysis, appear in symplectic geometry and their geodesic equations turn out to be related to several important partial differential equations in hydrodynamics (some more details are given in Sect. 1.1 below).

The basic setting is the following (see Sect. 2 for details): The diffeomorphism group $\text{Diff}(\mathcal{M})$ of a compact manifold \mathcal{M} is a Lie group, whose associated Lie algebra is the space of vector fields $\mathcal{X}(\mathcal{M})$. One can equip the diffeomorphism group with right-invariant Finsler metrics, by considering norms on the Lie algebra $\mathcal{X}(\mathcal{M})$; in this article we will focus on Sobolev norms $W^{s,p}$. For $p = 2$, these norms induce Riemannian metrics on the group $\text{Diff}(\mathcal{M})$; these are the most important metrics for hydrodynamics and shape analysis. The corresponding geodesic distance between $\varphi_0, \varphi_1 \in \text{Diff}(\mathcal{M})$ induced by this metric is then given by the variational problem

$$\text{dist}_{s,p}(\varphi_0, \varphi_1) := \inf_{\substack{\varphi: [0,1] \rightarrow \text{Diff}_c(\mathcal{M}) \\ \varphi(0) = \varphi_0, \varphi(1) = \varphi_1}} \int_0^1 \|\partial_t \varphi \circ \varphi^{-1}\|_{W^{s,p}}.$$

If the metric is weak enough, then this distance vanishes identically on every connected component; this phenomenon was first shown by Michor and Mumford [47], and was then analyzed in a series of works, culminating in a complete characterization in [32]. In this article we study a finer property of the geodesic distance, namely, the diameter of (connected components of) diffeomorphism groups with respect to the geodesic distance induced by these metrics:

$$\text{diam}_{s,p} \text{Diff}(\mathcal{M}) := \sup_{\varphi_0, \varphi_1 \in \text{Diff}(\mathcal{M})} \text{dist}_{s,p}(\varphi_0, \varphi_1).$$

Our main result is the following characterization of boundedness/unboundedness of this diameter:

Theorem 1.1 *Let \mathcal{M} be a compact manifold without boundary of dimension $n \geq 1$. Then the diameter of the connected component of the identity of the group of smooth diffeomorphisms with respect to the right-invariant $W^{s,p}$ -metric is*

1. Zero when $s \leq \frac{n}{p}$ and $s < 1$, i.e., $\text{diam}_{s,p} \text{Diff}(\mathcal{M}) = 0$;

2. For $M = S^n$, bounded but non-zero for $s \in (\frac{n}{p}, 1 + \frac{1}{p}) \cup [1, 1 + \frac{1}{p})$, i.e., $\text{diam}_{s,p} \text{Diff}(S^n) \in (0, \infty)$;
3. Unbounded for $s \geq 1 + \frac{n}{p}$, i.e., $\text{diam}_{s,p} \text{Diff}(M) = \infty$.

Here in $W^{s,p}$ we denote by s the number of derivatives and by p the exponent. The exact definition of these Sobolev norms, in particular for non-integer s , appears in Sect. 2. As mentioned above, the first part of this theorem is known from recent previous results on vanishing geodesic distance (see [32] and the references therein); the second and third parts are the main contributions of this paper. More generally, we prove the second part of the theorem for any closed manifold M that satisfies a certain uniform fragmentation property (see Definition 4.4), which holds in particular for spheres.

Note that for $M = S^1$ our theorem gives a complete characterization of boundedness (unboundedness, resp.) of the diameter of $\text{Diff}(S^1)$, i.e., we have that $\text{diam}_{s,p} \text{Diff}(S^1)$ is bounded for $s < 1 + \frac{1}{p}$ and unbounded otherwise. For higher dimensional spheres, there is a gap in the range $s \in [1 + \frac{1}{p}, 1 + \frac{n}{p})$, in which we do not know whether the diameter is finite or not. We believe (as explained in Sect. 4.4), that the diameter is finite in this range, that is, that the transition to infinite diameter happens at $s = 1 + \frac{n}{p}$.

If M is the non-compact space $M = \mathbb{R}^n$, we prove that the diameter only exhibits two different behaviors: it is either zero or unbounded. This leads to following complete characterization:

Theorem 1.2 *Let $p \geq 1$. The diameter of $\text{diam}_{s,p} \text{Diff}_c(\mathbb{R}^n)$ with respect to the right-invariant $W^{s,p}$ -metric is infinite if and only if $s \geq 1$ or $sp > n$. In any other case the diameter is zero.*

Here $\text{Diff}_c(\mathbb{R}^n)$ denotes the connected component of the identity of the group of compactly supported diffeomorphisms. Again, the zero diameter part is due to previous work, and the contribution of this paper is the infinite diameter part.

As described below, results on the boundedness/unboundedness of the diameter have been studied for a long time in the context of metrics on symplectomorphisms and volume-preserving diffeomorphisms. To the best of our knowledge, this work is the first to address this question for the full diffeomorphism group, and to show a transition of zero to finite to infinite diameter for a hierarchy of metrics.

1.1 Right-invariant Sobolev metrics on diffeomorphism groups: where they arise

The interest in right-invariant metrics on diffeomorphism groups originates from Arnold’s seminal observation [2] that Euler’s equation for the motion of an incompressible fluid admits a geometric interpretation in this setup: it is the geodesic equation of the right-invariant L^2 -metric on the group of all volume preserving diffeomorphisms (we will refer to this group also as volumorphism group)¹. Subsequent to Arnold’s geometric interpretations for Euler’s equation similar formulations have been found for several other partial differential equations that are of relevance in the field of mathematical physics; examples include

- the Camassa–Holm equation [18,40,49], which corresponds to the $W^{1,2}$ -metric on $\text{Diff}(S^1)$;

¹ This is, by no means, an excessive surveyy.

- the Hunter–Saxton equation [31,37,42,43], as the geodesic equation of the homogeneous $W^{1,2}$ -metric on $\text{Diff}(S^1)/S^1$; and more generally the p -Hunter–Saxton equation, the geodesic equation of the homogeneous $W^{1,p}$ -metric on $\text{Diff}(S^1)/S^1$, as introduced recently by Cotter et al. [20].
- the modified Constantin–Lax–Majda equation [11,19,25,62] corresponding to the homogeneous $W^{1/2,2}$ -metric on the same space. See, e.g., [57] and the references therein for further examples of Euler–Arnold equations, that are of relevance in mathematical physics.

An additional motivation for the study of (higher order) right-invariant metrics on the full diffeomorphism group stems from their central role in the field of mathematical shape analysis, where differences between objects such as point clouds, images, surfaces, or densities are encoded in the spirit of Grenander’s pattern analysis [27,28,51] by the cost of the minimal (diffeomorphic) transformation that (approximately) transports a source shape to a target shape. Using a right-invariant metric on the diffeomorphism group to measure the cost of these diffeomorphic transformations yields the so-called LDDMM-setting [7,12,34,48,63], which has proven successful in numerous applications in computational anatomy and medical imaging.

In yet another important line of research, right-invariant Sobolev metrics play a role in symplectic and contact geometry, starting from the Hofer metric on Hamiltonian symplectomorphisms [30], which is in this context a right-invariant $W^{-1,\infty}$ metric (which is also a bi-invariant metric).

1.2 Previous results on the geometry induced by right-invariant Sobolev metrics

The geodesic equations of right-invariant metrics, as they are related to many important partial differential equations, have been studied extensively, starting from Ebin and Marsden [21] who obtained local well-posedness and stability results for solutions to Euler’s equation by studying the geodesic spray of the right-invariant L^2 -metric on volumorphisms. Subsequently, local well-posedness results (and sometimes even global existence), have been obtained, using analogous methods, for geodesic equations on the diffeomorphism group as well [8,40,49,50,60]. See [16,39] for an overview on these results. Furthermore, Preston et al. [35,50] studied the curvature of the corresponding spaces and showed Fredholm properties of the exponential map for both volumorphisms and diffeomorphism groups.

In addition to the geodesic equation itself, right-invariant Sobolev metrics enable us to measure the lengths of curves, hence they give a structure of a length space on these diffeomorphism groups. A natural question is then—is this structure degenerate? That is, can the distance between two distinct diffeomorphisms be zero (meaning that there are arbitrarily short curves between them)? This is known as the vanishing geodesic distance phenomenon. On the other hand, one can ask—can we find two diffeomorphisms that are arbitrarily far away (i.e., the diameter is infinite)? This is the question we address in this paper. Note that these questions are of importance when this geodesic distance is explicitly used, e.g., as the regularization term in the LDDMM-setting.

The vanishing geodesic distance phenomenon was first shown for Hamiltonian symplectomorphisms [23] under $W^{-1,p}$ metrics for $p < \infty$ (in contrast to the Hofer $W^{-1,\infty}$ metric, which is non-degenerate). This was later extended to stronger metrics in [9]. Similar results were later obtained for contactomorphisms [54]. In the context of the full diffeomorphism group, the geodesic distance has been first investigated by Michor and Mumford in [47], where they showed the degeneracy (vanishing) of the geodesic distance for the L^2 -metric

and the non-degeneracy for metrics of order $W^{1,2}$ and above. These results have been later generalized to fractional $W^{s,p}$ -metrics and a complete characterization of vanishing (non-vanishing) geodesic distance for this class of metrics has been obtained [5,6,9,32,33]. The first part of Theorem 1.1 is essentially this characterization.

The diameter question was initiated by Shnirelman [55,56] who studied the diameter of the volumorphism group $\text{Diff}_\mu(\mathcal{M})$ with respect to the geodesic distance of the L^2 -metric. In particular he showed the boundedness of the diameter for contractible manifolds of dimension $\dim \mathcal{M} \geq 3$, and conjectured the unboundedness in the two-dimensional case.² For \mathcal{M} being either a (two-dimensional) surface with boundary or a closed surface of genus $g \geq 2$ this conjecture has been shown to be true by Eliashberg and Ratiu [24] for any L^p metric, $p \geq 1$. The case of the torus and the (significantly more complicated) case of S^2 were proved in [14], thus proving infinite diameter for any closed two dimensional surface with respect to the L^p metric. Eliashberg and Ratiu [24] also show that some higher dimensional manifolds with non-trivial topology have infinite diameter for this metric. So far, the analogue of Shnirelman's question regarding boundedness (unboundedness resp.) of the diameter of the full diffeomorphism group, has not been investigated.

1.3 Main ideas in the proofs

Interestingly, the techniques used in this paper are completely orthogonal to the ones used for studying the diameter of symplectomorphisms and volumorphisms, in the sense that all our proofs, both of boundedness and unboundedness of diameter, rely on volume change.

For sufficiently strong metrics, such that $W^{s,p}$ embeds in C^1 (that is, when $(s-1)p > n$), we show that we can bound from below the geodesic distance $\text{dist}_{s,p}(\text{Id}, \varphi)$ of a diffeomorphism φ to the identity by the logarithm of the Jacobian determinant of φ at any point. In particular, the distance from the identity to a diffeomorphism with an arbitrarily large volume change at a point is arbitrarily large. We call this the *supercritical case*.

The *critical case* is the one for which this embedding just fails, namely when $(s-1)p = n$. Here we extend an idea of Lenells [42] to construct an isometry from a degenerate $W^{1,q}$ -type metric that sees *only* volume changes to the space of smooth functions on \mathcal{M} , and from this we obtain a lower bound for the diameter of this metric, which diverges with q . This degenerate $W^{1,q}$ -type metric is weaker, for any $q < \infty$, than our critical $W^{s,p}$ metric, and by using the bound above and controlling the Sobolev embedding constants, we obtain the unboundedness of the diameter by letting $q \rightarrow \infty$.

In the *subcritical case*, when $(s-1)p < n$, we aim to prove that the diameter is bounded, at least for manifolds with nice fragmentation properties (see Sect. 4.3). In these cases the question can be reduced to a local question on the diameter of the diffeomorphism group of the Euclidean ball $\text{Diff}_c(B_1(\mathbb{R}^n))$. We show, by a rescaling argument, that for $(s-1)p < n$ the uniform boundedness of $\text{dist}_{s,p}(\text{Id}, \varphi)$ for any $\varphi \in \text{Diff}_c(B_1(\mathbb{R}^n))$ is equivalent to the uniform boundedness of $\text{dist}_{s,p}(\text{Id}, \psi_\lambda)$ for a class of diffeomorphisms $\psi_\lambda(x) \approx \lambda x$, as $\lambda \rightarrow \infty$. That is, boundedness of the diameter of an arbitrary, radially symmetric change of volume at a point implies the boundedness of the whole diffeomorphism group. We then show that when $(s-1)p < 1$, $\text{dist}_{s,p}(\text{Id}, \psi_\lambda)$ is indeed bounded, and give an indication of the fact that arbitrary changes of volume should be of bounded cost for the whole subcritical case $(s-1)p < n$.

² To be exact, Shnirelman proved the boundedness of the diameter of $\text{Diff}_\mu(\mathcal{M})$ when \mathcal{M} is the three-dimensional cube, but his proof can be modified to show the result for contractible manifolds of dimension $\dim \mathcal{M} \geq 3$, 0 see, e.g., [3,38].

1.4 Some open questions

- As mentioned, for spheres we have a gap in the range $s \in \left[1 + \frac{1}{p}, 1 + \frac{n}{p}\right)$; in order to bridge it and prove the boundedness of diameter in this range, we need to find a better way to transport the identity to the family of diffeomorphisms ψ_λ mentioned above.
- We do not know whether for other closed manifolds, that do not satisfy our fragmentation assumption, the diameter of $\text{Diff}(\mathcal{M})$ is finite or not in the subcritical case. That is, are there closed manifolds for which the diameter is either zero or infinity?
- Another open line of work is to extend the analysis to $W^{s,p}$ metrics on volumorphisms and symplectomorphisms, in particular for negative values of s . There it is not known what is the critical case below which the geodesic distance vanishes (it is known to vanish for $s \leq -1 + \frac{1}{p}$ and to not vanish for $s \geq 0$, see [9]). Also, to the best of our knowledge, it is not known if a similar phenomenon as seen here for $\text{Diff}(S^n)$, namely a transition zero \rightarrow finite \rightarrow infinite diameter, can occur for symplectomorphisms/volumorphisms on closed manifolds.
- More generally, it would be interesting to better understand the connections between the metric questions (vanishing geodesic distance, boundedness of diameter) to other geometric properties (having a smooth geodesic spray, Fredholm properties, etc.).

The structure of this paper In Sect. 2 we define the (fractional order) Sobolev norms we are considering in this paper, discuss some of their embedding properties, and define the right-invariant metrics they induce on the diffeomorphism group. In Sect. 3 we discuss the one dimensional case, namely, the full characterization of boundedness/unboundedness of the diameter for $\text{Diff}(S^1)$; this case already includes most of the key ingredients that are used in the higher dimensional case, which is the content of Sect. 4, in which we complete the proof of Theorem 1.1. Finally, in Sect. 5, we discuss the case of $\text{Diff}_c(\mathbb{R}^n)$ and prove Theorem 1.2.

2 Right-invariant $W^{s,p}$ -norms on diffeomorphism groups

Let \mathcal{N} be a finite dimensional manifold. We are interested in the connected component of the identity of the group $\text{Diff}_c(\mathcal{N})$ of all compactly supported, smooth diffeomorphisms on \mathcal{N} , where \mathcal{N} is either a closed manifold, the Euclidean space \mathbb{R}^n or the n -dimensional ball $B_r(\mathbb{R}^n)$ of radius r in \mathbb{R}^n . In the following, by a slight abuse of notation, we will denote the connected component of the identity by $\text{Diff}_c(\mathcal{N})$ as well.

For $\mathcal{N} = \mathcal{M}$ a closed manifold the requirement of a compact support is redundant, and we will simply write $\text{Diff}(\mathcal{M})$. The following classical result, see e.g., [4], summarizes the group and manifold structure of this infinite dimensional space:

Theorem 2.1 *The space of smooth, compactly supported diffeomorphisms $\text{Diff}_c(\mathcal{N})$ is a simple, Fréchet Lie-group whose Lie-algebra is the set of compactly supported vector fields $\mathcal{X}_c(\mathcal{N}) = \mathcal{C}_c^\infty(\mathcal{N}, T\mathcal{N})$.*

When \mathcal{N} is compact, then $\text{Diff}_c(\mathcal{N}) = \text{Diff}(\mathcal{N})$ is a Fréchet manifold and hence a Fréchet Lie-group [44]; in the non-compact case, this is no longer true but $\text{Diff}_c(\mathcal{N})$ can be modeled on an (LF)-space or a convenient vector space [41, Sect. 43]. This subtlety will not be relevant for the subsequent analysis.

In Sect. 3 we will be in addition interested in the homogeneous space of all smooth diffeomorphisms of the circle S^1 modulo translations, which we will identify with the set of

all diffeomorphisms that fix the point $0 \in S^1$, i.e.,

$$\text{Diff}(S^1)/S^1 \sim \{ \varphi \in \text{Diff}(S^1) : \varphi(0) = 0 \},$$

where we identified the circle S^1 with the interval $[0, 1]$.

2.1 Fractional order Sobolev spaces on \mathbb{R}^n

To introduce the class of right-invariant $W^{s,p}$ -norms on the diffeomorphism group we will start by introducing the fractional order Sobolev spaces $W^{s,p}(\mathbb{R}^n)$ for real valued functions on \mathbb{R}^n . There are several constructions of fractional order Sobolev spaces, which typically coincide in the important Hilbert case $p = 2$. Here we will use the Gagliardo-seminorm (also known as Slobodeckij seminorm) approach, resulting in the so called Sobolev–Slobodeckij spaces.

Let $p \in (1, \infty)$. For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^d$ and $s = k + \sigma$ with $k \in \mathbb{N}$ and $\sigma \in (0, 1)$ we define the homogeneous $\dot{W}^{s,p}$ -norm using the Gagliardo-seminorm via

$$\|f\|_{\dot{W}^{s,p}} = \left(\iint_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|D^k f(x) - D^k f(y)|^p}{|x - y|^{n+\sigma p}} dx dy \right)^{1/p}, \tag{2.1}$$

where $D^k f$ denotes the k -th differential of f . We extend this definition to the full $W^{s,p}$ -norm by adding the L^p -norm of the function, i.e.,

$$\|f\|_{W^{s,p}} = \|f\|_{W^{k,p}} + \|f\|_{\dot{W}^{s,p}}. \tag{2.2}$$

The fractional order Sobolev spaces $W^{s,p}(\mathbb{R}^n)$, as defined above, satisfy the Sobolev embedding theorem, i.e., $W^{s,p}(\mathbb{R}^n)$ embeds in $C^0(\mathbb{R}^n)$ iff $sp > n$ (see, e.g., [13, Theorem 3.7]). The following lemma deals with exact estimates for embeddings in L^q -spaces for the critical case $sp = n$:

Lemma 2.2 *Let $p \in (1, \infty)$. Then there exists $C = C(p, n)$ such that for every $f \in W^{n/p,p}(\mathbb{R}^n)$,*

$$\|f\|_{L^q(\mathbb{R}^n)} \leq C \|f\|_{W^{n/p,p}(\mathbb{R}^n)} q^{1-\frac{1}{p}}, \quad \forall q \in [p, \infty).$$

Proof For $s = n/p$ being an integer see [45, Theorem 12.33]. When s is not an integer, then the Gagliardo (Sobolev–Slobodeckij) spaces we consider here are equivalent to the Besov spaces $B^s_{p,p}(\mathbb{R}^n)$, i.e., the interpolation space $(L^p(\mathbb{R}^n), W^{N,p}(\mathbb{R}^n))_{\frac{s}{N},p}$ for $N > s$ (see e.g., [58, Section 2.5.1, Remark 4], or [13, Theorem 3.1]). The result for $(L^p(\mathbb{R}^n), W^{N,p}(\mathbb{R}^n))_{\frac{s}{N},p}$ is the content of [52, Theorem 9.1], in which a more general statement is shown and where the case treated here corresponds to $p = r$ in the notation of [52]. \square

Finally we state the behavior of the $W^{s,p}$ -norm with respect to scalings:

Lemma 2.3 *Let $p \in (1, \infty)$ and let $\lambda \in \mathbb{R}_{>0}$. For any $f \in W^{s,p}(\mathbb{R}^n)$ let f^λ denote the function*

$$f^\lambda(x) = \frac{1}{\lambda} f(\lambda x).$$

We then have

$$\|f^\lambda\|_{\dot{W}^{s,p}} = \lambda^{(s-1)-\frac{n}{p}} \|f\|_{\dot{W}^{s,p}}, \quad \|f^\lambda\|_{L^p} = \lambda^{-1-\frac{n}{p}} \|f\|_{L^p}.$$

Proof This follows immediately by the chain rule and changing variables in the standard Sobolev norm or the Gagliardo seminorm, depending on s . \square

2.2 Fractional order Sobolev norms on Riemannian manifolds

We now introduce the corresponding space of real valued functions $W^{s,p}(\mathcal{M})$ for \mathcal{M} a (compact) Riemannian manifold. Following [59, Sect. 7.2.1] let $B_\epsilon(x)$ denote the ball of radius ϵ with center x . We can then choose a finite cover of \mathcal{M} by balls $B_\epsilon(x_\alpha)$ with ϵ sufficiently small, such that normal coordinates are defined in the ball $B_\epsilon(x)$, and a partition of unity ρ_α , subordinated to this cover. Using this data we define the $W^{s,p}$ -norm of a function f on M via

$$\|f\|_{W^{s,p}(M,g)}^2 = \sum_\alpha \|(\rho_\alpha f) \circ \exp_{x_\alpha}\|_{W^{s,p}(\mathbb{R}^n)}^2$$

Changing the cover or the partition of unity leads to equivalent norms, see [59, Theorem 7.2.3] and thus this choice does not matter to us, as we are mainly interested in boundedness (unboundedness, resp.) of the diameter, a property which remains invariant under equivalent norms. For integer s and $p = 2$ we get norms which are equivalent to the Sobolev norms treated in [22, Chapter 2]. The norms depend on the choice of the Riemannian metric g , though again, different choices of metrics result in equivalent norms and thus are immaterial to this paper. This dependence is worked out in detail in [22].

2.3 Right-invariant fractional order Sobolev metrics on diffeomorphism groups

For vector fields we use the trivialization of the tangent bundle that is induced by the coordinate charts and define the norm in each coordinate as above. This leads to a well-defined $W^{s,p}$ -norm (up to the equivalence discussed above) on the Lie algebra $\mathcal{X}_c(\mathcal{M})$ of (compactly supported) vector fields on \mathcal{M} . This norm can be extended in the usual way to a right-invariant Finsler metric on the whole diffeomorphism group, i.e., for $\varphi \in \text{Diff}(\mathcal{M})$ and $h \in T_\varphi \text{Diff}(\mathcal{M})$,

$$F_\varphi^{s,p}(h) := \|h \circ \varphi^{-1}\|_{W^{s,p}}, \tag{2.3}$$

where in the right-hand side the norm is the $W^{s,p}$ norm on $\mathcal{X}(\mathcal{M})$. In the important case $p = 2$ this norm is equivalent to the standard H^s norm that is induced from the inner product $\langle \cdot, \cdot \rangle_{H^s}$, and therefore we obtain a right-invariant Riemannian metric

$$G_\varphi^s(h, k) := \langle h \circ \varphi^{-1}, k \circ \varphi^{-1} \rangle_{H^s}. \tag{2.4}$$

This is mentioned here for the sake of completeness—the Riemannian structure will not play a special role in this paper.

Equipping the diffeomorphism group with a Finsler metric gives rise to the corresponding geodesic distance, which is defined in the usual way via

$$\text{dist}_{s,p}(\varphi_0, \varphi_1) := \inf_\varphi \int_0^1 F_\varphi^{s,p}(\partial_t \varphi) dt,$$

where the infimum is taken over all paths $\varphi : [0, 1] \rightarrow \text{Diff}_c(\mathcal{M})$ with $\varphi(0) = \varphi_0$ and $\varphi(1) = \varphi_1$. Using this we can define the diameter of the diffeomorphism group with respect to the metric $F^{s,p}$ to be

$$\text{diam}_{s,p} \text{Diff}(\mathcal{M}) := \sup_{\varphi_0, \varphi_1 \in \text{Diff}(\mathcal{M})} \text{dist}_{s,p}(\varphi_0, \varphi_1) = \sup_{\varphi \in \text{Diff}(\mathcal{M})} \text{dist}_{s,p}(\text{id}, \varphi). \tag{2.5}$$

Here the second equality is due to the right invariance of the Finsler metric and thus of the geodesic distance function. Note, that all these definitions remain valid if the compact

manifold \mathcal{M} is replaced by the non-compact space \mathbb{R}^n or a connected open subset thereof (with $\text{Diff}_c(\mathbb{R}^n)$ and $\mathcal{X}_c(\mathbb{R}^n)$ instead of $\text{Diff}(\mathcal{M})$ and $\mathcal{X}(\mathcal{M})$).

The study of the geodesic distance—and thus of the diameter—is closely related to the study of the displacement energy [23], which in our context is defined as follows:

Definition 2.4 Given a manifold \mathcal{M} , the displacement energy of a set $A \subset \mathcal{M}$ with respect to the $W^{s,p}$ -metric is

$$E_{s,p}(A) = \inf \{ \text{dist}_{s,p}(\text{Id}, \varphi) : \varphi \in \text{Diff}(\mathcal{M}), \varphi(A) \cap A = \emptyset \}.$$

In fact it turns out that the geodesic distance collapses if and only if there exists an open set of zero displacement energy (see, e.g., [9,23,32,54]). This provides an important tool for studying vanishing distance phenomena. One could hope for a similar relation between bounded displacement energy and finite diameter, which is the geometric property that we aim to study in this article. A result of this type, in general settings, appears in “Appendix C”, however its assumptions are too restrictive for our applications to diffeomorphism groups (see Lemma C.2). Nevertheless, we do analyze the displacement energy, as it still provides some insight on the diameter.

3 The diameter of $\text{Diff}(S^1)$

The aim of this section is to prove the following complete characterization of boundedness (unboundedness, resp.) of the diameter of the diffeomorphism group of the circle S^1 with respect to right-invariant $W^{s,p}$ -norms:

Theorem 3.1 *Let $p \in (1, \infty)$. The diameter $\text{diam}_{s,p} \text{Diff}(S^1)$ of the diffeomorphism group of S^1 is zero for $s \leq 1/p$, bounded (but non-zero) for $1/p < s < 1 + 1/p$, and unbounded for $s \geq 1 + 1/p$.*

Note, that for the important special case $p = 2$ this shows that the diameter $\text{diam}_{s,2} \text{Diff}(S^1)$ of the H^s -metric is zero for $s \leq \frac{1}{2}$, bounded (but non-zero) for $s < 3/2$, and unbounded for $s \geq \frac{3}{2}$. For the case $p = 1$, the only change is that the diameter is finite but non-zero for $s = 1$ (due to the fact that $W^{1,1}$ embeds in C^0 in the one dimensional case).

Proof The zero diameter result for $s \leq \frac{1}{p}$ ($s < 1$ for $p = 1$) follows directly from the results on vanishing geodesic distance in [9,32,33]. We will split the proof of the remaining cases in three parts: the subcritical case $s < 1 + 1/p$, see Sect. 3.3, the critical case $s = 1 + 1/p$, see Sect. 3.2, and the supercritical case $s > 1 + 1/p$, see Sect. 3.1. \square

In the proof of the critical case in Sect. 3.2 we will show in addition that the diameter of the homogeneous $W^{1,p}$ -metric is bounded between p and $8p$. Before we present this analysis we want to point out an open question concerning the continuity of the diameter (in the parameter s).

Question 3.2 *Is the diameter continuous in the Sobolev index s at the critical exponents, i.e., do we have $\lim_{s \rightarrow 1/p+} \text{diam}_{s,p} \text{Diff}(S^1) = 0$ and $\lim_{s \rightarrow (1+1/p)-} \text{diam}_{s,p} \text{Diff}(S^1) = \infty$?*

3.1 The supercritical case $s > 1 + 1/p$

The unboundedness for the supercritical case $s \geq 1 + 1/p$ also follows directly from the analysis for the critical case, which is treated in Sect. 3.2. In the following we will present a

more elementary proof that in addition contains an explicit bound for the geodesic distance and will be of importance in the characterization of the displacement energy in Sect. 3.4.

Lemma 3.3 *Let $s > 1 + 1/p$. Then the geodesic distance of the right-invariant $W^{s,p}$ -norm on $\text{Diff}(S^1)$ satisfies*

$$\log \varphi'(x) \leq C \text{dist}_{s,p}(\text{Id}, \varphi), \tag{3.1}$$

where $C = C(s, p)$ depends on s and p . It follows that $\text{diam}_{s,p} \text{Diff}(S^1)$ is unbounded.

Proof Note that in this regime we have the Sobolev embedding $W^{s,p}(S^1) \subset C^1(S^1)$. Let φ_t be any curve starting at Id and ending at φ , and let u_t be the associated vector field, that is $\partial_t \varphi_t = u_t \circ \varphi_t$. Denote $\psi_t = \partial_x \varphi_t$. We then have $\partial_t \psi_t = \partial_x u_t \circ \varphi_t \cdot \psi_t$, or in other words, $\partial_t (\log \psi_t)(x) = \partial_x u_t(\varphi_t(x))$. Integrating this, and using the fact that $\log \psi_0 = 0$, we have for any $(s - 1)p > 1$ and any $x \in S^1$,

$$\log \varphi'(x) = \int_0^1 \partial_t (\log \psi_t(x)) dt \leq \int_0^1 \|\partial_x u_t\|_{L^\infty} dt \leq C \int_0^1 \|u_t\|_{W^{s,p}} dt, \tag{3.2}$$

where in the last inequality we used the above-mentioned Sobolev embedding. Since the above inequality holds for all paths φ_t connecting the identity to φ this yields Eq. (3.1). By choosing φ with φ' arbitrarily large at a point, we get an arbitrarily large lower bound to the diameter of $\text{Diff}(S^1)$ and thus the unboundedness follows. \square

Remark 3.4 For $p = 1$, the claim and its proof holds also for $s = 2$, since $W^{2,1}(S^1)$ embeds into $C^1(S^1)$.

3.2 The critical case $s = 1 + 1/p$

In the critical case, we do not have $W^{s,p}(S^1) \subset C^1(S^1)$ as before (unless $p = 1$), however we do have $W^{s,p}(S^1) \subset W^{1,q}(S^1)$ for $q < \infty$. Inspired by this, the proof for the critical case consists of two steps—first, we give a lower bound for the diameter with respect to the $\dot{W}^{1,q}$ -metric, which blows up as $q \rightarrow \infty$; second, we use Lemma 2.2 to show that the diameter bound blows up faster than the embedding constants of $W^{s,p}(S^1) \subset W^{1,q}(S^1)$, hence the diameter with respect to $W^{s,p}$ is infinite.

The following lower bound for the diameter with respect to the $\dot{W}^{1,q}$ -metric is based on a generalization of a result of Lenells [42], where he constructed an explicit solution formula for geodesics of the homogeneous $\dot{W}^{1,2}$ -metric. In the following lemma we will extend his construction to all homogeneous $\dot{W}^{1,q}$ -norms with $q \geq 1$:

Lemma 3.5 *Let*

$$\Phi : \text{Diff}(S^1)/S^1 \rightarrow C^\infty(S^1, \mathbb{R}), \quad \varphi \mapsto q(\varphi')^{1/q}. \tag{3.3}$$

We have:

1. *The mapping Φ is an isometric embedding, where $\text{Diff}(S^1)/S^1$ is equipped with the right-invariant homogeneous $\dot{W}^{1,q}$ -norm and $C^\infty(S^1, \mathbb{R})$ with the standard (i.e., non-invariant) L^q -norm.*
2. *The image $\Phi(\text{Diff}(S^1)) \subset C^\infty(S^1, \mathbb{R})$ is an open subset of the L^q -sphere of radius q given by*

$$\mathcal{U}_q := \{f \in C^\infty(S^1; \mathbb{R}) : f > 0, \|f\|_{L^q} = q\}.$$

3. For fixed $q \geq 1$ the diameter of the set \mathcal{U}_q is bounded from above and below by

$$q < \text{diam } \mathcal{U}_q \leq 8q. \tag{3.4}$$

As a consequence \mathcal{U}_q is unbounded for $q \rightarrow \infty$.

Proof The flat L^q metric on $C^\infty(S^1, \mathbb{R}_{>0})$ is given by

$$|\delta f|_f = \left(\int_0^1 |\delta f(\theta)|^q d\theta \right)^{1/q}, \quad \delta f \in T_f C^\infty(S^1, \mathbb{R}_{>0}) \cong C^\infty(S^1). \tag{3.5}$$

To see that the mapping Φ is an isometric embedding (where $\text{Diff}(S^1)/S^1$ is equipped with the right-invariant $\dot{W}^{1,q}$ -metric) we need to calculate the derivative of Φ . We have:

$$d\Phi(\varphi) \cdot h = (\varphi')^{1/q-1} h' \tag{3.6}$$

and thus

$$|d\Phi(\varphi) \cdot h| = \left(\int_0^1 (\varphi'(\theta))^{1-q} |h'(\theta)|^q d\theta \right)^{1/q} \tag{3.7}$$

which equals exactly the right-invariant, homogeneous $\dot{W}^{1,q}$ -metric. The characterization of the image of Φ follows directly from the definition of $\text{Diff}(S^1)/S^1$.

To calculate the lower bound for the diameter of \mathcal{U}_q we consider the functions $f = q$ and $g = cq(n\mathbb{1}_{(0,n-q)} + \varepsilon\mathbb{1}_{(n-q,1)})$ for $\varepsilon \ll 1$ and $c \approx 1$ such that $\|g\|_{L^q} = q$. Then

$$\text{diam } \mathcal{U}_q \geq \text{dist}_{\mathcal{U}_q}(f, g) \geq \|f - g\|_{L^q} \approx 2^{1/q}q,$$

where $\text{diam } \mathcal{U}_q$ and $\text{dist}_{\mathcal{U}_q}$ refer to the intrinsic distance in \mathcal{U}_q .

It remains to show that $\text{diam } \mathcal{U}_q$ is bounded from above for each fixed q . Towards this aim we will construct paths that connect given elements $f, g \in \mathcal{U}_q$ and are bounded independently of f and g . Let

$$f_t(\theta) = \frac{q}{\|\tilde{f}_t(\cdot)\|_{L^q}} \tilde{f}_t(\theta) \quad \text{with} \quad \tilde{f}_t(\theta) = (1-t)f(\theta) + tg(\theta). \tag{3.8}$$

It is easy to see that $f_t(\cdot) \in \mathcal{U}_q$ for any $t \in [0, 1]$. It remains to bound the L^q -length of $f_t(\theta)$. We have

$$\partial_t f_t(\theta) = \frac{q}{\|\tilde{f}_t(\cdot)\|_{L^q}} \partial_t \tilde{f}_t(\theta) - \frac{q}{\|\tilde{f}_t(\cdot)\|_{L^q}^2} \partial_t \|\tilde{f}_t(\cdot)\|_{L^q} \tilde{f}_t(\theta), \tag{3.9}$$

and thus

$$\text{dist}_{\mathcal{U}_q}(f, g) \leq \int_0^1 \|\partial_t f_t(\cdot)\|_{L^q} dt \leq q \left(\int_0^1 \frac{\|\partial_t \tilde{f}_t(\cdot)\|_{L^q}}{\|\tilde{f}_t(\cdot)\|_{L^q}} dt + \int_0^1 \frac{|\partial_t \|\tilde{f}_t(\cdot)\|_{L^q}|}{\|\tilde{f}_t(\cdot)\|_{L^q}} dt \right) \tag{3.10}$$

We will estimate the two integrals separately. For the first one we calculate

$$\|\partial_t \tilde{f}_t(\cdot)\|_{L^q} = \|g - f\|_{L^q} \leq \|g\|_{L^q} + \|f\|_{L^q} \leq 2q \tag{3.11}$$

and

$$\begin{aligned} \|\tilde{f}_t(\cdot)\|_{L^q}^q &= \int ((1-t)f(\theta) + tg(\theta))^q d\theta \\ &\geq \int (1-t)^q f(\theta)^q d\theta + \int t^q g(\theta)^q d\theta \\ &= (1-t)^q \|f\|_{L^q}^q + t^q \|g\|_{L^q}^q \geq \frac{2}{2^q} q^q. \end{aligned} \tag{3.12}$$

Thus the first term can be estimated by 4. In these estimates we made repeatedly use of the fact that all involved functions are positive. For the second term we calculate using the Hölder inequality

$$\begin{aligned} \partial_t \|\tilde{f}_t(\cdot)\|_{L^q} &= \partial_t \left(\int \tilde{f}_t(\theta)^q d\theta \right)^{1/q} = \left(\int \tilde{f}_t(\theta)^q d\theta \right)^{1/q-1} \int \tilde{f}_t(\theta)^{q-1} (g-f) d\theta \\ &\leq \left(\int \tilde{f}_t(\theta)^q d\theta \right)^{1/q-1} \left(\int \tilde{f}_t(\theta)^q d\theta \right)^{(q-1)/q} \left(\int (g-f)^q d\theta \right)^{1/q} \\ &= \left(\int (g-f)^q d\theta \right)^{1/q} = \|g-f\|_{L^q} \leq 2q, \end{aligned} \tag{3.13}$$

and thus the second term is bounded as well by 4. This, in turn, proves the desired bound for the diameter of \mathcal{U}_p . □

Note that the upper bound above implies directly the boundedness of the diameter of $\text{Diff}(S^1)/S^1$ with respect to the homogeneous $\dot{W}^{1,q}$ metric. We now use the lower bound, together with the Sobolev embedding theorem of Lemma 2.2 to show the unboundedness of $\text{diam}_{s,p} \text{Diff}(S^1)$ in the critical case:

Lemma 3.6 *Let $p \in (1, \infty)$. Then the diameter $\text{diam}_{1+1/p,p} \text{Diff}(S^1)$ of $\text{Diff}(S^1)$ with respect to the right-invariant $W^{1+1/p,p}$ -norm is unbounded.*

Proof In Lemma 3.5 we have shown that $\text{diam}_{1,q} \text{Diff}(S^1)/S^1 > q$, and thus we also have $\text{diam}_{1,q} \text{Diff}(S^1) > q$. In particular we have shown that for any $q \geq 1$ there is some $\varphi^q \in \text{Diff}(S^1)$ such that $\text{dist}_{1,q}(\text{Id}, \varphi^q) > q$. Therefore, using Lemma 2.2, we have

$$\begin{aligned} q < \text{dist}_{1,q}(\text{Id}, \varphi^q) &= \inf \int_0^1 \|\varphi_t^q \circ (\varphi_t^q)^{-1}\|_{W^{1,q}} dt \\ &< \inf Cq^{1-\frac{1}{p}} \int_0^1 \|\varphi_t^q \circ (\varphi_t^q)^{-1}\|_{W^{1+1/p,p}} dt \\ &= Cq^{1-\frac{1}{p}} \text{dist}_{1+1/p,p}(\text{Id}, \varphi^q). \end{aligned} \tag{3.14}$$

Thus $\text{dist}_{1+1/p,p}(\text{Id}, \varphi^q) \geq Cq^{1/p}$ and taking $q \rightarrow \infty$ completes the proof. □

Remark 3.7 It might be possible to use a similar argument to prove the second part of Question 3.2, i.e., that $\lim_{s \rightarrow (1+1/p)-} \text{diam}_{s,p} \text{Diff}(S^1) = \infty$, by controlling the embedding constants of $W^{s,p}$ into $W^{1,q}$ for $s \nearrow 1 + 1/p$ and an appropriate $q(s) \rightarrow \infty$.

3.3 The subcritical case $s < 1 + 1/p$

It remains to show the boundedness of the diameter for $s < 1 + 1/p$. Towards this aim we will first show that if (controlled) arbitrary change in length (volume) has a bounded cost, then the diameter is finite:

Lemma 3.8 *Identify S^1 with the unit interval, and let $(s - 1)p < 1$. For any $\lambda \in \mathbb{N}$ and $\delta \in (0, 1)$, denote by $\psi_{\lambda,\delta} \in \text{Diff}(S^1)$ a map satisfying $\psi_{\lambda,\delta}(x) = \lambda x$ for $x \in [0, \frac{1-\delta}{\lambda}]$. If there exists $C = C(s, p) > 0$, independent of λ and δ , such that*

$$\text{dist}_{W^{s,p}([0,1])}(\text{Id}, \psi_{\lambda,\delta}) < C \text{ for every } \lambda \in \mathbb{N} \text{ and } \delta \in (0, 1),$$

then

$$\text{diam}_{s,p} \text{Diff}(S^1) \leq \text{diam}_{s,p} \text{Diff}([0, 1]) + 1 < 4C + 1.$$

Proof Let $\varphi \in \text{Diff}(S^1)$; by translating, we can assume that $\varphi(0) = 0$. This translation costs at most 1, hence the “+1” in the statement of the theorem. We can always write $\varphi = \varphi_1 \circ \varphi_2$, where $\text{supp } \varphi_1 \subset [0, 1 - \delta]$, and $\text{supp } \varphi_2 \subset [\delta, 1]$ for some $\delta > 0$. Since $\text{dist}(\text{Id}, \varphi) \leq \text{dist}(\text{Id}, \varphi_1) + \text{dist}(\text{Id}, \varphi_2)$, it is enough to prove that both $\text{dist}(\text{Id}, \varphi_1)$ and $\text{dist}(\text{Id}, \varphi_2)$ are smaller than $2C$. Note that for $\varphi \in \text{Diff}(S^1)$ with $\varphi(0) = 0$ we have

$$\text{dist}_{W^{s,p}(S^1)}(\text{Id}, \varphi) \leq \text{dist}_{W^{s,p}([0,1])}(\text{Id}, \varphi),$$

hence it is enough to prove the statement for $W^{s,p}([0, 1])$. Henceforth in this proof, we will only refer to $[0, 1]$.

Following the decomposition above, from now on we will assume that $\text{supp } \varphi \subset [0, 1 - \delta]$ for some $\delta > 0$, and consider φ as a diffeomorphisms of \mathbb{R} . Denote

$$\varphi^\lambda(x) = \frac{1}{\lambda} \varphi(\lambda x) = \psi_{\lambda,\delta}^{-1} \circ \varphi \circ \psi_{\lambda,\delta}(x),$$

where the last equality holds because $\text{supp } \varphi \subset [0, 1 - \delta]$. Using our assumption, we have

$$\text{dist}_{s,p}(\text{Id}, \varphi) = \text{dist}_{s,p}(\text{Id}, \psi_{\lambda,\delta} \circ \varphi^\lambda \circ \psi_{\lambda,\delta}^{-1}) < 2C + \text{dist}_{s,p}(\text{Id}, \varphi^\lambda).$$

We now show that $\text{dist}_{s,p}(\text{Id}, \varphi^\lambda)$ can be controlled by $\text{dist}_{s,p}(\text{Id}, \varphi)$ times a small constant. A direct calculation shows that the map $\varphi(t, x) \mapsto \varphi^\lambda(t, x) = \lambda^{-1} \varphi(t, \lambda x)$ is a bijection between paths supported on $[0, 1]$ to paths supported on $[0, 1/\lambda]$, with the corresponding vector fields

$$u_t^\lambda(x) = \frac{1}{\lambda} u_t(\lambda x).$$

Note that

$$\|u_t^\lambda\|_{\dot{W}^{s,p}} = \lambda^{(s-1)-\frac{1}{p}} \|u_t\|_{\dot{W}^{s,p}}, \quad \|u_t^\lambda\|_{\dot{W}^{1,p}} = \lambda^{-\frac{1}{p}} \|u_t\|_{\dot{W}^{1,p}}, \quad \|u_t^\lambda\|_{L^p} = \lambda^{-1-\frac{1}{p}} \|u_t\|_{L^p}$$

where $\dot{W}^{s,p}$ refers to the $(s - 1)$ -Gagliardo seminorm on the derivative (if $s > 1$), c.f., Lemma 2.3. We therefore have

$$\|u_t^\lambda\|_{W^{s,p}} \leq \lambda^{(s-1)-\frac{1}{p}} \|u_t\|_{W^{s,p}},$$

and hence

$$\text{length}_{W^{s,p}}(\varphi_t^\lambda) \leq \lambda^{(s-1)-\frac{1}{p}} \text{length}_{W^{s,p}}(\varphi_t).$$

Therefore, taking the infimum over all possible paths between Id and φ , we have

$$\text{dist}_{s,p}(\text{Id}, \varphi^\lambda) \leq \lambda^{(s-1)-\frac{1}{p}} \text{dist}_{s,p}(\text{Id}, \varphi).$$

We conclude that

$$\text{dist}_{s,p}(\text{Id}, \varphi) < \frac{2C}{1 - \lambda^{(s-1)-\frac{1}{p}}}.$$

Since $(s - 1)p < 1$, taking $\lambda \rightarrow \infty$ concludes the proof. □

The following lemma shows that $\text{dist}_{W^{s,p}([0,1])}(\text{Id}, \psi_{\lambda,\delta})$ is indeed uniformly bounded, by showing that the affine homotopy is uniformly bounded (in λ and δ) in the subcritical regime. As the proof is a rather technical calculation, we postpone it to Appendix A.

Lemma 3.9 *Let $s < 1 + 1/p$. Then there exists a sequence of maps $\psi_{\lambda,\delta}$ with $\psi_{\lambda,\delta}(x) = \lambda x$ for $x \in [0, \frac{1-\delta}{\lambda}]$ such that*

$$\text{dist}_{W^{s,p}([0,1])}(\text{Id}, \psi_{\lambda,\delta}) < C \text{ for every } \lambda \in \mathbb{N} \text{ and } \delta \in (0, 1)$$

where $C = C(s, p)$ is independent of λ .

3.4 The displacement energy

Finally, we discuss boundedness properties of the displacement energy as introduced in Definition 2.4. Even though we do not have sufficiently strong result relating the boundedness of the diameter and the displacement energy (see ‘‘Appendix C’’, we will now show that indeed in our case boundedness (unboundedness resp.) of the displacement energy of arbitrarily large open subsets of S^1 is closely related to the boundedness of the diameter. While it is obvious that bounded diameter implies bounded displacement energy, we give below a direct, simpler proof for the boundedness of the displacement energy in the subcritical case.

Proposition 3.10 *Identify S^1 with the interval $[0, 1]$. We then have the following bounds*

1. For an interval I with length smaller than $1/2$, we have

$$E_{s,p}(I) \leq 1/2 \text{ for all } s, p.$$

2. For every $s < 1 + 1/p$, there exists $C = C(s, p) > 0$ such that

$$E_{s,p}(I) < C \text{ for every open interval } I \subset [0, 1].$$

3. If $s > 1 + 1/p$ then there exists $c = c(s, p) > 0$ such that

$$E_{s,p}((0, 1 - \delta)) > c|\log \delta|.$$

Remark 3.11 Note that we do not know whether the displacement energy is bounded or not in the critical case $s = 1 + 1/p$.

Proof The first assertion follows by flowing for time 1 along the constant vector field $u(t, x) = 1/2$, whose $W^{s,p}$ -norm is $\frac{1}{2}$.

The last assertion follows from (3.2). Indeed, if $\varphi \in \text{Diff}(S^1)$ such that $\varphi((0, 1 - \delta)) \cap (0, 1 - \delta) = \emptyset$, then $\varphi((1 - \delta, 1)) \supset (0, 1 - \delta)$, and therefore

$$\delta \max(\varphi') \geq \int_{1-\delta}^1 \varphi'(x) dx = \varphi(1) - \varphi(1 - \delta) \geq (1 - \delta) - 0 = 1 - \delta.$$

Therefore $\max(\varphi') > \frac{1-\delta}{\delta}$, and so by (3.2), $\text{dist}_{s,p}(\text{Id}, \varphi) \geq c|\log \delta|$.

The second assertion follows directly from Theorem 3.1. In the following we will sketch an alternative simpler proof. Consider the vector field

$$u_{\alpha,\varepsilon}(x) = \begin{cases} \varepsilon^{-\alpha} x & x \in [0, \varepsilon] \\ x^{1-\alpha} & x \in [\varepsilon, 3/4] \\ 4\left(\frac{3}{4}\right)^{1-\alpha} (1-x) & x \in [3/4, 1), \end{cases} \tag{3.15}$$

where $\alpha < 1 + \frac{1}{p} - s$ and $\varepsilon \ll 1$ to be determined (note that this vector field is simply $x^{1-\alpha}$ with linear interpolations to 0 at $0 \sim 1$). A direct calculation shows that $\|u_{\alpha,\varepsilon}\|_{s,p} < C(s, p, \alpha) < \infty$ when $(\alpha + (s - 1))p < 1$. Furthermore, we have that the flow φ_t along $u_{\alpha,\varepsilon}$ (that is, the solution to $\partial_t \varphi_t = u_{\alpha,\varepsilon} \circ \varphi_t$, $\varphi_0(x) = x$) satisfies, for $x \geq \varepsilon$, and as long as $\varphi_t(x) < 3/4$,

$$\varphi_t(x) = (x^\alpha + \alpha t)^{1/\alpha} > (\alpha t)^{1/\alpha},$$

hence in particular, for $t_0 = \frac{1}{\alpha 2^\alpha}$, $\varphi^{\alpha,\varepsilon} = \varphi_{t_0}$ satisfies

$$\varphi^{\alpha,\varepsilon}(0) = 0, \quad \varphi^{\alpha,\varepsilon}(1) = 1, \quad \varphi^{\alpha,\varepsilon}(x) > 1/2 \quad \text{for any } x > \varepsilon$$

and

$$\text{dist}_{s,p}(\text{Id}, \varphi^{\alpha,\varepsilon}) < \frac{1}{\alpha 2^\alpha} C(s, p, \alpha).$$

Consider now the interval $I = (\delta, 1)$ for some $\delta < 1/2$, and let $\varepsilon < \delta$. Then

$$\psi = (\varphi^{\alpha,\varepsilon})^{-1} \circ T_{1/2} \circ \varphi^{\alpha,\varepsilon}$$

where $T_{1/2}$ is the translation by $1/2$, satisfies

$$\psi(I) \subset (\varphi^{\alpha,\varepsilon})^{-1} \circ T_{1/2}((1/2, 1)) = (\varphi^{\alpha,\varepsilon})^{-1}(0, 1/2) \subset (0, \delta),$$

hence $\psi(I) \cap I = \emptyset$. Since

$$\text{dist}_{s,p}(\text{Id}, \psi) \leq 2 \text{dist}_{s,p}(\text{Id}, \varphi^{\alpha,\varepsilon}) + \text{dist}_{s,p}(\text{Id}, T_{1/2}) \leq \frac{2}{\alpha 2^\alpha} C(s, p, \alpha) + \frac{1}{2}$$

and the right-hand side is uniformly bounded in δ , the proof is complete. □

4 The diameter of $\text{Diff}(M)$ for compact manifolds in higher dimensions

In this section we prove Theorem 1.1 in its full generality. The main analytic ideas of the proof are similar to the one-dimensional case, however their adaptation to the higher dimensional settings is not always immediate; in particular, for the boundedness proof, we need the diffeomorphism group to have a localization property, which we call *uniform fragmentation property*, described in Sect. 4.3.

The structure of the section is as follows: in Sect. 4.1 we give a simple proof for the unboundedness of the diameter of $\text{Diff}(\mathcal{M})$ (for any manifold) when $s > 1 + \frac{\dim \mathcal{M}}{p}$, and in Sect. 4.2 we give a more elaborate proof for the unboundedness in the case $s \geq 1 + \frac{\dim \mathcal{M}}{p}$. We then present and discuss the uniform fragmentation property in Sect. 4.3, and prove boundedness of $\text{Diff}(\mathcal{M})$ for manifolds that satisfy this property (like spheres), when $s < 1 + \frac{1}{p}$. This completes the proof of Theorem 1.1, as the zero diameter result of item 1 follows directly from the results on vanishing geodesic distance in [5,32,33].

As noted in the introduction, when $\dim \mathcal{M} > 1$, we have a gap in the range $s \in \left[1 + \frac{1}{p}, 1 + \frac{\dim \mathcal{M}}{p}\right)$. We believe that in these cases the diameter is finite (assuming that the fragmentation property is satisfied); we show an indication for this in Sect. 4.4.

4.1 The supercritical case $s > 1 + \frac{\dim \mathcal{M}}{p}$

Lemma 4.1 *Let $s > 1 + \frac{\dim \mathcal{M}}{p}$. Then the geodesic distance of the right-invariant $W^{s,p}$ -norm on $\text{Diff}(\mathcal{M})$ satisfies*

$$\log |D\varphi| \leq C \text{dist}_{s,p}(\text{Id}, \varphi), \tag{4.1}$$

where $C = C(s, p, \dim \mathcal{M}, \mathfrak{g})$, and $|D\varphi|$ is the Jacobian determinant φ with respect to a chosen Riemannian metric \mathfrak{g} on \mathcal{M} . It follows that $\text{diam}_{s,p} \text{Diff}(\mathcal{M})$ is unbounded.

Proof In this regime we have the Sobolev embedding $W^{s,p}(\mathcal{M}) \subset C^1(\mathcal{M})$. Let φ_t be a curve starting at Id and ending at φ , and let u_t its associated vector field, that is $\partial_t \varphi_t = u_t \circ \varphi_t$. Denote $\psi_t = |D\varphi_t|$, the Jacobian determinant with respect to \mathfrak{g} . We have $\partial_t \psi_t = \text{div}(u_t) \circ \varphi \cdot \psi_t$, or in other words, $\partial_t (\log \psi_t)(x) = \text{div}(u_t)(\varphi_t(x))$. Integrating this, using the fact that $\log \psi_0 = 0$, we have for any $(s - 1)p > \dim \mathcal{M}$ and any $x \in \mathcal{M}$,

$$\log |D\varphi|(x) = \int_0^1 \partial_t (\log \psi_t(x)) dt \leq \int_0^1 \|\text{div}(u_t)\|_{L^\infty} dt \leq C \int_0^1 \|u_t\|_{W^{s,p}} dt,$$

where in the last inequality we used the above-mentioned Sobolev embedding. By choosing φ with $|D\varphi|$ arbitrarily large at a point, we get an arbitrarily large lower bound to the diameter. \square

Remark 4.2 As in the one dimensional case, for $p = 1$ this proof also works for the critical case $s = 1 + \dim \mathcal{M}$, as $W^{1+\dim \mathcal{M},1}(\mathcal{M}) \subset C^1(\mathcal{M})$.

4.2 The critical case $s = 1 + \frac{\dim \mathcal{M}}{p}$

Lemma 4.3 *Let $p \geq 1$ and $s = 1 + \frac{\dim \mathcal{M}}{p}$. Then the diameter $\text{diam}_{s,p} \text{Diff}(\mathcal{M})$ of $\text{Diff}(\mathcal{M})$ with respect to the right-invariant $W^{s,p}$ -norm is unbounded.*

Proof The proof of this result is inspired by connections between a homogeneous, degenerate H^1 -metric—called the information metric—on the group of diffeomorphisms and the Fisher–Rao metric on the space of probability densities, see [36]. Similarly as in the proof of the one-dimensional situation we aim to generalize this result to $W^{1,q}$ -metrics for general $q \geq 1$.

We start by introducing a right-invariant, degenerate Sobolev (Finsler) metric of order one on the diffeomorphism group:

$$F_{\text{Id}}(X) = \left(\int_{\mathcal{M}} |\text{div}(X)|^q \mu \right)^{1/q} \tag{4.2}$$

where μ is some fixed volume form on \mathcal{M} and div is the divergence with respect to μ . In the Riemannian case, $q = 2$, this metric is also called information metric due to its connections to the Fisher–Rao metric on the space of probability densities [10,36], which is the central object of interest in the area of information geometry [1].

In particular, we have for $h \in T_\varphi \text{Diff}(\mathcal{M})$

$$(F_{\varphi(h)})^q = \int_{\mathcal{M}} |\text{div}(h \circ \varphi^{-1})|^q \mu = \int_{\mathcal{M}} |\text{div}(h \circ \varphi^{-1}) \circ \varphi|^q \varphi^* \mu. \tag{4.3}$$

Consider the mapping

$$\Phi : \text{Diff}(\mathcal{M}) \rightarrow C^\infty(\mathcal{M}, \mathbb{R}), \quad \varphi \mapsto q|D\varphi|^{1/q}, \tag{4.4}$$

where $|D\varphi|$ the Jacobian determinant of φ with respect to μ (that is, $\varphi^*\mu = |D\varphi|\mu$). Denote the function $\varphi \mapsto |D\varphi|$ by $\tilde{\Phi}$. We have that

$$d\Phi(\varphi) \cdot h = |D\varphi|^{1/q-1} d\tilde{\Phi}(\varphi) \cdot h = |D\varphi|^{1/q} \operatorname{div}(h \circ \varphi^{-1}) \circ \varphi, \tag{4.5}$$

since if $\varphi(t)$ is a curve with $\varphi(0) = \varphi$ and $\dot{\varphi}(0) = h$, then

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} \varphi(t)^* \mu &= \frac{d}{dt} \Big|_{t=0} (\varphi(t) \circ \varphi^{-1} \circ \varphi)^* \mu = \frac{d}{dt} \Big|_{t=0} \varphi^* (\varphi(t) \circ \varphi^{-1})^* \mu \\ &= \varphi^* \left(\frac{d}{dt} \Big|_{t=0} (\varphi(t) \circ \varphi^{-1})^* \mu \right) \\ &= \varphi^* (\mathcal{L}_{h \circ \varphi^{-1}} \mu) = \varphi^* (\operatorname{div}(h \circ \varphi^{-1}) \mu) = \operatorname{div}(h \circ \varphi^{-1}) \circ \varphi \varphi^* \mu \\ &= \operatorname{div}(h \circ \varphi^{-1}) \circ \varphi |D\varphi| \mu. \end{aligned}$$

After equipping $C^\infty(M, \mathbb{R})$ with the flat L^q metric

$$|\delta f|_f = \left(\int_M |\delta f|^q \mu \right)^{1/q}, \tag{4.6}$$

Equations (4.3) and (4.5) imply that the mapping Φ is a Riemannian immersion onto the positive L^q -sphere in $C^\infty(M, \mathbb{R})$ with image

$$\mathcal{U}_q := \{f \in C^\infty(M; \mathbb{R}) : f > 0, \|f\|_{L^q} = q (\operatorname{Vol}_\mu(\mathcal{M}))^{1/q}\}.$$

The proof now continues as in the one-dimensional case (Sect. 3.2): The intrinsic diameter of \mathcal{U}_q with respect to the L^q metric is bounded below by a constant (depending on \mathcal{M}) times q —indeed, by choosing $f \equiv q$ and a function $g \in \mathcal{U}_q$ that is large on a small set and close to zero on the rest of M , we have that $\|f - g\|_{L^q} \geq Cq$. Since the $W^{1,q}$ -norm controls the degenerate metric F , and the map Φ is an immersion, we have that

$$\operatorname{diam}_{1,q} \operatorname{Diff}(\mathcal{M}) \geq Cq,$$

for some constant C independent of q . The infinite diameter with respect to the $W^{1+\frac{\dim \mathcal{M}}{p}, p}$ -norm follows in the same way as in Sect. 3.2, using the embedding $W^{1+\frac{\dim \mathcal{M}}{p}, p} \subset W^{1,q}$ as in Lemma 2.2 and taking $q \rightarrow \infty$. □

4.3 The subcritical case $s < 1 + \frac{1}{p}$

We start by introducing the geometric property of $\operatorname{Diff}(\mathcal{M})$ which we need to prove the boundedness:

Definition 4.4 Let \mathcal{M} be a compact finite dimensional manifold. The diffeomorphism group $\operatorname{Diff}(\mathcal{M})$ is said to satisfy the *uniform fragmentation property*, if there exists a constant $K > 0$ and a finite cover of \mathcal{M} by balls $B_{\epsilon_\alpha}(x_\alpha)$ on which normal coordinates are defined, such that any diffeomorphism $\varphi \in \operatorname{Diff}(M)$ can be written as a product of K diffeomorphisms φ_i , where each φ_i is only supported in one ball $B_{\epsilon_\alpha}(x_\alpha)$.

Remark 4.5 1. If we denote by $K(\varphi)$ the minimal number of diffeomorphisms φ_i needed in such a decomposition of $\varphi \in \operatorname{Diff}(\mathcal{M})$, then $K(\varphi) < \infty$ for any fixed cover of any compact manifold—this is the content of the well-known “fragmentation lemma” (see, e.g., [4, Lemma 2.1.8]). The quantity $K(\varphi)$ is sometimes referred to as the *fragmentation*

norm of φ with respect to this cover (see, e.g., in the context of homeomorphisms, [46, Section 2.3]). We ask for this fragmentation norm to be uniformly bounded, independent of φ .

2. If, instead of a fixed finite cover, we consider a cover by all open (topological) balls, then $K(\varphi)$ is a conjugation-invariant norm, in the sense of [17] (see Example 1.14 there). As such, it is known to be uniformly bounded for many manifolds, see, e.g., [26, Theorem VI] for a recent account on this. Unfortunately, we cannot allow for the balls to be arbitrary, as they are fixed a-priori in the definition of the norm, as discussed in Sect. 2.2 (see also the proof of Proposition 4.7 below).

Next we show that the n -dimensional sphere S^n satisfies the uniform fragmentation property:

Proposition 4.6 *Let $n \geq 1$ and let $A, B \subset S^n$ be open geodesic balls, $A \cup B = S^n$ and $\overline{A}, \overline{B} \neq S^n$. Then $\text{Diff}(S^n)$ has an uniform fragmentation property with respect to the cover $\{A, B\}$.*

Proof Let $\varphi \in \text{Diff}(S^n)$. The proof consists of two steps:

1. First we split $\varphi = \varphi_B \circ \varphi_A$, such that $S^n \setminus \text{supp } \varphi_A$ contains a ball U_A satisfying $\overline{U_A} \subset B^c \subset A$, and similarly to φ_B .
2. We then show that each of φ_A and φ_B can be written as a composition of at most 3 diffeomorphisms, each supported either in A or in B .

Step I Fix $x \in \varphi^{-1}(A^c)$, then there exists an open ball V_B , containing x , such that $\overline{V_B} \subset \varphi^{-1}(A^c)$, and therefore $B^c \cap \varphi(\overline{V_B}) = \emptyset$. We can choose V_B small enough such that $B^c \setminus \overline{V_B} \neq \emptyset$ as well. Then there exists $\varphi_A \in \text{Diff}(S^n)$ such that $\varphi_A|_{V_B} = \varphi|_{V_B}$ and $B^c \setminus \text{supp } \varphi_A \neq \emptyset$, hence there exists a ball U_A such that $\overline{U_A} \subset B^c \setminus \text{supp } \varphi_A$. Setting $\varphi_B = \varphi \circ \varphi_A^{-1}$, we have that $\varphi_B|_{\varphi(\overline{V_B})} = \text{Id}$, hence $\varphi(V_B) \subset A^c \setminus \text{supp } \varphi_B$. Choosing a ball U_B such that $\overline{U_B} \subset \varphi(V_B)$ completes this step.

Step II We prove the result for φ_A ; the case of φ_B is analogous. We decompose $\varphi_A = \varphi_3 \circ \varphi_2 \circ \varphi_1$, where each φ_i is supported in either A or B .

First, we construct φ_1 to make all the points that end in B start in B . Consider the set $\varphi_A^{-1}(B)$. Since $\overline{U_A} \cap \varphi_A^{-1}(B) = \emptyset$, we have

$$A \cap \varphi_A^{-1}(\overline{B}) \subset A \setminus \overline{U_A}. \tag{4.7}$$

Since $A \setminus \overline{U_A}$ is diffeomorphic to $A \cap B$, there exists a diffeomorphism φ_1 with $\varphi_1(A \setminus \overline{U_A}) = A \cap B$. Moreover, we can choose φ_1 such that it is supported on A (see Fig. 1).

We now have that

$$\varphi_1 \circ \varphi_A^{-1}(\overline{B}) \subset B. \tag{4.8}$$

Indeed, by (4.7) we have that

$$\varphi_1 \left(A \cap \varphi_A^{-1}(\overline{B}) \right) \subset A \cap B,$$

and since φ_1 is supported in A , we have also that

$$\varphi_1 \left(B \cap \varphi_A^{-1}(\overline{B}) \right) \subset B.$$

Next, we move all the points that end in a neighborhood of A^c to their final destination. Let D be an open ball containing A^c , such that $\overline{D} \subset B$. From (4.8) we have that $\varphi_1 \circ \varphi_A^{-1}(\overline{D})$




Fig. 1 A sketch of the construction of φ_1 . The sets A and B that cover the sphere are the blue and crossed one, respectively. On the left-hand figure, the orange domain is $\varphi_A^{-1}(\overline{B})$. Note that $\varphi_A^{-1}(\overline{B})$ does not wrap around A because of (4.7). The map φ_1 is supported in A , and maps the set $\varphi_A^{-1}(\overline{B})$ into B , as in the right-hand figure

is a closed ball contained in B . Therefore, there exists a diffeomorphism ψ , supported in B , such that $\psi|_{\overline{D}} = \varphi_1 \circ \varphi_A^{-1}|_{\overline{D}}$. Define $\varphi_2 := \psi^{-1}$.

Finally, define $\varphi_3 := \varphi_A \circ \varphi_1^{-1} \circ \varphi_2^{-1}$. We have that φ_3 is supported in A : indeed, for any $x \in D$,

$$\varphi_3(x) = \varphi_A \circ \varphi_1^{-1} \circ \psi(x) = \varphi_A \circ \varphi_1^{-1} \circ \varphi_1 \circ \varphi_A^{-1}(x) = x.$$

□

We will now continue with proving the boundedness of $\text{diam}_{s,p} \text{Diff}(\mathcal{M})$ by showing that if $\text{Diff}(\mathcal{M})$ satisfies the uniform fragmentation property, then the question of finiteness of $\text{diam}_{s,p} \text{Diff}(\mathcal{M})$ can be reduced to the finiteness of the diameter of diffeomorphisms groups of Euclidean balls.

Proposition 4.7 *Let \mathcal{M} be a closed n -dimensional manifold, such that $\text{Diff}(\mathcal{M})$ satisfies the uniform fragmentation property with respect to some cover. Assume that $\text{diam}_{s,p} \text{Diff}_c(B) < \infty$, where B is the unit ball of \mathbb{R}^n . Then, $\text{diam}_{s,p} \text{Diff}(\mathcal{M}) < \infty$.*

Proof Note that by scaling, our assumption $\text{diam}_{s,p} \text{Diff}_c(B) < \infty$ implies the finite diameter of the compactly-supported diffeomorphism group of any Euclidean ball of arbitrary radius.

Let $\{B_{\varepsilon_\alpha}(x_\alpha)\}_{\alpha \in A}$ be an open cover of \mathcal{M} by geodesic balls, with respect to which the uniform fragmentation property holds. Consider now the cover $\{B_{\eta_\alpha}(x_\alpha)\}_{\alpha \in A}$, where $\eta_\alpha > \varepsilon_\alpha$ for each $\alpha \in A$, such that normal coordinates are defined on $B_{\eta_\alpha}(x_\alpha)$ as well. To simplify notation, we denote $B^\alpha = B_{\varepsilon_\alpha}(x_\alpha)$ and $\tilde{B}^\alpha = B_{\eta_\alpha}(x_\alpha)$. We will henceforth consider $W^{s,p}$ -metrics on $\text{Diff}(\mathcal{M})$ with respect to $\{\tilde{B}^\alpha\}_{\alpha \in A}$ and a partition of unity ρ_α subordinate to this cover.

Let $\varphi \in \text{Diff}(\mathcal{M})$. By the uniform fragmentation property, there exists $\varphi_1, \dots, \varphi_K \in \text{Diff}(\mathcal{M})$, with $\text{supp } \varphi_i \subset B^\alpha$ for some $\alpha \in A$, such that $\varphi = \varphi_K \circ \dots \circ \varphi_1$. By right-invariance of the norm and the triangle inequality we have

$$\text{dist}_{s,p}(\text{Id}, \varphi) \leq \sum_{i=1}^K \text{dist}_{s,p}(\text{Id}, \varphi_i).$$

Therefore, in order to prove that $\text{diam}_{s,p} \text{Diff}(\mathcal{M}) < \infty$, it is enough to prove that $\text{dist}_{s,p}(\text{Id}, \varphi_i)$ is uniformly bounded. Therefore, we will henceforth assume that $\text{supp } \varphi \subset$

B^{α_0} for some $\alpha_0 \in A$. Since, by assumption, $\text{diam}_{s,p} \text{Diff}_c(B) < \infty$, there exists a path φ^t (with vector fields u^t) from Id to φ , supported on B^{α_0} , such that

$$\text{length}_{s,p}^{\mathbb{R}^n}(\varphi^t) < C$$

for some C independent of φ (we identified B^{α_0} with a ball in \mathbb{R}^n using normal coordinates as in the definition of the norm on $\mathcal{X}(\mathcal{M})$). This does not complete the proof as the support of u^t intersects other coordinate balls involved in the definition of the $W^{s,p}$ -norm on $\mathcal{X}(\mathcal{M})$, and therefore these balls also contribute to the length of the path. We now show that this contribution is uniformly bounded (and depends only on ε_{α_0} and η_{α_0}).

Let $\rho_{\alpha}^{\alpha_0}$ be another partition of unity subordinate to \tilde{B}^{α} , such that $\rho_{\alpha_0}^{\alpha_0}|_{B^{\alpha_0}} \equiv 1$. With respect to this partition of unity, the length of the path φ^t is the same as in the coordinate chart (since, by definition of $\rho_{\alpha}^{\alpha_0}$, the support of u^t intersect only the support of $\rho_{\alpha_0}^{\alpha_0}$), and is therefore bounded independent of φ .

The norm with respect to $\rho_{\alpha}^{\alpha_0}$ is equivalent to the one with respect to the original ρ_{α} , and therefore, since the choice of $\rho_{\alpha}^{\alpha_0}$ is independent of φ (depends only on ε_{α_0} and η_{α_0}) and the cover is finite, we obtain a uniform bound on the length of φ^t with respect to our original norm as well. □

Using Proposition 4.7, we now complete the proof of Theorem 1.1 by showing that $\text{diam}_{s,p} \text{Diff}_c(B) < \infty$ for $s < 1 + \frac{1}{p}$.

Lemma 4.8 *Let $s < 1 + \frac{1}{p}$ and let B denote the unit ball in \mathbb{R}^n . Then*

$$\text{diam}_{s,p} \text{Diff}_c(B) < C(s, p, n) < \infty.$$

Proof We will use polar coordinate on B_{ε} , in the standard notation

$$B = \{(r, \theta) : r \in [0, 1], \theta \in S^{n-1}\}. \tag{4.9}$$

Similar as in the one-dimensional situation the proof of this theorem will follow in two steps. *Step I* First we show that the boundedness of the diameter follows if (controlled) arbitrary change in volume has a bounded cost. To this end we denote by $\Psi_{\lambda,\delta} \in \text{Diff}(B)$ a map satisfying

$$\Psi_{\lambda,\delta}(r, \theta) = (\psi_{\lambda,\delta}(r), \theta), \tag{4.10}$$

where

$$\psi_{\lambda,\delta}(r) = \lambda r \quad \text{for } r \in \left[0, \frac{1-\delta}{\lambda}\right], \tag{4.11}$$

with $\lambda \in \mathbb{N}$ and $\delta \in (0, 1)$. In addition we assume that for some $C > 0$, independent of λ and δ , we have $\text{dist}_{s,p}(\text{Id}, \Psi_{\lambda,\delta}) < C$. Let now $\varphi \in \text{Diff}_c(B)$ be an arbitrary diffeomorphism. Since it has compact support, there exists $\delta > 0$ such that

$$\text{supp } \varphi \subset \{[0, 1 - \delta] \times S^{n-1}\} \subset B.$$

Denote

$$\varphi^\lambda(r, \theta) = \frac{1}{\lambda} \varphi(\lambda r, \theta) = \Psi_{\lambda,\delta}^{-1} \circ \varphi \circ \Psi_{\lambda,\delta}(r, \theta),$$

where the last equality holds because $\text{supp } \varphi \subset [0, 1 - \delta] \times S^{d-1}$ and since multiplication by a scalar in polar coordinates is given by $\lambda(r, \theta) = (\lambda r, \theta)$. Using the boundedness of $\text{dist}_{s,p}(\text{Id}, \Psi_{\lambda,\delta})$ we have

$$\text{dist}_{s,p}(\text{Id}, \varphi) = \text{dist}_{s,p}(\text{Id}, \Psi_{\lambda,\delta} \circ \varphi^\lambda \circ \Psi_{\lambda,\delta}^{-1}) < 2C + \text{dist}_{s,p}(\text{Id}, \varphi^\lambda).$$

By a direct calculation, similar as in Lemma 3.8, we have that the map $\varphi(t, x) \mapsto \varphi^\lambda(t, x) = \lambda^{-1}\varphi(t, \lambda x)$ is a bijection between paths supported on $[0, 1] \times S^{n-1}$ to paths supported on $[0, 1/\lambda] \times S^{n-1}$, with the corresponding vector fields

$$u_t^\lambda(r, \theta) = \frac{1}{\lambda}u_t(\lambda r, \theta).$$

From here the proof of the the above claims follows exactly as in Lemma 3.8, using Lemma 2.3 to obtain that

$$\text{dist}_{s,p}(\text{Id}, \varphi^\lambda) \leq \lambda^{(s-1)-\frac{n}{p}} \text{dist}_{s,p}(\text{Id}, \varphi).$$

Hence, by taking $\lambda \rightarrow \infty$, we have $\text{dist}_{s,p}(\text{Id}, \varphi) \leq 2C$. Note that it follows from Lemma 2.3 that this part of the proof holds whenever $s < 1 + \frac{n}{p}$, not merely when $s < 1 + \frac{1}{p}$.

Step II It remains to bound the distance from the identity to $\Psi_{\lambda,\delta}$ independently of λ and δ . Here we will rely on our construction from the one-dimensional case. From Lemma 3.9 we know that there exists a curve $\psi_{\lambda,\delta}^t$ in $\text{Diff}([0, 1])$ from $\text{Id}_{[0,1]}$ to $\psi_{\lambda,\delta}$ such that

$$\text{length}_{s,p}(\psi_{\lambda,\delta}^t) < C(s, p), \quad \text{for every } \lambda \in \mathbb{N} \text{ and } \delta \in (0, 1). \tag{4.12}$$

Let u_t be the vector field associated with the curve $\psi_{\lambda,\delta}^t$, and define a curve $\Psi_{\lambda,\delta}^t$ by flowing from the identity map, along the vector field

$$U_t(x) = u_t(|x|)\frac{x}{|x|}.$$

Obviously, $\Psi_{\lambda,\delta}^1 = \Psi_{\lambda,\delta}$ satisfies assumptions (4.10)–(4.11).

Since $u_t \in W_0^{s,p}(0, 1)$ we have, using Corollary B.3 and (4.12), that there exists $C = C(s, p, n) > 0$, independent of λ and δ , such that

$$\text{dist}_{W^{s,p}(B)}(\text{Id}, \Psi_{\lambda,\delta}) \leq \text{length}_{s,p}(\Psi_{\lambda,\delta}^t) < C, \quad \text{for every } \lambda \in \mathbb{N} \text{ and } \delta \in (0, 1),$$

which completes the proof. □

4.4 The displacement energy

Since step I of the proof of Lemma 4.8 holds for any $s < 1 + \frac{n}{p}$, the only ingredient needed for proving that $\text{diam}_{s,p} \text{Diff}(\mathcal{M}) < \infty$ for $s < 1 + \frac{n}{p}$, is a better vector field U_t in step II. That is, we need a better way of flowing from Id to $\Psi_{\lambda,\delta}$. An indication that this should be possible is the following proposition, which deals with the uniform boundedness of the displacement energy of sets in $\text{Diff}(S^n)$ (see Definition 2.4). Although, as discussed earlier, we do not know that bounded displacement energy is equivalent to bounded diameter, all our current examples are consistent with such a claim. Moreover, the proof shows that an arbitrary radial change of volume (which is what $\Psi_{\lambda,\delta}$ does) is possible at a bounded cost whenever $s < 1 + \frac{n}{p}$ (at least when $s \in \mathbb{N}$), although the change of volume in the proof is not as controlled as the one induced by $\Psi_{\lambda,\delta}$.

Proposition 4.9 *The following bounds on the displacement energy of subsets of S^n hold:*

1. *If $s > 1 + n/p$ then there exists $c = c(s, p) > 0$ such that*

$$E_{s,p}(S^n \setminus B_\delta) > c|\log \delta|,$$

where B_δ is a ball of radius δ in S^n .

2. For every integer $k < 1 + n/p$, there exists $C = C(k, p, n) > 0$ such that

$$E_{k,p}(A) < C \text{ for every closed set } A \subsetneq S^n.$$

Remark 4.10 We expect that the same line of proof below also yields the uniform boundedness for any non-integer $s < 1 + n/p$; we did not pursue the more involved estimates for non-integer values of s as the main point of this section is to indicate why we conjecture that $1 + n/p$ is the critical exponent. However, for some non-integer exponents (in particular when $1 + n/p$ is an integer) we could simply use the Sobolev embedding theorem (see, e.g., [13]):

$$\| \cdot \|_{W^{s,p}} \lesssim \| \cdot \|_{W^{r,q}}, \quad \frac{1}{q} - \frac{r}{n} = \frac{1}{p} - \frac{s}{n}, \quad r > s.$$

For example, consider $H^s = W^{s,2}$ in two dimensions. Then we know that for $s > 2$ the displacement energy is not bounded, while for $s < 2$, we can choose $q = \frac{2}{3-s} < 2$, and then

$$\| \cdot \|_{H^s} \lesssim \| \cdot \|_{W^{2,q}}.$$

The uniform boundedness of the displacement energy for $W^{2,q}(S^2)$, $q < 2$ therefore implies the boundedness for H^s , $s < 2$.

Proof We start by proving the unboundedness for large s :

Unboundedness for $s > 1 + n/p$. Let $\delta > 0$ be small enough. Denote $A_\delta = S^n \setminus B_\delta$. If $\varphi(A_\delta) \cap A_\delta = \emptyset$, then $\varphi(A_\delta) \subset B_\delta$, and therefore

$$\int_{A_\delta} |D\varphi| \, d\text{Vol} = \text{Vol}(\varphi(A_\delta)) \leq \text{Vol}(B_\delta).$$

Since $\text{Vol}(A_\delta)$ is of order one, and $\text{Vol}(B_\delta) = O(\delta^n)$, it follows that there exists a point $x \in S^n$ such that $|D\varphi(x)| = O(\delta^n)$. The first part of the proposition now follows immediately from the estimate 4.1.

Boundedness for $k < 1 + n/p$. For simplicity, we endow S^n with a round metric with diameter 1, and consider the cover of S^n with two balls of radius $3/4$, one centered at the south pole and the other at the north pole. Let $A \subsetneq S^n$ be a closed set. Then, there exists a ball of radius $\varepsilon > 0$, disjoint of A . Denote it by B_ε . Since an arbitrary rotation of S^n has a bounded cost, we can assume without loss of generality that B_ε is centered at the south pole.

We now construct φ such that $\varphi(A) \subset B_\varepsilon$ and $\text{dist}_{k,p}(\text{Id}, \varphi)$ is bounded independently of A and ε . Fix $\alpha \in (0, \frac{n}{p} + 1 - k)$. Let $u_{\alpha,\varepsilon} \in C_c^\infty((0, 3/4))$ be such that

$$u_{\alpha,\varepsilon}(x) = \begin{cases} 0 & x \in [0, \varepsilon/2) \\ x^{1-\alpha} & x \in [\varepsilon, 2/3) \end{cases}$$

and such that, for some $C > 0$ independent of ε ,

$$\left| u_{\alpha,\varepsilon}^{(j)}(x) \right| \leq C\varepsilon^{1-\alpha-j} \quad \forall x \in \left[\frac{\varepsilon}{2}, \varepsilon \right], \quad j = 0 \dots k. \tag{4.13}$$

Define now a vector field $U_{\alpha,\varepsilon}$ on the Euclidean ball of radius $3/4$ by

$$U_{\alpha,\varepsilon}(x) = u_{\alpha,\varepsilon}(|x|) \frac{x}{|x|}.$$

A straightforward calculation shows that

$$D^{(j)}U_{\alpha,\varepsilon}(x) = \sum_{i=0}^j \frac{u_{\alpha,\varepsilon}^{(i)}(|x|)}{|x|^{j-i}} G_{i,j} \left(\frac{x}{|x|} \right),$$

where $G_{i,j}$ is a tensor-valued polynomial (independent of $u_{\alpha,\varepsilon}$). We therefore have that

$$\left| D^{(j)} U_{\alpha,\varepsilon}(x) \right| \lesssim \sum_{i=0}^j \frac{|u_{\alpha,\varepsilon}^{(i)}(|x|)|}{|x|^{j-i}}. \tag{4.14}$$

We now evaluate $\|D^{(k)} U_{\alpha,\varepsilon}\|_p$, and show that it is independent of ε . By (4.14) it is enough to show that for every $i \leq k$,

$$\int_0^{3/4} \left| \frac{u_{\alpha,\varepsilon}^{(i)}(r)}{r^{k-i}} \right|^p r^{n-1} dr < C \tag{4.15}$$

for some C independent of ε . Indeed

$$\begin{aligned} \int_0^{3/4} \left| \frac{u_{\alpha,\varepsilon}^{(i)}(r)}{r^{k-i}} \right|^p r^{n-1} dr &= \int_{\varepsilon/2}^\varepsilon \left| \frac{u_{\alpha,\varepsilon}^{(i)}(r)}{r^{k-i}} \right|^p r^{n-1} dr + \int_\varepsilon^{2/3} \left| \frac{u_{\alpha,\varepsilon}^{(i)}(r)}{r^{k-i}} \right|^p r^{n-1} dr \\ &\quad + \int_{2/3}^{3/4} \left| \frac{u_{\alpha,\varepsilon}^{(i)}(r)}{r^{k-i}} \right|^p r^{n-1} dr \end{aligned}$$

The third addend on the right-hand side can obviously be bounded independently of ε , and therefore we can ignore it. The second addend can be evaluated explicitly, using the fact that $u_{\alpha,\varepsilon}(r) = r^{1-\alpha}$ in this region:

$$\begin{aligned} \int_\varepsilon^{2/3} \left| \frac{u_{\alpha,\varepsilon}^{(i)}(r)}{r^{k-i}} \right|^p r^{n-1} dr &= C_{\alpha,i} \int_\varepsilon^{2/3} \left| \frac{r^{1-\alpha-i}}{r^{k-i}} \right|^p r^{n-1} dr = C_{\alpha,i} \int_\varepsilon^{2/3} r^{n-1+(1-\alpha-k)p} dr \\ &< C_{\alpha,i} \int_0^1 r^{n-1+(1-\alpha-k)p} dr = C(\alpha, i, n, k, p), \end{aligned}$$

where in the last inequality we used the fact that $n + (1 - \alpha - k)p > 0$ since $\alpha < \frac{n}{p} + 1 - k$. As for the first addend, we have, using (4.13), that

$$\begin{aligned} \int_{\varepsilon/2}^\varepsilon \left| \frac{u_{\alpha,\varepsilon}^{(i)}(r)}{r^{k-i}} \right|^p r^{n-1} dr &\leq C \varepsilon^{(1-\alpha-i)p} \int_{\varepsilon/2}^\varepsilon r^{n-1-(k-i)p} dr \\ &\leq C' \varepsilon^{(1-\alpha-i)p} \varepsilon^{n-(k-i)p} = C' \varepsilon^{n+(1-\alpha-k)p}, \end{aligned}$$

which is uniformly bounded in ε since $n + (1 - \alpha - k)p > 0$. In the transition to the second line we use the fact that the lower bound of the integral is $\varepsilon/2$ rather than 0, and therefore we get boundedness even if $k p > n$ (for the case $i = 0$). This completes the proof of (4.15).

The proof for $j < k$ is similar, and therefore we obtain that there exists $C = C(k, p, n)$, independent of ε , such that

$$\|U_{\alpha,\varepsilon}\|_{W^{k,p}(\mathbb{R}^n)} < C. \tag{4.16}$$

Let ψ_t be a flow along $U_{\alpha,\varepsilon}$. Similar to the one dimensional case, after time $t_0 = \frac{1}{\alpha 2^\alpha}$ we have that

$$|\psi_{t_0}(x)| > 1/2 \quad \text{whenever } |x| > \varepsilon.$$

We now consider ψ_{t_0} as a diffeomorphism on S^n , using the normal coordinate chart centered at the south pole. $|\psi_{t_0}(x)| > 1/2$ for $|x| > \varepsilon$ implies then that ψ_{t_0} maps the

complement of B_ε to the northern hemisphere. Note that, by the same arguments as in Proposition 4.7, the bound (4.16) implies that

$$\text{dist}_{k,p}^{S^n}(\text{Id}, \psi_{t_0}) < C(k, n, p), \quad \text{independent of } \varepsilon.$$

Let R be a rotation of S^n that maps the south pole to the north pole, and consider

$$\varphi = \psi_{t_0}^{-1} \circ R \circ \psi_{t_0}.$$

Since ψ_{t_0} maps the complement of B_ε to the northern hemisphere, it follows that φ maps the complement of B_ε into B_ε , and therefore $\varphi(A) \subset B_\varepsilon$.

As the bound on $\text{dist}_{k,p}(\text{Id}, \psi_{t_0})$ implies that $\text{dist}_{k,p}(\text{Id}, \varphi)$ is bounded independent of ε , the proof is complete. □

5 The diameter of $\text{Diff}_c(\mathbb{R}^n)$

In the following we will consider the base manifold to be n -dimensional Euclidean space, i.e., $\mathcal{M} = \mathbb{R}^n$. In this case it will turn out, that the diameter of the the diffeomorphism group is either zero or unbounded (depending on the order s). We believe that the analogous results are also true for diffeomorphism groups on more general non-compact manifolds, but for simplicity, we will restrict ourselves here to the Euclidean case.

Proof of Theorem 1.2 The zero diameter result follows directly from the vanishing geodesic distance results of [5,32,33]. It remains to show that the diameter is unbounded otherwise.

For $sp > n$, the proof of positive geodesic distance [9,33] uses the Sobolev embedding $W^{s,p}(\mathbb{R}^n) \subset L^\infty(\mathbb{R}^n)$. It shows that for any $\varphi \in \text{Diff}_c(\mathbb{R}^n)$ and any $x \in \mathbb{R}^n$,

$$|\varphi(x) - x| \leq C \text{dist}_{s,p}(\text{Id}, \varphi). \tag{5.1}$$

Here $C = C(s, p, n) > 0$ is a constant depending on s, p and n . By choosing $\varphi(0)$ to be arbitrarily far away from the origin this shows that $\text{diam}_{s,p} \text{Diff}_c(\mathbb{R}^n) = \infty$ for $sp > n$.

For $s \geq 1$ a scaling argument yields the result independently from n and p . Fix $\lambda > 0$, and for $\varphi \in \text{Diff}_c(\mathbb{R}^n)$, define $\varphi^\lambda(x) := \lambda^{-1}\varphi(\lambda x)$. It is easy to see that $\text{supp } \varphi^\lambda = \lambda^{-1} \text{supp } \varphi$, hence $\varphi^\lambda \in \text{Diff}_c(\mathbb{R}^n)$. Similar arguments as in the proof of Lemma 3.8 show that given a path φ_t from Id to φ with a vector field u_t , φ_t^λ is a path from Id to φ^λ with a vector field

$$u_t^\lambda(x) = \frac{1}{\lambda} u_t(\lambda x).$$

It follows from Lemma 2.3 that

$$\|u_t^\lambda\|_{L^p}^p = \frac{1}{\lambda^{p+n}} \|u_t\|_{L^p}^p \quad \|u_t^\lambda\|_{W^{1,p}}^p = \frac{1}{\lambda^n} \|u_t\|_{W^{1,p}}^p,$$

and therefore, for $\lambda < 1$,

$$\|u_t^\lambda\|_{W^{1,p}} > \frac{1}{\lambda^{n/p}} \|u_t\|_{W^{1,p}}.$$

Since $\varphi_t \mapsto \varphi_t^\lambda$ is a bijection between the paths from Id to φ to the paths from Id to φ^λ , we have

$$\text{dist}_{1,p}(\text{Id}, \varphi^\lambda) \geq \frac{1}{\lambda^{n/p}} \text{dist}_{1,p}(\text{Id}, \varphi).$$

Taking $\lambda \rightarrow 0$, we obtain that $\text{diam}_{1,p} \text{Diff}_c(\mathbb{R}^n) = \infty$ (since from [5,33] we already know that $\text{dist}_{1,p}(\text{Id}, \varphi)$ is not zero). Since the $W^{s,p}$ norm for $s > 1$ controls the $W^{1,p}$ norm, we obtain that $\text{diam}_{s,p} \text{Diff}_c(\mathbb{R}^n) = \infty$ for any $s \geq 1$. \square

Acknowledgements We would like to thank to Stefan Haller, Philipp Harms, Stephen Preston, Tudor Ratiu and Josef Teichman for various discussions during the work on this paper, and to Meital Maor for her help with the figures. We are in particular grateful to Kathryn Mann and Tomasz Rybicki for introducing us to the literature on fragmentation and perfectness, and to Bob Jerrard for his continuous and valuable help throughout the work on this project. This project was initiated during the BIRS workshop ‘‘Shape Analysis, Stochastic Geometric Mechanics and Applied Optimal Transport’’ in December 2018; we are grateful to BIRS for their hospitality. M. Bauer was partially supported by NSF-grants 1912037 and 1953244. C. Maor was partially supported by ISF-grant 1269/19.

Author comment

The grant information should appear on the first page and not there.

Appendix A: Proof of Lemma 3.9

We consider the family of piecewise-linear maps $\psi_{\lambda+1,\delta}$

$$\psi_{\lambda+1,\delta} = \begin{cases} (\lambda + 1)x & x \in \left[0, \frac{1-\delta}{\lambda+1}\right] \\ \frac{\delta(\lambda+1)}{\lambda+\delta} \left(x - \frac{1-\delta}{\lambda+1}\right) + 1 - \delta & x \in \left[\frac{1-\delta}{\lambda+1}, 1\right] \end{cases} = \begin{cases} (\lambda + 1)x & x \in \left[0, \frac{1-\delta}{\lambda+1}\right] \\ \frac{\delta(\lambda+1)x + (1-\delta)\lambda}{\lambda+\delta} & x \in \left[\frac{1-\delta}{\lambda+1}, 1\right] \end{cases}.$$

Since piecewise-linear maps are not elements of the group of diffeomorphisms $\text{Diff}(S^1)$ we have to smoothen the maps around the break points $\frac{1-\delta}{\lambda+1}$ and $0 \sim 1$. However, since the $W^{s,p}$ -metric can be extended to the space of Lipschitz-maps (for $s < 1 + 1/p$) and since the smoothening can be done in such a way that the change in the distance to identity is arbitrarily small, we ignore this in the following.

In the following we will bound the length of the linear homotopy $\varphi_t(x)$ between Id and $\psi_{\lambda+1,\delta}$ which albeit being straightforward turns out to be a somewhat tedious calculation. We bound the length below with respect to the $\dot{W}^{s,p}$ norm, under the assumption that $s > 1$. Boundedness with respect to the lower order parts of $W^{s,p}$ norm, as well as for $W^{s,p}$ norm for $s \leq 1$, is similar, but simpler. We have

$$\varphi_t(x) = (1 - t)x + t\psi_{\lambda+1,\delta}(x) = \begin{cases} (1 + \lambda t)x & x \in \left[0, \frac{1-\delta}{\lambda+1}\right] \\ x + t \frac{(1-\delta)\lambda}{\lambda+\delta} (1 - x) & x \in \left[\frac{1-\delta}{\lambda+1}, 1\right] \end{cases}.$$

Its inverse is then given by

$$\varphi_t^{-1}(y) = \begin{cases} \frac{y}{1+\lambda t} & y \in \left[0, \frac{(1-\delta)(1+\lambda t)}{\lambda+1}\right] \\ \frac{y-1}{1-t \frac{(1-\delta)\lambda}{\lambda+\delta}} + 1 & y \in \left[\frac{(1-\delta)(1+\lambda t)}{\lambda+1}, 1\right] \end{cases},$$

and its time derivative is

$$\partial_t \varphi_t(x) = \psi_{\lambda+1,\delta}(x) - x = \begin{cases} \lambda x & x \in \left[0, \frac{1-\delta}{\lambda+1}\right] \\ \frac{(1-\delta)\lambda}{\lambda+\delta} (1 - x) & x \in \left[\frac{1-\delta}{\lambda+1}, 1\right] \end{cases}.$$

The vector field u_t defined by $\partial_t \varphi_t = u_t \circ \varphi_t$ is therefore

$$\begin{aligned}
 u_t(y) &= \partial_t \varphi_t(\varphi_t^{-1}(y)) = \begin{cases} \frac{\lambda y}{1+\lambda t} & y \in \left[0, \frac{(1-\delta)(1+\lambda t)}{\lambda+1}\right] \\ \frac{(1-\delta)\lambda}{\lambda+\delta} \frac{1-y}{1-t\frac{(1-\delta)\lambda}{\lambda+\delta}} & y \in \left[\frac{(1-\delta)(1+\lambda t)}{\lambda+1}, 1\right] \end{cases} \\
 &= \begin{cases} \frac{y}{t+\frac{1}{\lambda}} & y \in \left[0, \frac{(1-\delta)(1+\lambda t)}{\lambda+1}\right] \\ \frac{(1-\delta)(1-y)}{(1-t)(1-\delta)+\delta(1+\frac{1}{\lambda})} & y \in \left[\frac{(1-\delta)(1+\lambda t)}{\lambda+1}, 1\right] \end{cases},
 \end{aligned}$$

and therefore

$$u'_t(y) = \begin{cases} \frac{1}{t+\frac{1}{\lambda}} & y \in \left[0, \frac{(1-\delta)(1+\lambda t)}{\lambda+1}\right] \\ \frac{-(1-\delta)}{(1-t)(1-\delta)+\delta(1+\frac{1}{\lambda})} & y \in \left[\frac{(1-\delta)(1+\lambda t)}{\lambda+1}, 1\right] \end{cases}.$$

We now evaluate the $\dot{W}^{1+\sigma,p}$ -norm of u_t , for $\sigma p < 1$. That is, we evaluate the (σ, p) -Gagliardo seminorm of u'_t , whose p th power is

$$\begin{aligned}
 \iint_{\mathbb{R}^2} \frac{|u'_t(x) - u'_t(y)|^p}{|x - y|^{1+\sigma p}} dx dy &= 2 \iint_{y>x} \frac{|u'_t(x) - u'_t(y)|^p}{|x - y|^{1+\sigma p}} dx dy \\
 &= 2 \int_{-\infty}^{\infty} \int_0^{\infty} \frac{|u'_t(x) - u'_t(x+s)|^p}{s^{1+\sigma p}} ds dx.
 \end{aligned}$$

We split this double integral into different regions:

$$\begin{aligned}
 &\int_{-\infty}^{\infty} \int_0^{\infty} \frac{|u'_t(x) - u'_t(x+s)|^p}{s^{1+\sigma p}} ds dx \\
 &= \int_{-\infty}^0 \int_{-x}^{-x+\frac{(1-\delta)(1+\lambda t)}{\lambda+1}} \frac{\left(t + \frac{1}{\lambda}\right)^{-p}}{s^{1+\sigma p}} ds dx \\
 &\quad + \int_{-\infty}^0 \int_{-x+\frac{(1-\delta)(1+\lambda t)}{\lambda+1}}^{-x+1} \frac{\left(\frac{1-\delta}{(1-t)(1-\delta)+\delta(1+\frac{1}{\lambda})}\right)^p}{s^{1+\sigma p}} ds dx \\
 &\quad + \int_0^{\frac{(1-\delta)(1+\lambda t)}{\lambda+1}} \int_{-x+\frac{(1-\delta)(1+\lambda t)}{\lambda+1}}^{-x+1} \frac{\left(\frac{1}{t+\frac{1}{\lambda}} + \frac{1-\delta}{(1-t)(1-\delta)+\delta(1+\frac{1}{\lambda})}\right)^p}{s^{1+\sigma p}} ds dx \\
 &\quad + \int_{\frac{(1-\delta)(1+\lambda t)}{\lambda+1}}^1 \int_{-x+1}^{\infty} \frac{\left(\frac{1-\delta}{(1-t)(1-\delta)+\delta(1+\frac{1}{\lambda})}\right)^p}{s^{1+\sigma p}} ds dx.
 \end{aligned}$$

We now evaluate each of the four integrals in the right-hand side separately. We will use repeatedly the following: for $\alpha \in (0, 1)$ and $a > 0$,

$$\lim_{x \rightarrow \infty} (x+a)^\alpha - x^\alpha = 0,$$

and

$$(1-x)^\alpha \geq 1-x^\alpha \quad x \in [0, 1].$$

All the constants C below are $C = C(p, \sigma) > 0$, independent of λ, δ and t .

For the first integral we have:

$$\begin{aligned}
 & \left(t + \frac{1}{\lambda}\right)^{-p} \int_{-\infty}^0 \int_{-x}^{-x + \frac{(1-\delta)(1+\lambda t)}{\lambda+1}} \frac{1}{s^{1+\sigma p}} ds dx \\
 &= \left(t + \frac{1}{\lambda}\right)^{-p} \frac{1}{\sigma p} \int_{-\infty}^0 \left((-x)^{-\sigma p} - \left(-x + \frac{(1-\delta)(1+\lambda t)}{\lambda+1}\right)^{-\sigma p} \right) dx \\
 &= \left(t + \frac{1}{\lambda}\right)^{-p} \frac{1}{\sigma p} \int_0^{\infty} \left(x^{-\sigma p} - \left(x + \frac{(1-\delta)(1+\lambda t)}{\lambda+1}\right)^{-\sigma p} \right) dx \\
 &= \left(t + \frac{1}{\lambda}\right)^{-p} \frac{1}{(1-\sigma p)\sigma p} \left(x^{1-\sigma p} - \left(x + \frac{(1-\delta)(1+\lambda t)}{\lambda+1}\right)^{1-\sigma p} \right) \Big|_0^{\infty} \\
 &= \left(t + \frac{1}{\lambda}\right)^{-p} \frac{1}{(1-\sigma p)\sigma p} \left(\frac{(1-\delta)(1+\lambda t)}{\lambda+1} \right)^{1-\sigma p} \\
 &< C \left(t + \frac{1}{\lambda}\right)^{-p+(1-\sigma p)} < C t^{-p+(1-\sigma p)}.
 \end{aligned}$$

The second integral can be bounded via:

$$\begin{aligned}
 & \left(\frac{1-\delta}{(1-t)(1-\delta) + \delta(1 + \frac{1}{\lambda})} \right)^p \int_{-\infty}^0 \int_{-x + \frac{(1-\delta)(1+\lambda t)}{\lambda+1}}^{-x+1} \frac{1}{s^{1+\sigma p}} ds dx \\
 &= \frac{1}{\sigma p} \left(\frac{1-\delta}{(1-t)(1-\delta) + \delta(1 + \frac{1}{\lambda})} \right)^p \\
 & \quad \int_{-\infty}^0 \left(\left(\frac{(1-\delta)(1+\lambda t)}{\lambda+1} - x \right)^{-\sigma p} - (1-x)^{-\sigma p} \right) dx \\
 &= \frac{1}{\sigma p} \left(\frac{1-\delta}{(1-t)(1-\delta) + \delta(1 + \frac{1}{\lambda})} \right)^p \\
 & \quad \int_0^{\infty} \left(\left(\frac{(1-\delta)(1+\lambda t)}{\lambda+1} + x \right)^{-\sigma p} - (1+x)^{-\sigma p} \right) dx \\
 &= \frac{1}{(1-\sigma p)\sigma p} \left(\frac{1-\delta}{(1-t)(1-\delta) + \delta(1 + \frac{1}{\lambda})} \right)^p \\
 & \quad \left(\left(\frac{(1-\delta)(1+\lambda t)}{\lambda+1} + x \right)^{1-\sigma p} - (1+x)^{1-\sigma p} \right) \Big|_0^{\infty} \\
 &= \frac{1}{(1-\sigma p)\sigma p} \left(\frac{1-\delta}{(1-t)(1-\delta) + \delta(1 + \frac{1}{\lambda})} \right)^p \left(1 - \left(\frac{(1-\delta)(1+\lambda t)}{\lambda+1} \right)^{1-\sigma p} \right) \\
 &\leq \frac{1}{(1-\sigma p)\sigma p} \left(\frac{1-\delta}{(1-t)(1-\delta) + \delta(1 + \frac{1}{\lambda})} \right)^p \left(1 - \frac{(1-\delta)(1+\lambda t)}{\lambda+1} \right)^{1-\sigma p} \\
 &= \frac{1}{(1-\sigma p)\sigma p} \left(\frac{1-\delta}{(1-t)(1-\delta) + \delta(1 + \frac{1}{\lambda})} \right)^p \left(\delta + (1-\delta) \frac{\lambda}{\lambda+1} (1-t) \right)^{1-\sigma p}
 \end{aligned}$$

$$\begin{aligned}
 &< \frac{1}{(1-\sigma p)\sigma p} \left(\frac{1-\delta}{(1-t)(1-\delta)+\delta} \right)^p (\delta+(1-\delta)(1-t))^{1-\sigma p} \\
 &= \frac{(1-\delta)^p}{(1-\sigma p)\sigma p} (\delta+(1-\delta)(1-t))^{-p+(1-\sigma p)} < C(1-t)^{-p+(1-\sigma p)}.
 \end{aligned}$$

Similarly we calculate for the third integral:

$$\begin{aligned}
 &\left(\frac{1}{t+\frac{1}{\lambda}} + \frac{1-\delta}{(1-t)(1-\delta)+\delta(1+\frac{1}{\lambda})} \right)^p \int_0^{\frac{(1-\delta)(1+\lambda t)}{\lambda+1}} \int_{-x+\frac{(1-\delta)(1+\lambda t)}{\lambda+1}}^{-x+1} \frac{1}{s^{1+\sigma p}} ds dx \\
 &= p \left(\left(t + \frac{1}{\lambda} \right)^{-p} + \left(\frac{1-\delta}{(1-t)(1-\delta)+\delta(1+\frac{1}{\lambda})} \right)^p \right) \int_0^{\frac{(1-\delta)(1+\lambda t)}{\lambda+1}} \int_{-x+\frac{(1-\delta)(1+\lambda t)}{\lambda+1}}^{-x+1} \frac{1}{s^{1+\sigma p}} ds dx \\
 &= \frac{1}{\sigma} \left(\left(t + \frac{1}{\lambda} \right)^{-p} + \left(\frac{1-\delta}{(1-t)(1-\delta)+\delta(1+\frac{1}{\lambda})} \right)^p \right) \\
 &\quad \int_0^{\frac{(1-\delta)(1+\lambda t)}{\lambda+1}} \left(\left(\frac{(1-\delta)(1+\lambda t)}{\lambda+1} - x \right)^{-\sigma p} - (1-x)^{-\sigma p} \right) dx \\
 &= \frac{1}{(1-\sigma p)\sigma} \left(\left(t + \frac{1}{\lambda} \right)^{-p} + \left(\frac{1-\delta}{(1-t)(1-\delta)+\delta(1+\frac{1}{\lambda})} \right)^p \right) \\
 &\quad \left((1-x)^{1-\sigma p} - \left(\frac{(1-\delta)(1+\lambda t)}{\lambda+1} - x \right)^{1-\sigma p} \right) \Big|_0^{\frac{(1-\delta)(1+\lambda t)}{\lambda+1}} \\
 &= \frac{1}{(1-\sigma p)\sigma} \left(\left(t + \frac{1}{\lambda} \right)^{-p} + \left(\frac{1-\delta}{(1-t)(1-\delta)+\delta(1+\frac{1}{\lambda})} \right)^p \right) \\
 &\quad \left(\left(1 - \frac{(1-\delta)(1+\lambda t)}{\lambda+1} \right)^{1-\sigma p} - 1 + \left(\frac{(1-\delta)(1+\lambda t)}{\lambda+1} \right)^{1-\sigma p} \right) \\
 &\leq \frac{1}{(1-\sigma p)\sigma} \\
 &\quad \left[\left(t + \frac{1}{\lambda} \right)^{-p} \left(\frac{(1-\delta)(1+\lambda t)}{\lambda+1} \right)^{1-\sigma p} + \left(\frac{1-\delta}{(1-t)(1-\delta)+\delta(1+\frac{1}{\lambda})} \right)^p \left(1 - \frac{(1-\delta)(1+\lambda t)}{\lambda+1} \right)^{1-\sigma p} \right] \\
 &\leq \frac{1}{(1-\sigma p)\sigma} \\
 &\quad \left[\left(t + \frac{1}{\lambda} \right)^{-p} \left(\frac{(1-\delta)(1+\lambda t)}{\lambda+1} \right)^{1-\sigma p} + \left(\frac{1-\delta}{(1-t)(1-\delta)+\delta(1+\frac{1}{\lambda})} \right)^p \left(\delta + (1-\delta) \frac{\lambda}{\lambda+1} (1-t) \right)^{1-\sigma p} \right] \\
 &< \frac{1}{(1-\sigma p)\sigma} \\
 &\quad \left[\left(t + \frac{1}{\lambda} \right)^{-p} \left(\frac{1+\lambda t}{\lambda+1} \right)^{1-\sigma p} + \left(\frac{1}{(1-t)(1-\delta)+\delta} \right)^p (\delta+(1-\delta)(1-t))^{1-\sigma p} \right] \\
 &< \frac{1}{(1-\sigma p)\sigma} \left[\left(t + \frac{1}{\lambda} \right)^{-p+(1-\sigma p)} + (\delta+(1-\delta)(1-t))^{-p+(1-\sigma p)} \right] \\
 &< C \left(t^{-p+(1-\sigma p)} + (1-t)^{-p+(1-\sigma p)} \right)
 \end{aligned}$$

Finally the last integral can be bounded by:

$$\begin{aligned}
 & \left(\frac{1 - \delta}{(1 - t)(1 - \delta) + \delta(1 + \frac{1}{\lambda})} \right)^p \int_{\frac{(1-\delta)(1+\lambda t)}{\lambda+1}}^1 \int_{-x+1}^\infty \frac{1}{s^{1+\sigma p}} ds dx \\
 &= \frac{1}{\sigma p} \left(\frac{1 - \delta}{(1 - t)(1 - \delta) + \delta(1 + \frac{1}{\lambda})} \right)^p \int_{\frac{(1-\delta)(1+\lambda t)}{\lambda+1}}^1 (1 - x)^{-\sigma p} dx \\
 &= \frac{1}{(1 - \sigma p)\sigma p} \left(\frac{1 - \delta}{(1 - t)(1 - \delta) + \delta(1 + \frac{1}{\lambda})} \right)^p (1 - x)^{1-\sigma p} \Big|_1^{\frac{(1-\delta)(1+\lambda t)}{\lambda+1}} \\
 &= \frac{1}{(1 - \sigma p)\sigma p} \left(\frac{1 - \delta}{(1 - t)(1 - \delta) + \delta(1 + \frac{1}{\lambda})} \right)^p \left(1 - \frac{(1 - \delta)(1 + \lambda t)}{\lambda + 1} \right)^{1-\sigma p} \\
 &= \frac{1}{(1 - \sigma p)\sigma p} \left(\frac{1 - \delta}{(1 - t)(1 - \delta) + \delta(1 + \frac{1}{\lambda})} \right)^p \left(\delta + (1 - \delta) \frac{\lambda}{\lambda + 1} (1 - t) \right)^{1-\sigma p} \\
 &< \frac{(1 - \delta)^p}{(1 - \sigma p)\sigma p} \left(\frac{1}{(1 - t)(1 - \delta) + \delta} \right)^p (\delta + (1 - \delta)(1 - t))^{1-\sigma p} \\
 &= \frac{(1 - \delta)^p}{(1 - \sigma p)\sigma p} (\delta + (1 - \delta)(1 - t))^{-p+(1-\sigma p)} < C(1 - t)^{-p+(1-\sigma p)}.
 \end{aligned}$$

Overall we obtained

$$\|u'_t\|_{\dot{W}^{\sigma,p}(\mathbb{R})} < C \left((1 - t)^{-p+(1-\sigma p)} + t^{-p+(1-\sigma p)} \right)^{1/p} < C \left((1 - t)^{-1+\frac{1-\sigma p}{p}} + t^{-1+\frac{1-\sigma p}{p}} \right)$$

where we used the fact that $(1 + x)^\alpha < 1 + x^\alpha$ for $x > 0$ and $\alpha \in (0, 1)$.

We therefore have, using the fact that $1 - \sigma p > 0$, that

$$\int_0^1 \|u'_t\|_{\dot{W}^{\sigma,p}(\mathbb{R})} dt \leq C \int_0^1 C \left((1 - t)^{-1+\frac{1-\sigma p}{p}} + t^{-1+\frac{1-\sigma p}{p}} \right) dt = 2C \frac{p}{1 - \sigma p},$$

which is a bound independent of λ and δ .

Appendix B: Sobolev norms of radial functions

In this section we prove a technical lemma on Sobolev functions, which is used in Sect. 4.3.

Lemma B.1 *Let $n > 1$, and define the operator $T : C_c^\infty((0, 1)) \rightarrow C_c^\infty(B_1(\mathbb{R}^n))$ by*

$$Tf(x) = f(|x|).$$

Then for every $s \geq 0$ and $p \geq 1$, we have

$$\|Tf\|_{W^{s,p}} \leq C \|f\|_{W^{s,p}},$$

for some $C = C(s, p, n) > 0$ independent of f . That is, $T : W_0^{s,p}(0, 1) \rightarrow W_0^{s,p}(B_1(\mathbb{R}^n))$ is a bounded operator for every $s \geq 0$ and $p \geq 1$.

Proof *Step I: integer Sobolev spaces* We first prove the theorem for $W^{k,p}$ norms, where k is an integer. For $k = 0$, moving to polar coordinates, we have

$$\|Tf\|_{L^p}^p = \int_{B_1(\mathbb{R}^n)} |Tf(x)|^p dx = \omega_n \int_0^1 |f(r)|^p r^{n-1} dr \leq \omega_n \|f\|_{L^p}^p,$$

where ω_n is the measure of the $(n - 1)$ -dimensional unit sphere. For $k = 1$, we note that $D(Tf)(x) = f'(|x|) \frac{x}{|x|}$, hence $|D(Tf)(x)| = |f'(r)|$ and the estimate is similar. Differentiating further, we have for $k = 2$

$$D^2(Tf)(x) = \left(f''(|x|) - \frac{f'(|x|)}{|x|} \right) \frac{x}{|x|} \otimes \frac{x}{|x|} + \frac{f'(|x|)}{|x|} \text{Id},$$

and for higher derivatives we obtain

$$D^k(Tf)(x) = \sum_{j=1}^k \frac{f^{(j)}(|x|)}{|x|^{k-j}} G_j^k \left(\frac{x}{|x|} \right),$$

where G_j^k are smooth k -tensor-valued functions on S^{n-1} , which are independent of f .

In order to prove boundedness we need to prove that for $j \leq k$ we have that

$$\int_0^1 \left| \frac{f^{(j)}(r)}{r^{k-j}} \right|^p r^{n-1} dr \leq \int_0^1 |f^{(k)}(r)|^p dr.$$

This follows from Jensen’s inequality: For $k = 1$, we have

$$\begin{aligned} \int_0^1 \left| \frac{f'(r)}{r} \right|^p r^{n-1} dr &= \int_0^1 \left| \frac{1}{r} \int_0^r f''(t) dt \right|^p r^{n-1} dr \leq \int_0^1 \left(\frac{1}{r} \int_0^r |f''(t)|^p dt \right) r^{n-1} dr \\ &= \int_0^1 \int_0^r |f''(t)|^p dt r^{n-2} dr \leq \int_0^1 |f''(t)|^p dt \cdot \int_0^1 r^{n-2} dr \\ &= \frac{1}{n-1} \int_0^1 |f''(t)|^p dt. \end{aligned}$$

For $k = 2$ we have

$$\begin{aligned} \int_0^1 \left| \frac{f'(r)}{r^2} \right|^p r^{n-1} dr &= \int_0^1 \left| \frac{1}{r} \int_0^r \frac{1}{r} \int_0^t f^{(3)}(s) ds dt \right|^p r^{n-1} dr \\ &\leq \int_0^1 \frac{1}{r} \int_0^r \left| \frac{1}{r} \int_0^t f^{(3)}(s) ds \right|^p dt r^{n-1} dr \\ &= \int_0^1 \frac{1}{r} \int_0^r \frac{t^p}{r^p} \left| \frac{1}{t} \int_0^t f^{(3)}(s) ds \right|^p dt r^{n-1} dr \\ &\leq \int_0^1 \frac{1}{r} \int_0^r \frac{t^p}{r^p} \frac{1}{t} \int_0^t |f^{(3)}(s)|^p ds dt r^{n-1} dr \\ &\leq \int_0^1 |f^{(3)}(s)|^p ds \cdot \int_0^1 \frac{1}{r} \int_0^r \frac{t^p}{r^p} \frac{1}{t} dt r^{n-1} dr \\ &= \frac{1}{p(n-1)} \int_0^1 |f^{(3)}(s)|^p ds. \end{aligned}$$

The result for higher values of k follows in a similar manner.

Step II: Interpolation Assume for now that $s \in (0, 1)$. Since $B_1(\mathbb{R}^n)$ is a convex set, we have that the $W^{s,p}(\mathbb{R}^n)$ norm on functions supported on $B_1(\mathbb{R}^n)$ (the Gagliardo/Slobodeckij norm) is equivalent to the norm of the real interpolation space

$$\chi_0^{s,p}(B_1(\mathbb{R}^n)) = (L^p(B_1(\mathbb{R}^n)), W^{1,p}(B_1(\mathbb{R}^n)))_{s,p},$$

defined by

$$\begin{aligned} \|f\|_{\chi_0^{s,p}(B_1(\mathbb{R}^n))}^p &= \int_0^\infty \left(\frac{K(t, f)}{t^s} \right) \frac{dt}{t} \quad K(t, f) \\ &= \inf_{g \in C_0^\infty(B_1(\mathbb{R}^n))} (\|f - g\|_{L^p(B_1(\mathbb{R}^n))} + t\|g\|_{W^{1,p}(B_1(\mathbb{R}^n))}). \end{aligned}$$

See [15, Theorem 4.7].³ Since $\chi_0^{s,p}(B_1(\mathbb{R}^n))$ is an interpolation space, the map T is bounded as a map $L^p([0, 1]) \rightarrow L^p(B_1(\mathbb{R}^n))$ and as a map $W_0^{1,p}([0, 1]) \rightarrow W_0^{1,p}(B_1(\mathbb{R}^n))$ and thus is also bounded as a map between the corresponding interpolation spaces $\chi_0^{s,p}([0, 1]) \rightarrow \chi_0^{s,p}(B_1(\mathbb{R}^n))$ (see, e.g., [53, Section 2.3, Theorem 3]).

When $s = k + \sigma$, the proof is similar: T is bounded as a map of between the interpolation spaces $(\dot{W}^{k,p}(0, 1), \dot{W}^{k+1,p}(0, 1))_{\sigma,p} \rightarrow (\dot{W}^{k,p}(B_1(\mathbb{R}^n)), \dot{W}^{k+1,p}(B_1(\mathbb{R}^n)))_{\sigma,p}$, since by the previous step it is bounded as maps on the interpolating spaces; and the norm on these interpolation spaces is equivalent to the $\dot{W}^{s,p}(\mathbb{R}^n)$ -norm on $C_0^\infty(B_1(\mathbb{R}^n))$ functions, by the same results as for the $k = 0$ case. □

Remark B.2 This lemma could probably be proven, at least for low values of k , by brute force evaluation of the Gagliardo seminorm, using the Funk–Hecke theorem (see, e.g., [29]).

An immediate corollary is the analogous result for vector fields, instead of functions:

Corollary B.3 *Let $n > 1$, and define the operator $\tilde{T} : C_c^\infty((0, 1)) \rightarrow C_c^\infty(B_1(\mathbb{R}^n); \mathbb{R}^n)$ by*

$$\tilde{T}f(x) = f(|x|) \frac{x}{|x|}$$

Then for every $s \geq 0$ and $p \geq 1$, we have

$$\|\tilde{T}f\|_{W^{s,p}} \leq C\|f\|_{W^{s,p}},$$

for some $C = C(s, p, n) > 0$ independent of f . That is, $\tilde{T} : W_0^{s,p}(0, 1) \rightarrow W_0^{s,p}(B_1(\mathbb{R}^n); \mathbb{R}^n)$ is a bounded operator for any $s \geq 0$ and $p \geq 1$.

Proof Let $F \in C_c^\infty((0, 1))$ be an antiderivative of f . Then the corollary follows from Lemma B.1 since $\tilde{T}f = D(TF)$. □

Appendix C: Diameter and displacement energy

In this section we prove a general result relating bounded displacement energy and bounded diameter, inspired by previous results relating zero displacement energy and vanishing geodesic distance [9,23,54]. However, as shown below, compared with the vanishing case we need stronger assumptions on the norms involved, assumptions which are too restrictive to the applications in this paper; therefore we used other means to prove boundedness of the diameter.

Let G be a (possibly infinite dimensional) manifold and topological group with neutral element e , Lie algebra $\mathfrak{g} = T_eG$, and left and right translations L and R given by

$$g_1g_2 = L_{g_1}(g_2) = R_{g_2}(g_1), \quad \forall g_1, g_2 \in G. \tag{C.1}$$

³ In [15] the interpolation is defined with respect to the homogeneous $\dot{W}^{1,p}$ norm, but this does not matter as it is, by the Poincaré inequality, equivalent to the full $W^{1,p}$ norm on the space $W_0^{1,p}$ which we are considering. Similarly, the equivalence there is shown between the interpolation space and the homogeneous $\dot{W}^{s,p}$ norm, which is again equivalent to the full norm [15, Section 2.3].

Assume for each $g \in G$ that $R_g : G \rightarrow G$ is smooth, and let $\| \cdot \|$ be a norm on the Lie algebra \mathfrak{g} . This gives rise to the following right-invariant Riemannian metric on G :

$$\|h\|_g = \|TR_{g^{-1}}h\|, \quad \forall g \in G, \quad \forall h \in T_gG. \tag{C.2}$$

The corresponding geodesic distance function is defined as

$$\text{dist}(g_1, g_2) = \inf \int_0^1 \|\partial_t g(t)\|_{g(t)} dt, \quad \forall g_1, g_2 \in G, \tag{C.3}$$

where the infimum is taken over all smooth paths in G with $g(0) = g_1$ and $g(1) = g_2$.

Theorem C.1 *Let G be as above. Assume that*

1. *Any transformation g can be written as a product $g = g_1g_2$ where both g_1 and g_2 are supported on a proper closed subset of M .*
2. *For any proper closed subset $A \subset M$ the group $G_A \subset G$ of all transformations that have support in A is uniformly perfect, i.e., any $g \in G_A$ can be written as a product of n commutators, where n is independent of $g \in G$.*
3. *The geodesic distance to a commutator of g and h is uniformly controlled by the minimum of the distances to g and h , i.e.,*

$$\text{dist}(e, [g, h]) = \text{dist}(g \circ h, h \circ g) \leq C \min(\text{dist}(e, g), \text{dist}(e, h)), \quad \forall g, h \in G \tag{C.4}$$

where C is independent of both g and h .⁴

4. *The displacement energy is globally bounded, i.e., for any proper closed subset $A \subset M$ we have*

$$E(A) = \inf \{ \text{dist}(e, g) : g \in G, g(A) \cap A = \emptyset \} \leq D \tag{C.5}$$

where D is independent of the set A .

Then the diameter of the group G is bounded.

Proof Using Assumption C.1 and the right invariance of the geodesic distance we can reduce the boundedness of the diameter to consider only transformations that are supported on a proper closed subset of M , since

$$\begin{aligned} \text{dist}(e, g) &= \text{dist}(e, g_1g_2) = \text{dist}(g_2^{-1}, g_1) \leq \text{dist}(g_2^{-1}, e) + \text{dist}(e, g_1) \\ &= \text{dist}(e, g_2) + \text{dist}(e, g_1), \end{aligned} \tag{C.6}$$

where both g_1 and g_2 are supported in a proper subset of M .

Thus it remains to prove the boundedness of the distance from the identity to any transformation g with support in a proper closed subset A . Using Assumption C.1 we write any $g_1 = [h_1, h_2][h_3, h_4] \cdots [h_{2n-1}, h_{2n}]$ with $h_i \in G_A$. By the same argument as above we obtain

$$\text{dist}(e, g_1) \leq \sum_{i=1}^n \text{dist}(e, [h_{2i-1}, h_{2i-1}]). \tag{C.7}$$

To bound the distance from the identity to a commutator of transformations with support in A we proceed as in [9, Theorem 1] and use Assumption C.1 to obtain

$$\text{dist}(e, [h_{2i-1}, h_{2i-1}]) \leq (1 + C)^2 E(A). \tag{C.8}$$

⁴ Note that this holds if the left multiplication L_g is Lipschitz with Lipschitz constant that is independent of g , see [9, Theorem 1].

Putting all of this together we have for each $g \in G$ that

$$\text{dist}(e, g) \leq 2n(1 + C)^2 E(A) \tag{C.9}$$

and using assumption C.1 and the triangle inequality this yields

$$\text{dist}(g, h) \leq 4n(1 + C)^2 D \tag{C.10}$$

for any $g, h \in G$. □

Let now $M = S^n$ and let $G = \text{Diff}(S^n)$. Then Assumptions C.1 and C.1 are satisfied [17, 61]. Assumption C.1 is satisfied for $W^{s,p}$ -metrics of low enough order, see Proposition 4.9. In the following we will however show that already in the case $s = 1$ and $n = 1$ condition C.1 is too restrictive for our purposes as, e.g., the \dot{H}^1 metric on $\text{Diff}(S^1)$, which corresponds to bounded diameter, does not satisfy it:

Lemma C.2 *There exist sequences $\psi_n, \varphi_n \in \text{Diff}(S^1)$ such that $\text{dist}_{\dot{H}^1}(\varphi_n \circ \psi_n, \psi_n \circ \varphi_n) \rightarrow \pi/2$ but $\text{dist}_{\dot{H}^1}(\text{Id}, \varphi_n) \rightarrow 0$.*

Proof By the analysis of Lenells [42] we have an explicit formula for the geodesic distance of the homogeneous \dot{H}^1 -metric given by:

$$\text{dist}_{1,2}(\psi, \varphi) = \arccos \left(\int_0^1 \sqrt{\psi'} \sqrt{\varphi'} d\theta \right) \tag{C.11}$$

Now define the functions

$$\varphi_n : \begin{cases} 0 & 0 \leq \theta \leq \frac{1}{n} \\ 2x & \frac{1}{n} \leq \theta \leq \frac{2}{n} \\ x & \frac{2}{n} \leq \theta \leq 1 \end{cases} \quad \psi_n : \begin{cases} nx & 0 \leq \theta \leq \frac{1}{n} \\ 1 & \frac{1}{n} \leq \theta \leq 1 \end{cases} \tag{C.12}$$

The functions φ_n and ψ_n are not diffeomorphisms, but we can smooth them with an arbitrarily small change to the \dot{H}^1 distances considered. The claim now follows by a straightforward calculation. □

References

1. Amari, S., Nagaoka, H.: *Methods of Information Geometry*, vol. 191. American Mathematical Society, Providence (2007)
2. Arnold, V.: Sur la géométrie différentielle des groupes de Lie de dimension infinie et ses applications à l'hydrodynamique des fluides parfaits. *Ann. Inst. Fourier* **16**, 319–361 (1966)
3. Arnold, V.I., Khesin, B.A.: *Topological Methods in Hydrodynamics*, vol. 125. Springer, Berlin (1999)
4. Banyaga, A.: *The Structure of Classical Diffeomorphism Groups*, vol. 400. Springer, Berlin (1997)
5. Bauer, M., Bruveris, M., Harms, P., Michor, P.W.: Geodesic distance for right invariant Sobolev metrics of fractional order on the diffeomorphism group. *Ann. Glob. Anal. Geom.* **44**(1), 5–21 (2013)
6. Bauer, M., Bruveris, M., Michor, P.W.: Geodesic distance for right invariant Sobolev metrics of fractional order on the diffeomorphism group II. *Ann. Glob. Anal. Geom.* **44**(4), 361–368 (2013)
7. Bauer, M., Bruveris, M., Michor, P.W.: Overview of the geometries of shape spaces and diffeomorphism groups. *J. Math. Imaging Vis.* **50**(1–2), 60–97 (2014)
8. Bauer, M., Escher, J., Kolev, B.: Local and global well-posedness of the fractional order EPDiff equation on \mathbb{R}^d . *J. Differ. Equ.* **258**(6), 2010–2053 (2015)
9. Bauer, M., Harms, P., Preston, S.C.: Vanishing distance phenomena and the geometric approach to SQG. *Arch. Ration. Mech. Anal.* **235**, 1445–1466 (2020)
10. Bauer, M., Joshi, S., Modin, K.: Diffeomorphic density matching by optimal information transport. *SIAM J. Imaging Sci.* **8**(3), 1718–1751 (2015)

11. Bauer, M., Kolev, B., Preston, S.C.: Geometric investigations of a vorticity model equation. *J. Differ. Equ.* **260**(1), 478–516 (2016)
12. Beg, M.F., Miller, M.I., Trounev, A., Younes, L.: Computing large deformation metric mappings via geodesic flows of diffeomorphisms. *Int. J. Comput. Vis.* **61**(2), 139–157 (2005)
13. Behzadan, A., Holst, M.: Multiplication in Sobolev spaces, revisited. [arXiv:1512.07379](https://arxiv.org/abs/1512.07379) (2015)
14. Brandenbursky, M., Shelukhin, E.: The L^p -diameter of the group of area-preserving diffeomorphisms of S^2 . *Geom. Topol.* **21**(6), 3785–3810 (2017)
15. Brasco, L., Salort, A.: A note on homogeneous Sobolev spaces of fractional order. *Ann. Mate. Pura Appl.* (1923-) **198**, 1295–1330 (2019)
16. Bruveris, M., Vialard, F.-X.: On completeness of groups of diffeomorphisms. *J. Eur. Math. Soc.* **19**(5), 1507–1544 (2017)
17. Burago, D., Ivanov, S., Polterovich, L.: Conjugation-invariant norms on groups of geometric origin. In: *Groups of Diffeomorphisms: in Honor of Shigeyuki Morita on the Occasion of his 60th Birthday*, pp. 221–250. Mathematical Society of Japan (2008)
18. Camassa, R., Holm, D.D.: An integrable shallow water equation with peaked solitons. *Phys. Rev. Lett.* **71**(11), 1661–1664 (1993)
19. Constantin, P., Lax, P.D., Majda, A.: A simple one-dimensional model for the three-dimensional vorticity equation. *Commun. Pure Appl. Math.* **38**(6), 715–724 (1985)
20. Cotter, C.J., Deasy, J., Pryer, T.: The r -Hunter–Saxton equation, smooth and singular solutions and their approximation. *Nonlinearity* **33**(12), 7016 (2020)
21. Ebin, D.G., Marsden, J.: Groups of diffeomorphisms and the motion of an incompressible fluid. *Ann. Math.* **92**(1), 102–163 (1970)
22. Eichhorn, J.: *Global Analysis on Open Manifolds*. Nova Science Publishers Inc., New York (2007)
23. Eliashberg, Y., Polterovich, L.: Biinvariant metrics on the group of Hamiltonian diffeomorphisms. *Int. J. Math.* **4**(05), 727–738 (1993)
24. Eliashberg, Y., Ratiu, T.: The diameter of the symplectomorphism group is infinite. *Invent. Math.* **103**(1), 327–340 (1991)
25. Escher, J., Kolev, B., Wunsch, M.: The geometry of a vorticity model equation. *Commun. Pure Appl. Anal.* **11**(4), 1407–1419 (2012)
26. Fukui, K., Rybicki, T., Yagasaki, T.: The uniform perfectness of diffeomorphism groups of open manifolds. [arXiv:1905.07664](https://arxiv.org/abs/1905.07664)
27. Grenander, U.: *General Pattern Theory: A Mathematical Study of Regular Structures*. Clarendon Press, Oxford (1993)
28. Grenander, U., Miller, M.I.: Computational anatomy: an emerging discipline. *Q. Appl. Math.* **56**(4), 617–694 (1998)
29. Han, W., Atkinson, K., Zheng, H.: Some integral identities for spherical harmonics in an arbitrary dimension. *J. Math. Chem.* **50**(5), 1126–1135 (2012)
30. Hofer, H.: On the topological properties of symplectic maps. *Proc. R. Soc. Edinb. Sect. A Math.* **115**(1–2), 25–38 (1990)
31. Hunter, J.K., Saxton, R.: Dynamics of director fields. *SIAM J. Appl. Math.* **51**(6), 1498–1521 (1991)
32. Jerrard, R.L., Maor, C.: Geodesic distance for right-invariant metrics on diffeomorphism groups: critical Sobolev exponents. *Ann. Glob. Anal. Geom.* **56**(2), 351–360 (2019)
33. Jerrard, R.L., Maor, C.: Vanishing geodesic distance for right-invariant Sobolev metrics on diffeomorphism groups. *Ann. Glob. Anal. Geom.* **55**(4), 631–656 (2019)
34. Joshi, S.C., Miller, M.I.: Landmark matching via large deformation diffeomorphisms. *IEEE Trans. Image Process.* **9**(8), 1357–1370 (2000)
35. Khesin, B., Lenells, J., Misiolek, G., Preston, S.: Curvatures of Sobolev metrics on diffeomorphism groups. *Pure Appl. Math. Q.* **9**(2), 291–332 (2013)
36. Khesin, B., Lenells, J., Misiolek, G., Preston, S.: Geometry of diffeomorphism groups, complete integrability and geometric statistics. *Geom. Funct. Anal.* **23**(1), 334–366 (2013)
37. Khesin, B., Misiolek, G.: Euler equations on homogeneous spaces and Virasoro orbits. *Adv. Math.* **176**(1), 116–144 (2003)
38. Khesin, B., Wendt, R.: *The Geometry of Infinite-Dimensional Groups*, vol. 51. Springer, Berlin (2008)
39. Kolev, B.: Local well-posedness of the EPDiff equation: a survey. *J. Geom. Mech.* **9**(2), 167–189 (2017)
40. Kouranbaeva, S.: The Camassa–Holm equation as a geodesic flow on the diffeomorphism group. *J. Math. Phys.* **40**(2), 857–868 (1999)
41. Kriegl, A., Michor, P.W.: *The Convenient Setting of Global Analysis*. American Mathematical Society, Providence (1997)
42. Lenells, J.: The Hunter–Saxton equation describes the geodesic flow on a sphere. *J. Geom. Phys.* **57**(10), 2049–2064 (2007)

43. Lenells, J.: The Hunter–Saxton equation: a geometric approach. *SIAM J. Math. Anal.* **40**(1), 266–277 (2008)
44. Leslie, J.: On a differential structure for the group of diffeomorphisms. *Topology* **6**, 263–271 (1967)
45. Leoni, G.: *A First Course in Sobolev Spaces*, 2nd edn. American Mathematical Society, Providence (2017)
46. Mann, K., Rosendal, C.: Large-scale geometry of homeomorphism groups. *Ergod. Theory Dyn. Syst.* **38**(7), 2748–2779 (2018)
47. Michor, P.W., Mumford, D.: Vanishing geodesic distance on spaces of submanifolds and diffeomorphisms. *Doc. Math.* **10**, 217–245 (2005)
48. Miller, M.I., Trounev, A., Younes, L.: On the metrics and Euler-Lagrange equations of computational anatomy. *Ann. Rev. Biomed. Eng.* **4**(1), 375–405 (2002)
49. Misiołek, G.: A shallow water equation as a geodesic flow on the Bott–Virasoro group. *J. Geom. Phys.* **24**(3), 203–208 (1998)
50. Misiołek, G., Preston, S.C.: Fredholm properties of Riemannian exponential maps on diffeomorphism groups. *Invent. Math.* **179**(1), 191 (2010)
51. Mumford, D., Desolneux, A.: *Pattern Theory: the Stochastic Analysis of Real-World Signals*. AK Peters/CRC Press, Boca Raton (2010)
52. Peetre, J.: Espaces d’interpolation et théorème de Soboleff. *Ann. Inst. Fourier* **16**, 279–317 (1966)
53. Salo, M.: Function spaces—lecture notes. http://users.jyu.fi/~salomi/lecturenotes/fsp08_lectures.pdf (2008)
54. Shelukhin, E.: The Hofer norm of a contactomorphism. *J. Symp. Geom.* **15**(4), 1173–1208 (2017)
55. Shnirelman, A.I.: On the geometry of the group of diffeomorphisms and the dynamics of an ideal incompressible fluid. *Math. USSR-Sbornik* **56**(1), 79 (1987)
56. Shnirelman, A.I.: Generalized fluid flows, their approximation and applications. *Geom. Funct. Anal. GAFA* **4**(5), 586–620 (1994)
57. Tığlay, F., Vizman, C.: Generalized Euler–Poincaré equations on Lie groups and homogeneous spaces, orbit invariants and applications. *Lett. Math. Phys.* **97**(1), 45–60 (2011)
58. Triebel, H.: *Interpolation Theory, Function Spaces, Differential Operators*. North-Holland, Amsterdam (1978)
59. Triebel, H.: *Theory of Function Spaces II*, volume 84 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel (1992)
60. Trounev, A., Younes, L.: Local geometry of deformable templates. *SIAM J. Math. Anal.* **37**(1), 17–59 (2005). ((**electronic**))
61. Tsuboi, T.: On the uniform perfectness of diffeomorphism groups. In: *Groups of Diffeomorphisms: in Honor of Shigeyuki Morita on the Occasion of his 60th Birthday*, pp. 505–524. Mathematical Society of Japan (2008)
62. Wunsch, M.: On the geodesic flow on the group of diffeomorphisms of the circle with a fractional Sobolev right-invariant metric. *J. Nonlinear Math. Phys.* **17**(1), 7–11 (2010)
63. Younes, L.: *Shapes and Diffeomorphisms*, vol. 171. Springer, Berlin (2010)