

# Two-stage partial melting during the Variscan extensional tectonics (Montagne Noire, France)

Marc Poujol<sup>1</sup> · Pavel Pitra<sup>1</sup> · Jean Van Den Driessche<sup>1</sup> · Romain Tartèse<sup>2,3</sup> · Gilles Ruffet<sup>1</sup> · Jean-Louis Paquette<sup>4</sup> · Jean-Charles Poilvet<sup>1</sup>

Received: 2 December 2015 / Accepted: 28 June 2016 / Published online: 13 July 2016  
© Springer-Verlag Berlin Heidelberg 2016

**Abstract** One of the striking features that characterise the late stages of the Variscan orogeny is the development of gneiss and migmatite domes, as well as extensional Late Carboniferous and Permian sedimentary basins. It remains a matter of debate whether the formation of domes was related to the well-documented late orogenic extension or to the contractional tectonics that preceded. Migmatization and magmatism are expected to predate extension if the domes are compression-related regional anticlines, but they must both precede and be contemporaneous with extension if they are extensional core complexes. In the Montagne Noire area (southern French Massif Central), where migmatization, magmatism and the deformation framework are well documented, the age of the extensional event was unequivocally constrained to 300–290 Ma. Therefore, dating migmatization in this area is a key point for discriminating between the two hypotheses and understanding the

Late Palaeozoic evolution of this part of the Variscan belt. For this purpose, a migmatite and an associated anatectic granite from the Montagne Noire dome were dated by LA-ICP-MS (U–Th–Pb on zircon and monazite) and laser probe <sup>40</sup>Ar–<sup>39</sup>Ar (K–Ar on muscovite). Although zircon did not record any Variscan age unequivocally related to compression (380–330 Ma), two age groups were identified from the monazite crystals. A first event, at ca. 319 Ma (U–Th–Pb on monazite), is interpreted as a first stage of migmatization and as the emplacement age of the granite, respectively. A second event at ca. 298–295 Ma, recorded by monazite (U–Th–Pb) and by the muscovite <sup>40</sup>Ar–<sup>39</sup>Ar system in the migmatite and in the granite, could be interpreted as a fluid-induced event, probably related to a second melting event identified through the syn-extensional emplacement of the nearby Montalet leucogranite ca. 295 Ma ago. The ages of these two events post-date the Variscan compression and agree with an overall extensional context for the development of the Montagne Noire dome-shaped massif. Comparison of these results with published chemical (EPMA) dating of monazite from the same rocks demonstrates that the type of statistical treatment applied to EPMA data is crucial in order to resolve different monazite age populations.

**Electronic supplementary material** The online version of this article (doi:10.1007/s00531-016-1369-1) contains supplementary material, which is available to authorized users.

✉ Marc Poujol  
marc.poujol@univ-rennes1.fr

<sup>1</sup> Géosciences Rennes, UMR CNRS 6118, OSUR, Université Rennes 1, 35042 Rennes Cedex, France

<sup>2</sup> Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie, Muséum National d'Histoire Naturelle, Sorbonne Universités, CNRS, UPMC & IRD, 75005 Paris, France

<sup>3</sup> Planetary and Space Sciences, The Open University, Walton Hall, Milton Keynes MK7 6AA, UK

<sup>4</sup> UMR CNRS 6524, Laboratoire Magmas et Volcans, Université Blaise Pascal, 63038 Clermont-Ferrand Cedex, France

**Keywords** Monazite · LA-ICP-MS · U–Th–Pb dating · Muscovite · <sup>40</sup>Ar–<sup>39</sup>Ar dating · Variscan · Montagne Noire

## Introduction

Until the late 1980s, wrench tectonics was considered to control the late Palaeozoic tectonic evolution of the Variscan orogen (Arthaud and Matte 1977). It was interpreted as corresponding to an ultimate phase of N–S compression during the Late Carboniferous, and marked the end of the

Variscan continental collision. In such compressive context, the commonly coal-bearing continental sedimentary basins that developed during the very late Carboniferous (305–295 Ma) were interpreted as pull-apart basins or as related to horsetail splay faults at the termination of strike-slip systems faults (e.g. Arthaud and Matte 1977; Blès et al. 1989). The development of widespread continental sedimentary basins during the Permian was attributed to a subsequent pervasive N–S extension that occurred throughout the Variscan domain and was considered to result from a plate kinematics re-arrangement without any causal relation with the previous Variscan continental collision (e.g. Arthaud and Matte 1977; Blès et al. 1989).

More recently, the late Palaeozoic tectonic evolution has been compared to extensional tectonics of both the Tibetan plateau and the North American Cordillera during the Cenozoic (e.g. Ménard and Molnar 1988; Burg et al. 1994). E–W extension, nearly parallel to the belt took place between 330 and 305 Ma during escape tectonics driven by still active N–S compression forces. It was followed by a NE–SW to N–S extension between 300 and 260 Ma, which started during the waning shortening, and implies a radical change in extension direction induced by the modification of the boundary conditions and the collapse of the entire chain after continental convergence (e.g. Burg et al. 1994). Both episodes of extension are viewed as a consequence of the previous N–S shortening, extension being induced by the collapse of the crust, considerably thickened during the continental collision, after thermal relaxation. In this interpretation, and contrary to older “compressive” interpretations, the late Carboniferous basins are extensional in origin as well as the Permian basins (Becq-Giraudon and Van Den Driessche 1993).


In both interpretations, the pervasive high-temperature low-pressure (HT–LP) metamorphism and magmatism that occurred throughout the Variscan chain during the late Carboniferous resulted from crustal thickening. A striking feature of this period was the development of gneiss and migmatite domes, such as those found in the Montagne Noire and the Velay areas in the southern part of the French Massif Central. However, the two interpretations disagree on the cause of their development. In the compressive scenario, these structures are interpreted as regional anticlines that developed in response to the N–S shortening (Arthaud et al. 1966; Burg and Matte 1978; Matte et al. 1998), whereas they are interpreted as extensional gneiss domes, similar to the Cenozoic metamorphic core complexes of the Basin and Range province, in the extensional scenario (Van Den Driessche and Brun 1989, 1992; Echtler and Malavieille 1990; Brun and Van Den Driessche 1994). In the first case, migmatization and magmatism predate extension. The second interpretation requires thermal relaxation and related rheological softening, and change in boundary

conditions in order for the crust to collapse. Consequently, migmatization both precedes and is contemporaneous with the onset of extension, especially because extension can enhance partial melting by adiabatic decompression. In the Montagne Noire gneiss dome, the age of the extensional event was unequivocally constrained to ca. 295 Ma by dating a syntectonic leucogranite emplaced within the northward-dipping normal fault that bounds the gneiss dome to the north and controlled the development of Upper Carboniferous–Lower Permian sedimentary basins (Poilvet et al. 2011).

The Montagne Noire gneiss dome is a typical case where geochronology can yield critical constraints on tectonic models and where the precision obtained on each individual age is crucial. This becomes even more important in a region that underwent a complex polyphased metamorphic/magmatic history possibly characterised by several phases of mineral growth (involving datable minerals such as zircon, monazite and muscovite) in a relatively short period of time. For the purpose of this study, two samples (a migmatite and an associated granite) from the Montagne Noire dome were selected for geochronological investigations (U–Th–Pb on monazite and zircon and K–Ar on muscovite), in order to discriminate between the two contradictory tectonic interpretations.

## Geological setting

The Montagne Noire gneiss–migmatite massif is located in the southern French Massif Central (Fig. 1). It is composed of a high-grade gneissic core surrounded by mostly low-grade metasediments. The gneissic core is composed of migmatites and augen orthogneisses, with some fine-grained gneissic intercalations that have been interpreted as either metasediments or mylonitic zones (Bogdanoff et al. 1984; Van Den Driessche and Brun 1992, and references therein). Some of these intercalations contain HP/HT mafic and ultramafic metamorphic rocks, suggesting possible major early tectonic contacts (Bogdanoff et al. 1984; Alabouvette and Demange 1993; Demange et al. 1995). The migmatites resulted mostly from partial melting of sediments, but also of felsic augen orthogneisses (e.g. Bogdanoff et al. 1984; Demange 1982). Weakly deformed to undeformed anatectic granites intrude both the orthogneisses and the migmatites. The gneissic core is surrounded by weakly metamorphosed or unmetamorphosed lower to middle Palaeozoic sediments that are intensely deformed by southward verging folds and thrust faults (e.g. Arthaud 1970; Bogdanoff et al. 1984; Echtler 1990). Late Carboniferous to Early Permian detrital sediments unconformably overlie these tectonic units to the south. In contrast, to the North, the EW-trending north-dipping normal fault



**Fig. 1** Structural map of the southern French Massif Central (MC) showing the relationships between the Montagne Noire gneiss dome, the Stephanian-Permian basins, and the Variscan thrusts and nappes (modified after Brun and Van Den Driessche 1994). 7, 8: locations of samples ES7 and ES8, respectively; Af: St. Affrique basin, G Graissessac basin, L Laouzais granite, Lo Lodève basin, P Col de Picotalen (location of the sample of the Montalet syntectonic leucogranite,

ES5, Poilvet et al. 2011), V Vialais granite. *Inset* shows the location of the study area within the European Variscan belt (modified from Pitra et al. 2010). A Alps, AM Armorican Massif, BM Bohemian Massif, MC Massif Central. B Teplá-Barrandian, Mo Moldanubian, ST Saxothuringian, RH Rhenohercynian. L Lyon, M Montpellier, R Rennes

zone (Espinouse detachment) marks the tectonic contact between these sediments and the core units, and controls the development of the Lodève-Graissessac and Saint-Affrique basins (Fig. 1; Van Den Driessche and Brun 1989, 1992; Burg et al. 1994).

The foliation of the gneissic core developed within the lower crust during the thrusting event responsible for the deformation of the lower to middle Palaeozoic sedimentary cover. Its dome-shaped structure and tectonic evolution is still a matter of debate. Three types of models have been proposed ranging from a diapir (e.g. Schuiling 1960; Faure and Cottureau 1988), a double megafold (Arthaud 1970; Burg and Matte 1978; Bogdanoff et al. 1984) to a core complex (Van den Driessche and Brun 1989; Echtler and Malavieille 1990). Many recent models combine in a more or less complex way the processes responsible for these three types of structures to explain the final structure of the Montagne Noire: diapirism coeval with compression (Faure et al. 2010), compression and subsequent extension

(e.g. Cassard et al. 1993; Franke et al. 2011; Doublier et al. 2015; Rabin et al. 2015), compression during extension (Rey et al. 2011), or compression, diapirism and extension (e.g. Soula et al. 2001; Charles et al. 2009). The precise structure of the Montagne Noire dome-shaped massif is beyond the scope of the present paper. We just note that structural and metamorphic analyses have not allowed to reach a consensus, although these different models are built from mostly similar (especially structural) data (e.g. Rey et al. 2011; Van Den Driessche and Pitra 2012). We conclude that until now, discriminating data are lacking to arbitrate between these models.

On the scale of the Variscan belt, the tectonic origin of the Montagne Noire massif is emblematic of the two end-member interpretations discussed before (i.e. compressive or extensive). Eventually, the two main questions are the timing of (1) the development of the HT–LP metamorphism, including partial melting and (2) the initiation of the extensional tectonics, which predominates during Permian times.

Previous geochronological works on the protolith of the augen orthogneisses yielded Ordovician U–Pb zircon ages ( $456 \pm 3$  for Pont-de-Larn,  $450 \pm 6$  Ma for the Gorges d’Héric, Roger et al. 2004, 2015;  $455 \pm 2$  Ma for the Saint-Eutrope gneiss, Pitra et al. 2012). According to Faure et al. (2010), migmatization took place between 333 and 326 Ma (EPMA dating on monazite) while the emplacement of late anatectic granitoids took place between 325 and 316 Ma (including the Vialais granite at  $320 \pm 3$  Ma and Montalet leucogranite at ca. 330 Ma, Fig. 1). However, a recent study by Roger et al. (2015) documented an emplacement age of ca 303 Ma (U–Th–Pb on monazite) for the Vialais granite, while the Montalet leucogranite yielded monazite and zircon U–Th–Pb emplacement ages of ca 294 Ma (Poilvet et al. 2011). Three monazite grains from the Gorges d’Héric orthogneiss yielded  $^{206}\text{Pb}/^{238}\text{U}$  ID-TIMS dates around 310 Ma interpreted as a metamorphic age (Roger et al. 2015). Franke et al. (2011) reported a similar age of 313 Ma for monazite extracted from a foliated aplite dyke from the Gorges d’Héric, while Maluski et al. (1991) reported a biotite  $^{40}\text{Ar}$ – $^{39}\text{Ar}$  plateau age of  $316 \pm 4$  Ma for the Caroux massif. The undeformed garnet-bearing leucogranite of Ourtigas yielded a U–Th–Pb age of ca 298 Ma (Roger et al. 2015). Finally, a monazite Th–Pb age of  $294.4 \pm 4$  Ma (Pitra et al. 2012) as well as  $^{40}\text{Ar}$ – $^{39}\text{Ar}$  ages on muscovite and biotite of ca. 297 Ma (Maluski et al. 1991) have been obtained for orthogneiss samples sheared along the Espinouse detachment.

## Sampling and petrography

Two samples were selected for this study (Fig. 1): a cordierite-bearing granite located in the central part of the dome (sample ES7), and a migmatite (sample ES8), spatially associated with the cordierite-bearing granite. Both rocks were sampled at the same locations as their equivalent dated by Faure et al. (2010). One of the main reasons for this sampling strategy is linked to the fact that previous dating (EPMA on monazite) by Faure et al. (2010) on the Montalet granite returned an age  $327 \pm 7$  Ma that is very different from the age  $294 \pm 1$  Ma (U–Th–Pb on monazite) obtained by Poilvet et al. (2011) on the same granite. This age difference was also noticed by Roger et al. (2015) for the Vialais granite. Therefore, the age discrepancies cast some doubts on the EPMA ages obtained by Faure et al. (2010) on the Laouzas cordierite-bearing granite and the spatially associated migmatite. They have also been chosen because of the potentially complex history that these rocks underwent between 340 and 290 Ma.

### Laouzas granite, sample ES7

The Laouzas granite (Fig. 1) crops out in the west-central part of the axial zone of the Montagne Noire dome.


It was sampled near the Laouzas dam ( $43^{\circ}38'7.35''\text{N}$ ,  $2^{\circ}45'10.00''\text{E}$ ). The rock is an undeformed heterogeneous coarse-grained (1–5 mm in average, locally up to 3 cm) biotite-bearing granite containing numerous large clusters of cordierite (up to 5 cm), biotite-rich schlieren, tourmaline nodules and dark, foliated mica-rich enclaves (Fig. 2a). Although the schlieren and enclaves locally display a preferred orientation, no solid-state deformation is observed at the grain scale in the granite, with the exception of a weak undulose extinction of quartz crystals.

The sample is dominated by plagioclase, K-feldspar and quartz, and contains subordinate amounts of biotite, cordierite and muscovite (Fig. 2b). Dumortierite, tourmaline, andalusite and sillimanite are present locally. Plagioclase forms euhedral to subhedral stubby prismatic crystals, 1–3 mm long. Plagioclase cores are generally altered (saussuritised) and surrounded by a clear rim (Fig. 2c). The rims are similar to feldspar that also fills fractures in the plagioclase cores. Some plagioclase rims are intergrown with tiny crystals of quartz in a granophyric, myrmekite-like texture, in particular at the contact with K-feldspar. K-feldspar and quartz are anhedral, 1–5 mm in size. Cordierite forms anhedral crystals (up to 2 mm) that are slightly pinitised or replaced by fine-grained muscovite around the rims (Fig. 2b). Biotite crystals (0.1–3 mm) are subhedral and locally partly altered to chlorite in association with needles of rutile (sagenite). Rare minute needles of sillimanite are locally present in large quartz crystals. Although euhedral muscovite crystals (~1 mm) are locally present, muscovite mostly forms large subhedral poikiloblasts (up to 2 mm) or develops tiny crystals at the expense of K-feldspar, plagioclase or cordierite. Needles of dumortierite (pink to violet, strongly pleochroic acicular crystals, ~0.5 mm) or tourmaline (pale green, ~0.3 mm) are commonly associated with clear plagioclase overgrowths, anhedral pink andalusite (~0.5 mm) and muscovite (Fig. 2c).

These observations suggest a two-stage evolution. The crystallisation of the relatively coarse-grained granite (plagioclase cores, K-feldspar, quartz, biotite, cordierite,  $\pm$  muscovite  $\pm$  sillimanite) was followed by a second event resulting in the crystallisation of the plagioclase rims, dumortierite, tourmaline, andalusite and muscovite. The second stage was possibly associated with the alteration of the plagioclase cores and biotite. It could be either magmatic or, more probably, hydrothermal in origin. The hydrothermal origin is supported by the textures and in particular the position of the boron-bearing minerals. Indeed, tourmaline and dumortierite are known to be related to hydrothermal alteration (e.g. Taner and Martin 1993), but are liquidus rather than late-stage phases in leucogranitic magmatic systems (Benard et al. 1985).



**Fig. 2** **a** Outcrop photograph of the Laouzas granite (ES7). *Dark spots* are clusters of cordierite; the elongated object in the lower central part is a biotite-rich schlier. **b, c** Microphotographs of the granite. Note the presence of subhedral, partly altered crystals of plagioclase (pl) and K-feldspar (kfs), partly pinitised crystals of cordierite (cd), and late crystals of dumortierite (dum), andalusite (and) and muscovite (mu), and the clear rim around altered plagioclase core. **d** Outcrop photograph of the banded La Salvetat migmatite (ES8). Rare garnet crystals are shown by *arrows*. **e, f** Microphotographs of the migmatite leucosome. Interstitial brown-blue tourmaline is located between subhedral, partly altered crystals of feldspars and anhedral quartz (e). Plagioclase crystals are partly replaced by clinozoisite (tiny high-relief crystals), biotite contains prehnite (prh) and/or clay minerals parallel to the (001) cleavage; late muscovite is also present (f)



### La Salvetat migmatite, sample ES8

The La Salvetat migmatite crops out in the central part of the axial zone of the Montagne Noire gneiss dome, generally to the south of the Laouzas granite (Fig. 1). It was sampled ca. 2 km south of the Laouzas dam, close to the summit of a hill called Al Rec del Bosc (43°37'1.65"N, 2°45'23.70"E). The rock is a banded stromatic migmatite (Fig. 2d) and is locally garnet-bearing. The foliation is defined by a weak preferred orientation of biotite that is parallel to alternating layers of leucosome, mesosome and biotite-dominated melanosome. Leucosomes are relatively coarse-grained (0.5–3 mm), up to 1 cm thick, and are composed of subhedral plagioclase, quartz and subordinate K-feldspar, and locally contain cordierite, muscovite, biotite and tourmaline (Fig. 2e). Tourmaline commonly forms anhedral to subhedral interstitial, optically zoned grains. Plagioclase and K-feldspar have a dusty appearance, and plagioclase is partly replaced

by zoisite/clinozoisite and white mica (Fig. 2f). Cordierite is subhedral and partly altered to pinitite. Biotite is locally partly replaced by chlorite and contains lenses of minute prehnite (?) or clay minerals that are parallel to the (001) cleavage (Fig. 2f), in particular in the leucosome and in the adjacent melanosome. Euhedral muscovite is locally present in the leucosome. Subhedral muscovite, forming either fine-grained clusters or larger poikilitic crystals, commonly develops at the expense of biotite, cordierite and feldspars in the leucosomes.

These observations suggest at least two stages of evolution, where the migmatite textures, attributable to partial melting, are partly overprinted by hydrothermal alteration. The geographic proximity and the common petrographic character of the leucosomes and the Laouzas granite suggest a genetic relation between the two. Therefore, the La Salvetat migmatite is interpreted to be a likely source for the anatectic Laouzas granite.

## U–Th–Pb LA-ICP-MS and $^{40}\text{Ar}$ – $^{39}\text{Ar}$ dating

### Analytical techniques

#### *NanoSIMS analytical protocol*

The distribution of selected species was imaged in some monazite grains using the NanoSIMS 50 ion probe at the University of Rennes 1. Secondary ion images of  $^{89}\text{Y}$ ,  $^{139}\text{La}$ ,  $^{140}\text{Ce}$ ,  $^{149}\text{Sm}$ ,  $^{206}\text{Pb}$ ,  $^{208}\text{Pb}$ ,  $^{232}\text{Th}$  and  $^{238}\text{U}^{16}\text{O}$  were collected using the same primary O-beam of  $\sim 170$  pA over  $90\ \mu\text{m} \times 90\ \mu\text{m}$  areas. The mass resolving power was set to  $\sim 3500$ , sufficient to readily resolve isobaric interferences such as  $^{143,145}\text{NdPO}_2$  on  $^{206,208}\text{Pb}$ . A single plane of image data, divided in  $128\ \text{px} \times 128\ \text{pixels}$ , was collected using a dwell time of 40 ms/pixel, representing an acquisition time of  $\sim 11$  min for each area. The raw image data were processed using the L'image (Larry Nittler, Carnegie Institute of Washington, USA; [http://home.dtm.ciw.edu/users/nittler/limage/limage\\_manual.pdf](http://home.dtm.ciw.edu/users/nittler/limage/limage_manual.pdf)) and ImageJ softwares.

#### *U–Th–Pb dating technique*

A classic mineral separation procedure has been applied to concentrate minerals suitable for U–Th–Pb dating using the facilities available at Géosciences Rennes (see Poilvet et al. 2011). Zircon and monazite grains were carefully handpicked under a binocular microscope and embedded in epoxy mounts. The grains were then polished on a lap wheel with a 6- and  $1\text{-}\mu\text{m}$  diamond suspension successively. Zircon grains were imaged by cathodoluminescence (CL) using a Reliotron CL system equipped with a digital colour camera available in Géosciences Rennes and monazite grains by backscattered electron imaging on a JEOL JSM 6400 as well as by NanoSIMS. In addition, monazite grains have also been identified in thin sections in order to date them in context.

U–Th–Pb geochronology of zircon and monazite was conducted by in situ laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) at the Laboratoire Magmas et Volcans in Clermont-Ferrand, France. Ablation spot diameters of  $26\ \mu\text{m}$  and  $7\ \mu\text{m}$  with repetition rates of 3 Hz and 1 Hz were used for zircon and monazite, respectively. Data were corrected for U–Pb and Th–Pb fractionation and for the mass bias by standard bracketing with repeated measurements of the 91500 zircon (Wiedenbeck et al. 1995) or the Moacyr monazite standards (Gasquet et al. 2010). Repeated analyses of GJ-1 zircon ( $607 \pm 15$  Ma,  $N = 6$ ; Jackson et al. 2004) or Manangoutry monazite ( $554 \pm 23$  Ma,  $N = 6$ ; Paquette and Tiepolo 2007) standards treated as unknowns were used to control the reproducibility and accuracy of the corrections. Data reduction was carried out with the GLITTER<sup>®</sup> software

package developed by the Macquarie Research Ltd. (Jackson et al. 2004). Concordia ages and diagrams were generated using Isoplot/Ex (Ludwig 2001). All errors given in Tables 1 and 2 are listed at one sigma, but where data are combined for regression analysis or to calculate weighted means, the final results are provided with 95 % confidence limits. Further information on the instrumentation and the analytical technique is detailed in Hurai et al. (2010).

For each grain analysed (zircon and monazite), we also estimated the concentrations of U, Th and Pb as follows. First, the drift factor was calculated using the parameter  $a$  and  $b$  of a linear regression of the average  $^{206}\text{Pb}$  counts per second (cps) for all the standards measured during the course of the analyses as a function of their position during the acquisition.

$$^{206}\text{Pb}_{\text{cps}} = a \times N + b \quad (1)$$

with  $^{206}\text{Pb}_{\text{cps}}$  = average measured values for the standard in cps corrected from the blank,  $a$  = slope of the regression,  $b$  = ordinate at origin and  $N$  = analysis number.

The drift factor for each analysis ( $\text{DF}^N$ ) is then calculated as follows:

$$\text{DF}^N = \frac{b}{(a \times N + b)} \quad (2)$$

Then the Pb, Th and U concentrations are calculated using the known concentrations of these elements in the standards following:

$$\text{C}_{\text{ppm}}^N = \text{C}_{\text{meas}}^N \times \text{DF}^N \times \frac{\text{C}_{\text{std}_{\text{real}}}}{\text{C}_{\text{std}_{\text{aver}}}} \quad (3)$$

with  $N$  = analysis number,  $\text{C}_{\text{ppm}}^N$  = calculated concentration of the element in ppm,  $\text{C}_{\text{meas}}^N$  = measured values of the element in cps,  $\text{DF}^N$  = drift factor calculated for this analysis,  $\text{C}_{\text{std}_{\text{real}}}$  = known concentration of the standard and  $\text{C}_{\text{std}_{\text{aver}}}$  = drift-corrected average value for all the standards measured during the course of the analyses.

#### *Ar–Ar dating technique*

Single grains of muscovite used for the experiments were handpicked under a binocular microscope from 0.25 to 1.00 mm fractions of crushed rock samples. Care was taken to select inclusion-free crystals of about 1 mm in size, in order to avoid large poikilitic crystals as well as fine-grained clusters.

The samples were wrapped in Al foil to form small packets ( $11 \times 11$  mm) that were stacked up to form a pile within which packets of fluence monitors were inserted every 10 samples. Irradiation was performed at the HFR Petten reactor (Petten, the Netherlands) in the Cd-shielded Rodeo P3 facility and lasted 72 h ( $J/h \approx 2.54 \times 10^{-4}\ \text{h}^{-1}$ ). The irradiation standard was amphibole Hb3gr (Turner

**Table 1** U–Th–Pb LA-ICP-MS data for the zircon grains from sample ES7 and ES8

Grain (zircon)	[Pb] (ppm)	[U] (ppm)	Th/U	$^{207}\text{Pb}/^{235}\text{U}$	$\pm 1\sigma$	$^{206}\text{Pb}/^{238}\text{U}$	$\pm 1\sigma$	$^{207}\text{Pb}/^{206}\text{Pb}$	$\pm 1\sigma$	Rho	Ages		Conc. (%)		
											$^{207}\text{Pb}/^{235}\text{U}$	$^{206}\text{Pb}/^{238}\text{U}$	$^{207}\text{Pb}/^{206}\text{Pb}$	$\pm 1\sigma$	
<i>ES 7 Laouzas granite</i>															
1.2c	31	509	0.19	0.705	0.009	0.0736	0.0008	0.0695	0.0008	0.88	542	458	913	25	50
2.1c	39	376	0.43	1.166	0.014	0.1158	0.0013	0.0730	0.0008	0.92	785	707	1014	23	70
2.2c	104	2810	0.01	0.356	0.004	0.0480	0.0005	0.0539	0.0006	0.95	310	302	367	25	82
3c	110	2469	0.03	0.443	0.005	0.0570	0.0006	0.0563	0.0006	0.94	372	358	463	24	77
4.1c	102	988	0.85	1.028	0.012	0.1039	0.0011	0.0718	0.0008	0.94	718	637	979	22	65
4.2c	99	2374	0.02	0.438	0.005	0.0527	0.0006	0.0603	0.0007	0.94	369	331	613	24	54
6.1c	10	90	0.50	1.151	0.017	0.1255	0.0014	0.0665	0.0009	0.78	778	762	823	29	93
6.2c	15	212	0.17	0.723	0.010	0.0891	0.0010	0.0588	0.0008	0.84	552	550	561	27	98
7.2c	73	1834	0.02	0.387	0.005	0.0520	0.0006	0.0540	0.0006	0.92	332	327	371	25	88
8.1c	62	501	0.44	1.378	0.016	0.1413	0.0016	0.0708	0.0008	0.92	880	852	950	23	90
8.2c	29	317	0.20	0.958	0.012	0.1107	0.0012	0.0627	0.0008	0.88	682	677	699	25	97
10c	61	834	0.22	0.746	0.009	0.0880	0.0010	0.0615	0.0007	0.92	566	544	655	24	83
11.1c	7	87	0.78	0.691	0.012	0.0845	0.0010	0.0593	0.0010	0.69	533	523	579	36	90
11.2c	98	1572	0.05	0.635	0.008	0.0801	0.0009	0.0575	0.0006	0.93	499	497	511	24	97
12.1c	19	325	0.19	0.569	0.008	0.0719	0.0008	0.0574	0.0007	0.85	457	448	506	28	89
12.2c	157	4467	0.11	0.348	0.004	0.0454	0.0005	0.0557	0.0006	0.93	303	286	439	24	65
13c	46	1043	0.03	0.430	0.005	0.0568	0.0006	0.0550	0.0006	0.90	364	356	411	26	87
14.2c	87	2249	0.02	0.377	0.005	0.0504	0.0006	0.0542	0.0006	0.90	325	317	380	26	83
15c	67	1133	0.07	0.607	0.008	0.0760	0.0009	0.0579	0.0007	0.89	481	472	525	26	90
16.2c	86	1453	0.04	0.592	0.007	0.0762	0.0009	0.0564	0.0007	0.90	472	474	466	26	102
17c	106	1769	0.05	0.620	0.008	0.0773	0.0009	0.0582	0.0007	0.91	490	480	535	26	90
18c	75	1739	0.02	0.427	0.005	0.0563	0.0006	0.0550	0.0007	0.88	361	353	411	27	86
19c	117	2052	0.07	0.580	0.007	0.0734	0.0008	0.0573	0.0007	0.91	465	457	504	25	91
20c	72	1135	0.05	0.666	0.008	0.0817	0.0009	0.0591	0.0007	0.89	518	506	570	25	89
22c	74	1972	0.03	0.381	0.005	0.0486	0.0005	0.0568	0.0007	0.87	328	306	483	27	63
23c	99	1573	0.07	0.639	0.008	0.0808	0.0009	0.0574	0.0007	0.90	502	501	507	26	99
24c	102	1656	0.05	0.628	0.008	0.0798	0.0009	0.0571	0.0007	0.90	495	495	495	26	100
25c	112	2013	0.05	0.577	0.007	0.0711	0.0008	0.0589	0.0007	0.89	463	443	562	26	79
26c	94	1728	0.04	0.544	0.007	0.0702	0.0008	0.0562	0.0007	0.88	441	438	459	27	95
27c	133	2301	0.04	0.591	0.007	0.0753	0.0009	0.0570	0.0007	0.89	472	468	490	27	96
28c	114	1701	0.08	0.687	0.009	0.0861	0.0010	0.0579	0.0007	0.88	531	532	527	27	101
29c	96	1801	0.06	0.540	0.007	0.0689	0.0008	0.0568	0.0007	0.88	438	430	482	27	89
30c	95	1818	0.06	0.525	0.007	0.0677	0.0008	0.0562	0.0007	0.87	428	422	460	28	92

Table 1 continued

Grain (zircon)	[Pb] (ppm)	[U] (ppm)	Th/U	$^{207}\text{Pb}/^{235}\text{U}$	$\pm 1\sigma$	$^{206}\text{Pb}/^{238}\text{U}$	$\pm 1\sigma$	$^{207}\text{Pb}/^{206}\text{Pb}$	$\pm 1\sigma$	Rho	Ages	$^{206}\text{Pb}/^{238}\text{U}$	$^{207}\text{Pb}/^{238}\text{U}$	$^{207}\text{Pb}/^{206}\text{Pb}$	$\pm 1\sigma$	Conc. (%)
<i>ES 8 La Salvetat migmatite</i>																
1.1d	33	892	0.01	0.351	0.004	0.0478	0.0005	0.0532	0.0006	0.88	305	301	337	27	89	
1.2d	18	156	2.20	0.766	0.011	0.0932	0.0011	0.0596	0.0008	0.79	578	575	590	30	97	
2.1d	27	184	0.80	1.447	0.018	0.1533	0.0017	0.0685	0.0008	0.88	909	919	882	25	104	
2.2d	82	828	1.03	0.812	0.010	0.0995	0.0011	0.0592	0.0007	0.93	604	611	575	24	106	
3d	44	497	0.33	0.867	0.011	0.1047	0.0012	0.0601	0.0007	0.91	634	642	607	25	106	
4d	25	631	0.01	0.389	0.006	0.0504	0.0006	0.0560	0.0009	0.69	334	317	451	36	70	
5d	52	353	0.93	1.480	0.019	0.1512	0.0017	0.0710	0.0009	0.88	922	908	957	25	95	
6d	26	341	0.29	0.753	0.010	0.0886	0.0010	0.0616	0.0008	0.86	570	547	661	27	83	
7d	67	635	1.41	0.839	0.011	0.0983	0.0011	0.0619	0.0008	0.86	619	605	670	27	90	
8d	61	644	0.72	0.817	0.010	0.1014	0.0011	0.0585	0.0007	0.90	607	623	547	26	114	
9d	25	194	0.99	1.141	0.016	0.1276	0.0014	0.0648	0.0009	0.83	773	774	768	28	101	
10d	55	429	0.36	1.413	0.017	0.1496	0.0017	0.0685	0.0008	0.91	894	899	884	24	102	
11d	10	81	0.52	1.306	0.020	0.1414	0.0016	0.0670	0.0010	0.74	848	853	838	32	102	
12d	116	764	0.11	2.616	0.032	0.1827	0.0020	0.1039	0.0012	0.91	1305	1082	1694	21	64	
13d	201	2497	0.17	0.830	0.010	0.0990	0.0011	0.0608	0.0007	0.93	614	608	633	24	96	
14d	71	605	0.12	1.370	0.018	0.1463	0.0016	0.0679	0.0008	0.88	876	880	866	25	102	
15d	89	752	0.16	1.366	0.017	0.1454	0.0016	0.0681	0.0008	0.89	875	875	873	24	100	
16.1d	29	413	0.09	0.713	0.010	0.0872	0.0010	0.0593	0.0008	0.78	547	539	578	30	93	
16.2d	43	172	2.04	2.044	0.027	0.2038	0.0023	0.0728	0.0009	0.86	1130	1195	1007	25	119	
17d	40	255	0.73	1.693	0.023	0.1685	0.0019	0.0729	0.0010	0.83	1006	1004	1010	26	99	
18d	59	627	0.18	1.002	0.013	0.1162	0.0013	0.0625	0.0008	0.87	705	709	692	26	102	
19d	36	516	0.53	0.653	0.010	0.0779	0.0009	0.0608	0.0009	0.74	511	484	632	33	77	
20d	94	1132	0.29	0.824	0.011	0.0990	0.0011	0.0604	0.0007	0.87	611	608	619	26	98	
21d	53	597	0.50	0.852	0.011	0.1020	0.0012	0.0606	0.0008	0.87	626	626	627	27	100	
22d	295	653	0.48	11.792	0.149	0.4746	0.0053	0.1802	0.0022	0.89	2588	2504	2655	20	94	
23.1d	10	98	0.26	1.049	0.016	0.1205	0.0014	0.0631	0.0010	0.75	728	734	713	32	103	
23.2d	19	215	0.22	0.942	0.014	0.1084	0.0012	0.0630	0.0009	0.78	674	663	709	30	94	
24d	93	1001	0.69	0.927	0.012	0.0977	0.0011	0.0688	0.0009	0.84	666	601	894	26	67	
25d	31	352	0.69	0.764	0.011	0.0930	0.0011	0.0596	0.0008	0.81	576	573	588	29	98	
26d	113	653	0.28	2.378	0.031	0.2036	0.0023	0.0847	0.0011	0.86	1236	1195	1309	24	91	
27d	176	360	0.63	13.005	0.171	0.4959	0.0056	0.1902	0.0024	0.86	2680	2596	2744	20	95	
28d	13	138	0.53	0.867	0.014	0.1026	0.0012	0.0613	0.0010	0.73	634	630	649	33	97	

Errors are reported at 1 sigma



**Table 2** U–Th–Pb LA-ICP-MS data for the monazite grains from sample ES7 and ES8

Grain	[Pb] (ppm)	[U] (ppm)	[Th] (ppm)	Th/U	206 Pb/238 U ±(1σ)	207 Pb/235 U ±(1σ)	208 Pb/232 Th ±(1σ)	Ages						
								206 Pb/238 U	207 Pb/235 U	208 Pb/232 Th	±(1σ)			
<i>ES7 Laouzas granite</i>														
33a	1405	14522	72329	5	0.0459	0.0006	0.3167	0.0045	0.0143	0.0002	289	279	287	3
34a	1331	8983	82,279	9	0.0463	0.0006	0.3171	0.0046	0.0142	0.0002	292	280	284	3
35a	1561	13906	84,806	6	0.0469	0.0006	0.3241	0.0046	0.0144	0.0002	296	285	289	3
36a	1255	5650	86,272	15	0.0464	0.0006	0.3174	0.0050	0.0140	0.0002	292	280	282	3
37a	1526	7457	102,172	14	0.0469	0.0006	0.3195	0.0048	0.0141	0.0002	295	282	284	3
38a	1422	6446	95,482	15	0.0471	0.0006	0.3250	0.0050	0.0143	0.0002	296	286	287	3
39a	1223	9071	72,384	8	0.0464	0.0006	0.3141	0.0046	0.0143	0.0002	293	277	287	3
40a	1316	6291	88,790	14	0.0464	0.0006	0.3234	0.0050	0.0141	0.0002	292	285	283	3
41a	1869	5871	110,726	19	0.0510	0.0006	0.3636	0.0064	0.0169	0.0002	321	315	339	4
42a	1664	7306	95,010	13	0.0511	0.0006	0.3540	0.0056	0.0166	0.0002	321	308	332	4
43a	1427	5587	83,966	15	0.0506	0.0006	0.3511	0.0057	0.0165	0.0002	318	306	330	4
44a	1505	6351	89,387	14	0.0503	0.0006	0.3457	0.0056	0.0161	0.0002	316	302	323	3
45a	1421	3511	87,118	25	0.0515	0.0007	0.3737	0.0068	0.0168	0.0002	324	322	337	4
46a	1467	8153	80,517	10	0.0504	0.0006	0.3488	0.0053	0.0163	0.0002	317	304	328	4
47a	1385	12,594	62,010	5	0.0505	0.0006	0.3437	0.0050	0.0166	0.0002	317	300	332	4
48a	1402	13,808	60,267	4	0.0506	0.0006	0.3464	0.0050	0.0164	0.0002	318	302	330	4
50a	1343	12,757	60,159	5	0.0507	0.0006	0.3468	0.0051	0.0161	0.0002	319	302	322	3
51a	1254	5577	75,446	14	0.0514	0.0006	0.3570	0.0058	0.0156	0.0002	323	310	313	3
52a	1210	10,014	56,262	6	0.0506	0.0006	0.3483	0.0051	0.0166	0.0002	319	304	333	4
53a	1279	11,628	59,034	5	0.0506	0.0006	0.3470	0.0051	0.0160	0.0002	318	302	320	3
54a	1275	4595	79,000	17	0.0520	0.0006	0.3656	0.0063	0.0157	0.0002	327	316	315	3
55a	864	3086	52,825	17	0.0513	0.0006	0.3551	0.0065	0.0160	0.0002	323	309	320	3
56a	1224	5645	70,387	12	0.0510	0.0006	0.3458	0.0053	0.0162	0.0002	321	302	324	3
57a	1238	7421	68,419	9	0.0510	0.0006	0.3418	0.0054	0.0157	0.0002	320	299	315	3
58a	1312	5427	74,832	14	0.0522	0.0006	0.3527	0.0056	0.0166	0.0002	328	307	333	4
59a	1003	6587	54,512	8	0.0504	0.0006	0.3473	0.0057	0.0156	0.0002	317	303	312	3
60a	1107	6681	62,592	9	0.0502	0.0006	0.3376	0.0057	0.0154	0.0002	316	295	309	3
61a	1453	10,545	74,983	7	0.0499	0.0006	0.3476	0.0053	0.0158	0.0002	314	303	318	3
62a	1050	5823	58,804	10	0.0506	0.0006	0.3544	0.0057	0.0158	0.0002	318	308	318	3
63a	1061	3078	66,792	22	0.0509	0.0006	0.3549	0.0065	0.0159	0.0002	320	308	319	3
64a	1212	12,417	52,947	4	0.0497	0.0006	0.3419	0.0050	0.0158	0.0002	313	299	316	3
Mz1.1 (fsp)	346	3072	14,349	5	0.0506	0.0007	0.5039	0.0078	0.0167	0.0002	318	414	335	4
Mz1.2 (fsp)	1015	7710	50,744	7	0.0473	0.0006	0.3374	0.0048	0.0154	0.0002	298	295	309	4

Table 2 continued

Grain	[Pb] (ppm)	[U] (ppm)	[Th] (ppm)	Th/U	206 Pb/238 U	$\pm(1\sigma)$	207 Pb/235 U	$\pm(1\sigma)$	208 Pb/232 Th	$\pm(1\sigma)$	Ages			
											206 Pb/238 U	207 Pb/235 U	208 Pb/232 Th	
Mz2.1 (crd)	1072	8419	51,320	6	0.0489	0.0007	0.3815	0.0055	0.0157	0.0002	308	328	315	4
Mz2.2 (crd)	1218	7907	57,072	7	0.0519	0.0007	0.3798	0.0054	0.0169	0.0002	326	327	338	4
Mz3.1 (qtz)	1372	10,036	73,905	7	0.0458	0.0006	0.3173	0.0045	0.0148	0.0002	289	280	296	4
Mz3.2 (qtz)	1444	10,503	79,124	8	0.0460	0.0006	0.3171	0.0045	0.0145	0.0002	290	280	292	4
Mz4.1 (fsp)	1373	1157	55,048	48	0.0460	0.0006	0.3170	0.0046	0.0148	0.0002	290	280	297	4
Mz4.2 (fsp)	1423	1206	56,888	47	0.0462	0.0006	0.3264	0.0049	0.0149	0.0002	291	287	299	4
Mz5 (qtz)	880	732	35,685	49	0.0576	0.0008	0.4697	0.0089	0.0195	0.0003	361	391	390	5
Mz6 (qtz)	1247	1045	50,170	48	0.0508	0.0007	0.3525	0.0053	0.0175	0.0002	319	307	350	4
Mz7 (qtz)	1396	10,994	73,918	7	0.0474	0.0007	0.3261	0.0050	0.0157	0.0002	298	287	314	4
Mz8 (qtz)	713	5795	37,073	6	0.0462	0.0007	0.3156	0.0049	0.0148	0.0002	291	279	298	4
Mz9.1 (fsp)	667	7610	33,851	4	0.0478	0.0007	0.3389	0.0054	0.0152	0.0002	301	296	305	4
Mz9.2 (fsp)	136	1285	4509	4	0.0565	0.0008	0.3988	0.0065	0.0198	0.0003	354	341	396	5
Mz10.1 (qtz)	1305	6046	81,834	14	0.0465	0.0007	0.4293	0.0069	0.0148	0.0002	293	363	297	4
Mz10.2 (qtz)	1318	8843	61,053	7	0.0469	0.0007	0.3484	0.0057	0.0151	0.0002	295	304	303	4
Mz10.3 (qtz)	1344	10,119	76,298	8	0.0491	0.0007	0.3752	0.0062	0.0153	0.0002	309	324	306	4
Mz11.1 (qtz-j)	1121	8693	56,704	7	0.0466	0.0007	0.3239	0.0054	0.0148	0.0002	294	285	298	4
Mz11.2 (qtz-j)	1359	1165	55,336	48	0.0474	0.0007	0.3382	0.0056	0.0152	0.0002	298	296	304	4
Grain	[Pb] (ppm)	[U] (ppm)	[Th] (ppm)	Th/U	206 Pb/238 U	$\pm(1\sigma)$	207 Pb/235 U	$\pm(1\sigma)$	208 Pb/232 Th	$\pm(1\sigma)$	Ages			
											206 Pb/238 U	207 Pb/235 U	208 Pb/232 Th	$\pm(1\sigma)$
<i>E58 La Salvetat migmatite</i>														
65a	1230	5483	67,669	12	0.0518	0.0006	0.3603	0.0058	0.0163	0.0002	326	312	327	3
66a	1189	18,268	32,530	2	0.0496	0.0006	0.3407	0.0050	0.0160	0.0002	312	298	321	3
67a	853	8595	34,736	4	0.0511	0.0006	0.3507	0.0054	0.0160	0.0002	321	305	321	3
68a	942	12,580	30,216	2	0.0502	0.0006	0.3479	0.0052	0.0161	0.0002	316	303	324	3
69a	730	7226	30,260	4	0.0510	0.0006	0.3552	0.0055	0.0159	0.0002	321	309	319	3
70a	1097	16,108	32,500	2	0.0496	0.0006	0.3411	0.0051	0.0157	0.0002	312	298	315	3
71a	854	10,883	27,100	2	0.0522	0.0006	0.3715	0.0056	0.0164	0.0002	328	321	328	3
72a	1543	25,307	37,012	1	0.0502	0.0006	0.3449	0.0050	0.0160	0.0002	316	301	321	3
73a	1307	19,895	34,723	2	0.0504	0.0006	0.3497	0.0052	0.0162	0.0002	317	305	325	3
74a	1229	18,313	34,893	2	0.0499	0.0006	0.3445	0.0051	0.0159	0.0002	314	301	318	3
75a	1571	25,306	37,535	1	0.0508	0.0006	0.3528	0.0052	0.0161	0.0002	319	307	324	3
76a	781	8381	31,410	4	0.0501	0.0006	0.3488	0.0055	0.0156	0.0002	315	304	313	3

Table 2 continued

Grain	[Pb] (ppm)	[U] (ppm)	[Th] (ppm)	Th/U	206 Pb/238 U $\pm(1\sigma)$	207 Pb/235 U $\pm(1\sigma)$	208 Pb/232 Th $\pm(1\sigma)$	Ages					
								206 Pb/238 U	207 Pb/235 U	208 Pb/232 Th	$\pm(1\sigma)$		
77a	No data available				0.0500	0.0006	0.0052	0.0158	0.0002	314	299	317	3
78.1a					0.0503	0.0006	0.0052	0.0161	0.0002	317	304	322	3
78.2a					0.0505	0.0006	0.0052	0.0162	0.0002	317	303	325	3
78.3a					0.0510	0.0006	0.0055	0.0159	0.0002	321	307	318	3
79a					0.0485	0.0006	0.0053	0.0150	0.0002	305	294	300	3
80.1a					0.0475	0.0006	0.0052	0.0147	0.0002	299	291	295	3
80.2a					0.0475	0.0006	0.0053	0.0149	0.0002	299	295	298	3
81.1a					0.0474	0.0006	0.0050	0.0150	0.0002	299	289	300	3
81.2a					0.0467	0.0006	0.0049	0.0148	0.0002	294	285	296	3
82a					0.0466	0.0006	0.0049	0.0147	0.0002	294	284	296	3
83a					0.0477	0.0006	0.0052	0.0147	0.0002	300	289	294	3
84a					0.0469	0.0006	0.0049	0.0148	0.0002	295	285	297	3
2b	713	6409	34,056	5	0.0483	0.0006	0.0062	0.0152	0.0002	304	305	306	3
3.1b	929	9780	40,809	4	0.0480	0.0006	0.0059	0.0150	0.0002	302	298	301	3
3.2b	772	5903	39,878	7	0.0481	0.0006	0.0063	0.0150	0.0002	303	302	301	3
3.3b	890	7957	42,165	5	0.0484	0.0006	0.0061	0.0150	0.0002	305	302	301	3
4b	869	8361	39,428	5	0.0483	0.0006	0.0062	0.0149	0.0002	304	299	300	3
5.2b	924	10,015	39,179	4	0.0474	0.0006	0.0062	0.0147	0.0002	299	297	295	3
6b	676	6290	32,046	5	0.0472	0.0006	0.0063	0.0142	0.0002	297	284	285	3
7b	1186	19,504	31,251	2	0.0472	0.0006	0.0062	0.0145	0.0002	297	289	291	3
8.1b	1213	17,656	38,626	2	0.0468	0.0006	0.0064	0.0143	0.0002	295	292	288	3
8.2b	1300	20,670	37,530	2	0.0462	0.0006	0.0063	0.0141	0.0001	291	287	283	3
9b	1738	32,151	38,690	1	0.0455	0.0006	0.0063	0.0141	0.0001	287	282	282	3
10b	1074	16,844	30,899	2	0.0464	0.0006	0.0065	0.0141	0.0001	292	282	283	3
11b	1481	24,419	39,221	2	0.0461	0.0006	0.0066	0.0141	0.0001	291	285	283	3
12b	784	7353	35,558	5	0.0474	0.0006	0.0073	0.0140	0.0001	299	291	280	3
Mz12 (qtz)	731	9662	27,745	3	0.0465	0.0006	0.0054	0.0142	0.0002	293	309	285	3
Mz13 (qtz)	973	11,793	39,613	3	0.0455	0.0006	0.0047	0.0146	0.0002	287	286	294	3
Mz15 (bt)	137	1909	5184	3	0.0447	0.0006	0.0067	0.0142	0.0002	282	300	285	3
Mz17 (bt)	838	11,024	30,254	3	0.0454	0.0006	0.0048	0.0149	0.0002	286	280	300	3

Table 2 continued

Grain	[Pb] (ppm)	[U] (ppm)	[Th] (ppm)	Th/U	206 Pb/238 U $\pm(1\sigma)$	207 Pb/235 U $\pm(1\sigma)$	208 Pb/232 Th $\pm(1\sigma)$	Ages						
								206 Pb/238 U	207 Pb/235 U	208 Pb/232 Th	$\pm(1\sigma)$			
Mz20 (bt)	735	8040	29,039	4	0.0474	0.0006	0.3310	0.0053	0.0156	0.0002	299	290	312	4
Mz21.1 (bt)	1007	12,699	36,732	3	0.0495	0.0006	0.3700	0.0059	0.0155	0.0002	311	319	311	4
Mz21.2 (bt)	1138	12,832	44,059	3	0.0450	0.0006	0.3459	0.0067	0.0150	0.0002	284	302	300	4
Mz23 (bt)	463	4871	15,302	3	0.0468	0.0006	0.4067	0.0071	0.0147	0.0002	295	346	295	3

Errors are reported at 1 sigma

et al. 1971; Roddick 1983; Jourdan et al. 2006; Jourdan and Renne 2007), with an age of  $1081.0 \pm 1.2$  Ma (Renne et al. 2010, 2011).

Step-heating analyses of single grains were performed with a CO<sub>2</sub> laser probe coupled to a Map215<sup>®</sup> mass spectrometer. The experimental procedure is described in Ruffet et al. (1991) and Ruffet et al. (1995). The five argon isotopes and the background baselines were measured in eleven cycles, in peak-jumping mode. Blanks were performed routinely each first or third/fourth run, and subtracted from the subsequent sample gas fractions. All isotopic measurements are corrected for K, Ca and Cl isotopic interferences, mass discrimination and atmospheric argon contamination. Apparent age errors are plotted at the 1 $\sigma$  level and do not include the errors on the  $^{40}\text{Ar}^*/^{39}\text{Ar}_K$  ratio and age of the monitor and decay constant. The errors in the  $^{40}\text{Ar}^*/^{39}\text{Ar}_K$  ratio and age of the monitor and decay constant are included in the final calculation of the (pseudo-) plateau age error margins or for apparent ages individually cited. Details on the method and  $^{40}\text{Ar}/^{39}\text{Ar}$  analytical data are given in the Supporting Information.


It is commonly considered that a plateau is obtained when calculated  $^{40}\text{Ar}^*/^{39}\text{Ar}_K$  ratios of at least three consecutive steps, containing a minimum of 70 % of the  $^{39}\text{Ar}$  released, agree with the weighted mean calculated  $^{40}\text{Ar}^*/^{39}\text{Ar}_K$  ratio of the plateau segment. Pseudo-plateau ages can be defined with less than 70 % of the  $^{39}\text{Ar}$  released. All ages are displayed at the 1 $\sigma$  level. Analytical data, parameters used for calculations (isotopic ratios measured on K, Ca and Cl pure salts; mass discrimination; atmospheric argon ratios; J parameter; decay constants, etc.) and reference sources are available in a complementary data repository.

### U–Th–Pb LA-ICP-MS and $^{40}\text{Ar}$ – $^{39}\text{Ar}$ results

#### Sample ES7—Laouzas granite

Zircon and monazite grains were both recovered from this sample. Most of the zircon grains were pink in colour, euhedral, with very variable shapes from elongated to oval. Cathodoluminescence imaging revealed a rather heterogeneous population with anything from homogeneous low luminescent to heterogeneous (core + rim) grains (Fig. 3a). Thirty-seven analyses out of twenty-six zircon grains were made (Table 1). The heterogeneity of the grains is confirmed in a Tera-Wasserburg diagram (Fig. 4a) where data points are plotted in a concordant to very discordant position with apparent  $^{207}\text{Pb}/^{206}\text{Pb}$  ages ranging from 1010 Ma down to 370 Ma. Because of this heterogeneity, which is probably caused by a complex mixing of heterogeneous inheritance, plus variable degree of common Pb content and Pb loss, it is not possible to calculate any relevant ages. We





**Fig. 3** Cathodoluminescence images of some of the zircon grains dated in this study: **a** Laouzais granite; **b, c** La Salvetat migmatite. The white circle represents the spot analysis and the number corre-

sponds to the  $^{207}\text{Pb}/^{206}\text{Pb}$  age obtained. Zr number corresponds to the grain number in Table 1

therefore favoured plotting the data in a relative probability plot diagram (Fig. 4e) where only the  $^{207}\text{Pb}/^{206}\text{Pb}$  apparent ages for the more than 90 % concordant points were considered. One main peak can be defined at ca. 500 Ma, with minor peaks around 700 and 800 Ma.

Two types of monazite grains were found in this sample. Type 1 monazite comprises euhedral dark brown grains with sharp concentric zoning (Fig. 5), and type 2 monazite comprises euhedral to subhedral yellowish to orange grains characterised by complex patchy zoning (Fig. 6), as revealed by backscattered electron (BSE) imaging. Elemental imaging carried out using the NanoSIMS ion probe reveals that the concentric zoning for type 1 monazite (Fig. 5) is visible with all the elements imaged (Y, La, Ce, Sm, U, Th and Pb) with cores that are LREE-rich and poor in Y, U, Th and Pb. More importantly, the distributions of U and Th in this monazite type perfectly match each other. For type 2 monazite (Fig. 6), the complex zoning noticed in BSE images is well mimicked by the distribution of Y, REE, U and  $^{206}\text{Pb}$  while the distribution of Th and  $^{208}\text{Pb}$  is less disturbed.

These two types of monazite grains were therefore analysed separately. In addition, 19 analyses (11 grains) were performed directly in a thin section. In total, 50 analyses were carried out (Table 2). Plotted in a  $^{206}\text{Pb}/^{238}\text{U}$  versus  $^{208}\text{Pb}/^{232}\text{Th}$  concordia diagram (Fig. 4b), the two types plot in two distinct groups. For the monazite grains dated in context in the thin sections, there is no evident correlation between the mineral hosting the monazite grains and their apparent ages. Therefore, the location of type 1 and type 2 monazite is not related to any specific host mineral. Type 1 monazite ( $N = 26$ ; grains with sharp concentric zoning) plot in a concordant to slightly discordant position. A cluster of 14 concordant analyses (Fig. 4b) yields a concordia age (Ludwig 1998) of  $318.0 \pm 1.4$  Ma (MSWD = 0.87). This concordia age is equivalent within error to the average  $^{206}\text{Pb}/^{238}\text{U}$  date of  $318.8 \pm 1.5$  Ma ( $N = 26$ ; MSWD = 0.95) obtained for the 26 analyses

defining this first group. Interestingly, the  $^{208}\text{Pb}/^{232}\text{Th}$  apparent ages for this group display a bimodal distribution (see Fig. 4g), with one peak at  $331.5 \pm 2$  Ma and a second one at  $319.1 \pm 2.0$  Ma.









Data obtained on type 2 monazite (grains with complex patchy zoning;  $N = 20$ ) exhibit a slight reverse discordance (Fig. 4b). The mean  $^{206}\text{Pb}/^{238}\text{U}$  date obtained for these twenty analyses is consistent with a value of  $293.5 \pm 1.7$  Ma (MSWD = 0.78), while the  $^{208}\text{Pb}/^{232}\text{Th}$  apparent dates yield average dates of  $285.2 \pm 2.2$  Ma (MSWD = 0.66;  $N = 8$ ) for the grains analysed in the epoxy puck and of  $296.5 \pm 2.8$  Ma (MSWD = 0.42;  $N = 7$ ) for the grains dated in the thin section (Fig. 4g).

Muscovite single grain from sample ES7 yielded a flat  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  age spectrum (Fig. 4h) over most of  $^{39}\text{Ar}_K$  degassing (ca. 95 %), corresponding to a calculated plateau age of  $298.2 \pm 0.8$  Ma ( $2\sigma$  level).

#### Sample ES8—La Salvetat migmatite

Both monazite and zircon grains were extracted from this sample. Two types of zircon grains were found. The first type is characterised by elongate pinkish grains (Fig. 3b), while the second type is constituted by squat prismatic grains (Fig. 3c). Both types display complex zoning with core and rims apparent for most (Fig. 3b, c). Thirty-two analyses were performed on twenty-eight grains (Table 1). Plotted in a Tera–Wasserburg diagram (Fig. 4c), they are in a concordant to discordant position, with apparent  $^{207}\text{Pb}/^{206}\text{Pb}$  ages ranging from ca. 2650 down to 300 Ma. Once again, it is difficult to get any valuable geochronological constraint with this set of data. Plotted in a relative probability plot (Fig. 4e), two main peaks arise at ca. 610 Ma and 875 Ma, with minor peaks at 690, 1000, 1310 and 2450 Ma.

From a morphological point of view, all monazite crystals were yellow and euhedral to subhedral. Forty-six analyses were performed (38 on separated grains and 8 directly




**Fig. 4** Summary of the geochronological results. In all diagram,  $N$  refers to the number of analyses. Grey ellipses correspond to the data used to calculate the concordia ages. **a, c** Tera-Wasserburg  $^{207}\text{Pb}/^{206}\text{Pb}$  versus  $^{238}\text{U}/^{206}\text{Pb}$  concordia diagram for the zircon grains analysed in the Laouzas granite (**a**) and the La Salvetat migmatite (**c**). **b, d**:  $^{206}\text{Pb}/^{238}\text{U}$  versus  $^{208}\text{Pb}/^{232}\text{Th}$  diagrams for the monazite analysed in the Laouzas granite (**b**) and the La Salvetat migmatite (**d**). **e** Relative probability plot of  $^{207}\text{Pb}/^{206}\text{Pb}$  dates for all the more than 90 % concordant zircon grains obtained in this study. **f** Relative probability plots for all the monazite  $^{206}\text{Pb}/^{238}\text{U}$  dates obtained in this study. **g** Relative probability plots for all the monazite  $^{208}\text{Pb}/^{232}\text{Th}$  dates obtained in this study. **H**:  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  spectra of muscovite from samples ES5, ES7 and ES8. The error bars for each temperature steps are at the  $1\sigma$  level. The errors in the  $J$ -values are not included. Plateau age error is at the  $2\sigma$  level

in context in thin sections). Plotted in a  $^{206}\text{Pb}/^{238}\text{U}$  versus  $^{208}\text{Pb}/^{232}\text{Th}$  concordia diagram (Fig. 4d), they all plot in a concordant to slightly discordant position. A first group of 14 analyses defines a concordia age of  $318.5 \pm 0.7$  Ma


(MSWD = 1.3; Fig. 4d). A second cluster of 15 analyses yields a concordia age of  $298.8 \pm 1.3$  Ma (MSWD = 0.68; Fig. 4d). The remaining 17 analyses plot in a slightly reverse discordant position either between the two previous calculated concordia dates or are apparently younger than 299 Ma (Fig. 4d). The two analyses performed in monazite grains hosted by quartz yield dates close to 290 Ma, while the monazite grains hosted by biotite plot in a scattered position.

When looking at the BSE imaging, the first group (ca. 318.5 Ma) is characterised by fairly homogeneous monazite grains (Fig. 7), while monazite in the second group (ca. 299 Ma) is characterised by more patchy zoning (Fig. 8). The NanoSIMS imaging of the first group confirms the rather homogeneous distribution of the imaged elements with the exception of U and  $^{206}\text{Pb}$  that appear to be poorer in the core of the grain (Fig. 7). For the second group, the



**Fig. 5** Top picture backscattered electron image of the type 1 monazite from sample ES7 (i.e. ca. 318 Ma). White circle represents the spot analysis and has a diameter of 7 microns. Date refers to the

$^{208}\text{Pb}/^{232}\text{Th}$  individual date.  $M$  number refers to the grain number in Table 2. White square corresponds to the location of the NanoSIMS elemental images (8 bottom pictures)



**Fig. 6** Top picture backscattered electron image of the type 2 monazite from sample ES7 (i.e. ca. 294 Ma). White circle represents the spot analysis and has a diameter of 7 microns. Date refers to the

$^{208}\text{Pb}/^{232}\text{Th}$  individual date. *M* number refers to the grain number in Table 2. White square corresponds to the location of the NanoSIMS elemental images (8 bottom pictures)

elemental distributions are not as simple. The REE distribution seems to be homogeneous throughout the grain, whereas the Y, U, Th and Pb distributions are patchier (Fig. 8).

Similar to monazite in ES7, monazite data in ES8 are characterised by evident differences between the  $^{206}\text{Pb}/^{238}\text{U}$  and the  $^{208}\text{Pb}/^{232}\text{Th}$  dates (Fig. 4f, g). The  $^{208}\text{Pb}/^{232}\text{Th}$  dates fall into 3 distinct populations (Fig. 4g) at  $319.8 \pm 1.8$  Ma (MSWD = 1.3;  $N = 15$ ),  $298.2 \pm 1.5$  Ma (MSWD = 0.89;  $N = 19$ ) and  $284.7 \pm 2.1$  Ma (MSWD = 0.98;  $N = 8$ ), respectively.  $^{206}\text{Pb}/^{238}\text{U}$  dates define two different peaks at  $316.2 \pm 1.9$  Ma (MSWD = 0.68;  $N = 15$ ) and  $296.1 \pm 1.3$  Ma (MSWD = 2.8;  $N = 26$ ), respectively (Fig. 4f).

Muscovite single grain from sample ES8 yielded a flat  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  age spectrum (Fig. 4h) over most of  $^{39}\text{ArK}$  degassing (ca. 98 %) that correspond to a calculated plateau


age of  $298 \pm 1$  Ma ( $2\sigma$  level). This age is similar to the ES7 plateau age.

## Geological significance of the geochronological data

### Zircon dating

Zircon data from both the Laouzias granite and the La Salvetat migmatite show a considerable spread (Fig. 4e) and cannot be used to date either the emplacement of the granite or the age of migmatization. They rather demonstrate the existence of a complex polygenic history in the region, with dates ranging from the late Archean to the Ordovician. They are consistent with the data published by Faure et al. (2010), although more date populations were found





**Fig. 7** Top picture backscattered electron image of the first group of monazite grains (i.e. ca. 319 Ma) from sample ES8. White circle represents the spot analysis and has a diameter of 7 microns. Date

refers to the  $^{208}\text{Pb}/^{232}\text{Th}$  date. *M* number refers to the grain number in Table 2. White square corresponds to the location of the NanoSIMS elemental images (8 bottom pictures)

in our study. In the La Salvetat migmatite, the youngest point (grain 1.1d) that plots close to the concordia gives a  $^{207}\text{Pb}/^{206}\text{Pb}$  date of  $337 \pm 27$  Ma (1 sigma). This grain might have grown during the migmatization, but because of its large error, it does not help to constrain this event precisely.


### Monazite dating

#### The Laouzas granite

In the  $^{206}\text{Pb}/^{238}\text{U}$  versus  $^{208}\text{Pb}/^{232}\text{Th}$  concordia diagram (Fig. 4b), the monazite grains from the Laouzas granite plot in two clusters although the geochronological information brought, for each cluster, by the monazite grains  $^{206}\text{Pb}/^{238}\text{U}$  and  $^{208}\text{Pb}/^{232}\text{Th}$  respective dates is different. Indeed, within

each cluster, the  $^{206}\text{Pb}/^{238}\text{U}$  dates are comparable, while they are more scattered in the case of the  $^{208}\text{Pb}/^{232}\text{Th}$  ones.

To first order, four dates can be defined for the Laouzas granite. The oldest date at ca. 330 Ma can be calculated using the older apparent  $^{208}\text{Pb}/^{232}\text{Th}$  dates (Fig. 4g). The second one around 319 Ma is given by a concordia age of  $318.0 \pm 1.4$  Ma (Fig. 4b), the mean  $^{206}\text{Pb}/^{238}\text{U}$  date of  $318.8 \pm 1.5$  Ma obtained for all the analyses from the older cluster (Fig. 4b) and from one of the peaks defined by the  $^{208}\text{Pb}/^{232}\text{Th}$  dates at  $319.1 \pm 2.0$  Ma (Fig. 4g). The third date of around 298–292 Ma is defined by one peak in the  $^{208}\text{Pb}/^{232}\text{Th}$  dates distribution at  $298.2 \pm 1.5$  Ma (Fig. 4g) and the mean  $^{206}\text{Pb}/^{238}\text{U}$  date of  $293.5 \pm 1.7$  Ma calculated for the second group (Fig. 4b). This third date is in a good agreement with the  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  age of  $298.2 \pm 0.8$  Ma ( $2\sigma$  level) obtained on muscovite from the same sample



**Fig. 8** Top picture backscattered electron image of the second group of monazite grains (i.e. ca. 298 Ma) from sample ES8. White circle represents the spot analysis and has a diameter of 7 microns. Date

refers to the  $^{208}\text{Pb}/^{232}\text{Th}$  date. M number refers to the grain number in Table 2. White square corresponds to the location of the NanoSIMS elemental images (8 bottom pictures)

(Fig. 4h). Finally, the youngest date around 285 Ma is defined by the youngest  $^{208}\text{Pb}/^{232}\text{Th}$  dates (Fig. 4g).

It seems evident that the dates of ca. 319 Ma and ca. 298 Ma are representative of specific events as they are common to both chronometers (U/Pb and Th/Pb) and because the second one is also defined by  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  dating. Therefore, two scenarios can be suggested. Either the date of ca. 319 Ma yields the emplacement age of the Laouzaz granite and the date of ca. 298 Ma is related to a post-emplacement event, or the granite was emplaced ca. 298 Ma ago, in which case the date of ca. 319 Ma should be regarded as “inherited”. Three observations are helpful to discriminate between these two scenarios. (1) The backscattered imaging shows that the ca. 319 Ma monazite population (type 1) is characterised by a rather simple concentric zoning, whereas the ca. 298 Ma old population (type 2) systematically displays a complex patchy zoning.

Concentric zoning is consistent with a magmatic origin, whereas patchy zoning suggests the involvement of a post-crystallisation perturbing event, such as fluid-related dissolution/recrystallisation (Williams et al. 2011; Tartèse et al. 2011; Didier et al. 2013). (2) As noticed on the NanoSIMS imaging (Figs. 5, 6), the elemental distributions perfectly match for the type 1 monazite while they differ for type 2. (3) Finally, petrographic observations suggest an overprint of the primary magmatic assemblage associated with a later circulation of hydrothermal fluids. In the light of these observations, we propose that the Laouzaz granite was emplaced ca. 319 Ma ago and was affected by a post-emplacement, fluid-related event, ca. 298 Ma ago. In this case, the  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  age at  $298.2 \pm 0.8$  Ma yielded by muscovite from the same sample would also characterise this late hydrothermal event, since it has been shown that the K–Ar geochronometer in muscovite can be highly

sensitive to fluid circulations in granites in a similar context (Questembert leucogranite, Armorican massif, France by Tartèse et al. 2011).

The oldest date at ca. 330 Ma is only evidenced by the  $^{208}\text{Pb}/^{232}\text{Th}$  dates. A similar phenomenon has also been noticed by Tartèse et al. (2012) for monazite grains from a mylonitised granite from the South Armorican Shear Zone, in which the  $^{208}\text{Pb}/^{232}\text{Th}$  dates (defining an average date of  $313 \pm 3$  Ma) were systematically older than the U–Pb dates (defining an average date of  $299 \pm 4$  Ma). A recent study by Didier et al. (2013) has demonstrated that F-rich fluids can be responsible for the disturbance of the Th/Pb ratios in monazite and the incorporation of excess Pb, leading to a large spread of the  $^{208}\text{Pb}/^{232}\text{Th}$  dates. Disturbed  $^{208}\text{Pb}/^{232}\text{Th}$  and  $^{206}\text{Pb}/^{238}\text{U}$  data were also obtained by Poitrasson et al. (2000) and were attributed to variable inputs and/or depletions in U, Th and Pb in the monazite crystals during hydrothermal alteration. As detailed earlier, the Laouzas anatectic granite bears petrographic evidence of hydrothermal fluid circulation. We therefore believe that this date of ca. 330 Ma is meaningless and is attributed to fluid perturbation of some of the monazite grains leading to a fractionation of their Th/Pb ratios (cf. Didier et al. 2013).

The other date at ca. 285 Ma is also obtained with only the  $^{208}\text{Pb}/^{232}\text{Th}$  dates. Once again, either this date is related to a fluid-induced perturbation of the monazite Th–Pb isotope system, and is, therefore, meaningless or it reflects the age of a yet unknown event in the region.

#### *La Salvetat migmatite*

Monazite from the La Salvetat migmatite also yields different  $^{208}\text{Pb}/^{232}\text{Th}$  and  $^{206}\text{Pb}/^{238}\text{U}$  dates, but here again two main dates can be proposed. A first one at ca. 319 Ma is given by a concordia age of  $318.5 \pm 0.7$  Ma (Fig. 4d), a mean  $^{208}\text{Pb}/^{232}\text{Th}$  date of  $319.8 \pm 1.8$  Ma (Fig. 4g) and a mean  $^{206}\text{Pb}/^{238}\text{U}$  date of  $316.2 \pm 1.9$  Ma (Fig. 4f). The second one, around 298 Ma, is given by a concordia age of  $298.8 \pm 1.3$  Ma (Fig. 4d), a mean  $^{208}\text{Pb}/^{232}\text{Th}$  date of  $298.2 \pm 1.5$  Ma (Fig. 4g), a mean  $^{206}\text{Pb}/^{238}\text{U}$  date of  $296.1 \pm 1.3$  Ma (Fig. 4f) and a muscovite  $^{40}\text{Ar}$ – $^{39}\text{Ar}$  plateau age of  $298 \pm 1$  Ma (Fig. 4h).

Lastly, the youngest  $^{208}\text{Pb}/^{232}\text{Th}$  dates define a mean date of  $284.7 \pm 2.1$  Ma. Monazite is known to be very resistant to diffusional reequilibration (e.g. Seydoux-Guil-laume et al. 2002; Gardés et al. 2007), preserving the age of their crystallisation. On the other hand, they recrystallise readily by dissolution/precipitation processes, when fluids or magmas are involved (Williams et al. 2011; Tartèse et al. 2011; Didier et al. 2013). As a general rule, two major “pulses” of monazite growth are predicted in metapelitic rocks: subsolidus growth in the upper amphibolite facies and growth during the cooling of leucosomes (rather than

partial melting) following migmatization (e.g. Foster et al. 2000; Rubatto et al. 2001; Kelsey et al. 2008; Spear and Pyle 2010). Furthermore, it is now well established that monazite may (re)crystallise due to fluid–rock interactions relatively late in the metamorphic history (e.g. Bosse et al. 2009; Tartèse et al. 2011, 2012; Didier et al. 2013).

Consequently, the three dates of ca. 319, ca. 298 and ca. 284 Ma obtained for the monazite grains from the La Salvetat migmatite may represent the ages of: (1) crystallisation during the prograde metamorphism in the upper amphibolite facies conditions; (2) crystallisation of the leucosomes following partial melting and (3) recrystallisation due to late fluid circulations, respectively. On the other hand, the close spatial and inferred genetic association of the migmatites with the Laouzas granite suggests that monazite grains of the same age should be found in both rock types. The crystallisation of the Laouzas granite is inferred to have taken place at ca. 319 Ma (see earlier), suggesting that the date of 319 Ma in the La Salvetat migmatite should be interpreted as the age of the crystallisation of the leucosomes, rather than that of the prograde amphibolite–facies metamorphism. The second date of ca. 298 Ma found in the migmatite is identical within error to the date of ca. 298 Ma found in the Laouzas granite, but also to the emplacement age of  $294 \pm 3$  Ma (U–Pb on zircon) obtained for the syntectonic Montalet leucogranite (Poilvet et al. 2011), situated about 20 km to the NW (Fig. 1). It is conceivable that a second phase of partial melting of the La Salvetat migmatite was the source of this leucogranite, but the poor outcrop conditions do not allow to validate such hypothesis. Alternatively and more probably, partial melting of other deeper formations formed the Montalet magmas, which percolated with associated fluids through the crust, resulting in the recrystallisation of some monazite grains in the migmatite–granite dome, as suggested by the late fluid circulations inferred from the petrographic observations of the La Salvetat migmatite. A detailed study of the geochemical affinities between the migmatites and the various granites would be necessary to answer this question.

Finally, the 284 Ma date, identical within uncertainty to the date of ca. 285 Ma obtained in the Laouzas granite is, as argued above, either an artefact due to fluid-enhanced modification of some of the monazite crystals or an evidence of a younger, although unidentified, event in the region. Similar Permian ages are known elsewhere in the European Variscan belt. Mougeot et al. (1997) reported an U–Pb apatite age ca. 289 Ma for the Velay granite. Cathelineau et al. (1990) obtained Permian ages on vein-type deposits from the Mortagne district in the South Armorican Massif and the French Massif Central, with a major stage of uranium mobilization between 290 and 260 Ma. In the Erzgebirge (Germany), the emplacement of vein-type deposits is also

Permian in age and post-dates the emplacement of the youngest Variscan granites by no less than 20–25 Ma (e.g. Velichkin and Vlasov 2011 and references therein). More recently, Boutin et al. (2015) reported some Permian U–Pb ages obtained on titanite associated with chlorite-talc mineralization in the Pyrenees. This non-exhaustive list demonstrates that this date of ca. 285 Ma is not unique at the scale of the Variscan belt as numerous ore deposits linked to fluid circulations are contemporaneous, and could, therefore, be considered as meaningful for the Montagne Noire. Finally, the remaining data that plot in a scattered position (Fig. 4d) could be attributed to an incomplete resetting of their U–Th–Pb system during the subsequent fluid circulation events.

### Partial melting and regional correlations

In summary, the Laouzas granite and the La Salvetat migmatite are spatially close and petrographically similar. The Laouzas granite is therefore interpreted as a product of the partial melting recorded in the La Salvetat migmatite. Monazite grains from both rocks recorded a date of ca. 319 Ma, which is interpreted as the emplacement age of the Laouzas granite, and hence also that of a first stage of migmatization (or rather the crystallisation of the leucosomes resulting from this partial melting). The date of ca. 298 Ma is either tentatively attributed to a second stage of migmatization or more probably to a pervasive percolation of magmas and associated fluids coming from a deeper source and resulting in the syntectonic crystallisation of the more superficial Montalet leucogranite.

It is interesting to draw a parallel with the scenario described for the Velay gneiss dome, located some 150 km ENE of the Montagne Noire. In this area, Montel et al. (1992) described two successive stages of migmatization that have been dated at  $314 \pm 5$  Ma and  $301 \pm 5$  Ma (U–Pb ID-TIMS on monazite, Mougeot et al. 1997). Barbey et al. (2015) suggest three melting events, estimated to have occurred at 325–315 Ma, ca. 305 Ma, and 305–295 Ma. In addition, Roger et al. (2015) bracketed the high-temperature deformation and metamorphism both in the Gorges d’Héric and the Vialais granite between 310 and 300 Ma.

### Tectonic implications for the formation of the Montagne Noire dome


Two major hypotheses are proposed at present to explain the origin of the Montagne Noire dome. Both agree on the presence of compressional and extensional features, but disagree on their timing and their relative importance. The first hypothesis considers that the domal structure developed as a regional anticline during the collisional stage of

the Variscan orogeny (e.g. Arthaud et al. 1966; Burg and Matte 1978; Charles et al. 2009) and interpret the extensional features as second-order and late with respect to the formation of the dome. For the second hypothesis, dome-like exhumation of the lower continental crust beneath a major crustal-scale extensional detachment is at the origin of the Montagne Noire dome (Van Den Driessche and Brun 1989; Echtler and Malavieille 1990; Van Den Driessche and Brun 1992). Beyond the structural record, this second hypothesis is supported and constrained in time by the syntectonic emplacement of a leucogranite at ca. 295 Ma (Poilvet et al. 2011) and by monazite and mica ages of ca. 295 Ma in sheared metasediments and orthogneisses (Maluski et al. 1991; Pitra et al. 2012) along the detachment that is contemporaneous with and controls the development of the Stephanian to Permian Graissessac and Lodève basins (Fig. 1; Van Den Driessche and Brun 1989; Bruguier et al. 2003). In contrast, the advocates of the compressional origin of the dome either associate the extensional structures exclusively with the compressional phase (Brunel and Languet 1997) or relegate it to a secondary role in the brittle domain (Matte et al. 1998; Charles et al. 2009). The principal argument was the age of the late- to post-kinematic Vialais granite, supposedly dated at  $327 \pm 5$  Ma (TIMS on zircon and monazite fractions; Matte et al. 1998), emplaced in the central-eastern part of the dome (Fig. 1), and the ca. 330 Ma dates obtained by electron probe micro-analysis (EPMA) monazite dating from migmatites and anatectic granites summarised in Charles et al. (2009) and Faure et al. (2010).

In the “compressional” interpretation, migmatization and magmatism predate extension. In the “extensional” interpretation, which requires thermal relaxation and related rheological softening in order for the crust to collapse, migmatization is also contemporaneous with the onset of extension, especially because extension can enhance partial melting by adiabatic decompression (e.g. Hollister 1993; Holtz and Johannes 1994; Holtz et al. 2001; Thompson 2001). Clearly, the clue is to be sought in the absolute timing of regional migmatization and related granite emplacement.

First, the Vialais granite and an associated post-kinematic leucogranite have been recently re-dated at  $303 \pm 4$  Ma and  $298 \pm 2$  Ma, respectively (U–Pb monazite ICP-MS ages, Roger et al. 2015). Second, our data support a genetic link between the migmatites and granites in the “axial zone” of the Montagne Noire gneiss dome. Three age groups were identified from the monazite U–Th–Pb data. A first event, at ca. 319 Ma, is recorded in both the La Salvetat migmatite and the Laouzas granite, and is interpreted as the end of a first stage of migmatization and as the emplacement age of the Laouzas granite, respectively. A second event, at ca. 298 Ma, is recorded in the migmatite





**Fig. 9** Th/Pb versus U/Pb plots using data for the monazite grains from the Montalet granite (a; from Poilvet et al. 2011), the Laouzas granite (b) and the La Salvetat migmatite (c). For all the diagrams,

ages obtained by LA-ICP-MS (this study) are given for comparison. *n* refers to the number of analyses plotted in the diagrams. For more explanation on these plots see Cocherie and Albarède (2001)

and in the Laouzas granite, and could be interpreted as a fluid-induced event, probably related to a second melting event identified through the emplacement of the Montalet and Vialais leucogranites. The third event, dated around 285 Ma, although not clear, could be linked to Permian fluid circulations. The presence of two stages of partial melting at ca. 320 Ma and ca. 300 Ma confirms (1) the interpretation of the Late Carboniferous–Early Permian evolution of the Variscan belt dominated by the extensional collapse, which predicts migmatization both preceding and contemporaneous of the onset of extension and (2) the interpretation of the Montagne Noire dome as an extensional gneiss dome.

### EPMA versus LA-ICP-MS dating

In recent years, some studies comparing monazite ages obtained by the LA-ICP-MS and EPMA techniques encountered discrepancies between both sets of ages (Paquette and Tiepolo 2007; Poilvet et al. 2011 and references therein). Other studies also pointed out that EPMA dating should be acquired with caution as, for example, incorrect determination of background intensities could result in artificially older ages (Jercinovic and Williams 2005; Spear et al. 2009).


This study offers, therefore, the opportunity to compare the results obtained through EPMA chemical dating and those obtained by LA-ICP-MS. Indeed, two aspects differ between our data and those of Faure et al. (2010)—(1) a minimum of two generations of monazite are distinguished in our data set whereas only one was identified in Faure et al. (2010) and (2) there is a significant difference in the absolute ages obtained by both approaches.

Although Faure et al. (2010) identified “three groups of composition according to the Th/U ratio” (p. 660) for monazite grains from the La Salvetat migmatite, they interpreted them in terms of only one chemical date of

327 ± 7 Ma, reflecting the crystallisation age of these monazite grains. This age is barely within error of the oldest age of 319.8 ± 1.8 Ma found in this study, although the latter is more precise. However, one more age at 298 Ma was obtained by LA-ICP-MS (with another possible event at ca. 285 Ma). In the case of the monazite from the Laouzas granite, EPMA dating yielded only one date at 336 ± 6 Ma, whereas at least two (potentially four) were obtained by LA-ICP-MS. In this case, Faure et al. (2010) describe their monazite as patchy zoned, a feature encountered only in our younger monazite population dated at ca. 298 Ma. The fact that they did not find two age populations in their data set could be explained if they did not encounter the concentrically zoned monazite in their samples.

These differences could also be linked to the statistical treatment applied to EPMA dating. Indeed, each individual analysis bears fairly high error, but once they are all combined to calculate a total U–Th–Pb date, the resulting error becomes relatively small ( $\leq 2\%$ ). One could therefore argue that this statistic treatment is not able to resolve different populations that are relatively close in age.

In order to test this hypothesis, we took the U, Th and Pb contents calculated for three of our samples (ES7 and ES8, this study Table 2; ES5 of Poilvet et al. 2011) and ran them into the EPMA dating add-in developed by Pommeret et al. (2002) following the data treatment described in Cocherie et al. (1998) and Cocherie and Albarède (2001). In order to test the viability of our approach, we first took the data from the Montalet granite in Poilvet et al. (2011). The monazite grains in this sample yielded a single concordia age of 294 ± 1 Ma, identical to the concordia age of 294 ± 3 Ma obtained on zircon. Plotted in a Th/Pb versus U/Pb diagram (Fig. 9a), they define a similar U–Th–Pb age of 295 ± 10 Ma. This demonstrates that when dating a simple (i.e. single age) population of monazite, the results obtained by both the LA-ICP-MS and EPMA dating techniques are comparable. We then tested this approach with two other samples from this study, which gave several age



**Fig. 10** Weighted histogram representation of the data acquired in this study for sample ES7 following the statistical 1D data treatment of Montel et al. (1996) for EPMA dating. For more explanation on this plot see Montel et al. (1996)

populations. For sample ES7, the resulting Th/Pb versus U/Pb isochron diagram (Fig. 9b) allows to calculate a single U–Th–Pb date of  $297 \pm 15$  Ma (MSWD = 0.28) at the centroid of the population. In this diagram, the regression line lies fairly close to the theoretical isochron; therefore, this age of ca. 297 Ma would have been considered as reliable and, therefore, unique. Yet, we know that in fact at least two ages can be calculated using the LA-ICP-MS data (ca. 319 Ma and ca. 298 Ma, respectively). Although these two ages are ~20 Ma apart, the Th/Pb versus U/Pb isochron diagram is not able to distinguish them. If we do the same operation with monazite data from sample ES8, we also end up with a U–Th–Pb chemical age of  $291 \pm 8$  Ma (MSWD = 0.18) at the centroid of the population (Fig. 9c) although three ages can be calculated in a conventional concordia diagram (ca. 319 Ma, 298 Ma and, maybe, 284 Ma). In this case, we reached the limits of the technique as the theoretical isochron fits just within the limits of the error envelope. This does not explain, however, why the EPMA ages found by Faure et al. (2010) are significantly older than the ages found by LA-ICP-MS.

It is interesting to note that, with the same data set, but using the procedure described in Montel et al. (1996) where, for each individual age, the 95 % confidence interval is estimated by a Monte Carlo procedure assuming U, Th and Pb content to obey a Gaussian distribution, we end up with completely different results, as the age distributions for sample ES7 yield three different date peaks at ca. 318 Ma, 302 Ma and 270 Ma (Fig. 10), which is in a good agreement with the ages found in this study.

## Conclusion

The presence of two stages of partial melting, at ca. 320 Ma and ca. 300 Ma, confirms (1) the interpretation of the Late Carboniferous–Early Permian evolution of the Variscan belt dominated by extensional collapse, which predicts migmatization both preceding and contemporaneous with the onset of extension and (2) the interpretation of the Montagne Noire dome as an extensional gneiss dome.

We also demonstrate that EPMA dating of monazite in this type of complex polyphased environment should be used with extreme caution. Indeed, as illustrated here, the Pb/U and Th/U isochron statistical data treatment (Pommier et al. 2002) can sometimes fail to resolve different age populations and can, therefore, produce erroneous results.

**Acknowledgments** Xavier Le Coz is acknowledged for making promptly the thin sections required for our study, Yann Lepagnet for the efficient rock crushing, and Thomas Delhaye for help with NanoSIMS analysis. Jean-Marc Montel and an anonymous reviewer are thanked for their constructive remarks on a previous version of this manuscript. The reviews of P. Rey and E. Oliot are acknowledged.

## References

- Alabouvette B, Demange M (1993) Notice explicative, Carte géol. France (1/50 000), feuille Saint-Pons (1013). BRGM, Orléans
- Arthaud F (1970) Etude tectonique comparée de deux domaines hercyniens: les nappes de la Montagne Noire (France) et l'anticlinorium de l'Iglesiente (Sardaigne), Thesis, University of Montpellier, Publications USTELA Géol. Struct
- Arthaud F, Matte P (1977) Late Paleozoic strike slip faulting in southern Europe and northern Africa: result of a right-lateral shear zone between the Appalachians and the Urals. *Geol Soc Am Bull* 88:1305–1320
- Arthaud F, Mattauer M, Proust F (1966) La structure et la microtectonique des nappes hercyniennes de la Montagne Noire. In: Etages tectoniques: colloque de Neuchâtel, 18–19 avril 1966, Université de Neuchâtel, pp 231–243
- Barbey P, Villaros A, Marignac C, Montel J-M (2015) Multiphase melting, magma emplacement and P-T-time path in late-collisional context: the Velay example (Massif Central, France). *Bull Soc Géol Fr* 186(2–3):93–116
- Becq-Giraudon JF, Van Den Driessche J (1993) Continuité de la sédimentation entre le Stéphanois et l'Autunien dans le bassin de Graissessac-Lodève (sud du Massif Central): implications tectoniques. *Comptes Rendus de l'Acad des Sci sér 2(317)*:939–945
- Benard F, Moutou P, Pichavant M (1985) Phase relations of tourmaline leucogranites and the significance of tourmaline in silicic magmas. *J Geol* 93:271–291
- Blès J-L, Bonijoly C, Castaing C, Gros Y (1989) Successive post-Variscan stress fields in the French Massif Central and its borders (Western European plate): comparison with geodynamical data. *Tectonophysics* 169:79–111
- Bogdanoff S, Donnot F, Ellenberger F (1984) Note explicative, Carte géol. France (1/50 000), feuille Bédarieux (988). BRGM, Orléans
- Bosse V, Boulvais P, Gautier P, Tiepolo M, Ruffet G, Devidal J-L, Cherneva Z, Gerdjikov I, Paquette J-L (2009) Fluid-induced disturbance of the monazite Th–Pb chronometer: In situ dating and

- element mapping in pegmatites from the Rhodope (Greece, Bulgaria). *Chem Geol* 261:286–302
- Boutin A, de Saint Blanquat M, Poujol M, Boulvais P, de Parseval P, Rouleau C, Robert JF (2015) Succession of Permian and Mesozoic metasomatic events in the eastern Pyrenees with emphasis on the Trimouns talc-chlorite deposit. *Int J Earth Sci*. doi:10.1007/s00531-015-1223-x
- Bruguier O, Becq-Giraudon JF, Champenois M, Deloule E, Ludden J, Mangin D (2003) Application of in situ zircon geochronology and accessory phase chemistry to constraining basin development during post-collisional extension: a case study from the French Massif Central. *Chem Geol* 201:319–336
- Brun J-P, Van Den Driessche J (1994) Extensional gneiss domes and detachment fault systems: structures and kinematics. *Bull de la Soc Géol de Fr* 165:519–530
- Brunel M, Lansigu C (1997) Déformation et cinématique de mise en place du dôme de la zone axiale de la Montagne Noire: signification des nodules à quartz-sillimanite (Massif central français) [Deformation and kinematics of emplacement of the axial dome of the Montagne Noire: implications of quartz-sillimanite nodule attitudes (French Massif Central)]. *Comptes Rendus de l'Acad des Sci Paris Ser IIA Earth Planet Sci* 325:517–523
- Burg J-P, Matte PJ (1978) A Cross section through the French Massif central and the scope of its Variscan geodynamic evolution. *Z der Dtsch Geol Ges* 129:429–460
- Burg J-P, Van Den Driessche J, Brun J-P (1994) Syn- to post-thickening extension in the Variscan Belt of Western Europe: modes and structural consequences. *Géol Fr* 3:33–51
- Cassard D, Feybesse J-L, Lescuyer J-L (1993) Variscan crustal thickening, extension and late overstacking during the Namurian-Westphalian in the western Montagne Noire. *Tectonophysics* 222:33–53
- Cathelineau M, Boiron M-C, Holliger P, Poty B (1990) Metallogensis of the French part of the Variscan orogen. Part II: time-space relationships between U, Au and Sn-W ore deposition and geodynamic events—mineralogical and U–Pb data. *Tectonophysics* 177:59–79
- Charles N, Faure M, Chen Y (2009) The Montagne Noire migmatitic dome emplacement (French Massif Central): new insights from petrofabric and AMS studies. *J Struct Geol* 31:1423–1440
- Cocherie A, Albarède F (2001) An improved U–Th–Pb age calculation for electron microprobe dating of monazite. *Geochim Cosmochim Acta* 65:4509–4522
- Cocherie A, Legendre O, Peucat J-J, Kouamelan AN (1998) Geochronology of polygenetic monazites constrained by in situ electron microprobe Th–U–total lead determination; implications for lead behaviour in monazite. *Geochim Cosmochim Acta* 62:2475–2497
- Demange M (1982) Etude géologique du massif de l'Agout, Montagne Noire. France, Thèse d'Etat, University Paris 6
- Demange M, Guérangé-Lozes J, Guérangé B (1995) Notice explicative, Carte géol. France (1/50 000), feuille Lacaune (987). Orléans, BRGM, p 153
- Didier A, Bosse V, Boulvais P, Bouloton J, Paquette J-L, Montel J-M, Devidal J-L (2013) Disturbance versus preservation of U–Th–Pb ages in monazite during fluid-rock interaction: textural, chemical and isotopic in situ study in microgranites (Velay Dome, France). *Contrib Miner Petrol* 165:1051–1072
- Doublier MP, Potel S, Wemmer K (2015) The tectono-metamorphic evolution of the very low-grade hangingwall constrains two-stage gneiss dome formation in the Montagne Noire (Southern France). *J Metamorph Geol* 33:71–89
- Echtler H (1990) Geometry and kinematics of recumbent folding and low-angle detachment in the Pardailhan nappe (Montagne Noire, Southern French Massif Central). *Tectonophysics* 177:109–123
- Echtler H, Malavieille J (1990) Extensional tectonics, basement uplift and Stephano-Permian collapse basin in a late Variscan metamorphic core complex (Montagne Noire, Southern Massif Central). *Tectonophysics* 177:125–138
- Faure M, Cottreau N (1988) Données cinématiques sur la mise en place du dôme migmatitique carbonifère moyen de la zone axiale de la Montagne noire (Massif central français). *Comptes rendus de l'Acad des Sci* 307:1787–1794
- Faure M, Cocherie A, Bé Mézème E, Charles N, Rossi P (2010) Middle Carboniferous crustal melting in the Variscan belt: new insights from U–Th–Pb<sub>cor</sub> monazite and U–Pb zircon ages of the Montagne Noire Axial Zone (southern French Massif Central). *Gondwana Res* 18:653–673
- Foster G, Kinny P, Vance D, Prince C, Harris N (2000) The significance of monazite U–Th–Pb age data in metamorphic assemblages; a combined study of monazite and garnet chronometry. *Earth Planet Sci Lett* 181:327–340
- Franke W, Doublier MP, Klama K, Potel S, Wemmer K (2011) Hot metamorphic core complex in a cold foreland. *Int J Earth Sci* 100:753–785
- Gardés E, Montel J-M, Seydoux-Guillaume A-M, Wirth R (2007) Pb diffusion in monazite: New constraints from the experimental study of Pb<sup>2+</sup> ↔ Ca<sup>2+</sup> interdiffusion. *Geochim Cosmochim Acta* 71:4036–4043
- Gasquet D, Bertrand J-M, Paquette J-L, Lehmann J, Ratzov G, De Ascensão Guedes R, Tiepolo M, Boullier AM, Scaillet S, Nomade S (2010) Miocene to Messinian deformation and hydrothermalism in the Lauzière Massif (French Western Alps): New U–Th–Pb and Argon ages. *Bull de la Soc Géol de Fr* 181:227–241
- Hollister LS (1993) The role of melt in the uplift and exhumation of orogenic belts. *Chem Geol* 108:31–48
- Holtz F, Johannes W (1994) Maximum and minimum water contents of granitic melts: implications for chemical and physical properties of ascending magmas. *Lithos* 32:149–159
- Holtz F, Johannes W, Tamic N, Behrens H (2001) Maximum and minimum water contents of granitic melts generated in the crust: reevaluation and implications. *Lithos* 56:1–14
- Hurav V, Paquette J-L, Huraiová M, Konečný P (2010) U–Th–Pb geochronology of zircon and monazite from syenite and pincinite xenoliths in Pliocene alkali basalts of the intra-Carpathian back-arc basin. *J Volcanol Geoth Res* 198:275–287
- Jackson SE, Pearson NJ, Griffin WL, Belousova EA (2004) The application of laser ablation-inductively coupled plasma-mass spectrometry to in situ U–Pb zircon geochronology. *Chem Geol* 211:47–69
- Jercinovic MJ, Williams ML (2005) Analytical perils (and progress) in electron microprobe trace elements analysis applied to geochronology: background, interferences and beam irradiation effects. *Am Miner* 90:526–546
- Jourdan F, Renne PR (2007) Age calibration of the Fish Canyon sanidine <sup>40</sup>Ar/<sup>39</sup>Ar dating standard using primary K–Ar standards. *Geochim Cosmochim Acta* 71:387–402
- Jourdan F, Verati C, Féraud G (2006) Intercalibration of the Hb3gr <sup>40</sup>Ar/<sup>39</sup>Ar dating standard. *Chem Geol* 231:77–189
- Kelsey DE, Clark C, Hand M (2008) Thermobarometric modelling of zircon and monazite growth in melt-bearing systems: examples using model metapelitic and metapsammitic granulites. *J Metamorph Geol* 26:199–212
- Ludwig KR (1998) On the treatment of concordant uranium-lead ages. *Geochim Cosmochim Acta* 62:665–676
- Ludwig KR (2001) User's manual for Isoplot/Ex Version 2.49, a geochronological toolkit for Microsoft Excel. Spec Publ. 1a. Berkeley Geochronological Center, Berkeley, USA
- Maluski H, Costa S, Echtler H (1991) Late Variscan tectonic evolution by thinning of earlier thickened crust. An <sup>40</sup>Ar–<sup>39</sup>Ar study of the Montagne Noire, southern Massif Central, France. *Lithos* 26:287–304

- Matte P, Lancelot J, Mattauer M (1998) The Montagne Noire Axial Zone is not an extensional metamorphic core complex but a compressional post-nappe anticline with an anatectic core. *Geodin Acta* 11:3–22
- Ménard G, Molnar P (1988) Collapse of a Hercynian Tibetan plateau into a late Paleozoic European Basin and Range province. *Nature* 334:235–237
- Montel J-M, Marignac C, Barbey P, Pichavant M (1992) Thermo-barometry and granite genesis: the Hercynian low-P high-T Velay anatectic Dome (French Massif Central). *J Metamorph Geol* 10:1–15
- Montel J, Foret S, Veschambre M, Nicollet C, Provost A (1996) Electron microprobe dating of monazite. *Chem Geol* 131:37–53
- Mougeot R, Respaut J-P, Ledru P, Marignac C (1997) U–Pb geochronology on accessory minerals of the Velay anatectic Dome (French Massif Central). *Eur J Miner* 9:141–156
- Paquette J-L, Tiepolo M (2007) High resolution (5  $\mu\text{m}$ ) U–Th–Pb isotopes dating of monazite with excimer laser ablation (ELA)-ICPMS. *Chem Geol* 240:222–237
- Pitra P, Ballèvre M, Ruffet G (2010) Inverted metamorphic field gradient towards a Variscan suture zone (Champtoceaux Complex, Armorican Massif, France). *J Metamorph Geol* 28:183–208
- Pitra P, Poujol M, Van Den Driessche J, Poilvet J-C, Paquette J-L (2012) Early Permian extensional shearing of an Ordovician granite: the Saint-Eutrope, “C/S-like” orthogneiss (Montagne Noire, French Massif Central). *CR Geosci* 344:377–384
- Poilvet J-C, Poujol M, Pitra P, Van Den Driessche J, Paquette J-L (2011) The Montalet granite, Montagne Noire, France: an early Permian syn-extensional pluton as evidenced by new U–Th–Pb data on zircon and monazite. *CR Geosci* 343:454–461
- Poitrasson F, Chenery S, Shepherd TJ (2000) Electron microprobe and LA-ICP-MS study of monazite hydrothermal alteration: implications for U–Th–Pb geochronology and nuclear ceramics. *Geochimica and Cosmochimica Acta* 64:3283–3297
- Pommier A, Cocherie A, Legendre O (2002) EPMA dating user’s manual: age calculation from electron probe microanalyser measurements of U–Th–Pb. BRGM internal report, p 9
- Rabin M, Trap P, Carry N, Fréville K, Centi-Tok B, Lobjoie C, Goncalves P, Marquer D (2015) Strain partitioning along the anatectic front in the Variscan Montagne Noire massif (southern French Massif Central). *Tectonics*. doi:10.1002/2014TC003790
- Renne PR, Mundil R, Balco G, Min K, Ludwig RL (2010) Joint determination of  $^{40}\text{K}$  decay constants and  $^{40}\text{Ar}^*/^{40}\text{K}$  for the Fish Canyon sanidine standard, and improved accuracy for  $^{40}\text{Ar}/^{39}\text{Ar}$  geochronology. *Geochim Cosmochim Acta* 74:5349–5367
- Renne PR, Balco G, Ludwig RL, Mundil R, Min K (2011) Response to the comment by W.H. Schwarz et al. on “Joint determination of  $(^{40}\text{K})$  decay constants and  $(^{40}\text{Ar})^*/(^{40}\text{K})$  for the Fish Canyon sanidine standard, and improved accuracy for  $(^{40}\text{Ar})/(^{39}\text{Ar})$  geochronology” by PR Renne et al. (2010). *Geochim Cosmochim Acta* 75:5097–5100
- Rey PF, Teyssier C, Whitney DL (2011) Viscous collision in channel explains double domes in metamorphic core complexes. *Geology* 39:387–390
- Roddick JC (1983) High precision intercalibration of  $^{40}\text{Ar}/^{39}\text{Ar}$  standards. *Geochim Cosmochim Acta* 47:887–898
- Roger F, Respaut J-P, Brunel M, Matte P, Paquette J-L (2004) U–Pb dating of Augen orthogneisses from the Axial Zone of the Montagne Noire (Southern Massif Central): new witness of Ordovician magmatism into the Variscan Belt. *CR Geosci* 336:19–28
- Roger F, Teyssier C, Respaut J-P, Rey PF, Jolivet M, Whitney DL, Paquette J-L, Brunel M (2015) Timing of formation and exhumation of the Montagne Noire double dome, French Massif Central. *Tectonophysics* 640–641:53–69
- Rubatto D, Williams IS, Buick IS (2001) Zircon and monazite response to prograde metamorphism in the Reynolds Range, central Australia. *Contrib Miner Petrol* 140:458–468
- Ruffet G, Féraud G, Amouric M (1991) Comparison of  $^{40}\text{Ar}/^{39}\text{Ar}$  conventional and laser dating of biotites from the North Trégor Batholith. *Geochim Cosmochim Acta* 55:1675–1688
- Ruffet G, Féraud G, Ballèvre M, Kiénast JR (1995) Plateau ages and excess argon on phengites: a  $^{40}\text{Ar}/^{39}\text{Ar}$  laser probe study of alpine micas (Sesia zone). *Chem Geol* 121:327–343
- Seydoux-Guillaume A-M, Paquette J-L, Wiedenbeck M, Montel J-M, Heinrich W (2002) Experimental resetting of the U–Th–Pb systems in monazite. *Chem Geol* 191:165–181
- Schuiling RD (1960) Le dôme gneissique de l’Agout (Tarn et Hérault). *Mém Soc géol France* 39(91):1–59
- Soula JC, Debat P, Brusset S, Bessière G, Christophoul F, Déramond J (2001) Thrust-related, diapiric, and extensional doming in a frontal orogenic wedge: example of the Montagne noire, Southern French Hercynian Belts. *J Struct Geol* 23:1677–1699
- Spear FS, Pyle JM (2010) Theoretical modeling of monazite growth in a low-Ca metapelite. *Chem Geol* 273:111–119
- Spear FS, Pyle JM, Cherniak D (2009) Limitations of chemical dating of monazite. *Chem Geol* 266:218–230
- Taner MF, Martin RF (1993) Significance of dumortierite in an aluminosilicate-rich alteration zone, Louvicourt, Quebec. *Can Miner* 31:137–146
- Tartèse R, Ruffet G, Poujol M, Boulvais P, Ireland TR (2011) Simultaneous resetting of the muscovite K–Ar and monazite U–Pb geochronometers: a story of fluids. *Terra Nova* 23:390–398
- Tartèse R, Boulvais P, Poujol M, Chevalier T, Paquette J-L, Ireland TR, Deloule E (2012) Mylonites of the South Armorican Shear Zone: insights for crustal-scale fluid flow and water-rock interaction processes. *J Geodyn* 56–57:86–107
- Thompson AB (2001) Clockwise P–T paths for crustal melting and  $\text{H}_2\text{O}$  recycling in granite source regions and migmatite terrains. *Lithos* 56:33–45
- Turner G, Huneke JC, Podosek FA, Wasserburg GJ (1971)  $^{40}\text{Ar}/^{39}\text{Ar}$  ages and cosmic ray exposure age of Apollo 14 samples. *Earth Planet Sci Lett* 12:19–35
- Van Den Driessche J, Brun J-P (1989) Un modèle cinématique de l’extension paléozoïque supérieur dans le sud du Massif Central. *Comptes Rendus de l’Acad des Sci Sér 2*(309):1607–1613
- Van Den Driessche J, Brun J-P (1992) Tectonic evolution of the Montagne Noire (French Massif Central): a model of extensional dome. *Geodin Acta* 5:85–99
- Van Den Driessche J, Pitra P (2012) Viscous collision channel explains double dome in metamorphic core complexes: comment. *Geology* 40:e279
- Velichkin VI, Vlasov BP (2011) Domal structures and hydrothermal uranium deposits of the Erzgebirge, Saxony, Germany. *Geol Ore Depos* 53:74–83
- Wiedenbeck M, Allé P, Corfu F, Griffin WL, Meier M, Oberli F, von Quadt A, Roddick JC, Spiegel W (1995) Three natural zircon standards for U–Th–Pb, Lu–Hf, trace element and REE analyses. *Geostand Newslett* 19:1–23
- Williams ML, Jercinovic MJ, Harlov DE, Budzyn Hetherington CJ (2011) Resetting monazite ages during fluid-related alteration. *Chem Geol* 283:218–225