

# The impact of patatin-like phospholipase domain-containing protein 3 polymorphism on hepatocellular carcinoma prognosis

Yasuto Takeuchi · Fusao Ikeda · Yuki Moritou · Hiroaki Hagihara · Tetsuya Yasunaka · Kenji Kuwaki · Yasuhiro Miyake · Hideki Ohnishi · Shinichiro Nakamura · Hidenori Shiraha · Akinobu Takaki · Yoshiaki Iwasaki · Kazuhiro Nouse · Kazuhide Yamamoto

Received: 3 June 2012 / Accepted: 9 July 2012 / Published online: 7 August 2012  
© Springer 2012

## Abstract

**Background** The single nucleotide polymorphism (SNP) rs738409 in patatin-like phospholipase domain-containing protein 3 (PNPLA3) is associated with hepatic fat accumulation and disease progression in patients with non-alcoholic fatty liver disease and alcoholic liver disease (ALD). This study was conducted to determine whether PNPLA3 rs738409 SNPs affect development and prognosis of hepatocellular carcinoma (HCC) in patients with various liver diseases.

**Methods** We enrolled 638 consecutive Japanese patients newly diagnosed with HCC between 2001 and 2010: 72 patients with hepatitis B virus (HBV), 462 with hepatitis C virus (HCV), and 104 with non-B non-C (NBNC).

**Results** NBNC patients exhibited large tumors of advanced TNM stages at HCC diagnosis, and had significantly poorer prognosis than HBV or HCV patients

( $P < 0.001$  and  $< 0.001$ , respectively; log-rank test). The G/G genotype of PNPLA3 rs738409 SNP had significantly higher distribution in NBNC patients ( $P < 0.001$ ) and was significantly associated with higher body mass index (BMI) and an increased aspartate aminotransferase to platelet ratio index. No significant differences were observed in survival with differences in PNPLA3 SNP genotypes among the patients, although ALD patients with the G/G genotype of PNPLA3 SNP and low BMI had significantly poorer survival than those with high BMI ( $P = 0.028$ ).

**Conclusions** The G/G genotype of PNPLA3 rs738409 SNP was more frequently distributed, and associated with BMI and fibrosis among NBNC-HCC patients but not among HBV or HCV patients. These genotypes might affect HCC prognosis in ALD patients, but not in HBV, HCV, or NAFLD patients.

**Keywords** Hepatocellular carcinoma · PNPLA3 SNP · Survival

Y. Takeuchi · F. Ikeda (✉) · Y. Moritou · H. Hagihara · T. Yasunaka · K. Kuwaki · Y. Miyake · H. Ohnishi · S. Nakamura · H. Shiraha · A. Takaki · K. Nouse · K. Yamamoto

Department of Gastroenterology and Hepatology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, 2-5-1 Shikata-cho, Okayama 700-8558, Japan  
e-mail: fikeda@md.okayama-u.ac.jp

F. Ikeda · Y. Miyake · H. Ohnishi · K. Nouse · K. Yamamoto  
Department of Molecular Hepatology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, Japan

Y. Iwasaki  
Health and Environment Center, Okayama University, Okayama, Japan

## Abbreviations

AFP	Alpha-fetoprotein
ALD	Alcoholic liver disease
APRI	Aspartate aminotransferase to platelet ratio index
BMI	Body mass index
DCP	Des-gamma-carboxy prothrombin
HBV	Hepatitis B virus
HCC	Hepatocellular carcinoma
HCV	Hepatitis C virus
NAFLD	Non-alcoholic fatty liver disease
NBNC	Non-B non-C
PNPLA3	Patatin-like phospholipase domain-containing protein 3
SNP	Single nucleotide polymorphism

## Introduction

Hepatocellular carcinoma (HCC) is the fifth most common cancer and the third most common cause of cancer-related deaths worldwide [1–3]. Various factors, including chronic viral infection, age, sex, alcoholic liver disease (ALD), and diabetes mellitus, are involved in HCC development and progression. Hepatitis C virus (HCV) and/or hepatitis B virus (HBV) infections are the major cause of HCC development in Japan [4–6]. Recently, the incidence of HCC without HBV or HCV infection has increased, accounting for 12–16 % of HCC patients in Japan [7, 8]. Obesity is the leading cause of non-alcoholic fatty liver disease (NAFLD) [9] and represents the second major cause of cancer development in western countries and Japan [10, 11]. Almost 25 % of all HCC cases in western countries, and 8–15 % in Japan are estimated to originate from NAFLD [12, 13]. Because effective predictive biomarkers for non-B non-C (NBNC) HCC development in NAFLD patients have not been clarified, NBNC-HCC is often diagnosed in advanced stages with poor prognosis [14].

There is compelling evidence that host genetic polymorphisms are involved in HCC pathogenesis and may be useful as predictive biomarkers [15]. The associations of functional genetic polymorphisms of interleukin 6 and tumor necrosis factor alpha with HCC development have been reported [16, 17]. In the present study, we focused on the association of patatin-like phospholipase domain-containing protein 3 (PNPLA3) genetic polymorphisms with HCC prognosis. PNPLA3, also called as adiponutrin, is a member of the patatin-like phospholipase family and its activity is regulated by the same hormonal pathways that regulate fat deposition in the liver [18–20]. Non-synonymous single nucleotide polymorphism (SNP) rs738409 C>G transversion (Ile148Met) in exon 3 of the gene encoding for PNPLA3 has been reported to be associated with NAFLD development [21]. Moreover, the PNPLA3 rs738409 SNP is related to the degree of fibrosis observed in ALD and NAFLD [22–24]; however, it is not clear whether this polymorphism affects HCC development and prognosis in NAFLD and ALD. The present study was conducted to determine whether rs738409 C>G non-synonymous PNPLA3 SNP is associated with HCC development and prognosis in HBV, HCV, or NBNC patients.

## Methods

### Patients

The present study enrolled 638 consecutive Japanese patients newly diagnosed with HCC at the Okayama University Hospital between 2001 and 2010. Liver disease

etiologies were as follows: 462 patients with HCV-related disease, 72 with HBV-related disease, and 104 with non-B non-C (NBNC) disease. NBNC etiologies included metabolic disorders such as NAFLD in 57 patients, ALD in 34 patients, and autoimmune liver diseases and cryptogenic liver injury in 13 patients. Heavy drinking behavior was defined as daily alcohol intake of at least 70 g for ALD patients, and non-drinking behavior was defined as daily alcohol intake of less than 20 g for NAFLD patients. Fifty NAFLD patients without HCC were also included in the study. The study was conducted in accordance with the Declaration of Helsinki, and all protocols were approved by the ethics committees of the institutes. All patients provided informed consent before study enrollment.

### HCC diagnosis and follow-up

In accordance with the clinical practice manual from the Japan Society of Hepatology [25, 26], HCC diagnosis was accomplished using imaging techniques, including ultrasonography, computed tomography, magnetic resonance imaging, hepatic angiography, and/or tumor biopsy, in combination with the detection of serum levels of alpha-fetoprotein (AFP), AFP-L3, and des-gamma-carboxy prothrombin (DCP). HCC staging conformed to the standards of general rules for clinical and pathological study of primary liver cancer by the Liver Cancer Study Group of Japan [27]. The cancer was treated following suitable measures according to the treatment algorithm for HCC in the clinical practice manual from the Japan Society of Hepatology [18, 19]. After curative treatment of primary HCC, all patients underwent liver function tests every 1–3 months and ultrasonography or dynamic study every 3–4 months. Serum levels of AFP, AFP-L3, and DCP were also determined every 1–3 months. HCC recurrence was diagnosed using the same criteria as those used for primary HCC diagnosis. HCC recurrence was treated following suitable measures.

### SNP genotyping

Genomic DNA was extracted from whole blood samples using the QIAamp DNA Mini Kit according to the manufacturer's protocol (Qiagen, Tokyo, Japan). PNPLA3 SNP was genotyped using the TaqMan predesigned SNP genotyping assays, as recommended by the manufacturer (Applied Biosystems, Tokyo, Japan).

### Statistical analysis

Data are expressed as mean  $\pm$  standard deviation or median (range). Patient laboratory data were compared among the groups using the Kruskal–Wallis test. Survival

rates of patients were estimated using the Kaplan–Meier method and compared with the log-rank test.  $P < 0.05$  was considered significant. Statistical analysis was performed using JMP software (SAS Institute, Cary, NC, USA).

**Results**

**Patient characteristic at diagnosis**

HCC development occurred at a much younger age in HBV patients than in HCV or NBNC patients ( $P < 0.001$ , Kruskal–Wallis test, Table 1). The majority of NBNC and HBV patients were male ( $P < 0.001$ ). Body mass index (BMI) of NBNC patients was significantly higher than that of HBV or HCV patients ( $P = 0.001$ ). The NBNC patients also exhibited the highest incidence of diabetes mellitus (38.5 %). The incidence of diabetes mellitus was much lower among HCV (18.8 %) and HBV patients (4.0 %, lowest,  $P < 0.001$ ). In terms of drinking behavior, 32.7 % of NBNC patients drank heavily, whereas only 5.5 % of HBV patients and 11.0 % of HCV patients ( $P < 0.001$ ) were heavy drinkers. No significant differences were observed among the experimental groups regarding the ratio of patients with Child–Pugh A ( $P = 0.44$ ). Significantly higher serum levels of aspartate aminotransferase and alanine aminotransferase as well as lower platelet counts were observed in HCV patients ( $P < 0.001$ , 0.003, and  $<0.001$ , respectively). Gamma-glutamyl transpeptidase

was significantly higher in NBNC patients than in HBV or HCV patients ( $P < 0.001$ ).

**HCC diagnosis and prognosis**

At diagnosis, NBNC patients exhibited a higher incidence of large HCC tumors of advanced TNM stages as well as higher serum DCP levels than HBV or HCV patients ( $P < 0.001$ , Table 1). After curative treatment of primary HCC, all patients underwent periodical liver function and imaging tests at follow-up. The mean follow-up period was approximately 3.0 years (range 0.1–9.5 years). This period was defined as the time between diagnosis and death or the latest confirmation of survival. The accumulated observation time was 1,914 person-years. The survival rate of NBNC patients was significantly lower than that of HBV or HCV patients ( $P = 0.011$ , log-rank test, Fig. 1), with an estimated 68.7 % survival at 3 years and 41.8 % survival at 5 years. The survival of HCV or NBNC patients was not dependent on TNM stage (Fig. 2). The survival of HBV patients on TNM stage II was significantly better than that of HCV or NBNC patients ( $P = 0.044$ ).

**Subgroup analysis based on PNPLA3 rs738409 SNP**

The ratio of patients with the G/G genotype of PNPLA3 SNP was significantly greater among NBNC patients (46.2 %) than among HBV or HCV patients (16.7 and 24.0 %, respectively,  $P < 0.001$ , Table 1). Age and gender

**Table 1** Patient characteristics


	HBV (n = 72)	HCV (n = 462)	NBNC (n = 104)	P
Age (years) <sup>a</sup>	61 (38–87)	75 (42–93)	71 (27–91)	<0.0001
Gender (female/male)	11/61	151/311	22/82	<0.0001
Diabetes mellitus (yes)	3	87	40	<0.0001
Heavy drinking behavior (yes)	4	51	34	<0.0001
Body mass index <sup>b</sup>	23.2 ± 3.5	22.7 ± 3.1	24.6 ± 4.3	0.0001
PNPLA rs738409 (CC/CG/GG)	20/40/12	125/226/111	17/39/48	<0.0001
Child–Pugh score (A/B/C)	57/13/2	332/118/12	83/19/2	0.14
APRI <sup>b</sup>	4.5 ± 3.8	7.0 ± 7.0	4.6 ± 4.6	<0.0001
TNM stage (1/2/3/4)	24/24/13/11	161/155/111/35	22/32/24/26	<0.0001
Tumor size (mm) <sup>b</sup>	22 ± 3.6	20 ± 0.9	32 ± 4.0	<0.0001
Tumor number (solitary)	36	260	52	0.36
Laboratory data				
ALT (IU/l) <sup>b</sup>	40 ± 23	57 ± 39	46 ± 44	0.003
γGTP (IU/l) <sup>b</sup>	103 ± 129	85 ± 95	143 ± 146	<0.0001
Albumin (g/dl) <sup>b</sup>	3.9 ± 0.6	3.6 ± 0.5	3.7 ± 0.5	<0.0001
Total bilirubin (mg/dl) <sup>b</sup>	1.1 ± 0.7	1.0 ± 0.6	1.1 ± 0.7	0.15
Platelet count (×10 <sup>4</sup> /μl) <sup>b</sup>	16.2 ± 8.7	12.4 ± 5.9	13.4 ± 6.2	<0.0001
AFP (ng/ml) <sup>a</sup>	10 (0.9–67,951)	19 (0.5–85,292)	26 (2–455,560)	0.54
DCP (mAU/ml) <sup>a</sup>	46 (2–410,500)	36 (10–708,400)	126 (11–1,323,600)	<0.0001
Anti-HBc antibody (yes)		135	35	0.44

TNM tumor–node–metastasis stage according to the Liver Cancer Study Group of Japan, APRI aspartate aminotransferase to platelet ratio index, ALT alanine aminotransferase, γGTP γ-glutamyl transpeptidase, AFP alpha-fetoprotein, DCP des-gamma-carboxy prothrombin


<sup>a</sup> Median (range)

<sup>b</sup> Mean ± standard deviation

did not significantly correlate with PNPLA3 SNP genotype distribution among HBV, HCV, or NBNC patients. Among NBNC patients, patients with the G/G genotype of PNPLA3 SNP had significantly higher BMI than those with



**Fig. 1** Patient survival rates in the three groups. The patient survival rate was presumed for each group by the Kaplan–Meier method. The survival of NBNC patients was significantly poorer than that of HBV or HCV patients ( $P = 0.011$ , log-rank test), and it was estimated as 68.7 % at 3 years and 41.8 % at 5 years



**Fig. 2** Comparison of survival rates among the three groups in different TNM stages. By dividing the patients on the basis of the TNM stage at diagnosis, the patient survival rate was compared among the three groups of HBV, HCV, and NBNC etiology for each TNM stage. No prognostic differences were observed among the three

the C/C or C/G genotype ( $P < 0.001$ , Table 2), whereas no significant correlation was observed between HBV or HCV patients regarding BMI and PNPLA3 SNP genotypes. As for laboratory data, NBNC patients with the G/G genotype exhibited a slight tendency toward smaller tumors ( $P = 0.047$ ), and higher aspartate aminotransferase to platelet ratio index (APRI) ( $P = 0.054$ ) than those with C/C or C/G genotype; however, no significant differences were observed in the TNM stage and serum levels of AFP and DCP. We also compared patient survival to the distribution of the different PNPLA3 SNP genotypes (Fig. 3). No significant correlation was observed between survival and the genotypes among HBV, HCV, or NBNC patients. For HBV patients, the viral status including HBe antigen, HBs antigen, and HBV DNA at diagnosis did not differ on the basis of the different PNPLA3 SNP genotypes. We further investigated ALD or NAFLD patients among NBNC patients. The ratio of patients with the G/G genotype of PNPLA3 SNP was significantly greater among NAFLD patients (56.1 %) than the other groups with ALD

groups for TNM stage I, III, or IV ( $P = 0.13$ , 0.36, and 0.64, respectively, log-rank test). The survival of HBV patients on TNM stage II was significantly better than that of HCV or NBNC patients ( $P = 0.044$ )


**Table 2** Characteristics of NBNC patients with PNPLA3 rs738409 polymorphisms

	rs738409 SNP genotype		P
	C/C or C/G (n = 56)	G/G (n = 48)	
Age (years) <sup>a</sup>	70 (27–91)	72 (46–83)	0.85
Gender (female/male)	10/46	12/36	0.26
Diabetes mellitus (yes)	20	20	0.34
Heavy drinking behavior (yes)	22	12	0.09
Body mass index <sup>b</sup>	23.6 ± 3.2	26.3 ± 3.8	<0.0001
Child–Pugh score (A/B/C)	44/8/4	39/6/3	0.46
APRI <sup>b</sup>	2.4 ± 2.2	3.6 ± 3.1	0.054
TNM stage (1/2/3/4)	13/15/9/19	7/14/14/13	0.21
Tumor size (mm)	38 (10–200)	28 (5–100)	0.047
Laboratory data			
ALT (IU/l) <sup>b</sup>	38 ± 34	40 ± 22	0.66
γGTP (IU/l) <sup>b</sup>	101 ± 95	85 ± 82	0.68
Albumin (g/dl) <sup>b</sup>	3.7 ± 0.6	3.7 ± 0.5	0.92
Total bilirubin (mg/dl) <sup>b</sup>	0.9 ± 0.5	1.0 ± 0.7	0.35
Platelet count (×10 <sup>4</sup> /μl) <sup>b</sup>	16.1 ± 6.9	13.7 ± 6.7	0.08
AFP (ng/ml) <sup>a</sup>	8.1 (0.9–67,951)	10.1 (2–37,387)	0.34
DCP (mAU/ml) <sup>a</sup>	183.5 (16–472,100)	100 (11–1,323,600)	0.33
Anti-HBc antibody (yes)	17	18	0.29

TNM tumor–node–metastasis stage according to the Liver Cancer Study Group of Japan, APRI aspartate aminotransferase to platelet ratio index, ALT alanine aminotransferase, γGTP γ-glutamyl transpeptidase, AFP alpha-fetoprotein, DCP des-gamma-carboxy prothrombin

<sup>a</sup> Median (range)

<sup>b</sup> Mean ± standard deviation



**Fig. 3** Comparison of survival rates based on the different PNPLA3 SNP genotypes in the three groups of HBV, HCV, and NBNC etiology. The PNPLA3 SNP genotype was not significantly related to survival rates among HBV, HCV, or NBNC patients

or other etiologies (35.3 and 30.8 %, respectively,  $P = 0.046$ , Table 3). As presented in Fig. 4, ALD patients with the G/G genotype of PNPLA3 SNP who had BMI of

less than 25 kg/m<sup>2</sup> exhibited significantly lower survival rates than those who had BMI of at least 25 kg/m<sup>2</sup> ( $P = 0.028$ , log-rank test). A similar association was not

observed for ALD patients with the C/C or C/G genotype or NAFLD patients. To investigate the association of PNPLA3 SNP with carcinogenesis, the NAFLD patients with HCC were compared to those without HCC in terms of the prevalence of the G/G genotype of PNPLA3 SNP. The genotypes of PNPLA3 SNP in the NAFLD patients without HCC were C/C in 3 patients (6.0 %), C/G in 22 patients (44.0 %), and G/G in 25 patients (50.0 %), which

did not differ significantly from those in the NAFLD patients with HCC ( $P = 0.61$ ).


## Discussion

HCC is commonly associated with HBV or HCV infections; however, the incidence of NBNC-HCC with NAFLD or ALD has increased significantly in recent years [12, 13]. The incidence of NBNC-HCC in the present study supports the trend that non-viral-associated HCC is increasing as the ratio of NBNC-HCC has increased from 11.8 % between 2001 and 2005 to 15.1 % between 2006 and 2010. Several studies have attempted to elucidate useful predictors for NBNC-HCC; however, these studies were not completely successful. Recent genome-wide association studies revealed that PNPLA3 rs738409 SNPs are associated with hepatic steatosis and fibrosis [21, 28]. PNPLA3 (rs738409 C>G) SNP can modulate hepatic inflammation and fibrosis progression [29]. The G allele of PNPLA3 SNP is associated with increased steatosis [30–32] and severe fibrosis [22, 27, 33–35]. Liver cirrhosis and steatohepatitis are

**Table 3** Frequency of PNPLA3 rs738409 polymorphisms in NBNC patients

	rs738409 SNP genotype		
	C/C (n = 17)	C/G (n = 39)	G/G (n = 48)
NAFLD	5 (8.8 %)	20 (35.1 %)	32 (56.1 %)
Alcoholic liver disease	7 (20.6 %)	15 (44.1 %)	12 (35.3 %)
Disease for other etiologies	5 (38.4 %)	4 (30.8 %)	4 (30.8 %)

PNPLA3 patatin-like phospholipase domain-containing protein 3, NBNC non-B non-C, SNP single nucleotide polymorphism, NAFLD non-alcoholic fatty liver disease



**Fig. 4** Comparison of survival rates based on different BMI for NBNC patients with ALD or NAFLD showing the impact of PNPLA3 SNP on survival. Among ALD patients with the G/G genotype, patients with BMI of less than 25 kg/m<sup>2</sup> had significantly poorer

survival than those with BMI of at least 25 kg/m<sup>2</sup> ( $P = 0.028$ , log-rank test). ALD patients with the C/C or C/G genotype of PNPLA3 or NAFLD patients showed no prognostic difference in BMI

associated with carcinogenesis and increased susceptibility to HCC development; however, no published studies have focused on the role of PNPLA3 SNP genotypes in HCC development and prognosis. Therefore, we investigated PNPLA3 SNP genotypes in HCC patients to clarify the importance of this polymorphism as a predictor for HCC development and prognosis.

Our results of patient characteristics at HCC diagnosis revealed that NBNC-HCC patients exhibited a significantly higher incidence of the G/G genotype than HCV or HBV patients with a higher incidence of diabetes mellitus, obesity, and alcoholic liver injury, supporting the results of previous studies [36–38]. Furthermore, subgroup analysis revealed that the NBNC-HCC patients with the G/G genotype had higher BMI and APRI, reflecting obesity and severe fibrosis, than those with the other genotypes, and that the patients with NAFLD had a higher frequency of the G/G genotype than those with ALD. In contrast, among HBV or HCV patients with HCC, the presence of obesity or severe hepatic fibrosis showed no significant correlation with the PNPLA3 SNP genotypes. Therefore, these results indicated that among the patients with HCC, the G/G genotype of PNPLA3 SNP is strongly associated with NAFLD with obesity and severe fibrosis, which is a similar situation in those without HCC in the previous studies.

Our results of HCC prognosis clearly showed that no significant association was observed between HCC prognosis and the PNPLA3 SNP genotypes in HCV or NAFLD patients. Interestingly, subgroup analysis of ALD patients indicated prognostic differences according to the genotypes of PNPLA3 SNP; the ALD patients with the G/G genotype of PNPLA3 SNP and high BMI might maintain body weight against HCC progression and malnutrition, and resulted in better prognosis, by comparing to the ALD patients with low BMI, although the number of ALD patients might be relatively small.

In conclusion, our results indicate that differences in the distribution of PNPLA3 SNP genotypes have important and independent roles in the regulation of various liver disease etiologies and HCC prognosis. While the G/G genotype showed greater distribution in NBNC-HCC patients, only among ALD patients was the G/G genotype, but not the C/C and C/G genotypes, associated with HCC prognosis. More research is necessary to further clarify the exact manner in which differences in expression of PNPLA3 SNP genotypes may regulate liver etiology as well as HCC development and prognosis. Longitudinal study of HCC development in patients without HCC is required.

**Acknowledgments** This work was supported in part by the research program of intractable disease provided by the Ministry of Health, Labor, and Welfare of Japan, and by a grant-in-aid (22590733) for

scientific research from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to Y. I.).

**Conflict of interest** The authors have no conflict of interest.

## References

1. El-Serag HB, Marrero JA, Reddy KR, et al. Diagnosis and treatment of hepatocellular carcinoma. *Gastroenterology*. 2008;134:1752–63.
2. Kawaguchi T, Taniguchi E, Sata M, et al. The pathogenesis, complications and therapeutic strategy for hepatitis C virus-associated insulin resistance in the era of anti-viral treatment. *Rev Recent Clin Trials*. 2010;5:147–57.
3. Starley BQ, Calcagno CJ, Harrison SA. Nonalcoholic fatty liver disease and hepatocellular carcinoma: a weighty connection. *Hepatology*. 2010;51:1820–32.
4. Fattovich G, Stroffolini T, Donato F, et al. Hepatocellular carcinoma in cirrhosis: incidence and risk factors. *Gastroenterology*. 2004;127:S35–50.
5. Hassan MM, Hwang LY, Patt YZ, et al. Risk factors for hepatocellular carcinoma: synergism of alcohol with viral hepatitis and diabetes mellitus. *Hepatology*. 2002;36:1206–13.
6. Kiyosawa K, Umemura T, Tanaka E, et al. Hepatocellular carcinoma: recent trends in Japan. *Gastroenterology*. 2004;127:S17–26.
7. Taura N, Fukushima N, Taketomi A, et al. The incidence of hepatocellular carcinoma associated with hepatitis C infection decreased in Kyushu area. *Med Sci Monit*. 2010;17:PH7–11.
8. Abe H, Yoshizawa K, Aizawa Y, et al. Etiology of non-B non-C hepatocellular carcinoma in the eastern district of Tokyo. *J Gastroenterol*. 2008;43:967–74.
9. Marchesini G, Bugianesi E, Rizzetto M, et al. Nonalcoholic fatty liver, steatohepatitis, and the metabolic syndrome. *Hepatology*. 2003;37:917–23.
10. James WP. The epidemiology of obesity: the size of the problem. *J Intern Med*. 2008;263:336–52.
11. Wolin KY, Carson K, Colditz GA. Obesity and cancer. *Oncologist*. 2010;15:556–65.
12. Hui JM, Kench JG, George J, et al. Long-term outcomes of cirrhosis in nonalcoholic steatohepatitis compared with hepatitis C. *Hepatology*. 2003;38:420–7.
13. Sugimoto K, Takei Y. Clinicopathological features of non-alcoholic fatty liver disease. *Hepatol Res*. 2011;41:911–20.
14. Nouse K, Kobayashi Y, Yamamoto K, et al. Evolution of prognostic factors in hepatocellular carcinoma in Japan. *Aliment Pharmacol Ther*. 2010;31:407–14.
15. Bosch FX, Ribes J, Diaz M, Cleries R. Primary liver cancer: worldwide incidence and trends. *Gastroenterology*. 2004;127:S5–6.
16. Falletti E, Fabris C, Pirisi M, et al. Interleukin-6 polymorphisms and gender: relationship with the occurrence of hepatocellular carcinoma in patients with end-stage liver disease. *Oncology*. 2009;77:304–13.
17. Yang Y, Luo C, Feng R, Bi S. The TNF-alpha, IL-1B and IL-10 polymorphisms and risk for hepatocellular carcinoma: a meta-analysis. *J Cancer Res Clin Oncol*. 2010. doi:10.1007/s00432-010-0959-8.
18. Wilson PA, Gardner SD, Crowther DJ, et al. Characterization of the human patatin-like phospholipase family. *J Lipid Res*. 2006;47:1940–9.
19. Huang Y, He S, Hobbs HH, et al. A feed-forward loop amplifies nutritional regulation of PNPLA3. *Proc Natl Acad Sci USA*. 2010;107:7892–7.

20. Kershaw EE, Hamm JK, Flier JS, et al. Adipose triglyceride lipase: function, regulation by insulin, and comparison with adiponutrin. *Diabetes*. 2006;55:148–57.
21. Romeo S, Kozlitina J, Hobbs HH, et al. Genetic variation in PNPLA3 confers susceptibility to nonalcoholic fatty liver disease. *Nat Genet*. 2008;40:1461–5.
22. Rotman Y, Koh C, Liang TJ, et al. The association of genetic variability in patatin-like phospholipase domain-containing protein 3 (PNPLA3) with histological severity of nonalcoholic fatty liver disease. *Hepatology*. 2010;52:894–903.
23. Valenti L, Al-Serri A, Day CP, et al. Homozygosity for the patatin-like phospholipase-3/adiponutrin I148M polymorphism influences liver fibrosis in patients with nonalcoholic fatty liver disease. *Hepatology*. 2010;51:1209–17.
24. Tian C, Stokowski RP, Hinds DA, et al. Variant in PNPLA3 is associated with alcoholic liver disease. *Nat Genet*. 2010;42:21–3.
25. Kudo M, Okanoue T. Management of hepatocellular carcinoma in Japan: consensus-based clinical practice manual proposed by the Japan Society of Hepatology. *Oncology*. 2007;72(Suppl 1):2–15.
26. Makuuchi M, Kokudo N, Aii S. Clinical practice guidelines for hepatocellular carcinoma—the Japan Society of Hepatology 2009 update. *Hepatol Res*. 2010;40(Suppl 1):2–144.
27. Liver Cancer Study Group of Japan. The general rules for the clinical and pathological study of primary liver cancer. 5th ed. Tokyo: Kanehara; 2009. p. 19 (in Japanese).
28. Valenti L, Al-Serri A, Day CP, et al. Homozygosity for the patatin-like phospholipase-3/adiponutrin I148M polymorphism influences liver fibrosis in patients with nonalcoholic fatty liver disease. *Hepatology*. 2010;51:1209–17.
29. Kollerits B, Coassin S, Kronenberg F, et al. A common variant in the adiponutrin gene influences liver enzyme values. *J Med Genet*. 2010;47:116–9.
30. Kotronen A, Johansson LE, Yki-Järvinen H, et al. A common variant in PNPLA3, which encodes adiponutrin, is associated with liver fat content in humans. *Diabetologia*. 2009;52:1056–60.
31. Kotronen A, Peltonen M, Yki-Järvinen H, et al. Prediction of non-alcoholic fatty liver disease and liver fat using metabolic and genetic factors. *Gastroenterology*. 2009;137:865–72.
32. Kantartzis K, Peter A, Stefan N, et al. Dissociation between fatty liver and insulin resistance in humans carrying a variant of the patatin-like phospholipase 3 gene. *Diabetes*. 2009;58:2616–23.
33. Sookoian S, Castaño GO, Pirola CJ, et al. A nonsynonymous gene variant in the adiponutrin gene is associated with nonalcoholic fatty liver disease severity. *J Lipid Res*. 2009;50:2111–6.
34. Speliotes EK, Butler JL, Hirschhorn JN, et al. PNPLA3 variants specifically confer increased risk for histologic nonalcoholic fatty liver disease but not metabolic disease. *Hepatology*. 2010;52:904–12.
35. Kawaguchi T, Sumida Y, Matsuda F, et al. Genetic polymorphisms of the human PNPLA3 gene are strongly associated with severity of non-alcoholic fatty liver disease in Japanese. *PLoS One*. 2012;7:e38322.
36. Inoue M, Iwasaki M, Tsugane S, et al. Diabetes mellitus and the risk of cancer: results from a large-scale population-based cohort study in Japan. *Arch Intern Med*. 2006;166:1871–7.
37. Hara M, Sasaki S. Japan Public Health Center Study Group. Effect of smoking on the association between alcohol consumption and cancer mortality among middle-aged Japanese men: JPHC Study Cohort I. *IARC Sci Publ*. 2002;156:165–8.
38. Ishiguro S, Inoue M, Tsugane S, JPHC Study Group, et al. Serum aminotransferase level and the risk of hepatocellular carcinoma: a population-based cohort study in Japan. *Eur J Cancer Prev*. 2009;18:26–32.