




Correction to: The characteristics of gastrointestinal symptoms in patients with severe COVID-19: a systematic review and meta-analysis

Yuki Hayashi¹ · Kohei Wagatsuma¹ · Masanori Nojima² · Tsukasa Yamakawa¹ ·
Tadashi Ichimiya¹ · Yoshihiro Yokoyama¹ · Tomoe Kazama¹ · Daisuke Hirayama¹ ·
Hiroshi Nakase¹ 

Published online: 17 April 2021
© Japanese Society of Gastroenterology 2021

Correction to: J Gastroenterol
<https://doi.org/10.1007/s00535-021-01778-z>

In the original publication of the article, the second sentence in second paragraph under the heading “What are the symptoms characteristic of severe COVID-19?” in page 4, should read as “Patients with severe COVID-19 had a significantly higher incidence of abdominal pain than those

non-severe illness (OR 2.70; [95% CI 1.17–6.27; $p = 0.02$; $I^2 = 0\%$]) (Fig. 2).”

In Table 2, the value of “Effect estimate” for “2.4 Vomiting” should read as “0.53 [0.31, 0.92]”.

The original article can be found online at <https://doi.org/10.1007/s00535-021-01778-z>.

✉ Hiroshi Nakase
hiropynakase@gmail.com

- ¹ Department of Gastroenterology and Hepatology, Sapporo Medical University of Medicine, S-1, W16, Chuoku, Sapporo, Hokkaido 060-8543, Japan
- ² Center for Translational Research, The Institute of Medical Science, The University of Tokyo, Tokyo 108-8639, Japan