

One-sided band-limited approximations of some radial functions

Felipe Gonçalves, Michael Kelly* and Jose Madrid

Abstract. We construct majorants and minorants of a Gaussian function in Euclidean space that have Fourier transforms supported in a box. The majorants that we construct are shown to be extremal and our minorants are shown to be *asymptotically extremal* as the sides of the box become uniformly large. We then adapt the Distribution and Gaussian Subordination methods of [12] to the multidimensional setting to obtain majorants and minorants for a class of radial functions. Periodic analogues of the main results are proven and applications to Hilbert-type inequalities are given.

Keywords: Beurling-Selberg problem, band-limited, low-pass filter, Gaussian, Gaussian subordination, distribution method, one-sided approximation, entire function of exponential type.

Mathematical subject classification: 41A30, 46E22, 41A05, 42A05.

1 Introduction

In this paper we address a class of problems that have come to be known as *Beurling-Selberg extremal problems*. The most well-known example of such a problem is due to Selberg himself [39, 45, 51]. Given an interval $I \subset \mathbb{R}$ and $\delta > 0$, Selberg constructed an integrable function $M(x)$ that satisfies

- (i) $\hat{M}(\xi) = 0$ if $|\xi| > \delta$ where $\hat{M}(\xi)$ is the Fourier transform (see §2) of $M(x)$,
- (ii) $M(x) \geq \chi_I(x)$ for each $x \in \mathbb{R}$, and
- (iii) $M(x)$ has the smallest integral¹ among all functions satisfying (i) and (ii).

Received 26 August 2014.

*Corresponding author.

¹This was shown by Selberg in the case when $\delta \text{Length}(I) \in \mathbb{Z}$. If $\delta \text{Length}(I) \notin \mathbb{Z}$, then the minimal integral was found by BF Logan ([37], unpublished) and Littmann [36].

The key constraint (which is common in Beurling-Selberg problems) is condition (ii), that $M(x)$ majorizes the characteristic function of I . Such problems are sometimes called *one-sided* approximation problems.

Most work on these sorts of problems have been focused on solving Selberg's problem but with $\chi_I(x)$ replaced by a different single variable function, such as $e^{-\lambda|x|}$. Such problems are considered in [5, 6, 11, 12, 13, 14, 34, 35, 36, 51]. Some work has also been done in the several variables setting. Shortly after his construction of his function $M(x)$, Selberg was able to construct majorants and minorants of the characteristic function of a box whose Fourier transforms are supported in a (possibly different) box. His majorant can be shown to be extremal for certain configurations of boxes, and it is unknown if the minorant is ever extremal for any given configuration of boxes. Nevertheless, such approximations have proven to be useful in applications [1, 2, 17, 20, 25, 26, 27, 31] because the best known approximations are *asymptotically extremal* as the Fourier support becomes uniformly large. We will use Selberg's method of constructing minorants in what follows (see Proposition 6).

We first study the Beurling-Selberg problem for multivariate Gaussian functions.² We then generalize the *Gaussian subordination* and *distribution method*, originally developed in [12], to higher dimensions and apply the method to study the Beurling-Selberg problem for a class of radial functions. We conclude our investigations with some applications to Hilbert-type inequalities and adapt the construction to periodic functions.

To state the first of our main results we will use the following notation (see §2 for additional information). For $c > 0$, let $g_c(t) = e^{-c\pi t^2}$ and for $a, \lambda \in \Lambda$ (where $\Lambda = (0, \infty)^d$) let

$$x \in \mathbb{R}^d \mapsto G_\lambda(x) = \prod_{j=1}^d g_{\lambda_j}(x_j) = \exp \left\{ - \sum_{j=1}^d \lambda_j \pi x_j^2 \right\}.$$

and let

$$Q(a) = \prod_{j=1}^d [-a_j, a_j].$$

Our first result is a solution to the Beurling-Selberg extremal problem of determining optimal *majorants* of the Gaussian function $G_\lambda(x)$ that have Fourier transform supported in $Q(a)$.

Theorem 1. *Let $a, \lambda \in \Lambda$. If $F : \mathbb{R}^d \rightarrow \mathbb{R}$ is an integrable function that satisfies*

²This problem, for a single variable, was solved in [12].

- (i) $F(x) \geq G_\lambda(x)$ for each $x \in \mathbb{R}^d$, and
- (ii) $\hat{F}(\xi) = 0$ for each $\xi \notin Q(a)$,

then

$$\int_{\mathbb{R}^d} F(x)dx \geq \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} \Theta(0; ia_j^2/\lambda_j). \tag{1.1}$$

where $\Theta(v; \tau)$ is Jacobi's theta function (see §2). Moreover, equality holds if $F(z) = M_{\lambda,a}(z)$ where $M_{\lambda,a}(z)$ is defined by (3.3).

This theorem is essentially a corollary of Theorem 3 of [12]. The proof simply uses the product structure and positivity of $G_\lambda(x)$ in conjunction with Theorem 3 of [12]. It would be interesting to determine the analogue of the above theorem for *minorants* of $G_\lambda(x)$ (i.e. the high dimensional analogue of Theorem 2 of [12]) where the extremal functions cannot be obtained by a tensor product of lower dimensional extremal functions.

In our second result we address this problem by constructing minorants of the Gaussian function $G_\lambda(x)$ that have Fourier transform supported in $Q(a)$ and that are asymptotically extremal as a becomes uniformly large in each coordinate.

Theorem 2. Let $a, \lambda \in \Lambda$. If $F : \mathbb{R}^d \rightarrow \mathbb{R}$ is an integrable function that satisfies

- (i) $F(x) \leq G_\lambda(x)$ for each $x \in \mathbb{R}^d$, and
- (ii) $\hat{F}(\xi) = 0$ for each $\xi \notin Q(a)$,

then

$$\int_{\mathbb{R}^d} F(x)dx \leq \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} \Theta(\frac{1}{2}; ia_j^2/\lambda_j). \tag{1.2}$$

Furthermore, there exists a positive constant $\gamma_0 = \gamma_0(d)$ such that if $\gamma := \min\{a_j^2/\lambda_j : 1 \leq j \leq d\} \geq \gamma_0$, then

$$\prod_{j=1}^d \lambda_j^{-\frac{1}{2}} \Theta(\frac{1}{2}; ia_j^2/\lambda_j) \leq (1 + 5de^{-\pi\gamma}) \int_{\mathbb{R}^d} L_{\lambda,a}(x)dx \tag{1.3}$$

where $L_{\lambda,a}(x)$ is defined by (3.2).

Remark 3. For a fixed a , if λ is large enough then the right hand side (RHS) of (1.3) would be negative and the inequality would not hold, but this is not true for large values of γ . In fact, this happens because the zero function would be a better minorant.

Remark 4. Inequality (1.3) implies that if λ is fixed and a is large, then $\int_{\mathbb{R}^d} L_{\lambda,a}(x)dx$ approximates exponentially fast the optimal answer. In this sense, we say that $L_{\lambda,a}(x)$ is asymptotically optimal with respect to the type.

Our next set of results (which are too lengthy to state here) are Theorems 9, 10, and 11. See §4 for the statements of the theorems. These theorems generalize the so-called *distribution* and *Gaussian subordination* methods of [12]. The main idea behind these methods goes back to the paper of Graham and Vaaler [24]. We will describe a “watered down” version of the approach here. Let us begin with the inequality

$$G_\lambda(x) \leq F_\lambda(x)$$

where $F_\lambda(x)$ is defined by (3.3). The idea is to *integrate the free parameter* λ in the function $G_\lambda(x)$ with respect to a (positive) measure ν on $\Lambda = (0, \infty)^d$ to obtain a pair of new functions of x :

$$g(x) = \int_\Lambda G_\lambda(x) d\nu(\lambda) \leq \int_\Lambda F_\lambda(x) d\nu(\lambda) = f(x).$$

The process simultaneously produces a function $g(x)$ and a majorant $f(x)$ having $\hat{f}(\xi)$ supported in $Q(a)$. The difference of the functions in L^1 -norm is similarly obtained by integrating against ν . The method that we present allows us to produce majorants and minorants for $g(x)$ equal to the one of the following functions (among others):

$$\begin{aligned} g(x) &= e^{-\alpha|x|^r}, \text{ for } \alpha > 0 \text{ and } 0 < r < 2, \\ g(x) &= (|x|^2 + \alpha^2)^{-\beta}, \text{ for } \alpha > 0 \text{ and } \beta > 0, \\ g(x) &= -\log\left(\frac{|x|^2 + \alpha^2}{|x|^2 + \beta^2}\right), \text{ for } 0 < \alpha < \beta, \text{ and} \\ g(x) &= |x|^\sigma, \text{ for } \sigma \in (0, \infty) \setminus 2\mathbb{Z}_+ \end{aligned}$$

where $|x|$ is the Euclidean norm of x (see §2). In §5.2, we will discuss the full class of functions for which our method produces majorants and minorants.

One dimensional extremal functions have proven to be useful in several problems in analytic number theory [4, 7, 9, 10, 12, 15, 21, 22, 24, 26, 39, 40, 44, 45, 51]. The most well known examples make use of Selberg’s extremal functions for the characteristic function of the interval. These include sharp forms of the large sieve and the Erdős-Turán inequality. Recent activity include estimates of quantities related to the Riemann zeta function. These include estimates of the zeta function on vertical lines in the critical strip and estimates of the pair correlation of the zeros of the zeta function. It would be interesting to see similar applications in the multidimensional setting.

Organization of the paper

In §2 we introduce notation and gather the necessary background material for the remainder of the paper. Then in §3 we will discuss and prove Theorems 1 and 2. Next, in §4 we present and prove a generalization of the *distribution method* introduced in [12] as well as a generalization of their Gaussian subordination result. In §5.1 we study periodic analogues of Theorems 1, 2, 9, 10, and 11. We conclude with some applications to Hilbert-type inequalities §5.3 and some final remarks in §6

2 Preliminaries

Let us first have a word about our notation. Throughout this manuscript vectors in \mathbb{R}^d will be denoted by lowercase letters such as $x = (x_1, \dots, x_d)$, and the Euclidean norm of x is given by

$$|x| = \{x_1^2 + \dots + x_d^2\}^{1/2}.$$

We will write $\mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$, $\Lambda = (0, \infty)^d$ and for each $a \in \Lambda$ we let

$$Q(a) = \{x \in \mathbb{R}^d : |x_j| \leq a_j\}. \quad (2.1)$$

The vector $u = (1, \dots, 1) \in \Lambda$ will be called the *unitary* vector and we will use the notation $Q(R) = Q(Ru)$ whenever $R > 0$ is a positive real number.

Now we will introduce some notation that is not standard, but convenient for our purposes. Given vectors $x, y \in \mathbb{R}^d$ we will write $xy = (x_1y_1, \dots, x_dy_d)$ and if $y_j \neq 0$ for each j , then we will write $x/y = (x_1/y_1, \dots, x_d/y_d)$. We say that $x < y$ ($x \leq y$) if $x_j < y_j \forall j$ ($x_j \leq y_j \forall j$). We will always denote the inner product of $x, y \in \mathbb{R}^d$ by a central dot, that is, $x \cdot y$.

One of the main objects of study in this paper is the *Fourier transform*. Given an integrable function $F(x)$ on \mathbb{R}^d , we define the Fourier transform of $F(x)$ by

$$\hat{F}(\xi) = \int_{\mathbb{R}^d} e(x \cdot \xi) F(x) dx$$

where $\xi \in \mathbb{R}^d$ and $e(\theta) = e^{-2\pi i\theta}$. We extend the definition in the usual way to tempered distributions (see for instance [49]). We will mainly be considering functions whose Fourier transforms are supported in a bounded subset of \mathbb{R}^d . Such functions are called *band-limited*. It is well-known that band-limited functions can be extended to an entire functions on \mathbb{C}^d satisfying an exponential growth condition. An entire function $F : \mathbb{C}^d \rightarrow \mathbb{C}$ is said to be of *exponential*

type if there exists a number $\tau \geq 0$ such that for every $\epsilon > 0$ there exists a constant $C_\epsilon > 0$ such that

$$|F(z)| \leq C_\epsilon e^{(\tau+\epsilon)|z|}.$$

If $F(z)$ satisfies such a growth estimate, then $F(z)$ is said to be of exponential type at most τ .

There is a refinement of the definition of exponential type due to Stein [46, 49]. Given an origin symmetric convex body K with supporting function

$$H(z) = H_K(z) = \sup_{\xi \in K} z \cdot \xi, \quad (2.2)$$

an entire function $F : \mathbb{C}^d \rightarrow \mathbb{C}$ is said to be of exponential type with respect to K if

$$|F(z)| \leq C_\epsilon e^{(1+\epsilon)H(z)}.$$

This is the natural generalization of exponential type used by Stein in his generalization of the Paley-Wiener theorem [46, 49]. The Paley-Wiener theorem has another generalization to higher dimensions that is formulated for certain tempered distributions. In this formulation, it is not necessary that the body K be symmetric. We will now state this generalization of the Paley-Wiener theorem which can be found in [29], Theorem 7.3.1.

Theorem 5 (Paley-Wiener-Schwartz). *Let K be a convex compact subset of \mathbb{R}^d with supporting function $H(x)$ given by (2.2). If F is a tempered distribution such that the support of \widehat{F} is contained in K , then $F : \mathbb{C}^d \rightarrow \mathbb{C}$ is an entire function and exist $N, C > 0$ such that*

$$|F(x + iy)| \leq C(1 + |x + iy|)^N e^{2\pi H(y)}.$$

for every $x + iy \in \mathbb{C}^d$.

Conversely, every entire function $F : \mathbb{C}^d \rightarrow \mathbb{C}$ satisfying an estimate of this form defines a tempered distribution with Fourier transform supported on K .

We will now define and compile some results about Gaussians and theta functions that we will need in the sequel. Given a positive real number $\delta > 0$ the Gaussian $g_\delta : \mathbb{R} \rightarrow \mathbb{R}$ is defined by

$$g_\delta(t) = e^{-\delta\pi t^2},$$

and its Fourier transform is given by $\widehat{g}_\delta(\xi) = \delta^{-1/2} g_{1/\delta}(\xi)$.

For a $\tau = \sigma + it$ with $t > 0$, if $q = e^{\pi i \tau}$, then Jacobi's theta function (see [16]) is defined by

$$\Theta(v; \tau) = \sum_{n \in \mathbb{Z}} e(nv)q^{n^2}. \tag{2.3}$$

These functions are related through the Poisson summation formula by

$$\sum_{m \in \mathbb{Z}} g_\delta(v + m) = \sum_{n \in \mathbb{Z}} e(nv)\hat{g}_\delta(n) = \delta^{-1/2}\Theta(v; i\delta^{-1}). \tag{2.4}$$

The one dimensional case of Theorems 1 and 2 are proven in [12]. There it is proved that the functions

$$l_\delta(z) = \left(\frac{\cos \pi z}{\pi}\right)^2 \left\{ \sum_{k \in \mathbb{Z}} \frac{g_\delta(k + \frac{1}{2})}{(z - k - \frac{1}{2})^2} + \sum_{k \in \mathbb{Z}} \frac{g'_\delta(k + \frac{1}{2})}{(z - k - \frac{1}{2})} \right\} \tag{2.5}$$

and

$$m_\delta(z) = \left(\frac{\sin \pi z}{\pi}\right)^2 \left\{ \sum_{k \in \mathbb{Z}} \frac{g_\delta(k)}{(z - k)^2} + \sum_{k \in \mathbb{Z}} \frac{g'_\delta(k)}{(z - k)} \right\} \tag{2.6}$$

are entire functions of exponential type at most 2π and they satisfy

$$l_\delta(x) \leq g_\delta(x) \leq m_\delta(x) \tag{2.7}$$

for all real x . Moreover,

$$\int_{-\infty}^{\infty} m_\delta(x)dx = \delta^{-\frac{1}{2}}\Theta(0; i/\delta) \tag{2.8}$$

and

$$\int_{-\infty}^{\infty} l_\delta(x)dx = \delta^{-\frac{1}{2}}\Theta(\frac{1}{2}; i/\delta). \tag{2.9}$$

In view of (2.4), (2.8), and (2.9), the functions $l_\delta(z)$ and $m_\delta(z)$ are the best one-sided L^1 -approximations of g_δ having exponential type at most 2π .

3 The Multidimensional Gaussian Function

In this section we will prove Theorem 1 and 2. To construct a minorant of the Gaussian, we begin with the functions $m_\delta(z)$ and $l_\delta(z)$ defined by (2.5) and (2.6) and use Selberg's bootstrapping technique to obtain multidimensional minorants. The majorant is constructed by taking $m_\delta(z)$ tensored with itself d -times.

For every $\lambda = (\lambda_1, \dots, \lambda_n) \in \Lambda$ we define the function $G_\lambda : \mathbb{R}^d \rightarrow \mathbb{R}$ by

$$G_\lambda(x) = \prod_{j=1}^d g_{\lambda_j}(x_j) = \prod_{j=1}^d e^{-\lambda_j \pi x_j^2}. \quad (3.1)$$

The following proposition is due to Selberg, but it was never published [50]. We called it *Selberg's bootstrapping method* because it enables us to construct a minorant for a tensor product of functions provided that we have majorants and minorants of each component at our disposal. The method has been used in one form or the other in [2, 20, 25, 26, 27].

Proposition 6. *Let $d > 0$ be natural number and $f_j : \mathbb{R} \rightarrow (0, \infty)$ be functions for every $j = 1, \dots, d$. Let $l_j, m_j : \mathbb{R} \rightarrow \mathbb{R}$ be real-valued functions such that*

$$l_j(x) \leq f_j(x) \leq m_j(x)$$

for every x and j . Then

$$-(d-1) \prod_{k=1}^d m_k(x_k) + \sum_{k=1}^d l_k(x_k) \prod_{\substack{j=1 \\ j \neq k}}^d m_j(x_j) \leq \prod_{k=1}^d f_k(x_k).$$

This proposition is easily deduced from the following inequality.

Lemma 7. *If $\beta_1, \dots, \beta_d \geq 1$, then*

$$\sum_{k=1}^d \prod_{\substack{j=1 \\ j \neq k}}^d \beta_j \leq 1 + (d-1) \prod_{k=1}^d \beta_k.$$

Proof. We give a proof by induction, starting with the inductive step since the base case is simple.

Suppose that the claim is true for $d = 1, \dots, L$. Let $\beta_1, \dots, \beta_L, \beta_{L+1}$ be a sequence of real numbers not less than one and write $\beta_j = 1 + \epsilon_j$. We obtain

$$\begin{aligned} \sum_{k=1}^{L+1} \prod_{\substack{j=1 \\ j \neq k}}^{L+1} \beta_j &= \prod_{j=1}^L \beta_j + (1 + \epsilon_{L+1}) \sum_{k=1}^L \prod_{\substack{j=1 \\ j \neq k}}^L \beta_j \\ &\leq \prod_{j=1}^L \beta_j + (1 + \epsilon_{L+1}) \left\{ 1 + (L-1) \prod_{j=1}^L \beta_j \right\} \end{aligned}$$

$$\begin{aligned}
 &= 1 + \epsilon_{L+1} + \prod_{j=1}^L \beta_j + (L - 1) \prod_{j=1}^{L+1} \beta_j \\
 &\leq 1 + \epsilon_{L+1} \prod_{j=1}^L \beta_j + \prod_{j=1}^L \beta_j + (L - 1) \prod_{j=1}^{L+1} \beta_j \\
 &= 1 + L \prod_{j=1}^{L+1} \beta_j \quad \square
 \end{aligned}$$

Now we can define our candidates for majorant and minorant of $G_\lambda(x)$. For a given $\lambda \in \Lambda$ define the functions

$$z \in \mathbb{C}^d \mapsto L_\lambda(z) = -(d - 1) \prod_{j=1}^d m_{\lambda_j}(z_j) + \sum_{k=1}^d l_{\lambda_k}(z_k) \prod_{\substack{j=1 \\ j \neq k}}^d m_{\lambda_j}(z_j) \quad (3.2)$$

and

$$z \in \mathbb{C}^d \mapsto M_\lambda(z) = \prod_{j=1}^d m_{\lambda_j}(z_j). \quad (3.3)$$

It follows from Proposition 6 and (2.7) that

$$L_\lambda(x) \leq G_\lambda(x) \leq M_\lambda(x) \text{ for all } x \in \mathbb{R}^d. \quad (3.4)$$

Moreover, since $l_\delta(x)$ and $m_\delta(x)$ have exponential type at most 2π , we conclude that the Fourier transforms of $L_\lambda(x)$ and $M_\lambda(x)$ are supported on Q . We modify $L_\lambda(z)$ and $M_\lambda(z)$ to have exponential type with respect to $Q(a)$ in the following way. Given $a, \lambda \in \Lambda$ we define the functions

$$L_{\lambda,a}(z) = L_{\lambda/a^2}(az) \quad (3.5)$$

and

$$M_{\lambda,a}(z) = M_{\lambda/a^2}(az). \quad (3.6)$$

By (3.4) we obtain

$$L_{\lambda,a}(x) \leq G_\lambda(x) \leq M_{\lambda,a}(x) \text{ for all } x \in \mathbb{R}^d \quad (3.7)$$

and using the scaling properties of the Fourier transform, we conclude that $L_{\lambda,a}(x)$ and $M_{\lambda,a}(x)$ have exponential type with respect to $Q(a)$. By formula (2.6), we have $m_\delta(k) = g_\delta(k)$ for all integers k , hence we obtain

$$M_{\lambda,a}(k/a) = G_\lambda(k/a) \quad (3.8)$$

for all $k \in \mathbb{Z}^d$ (recall that $k/a = (k_1/a_1, \dots, k_d/a_d)$).

We are now in a position to prove Theorems 1 and 2.

Proof of Theorem 1. It follows from (3.4) and (3.6) that the function $M_{\lambda,a}(x)$ is majorant of $G_\lambda(x)$ of exponential type with respect to $Q(a)$. Define $\alpha = a_1 \cdots a_d$. Using definition (3.3) and (3.6) we obtain

$$\begin{aligned} \int_{\mathbb{R}^d} M_{\lambda,a}(x) dx &= \alpha^{-1} \int_{\mathbb{R}^d} M_{\lambda/a^2}(x) dx \\ &= \alpha^{-1} \prod_{j=1}^d \int_{\mathbb{R}^d} m_{\lambda_j/a_j^2}(x_j) dx_j \\ &= \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} \Theta(0; ia_j^2/\lambda_j), \end{aligned}$$

where the first equality is due to a change of variables, the second one due to the product structure and the third one due to (2.8).

Now we will prove that (3.6) is extremal. Suppose that $F(z)$ is an entire majorant of $G_\lambda(x)$ of exponential type with respect to $Q(a)$ and integrable on \mathbb{R}^d (and therefore absolutely integrable on \mathbb{R}^d). By the Poisson summation formula we have

$$\begin{aligned} \int_{\mathbb{R}^d} F(x) dx &= \alpha^{-1} \sum_{k \in \mathbb{Z}^d} F(k/a) \\ &\geq \alpha^{-1} \sum_{k \in \mathbb{Z}^d} G_\lambda(k/a) \\ &= \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} \Theta(0; ia_j^2/\lambda_j) \end{aligned} \quad (3.9)$$

because $M_{\lambda,a}(x)$ majorizes G_λ , and the rightmost equality is given by (2.4). \square

To prove Theorem 2 we will need the following lemma.

Lemma 8. *For all $t > 0$ we have*

$$1 - 4q/(1 - q)^2 < \frac{\Theta(\frac{1}{2}; it)}{\Theta(0; it)} < e^{-2q}, \quad (3.10)$$

where $q = e^{-\pi t}$.

Proof of Theorem 2. Suppose that $F(z)$ is an entire minorant of exponential type with respect to $Q(a)$ and absolutely integrable on \mathbb{R}^d . Recalling that $u = (1, \dots, 1)$ and applying the Poisson summation formula, we obtain

$$\begin{aligned} \int_{\mathbb{R}^d} F(x)dx &= \alpha^{-1} \sum_{k \in \mathbb{Z}^d} F(k/a + u/2a) \\ &\leq \alpha^{-1} \sum_{k \in \mathbb{Z}^d} G_\lambda(k/a + u/2a) \\ &= \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} \Theta\left(\frac{1}{2}; ia_j^2/\lambda_j\right). \end{aligned}$$

where the last equality is given by (2.4). This proves (1.2). By construction, $L_{\lambda,a}(z)$ is an entire minorant of exponential type with respect to $Q(a)$. Using definitions (3.2) and (3.5) we conclude that

$$\begin{aligned} \int_{\mathbb{R}^d} L_{\lambda,a}(x)dx &= \left\{ \sum_{j=1}^d \frac{\Theta\left(\frac{1}{2}; ia_j^2/\lambda_j\right)}{\Theta(0; ia_j^2/\lambda_j)} - (d-1) \right\} \\ &\quad \times \prod_{j=1}^d \Theta(0; ia_j^2/\lambda_j) \lambda_j^{-\frac{1}{2}}. \end{aligned} \tag{3.11}$$

Thus, to deduce (1.3), we only need to prove that

$$(1 + 5de^{-\pi\gamma}) \left\{ \sum_{j=1}^d \frac{\Theta\left(\frac{1}{2}; ia_j^2/\lambda_j\right)}{\Theta(0; ia_j^2/\lambda_j)} - (d-1) \right\} \geq \prod_{j=1}^d \frac{\Theta\left(\frac{1}{2}; ia_j^2/\lambda_j\right)}{\Theta(0; ia_j^2/\lambda_j)}. \tag{3.12}$$

for large γ (recall that $\gamma = \min\{\lambda_j/a_j^2\}$). If we let $q_j = e^{-\pi a_j^2/\lambda_j}$ and γ sufficient large such that $(1 - e^{-\pi\gamma})^2 > 4/5$, we can use Lemma 8 to obtain

$$\begin{aligned} \sum_{j=1}^d \frac{\Theta\left(\frac{1}{2}; ia_j^2/\lambda_j\right)}{\Theta(0; ia_j^2/\lambda_j)} - (d-1) &\geq 1 - 4 \sum_{j=1}^d q_j / (1 - q_j)^2 \\ &\geq 1 - 5 \sum_{j=1}^d q_j. \end{aligned} \tag{3.13}$$

Applying Lemma 8 for a sufficient large γ we obtain

$$\prod_{j=1}^d \frac{\Theta\left(\frac{1}{2}; ia_j^2/\lambda_j\right)}{\Theta(0; ia_j^2/\lambda_j)} \leq \exp \left\{ -2 \sum_{j=1}^d q_j \right\} \leq 1 - \sum_{j=1}^d q_j. \tag{3.14}$$

where the last inequality holds, say, if $\sum_{j=1}^d q_j < \log 2$. Write $\beta = \sum_{j=1}^d q_j$ and note that

$$1 - \beta \leq (1 - 5\beta)(1 + 5\beta)$$

if β is sufficiently small, for instance if $\beta \in [0, 1/25]$. If γ is sufficiently large such that $\sum_{j=1}^d q_j < 1/25$ we obtain

$$\begin{aligned} 1 - \sum_{j=1}^d q_j &\leq (1 - 5 \sum_{j=1}^d q_j)(1 + 5 \sum_{j=1}^d q_j) \\ &< (1 - 5 \sum_{j=1}^d q_j)(1 + 5de^{-\pi\gamma}). \end{aligned} \tag{3.15}$$

By (3.13), (3.14) and (3.15) we conclude that there exists an $\gamma_0 = \gamma_0(d) > 0$ such that if $\gamma \geq \gamma_0$, then (3.12) holds. □

Proof of Lemma 8. Recall that $e(v) = e^{2\pi i v}$ and $q = e^{\pi i \tau}$. By Theorem 1.3 of chapter 10 of [48] the Theta function has the following product representation

$$\Theta(v; \tau) = \prod_{n=1}^{\infty} (1 - q^{2n})(1 + q^{2n-1}e(v))(1 + q^{2n-1}e(-v)). \tag{3.16}$$

It follows from (3.16) that

$$\frac{\Theta(\frac{1}{2}; it)}{\Theta(0; it)} = \prod_{n=1}^{\infty} \left(\frac{1 - q^{2n-1}}{1 + q^{2n-1}} \right)^2 = \exp \left\{ 2 \sum_{n=0}^{\infty} \log \left(1 - \frac{2q^{2n+1}}{1 + q^{2n+1}} \right) \right\}.$$

Using the inequality $\log(1 - x) \geq -x/(1 - x)$ for all $x \in [0, 1)$ we obtain

$$\begin{aligned} \frac{\Theta(\frac{1}{2}; it)}{\Theta(0; it)} &\geq \exp \left\{ -4 \sum_{n=0}^{\infty} \frac{q^{2n+1}}{1 - q^{2n+1}} \right\} \\ &\geq \exp \left\{ -\frac{4}{1 - q} \sum_{n=0}^{\infty} q^{2n+1} \right\} \\ &= \exp \left\{ -\frac{4q}{(1 - q)(1 - q^2)} \right\} \\ &> e^{-4q/(1-q)^2} \\ &> 1 - 4q/(1 - q)^2 \end{aligned}$$

and this proves the left hand side (LHS) inequality in (3.10). The (RHS) inequality in (3.10) is deduced by a similar argument using the inequality $\log(1 - x) \leq -x$ for all $x \in [0, 1)$. □

4 Gaussian Subordination Method

In this section we adapt the distribution method developed in [12] to the several variables setting, and we apply the Gaussian subordination method to the majorant and minorant for the multidimensional Gaussian defined in the previous section. This method allow us to extend the class of functions for which we can solve the corresponding Beurling-Selberg extremal problems. The central idea is to integrate the functions $G_\lambda(x)$, $L_{\lambda,a}(z)$ and $M_{\lambda,a}(z)$, in a distributional sense, with respect to a non-negative measure ν defined on $\Lambda = (0, \infty)^d$.

We begin with a generalization of the distribution method developed in [12] for existence of majorants and minorants, we will deal with extremality later.

Theorem 9 (Distribution Method – (Existence)). *Let $K \subset \mathbb{R}^d$ be a compact convex set, Λ be a measurable space of parameters, and for each $\lambda \in \Lambda$ let $G(x; \lambda) \in L^1(\mathbb{R}^d)$ be a real-valued function. For each λ let $F(z; \lambda)$ be a entire function defined for $z \in \mathbb{C}^d$ of exponential type with respect to K . Let ν be a non-negative measure on Λ that satisfies*

$$\int_\Lambda \int_{\mathbb{R}^d} |F(x; \lambda) - G(x; \lambda)| dx d\nu(\lambda) < \infty. \tag{4.1}$$

and

$$\int_\Lambda \int_{\mathbb{R}^d} |\widehat{G}(x; \lambda)\varphi(x)| dx d\nu(\lambda) < \infty \tag{4.2}$$

for all $\varphi \in S(\mathbb{R}^d)$ supported in K^c .

Let $\mathcal{G} \in S'(\mathbb{R}^d)$ be a real-valued continuous function and

$$\widehat{\mathcal{G}}(\varphi) = \int_{\mathbb{R}^d} \int_\Lambda \widehat{G}(x; \lambda) d\nu(\lambda) \varphi(x) dx \tag{4.3}$$

for all $\varphi \in S(\mathbb{R}^d)$ supported in K^c .

- (i) *If $G(x; \lambda) \leq F(x; \lambda)$ for each $x \in \mathbb{R}^d$ and $\lambda \in \Lambda$, then there exists a real entire majorant $\mathcal{M}(x)$ for $\mathcal{G}(x)$ of exponential type with respect to K and*

$$\int_{\mathbb{R}^d} \{\mathcal{M}(x) - \mathcal{G}(x)\} dx$$

is equal to the quantity at (4.1).

- (ii) *If $F(x; \lambda) \leq G(x; \lambda)$ for each $x \in \mathbb{R}^d$ and $\lambda \in \Lambda$, then there exists a real entire minorant $\mathcal{L}(x)$ for $\mathcal{G}(x)$ of exponential type with respect to K and*

$$\int_{\mathbb{R}^d} \{\mathcal{G}(x) - \mathcal{L}(x)\} dx$$

is equal to the quantity at (4.1).

With the exception of Theorem 9, Λ will always stand for $(0, \infty)^d$. For a given $a \in \Lambda$ define $\mathfrak{G}_+^d(a)$ as the set of ordered pairs (\mathcal{G}, ν) where $\mathcal{G} : \mathbb{R}^d \rightarrow \mathbb{R}$ is a continuous function and ν is a non-negative Borel measure in Λ such that

(C1) $\mathcal{G} \in S'(\mathbb{R}^d)$ is a tempered distribution.

(C2) For all $\varphi \in S(\mathbb{R}^d)$ supported in $Q(a)^c$ we have

$$\int_{\mathbb{R}^d} \int_{\Lambda} |\widehat{\mathcal{G}}_{\lambda}(x)\varphi(x)|d\nu(\lambda)dx < \infty.$$

(C3) For all $\varphi \in S(\mathbb{R}^d)$ supported in $Q(a)^c$ we have

$$\int_{\mathbb{R}^d} \mathcal{G}(x)\widehat{\varphi}(x) = \int_{\mathbb{R}^d} \int_{\Lambda} \widehat{\mathcal{G}}_{\lambda}(x)d\nu(\lambda)\varphi(x)dx. \tag{4.4}$$

(C4+) The following integrability condition holds

$$\int_{\Lambda} \prod_{k=1}^d \lambda_k^{-\frac{1}{2}} \left\{ \prod_{j=1}^d \Theta(0; ia_j^2/\lambda_j) - 1 \right\} d\nu(\lambda) < \infty. \tag{4.5}$$

In a analogous way, we define the class $\mathfrak{G}_-^d(a)$ replacing condition (C4+) by

(C4-) The following integrability condition holds

$$\int_{\Lambda} \left\{ 1 - \left(\sum_{j=1}^d \frac{\Theta(\frac{1}{2}; ia_j^2/\lambda_j)}{\Theta(0; ia_j^2/\lambda_j)} - (d-1) \right) \prod_{j=1}^d \Theta(0; ia_j^2/\lambda_j) \right\} \times \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} d\nu(\lambda) < \infty. \tag{4.6}$$

Theorem 10 offers a optimal resolution of the Majorization Problem for the class functions $\mathfrak{G}_+^d(a)$ and the Theorem 11 offers asymptotically optimal resolution of the Minorization Problem for the class of functions $\mathfrak{G}_-^d(a)$.

Theorem 10 (Gaussian Subordination – Majorant). *For a given $a \in \Lambda$, let $(\mathcal{G}, \nu) \in \mathfrak{G}_+^d(a)$. Then there exists an extremal majorant $\mathcal{M}_a(z)$ of exponential type with respect to $Q(a)$ for $\mathcal{G}(x)$. Furthermore, $\mathcal{M}_a(x)$ interpolates $\mathcal{G}(x)$ on \mathbb{Z}^d/a and satisfies*

$$\int_{\mathbb{R}^d} \mathcal{M}(x) - \mathcal{G}(x)dx = \int_{\Lambda} \prod_{k=1}^d \lambda_k^{-\frac{1}{2}} \left\{ \prod_{j=1}^d \Theta(0; ia_j^2/\lambda_j) - 1 \right\} d\nu(\lambda). \tag{4.7}$$

Theorem 11 (Gaussian Subordination – Minorant). *For a given $a \in \Lambda$, let $(\mathcal{G}, \nu) \in \mathfrak{G}^d(a)$. Then, if $\mathcal{F}(z)$ is a real entire minorant of $\mathcal{G}(x)$ of exponential type with respect to $Q(a)$, we have*

$$\int_{\mathbb{R}^d} \mathcal{G}(x) - \mathcal{F}(x) dx \geq \int_{\Lambda} \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} \left\{ 1 - \prod_{j=1}^d \Theta\left(\frac{1}{2}; ia_j^2/\lambda_j\right) \right\} d\nu(\lambda). \tag{4.8}$$

Furthermore, there exists a family of minorants $\{\mathcal{L}_a(z) : a \in \Lambda\}$ where $\mathcal{L}_a(z)$ is of exponential type with respect to $Q(a)$ such that

$$\int_{\mathbb{R}^d} \mathcal{G}(x) - \mathcal{L}_a(x) dx$$

is equal to the (LHS) of (4.6). Also

$$\lim_{a \uparrow \infty} \int_{\mathbb{R}^d} \mathcal{G}(x) - \mathcal{L}_a(x) dx = 0, \tag{4.9}$$

where $a \uparrow \infty$ means $a_j \uparrow \infty$ for each j .

Corollary 12. *Under the hypothesis of Theorem 11, suppose also that exists an $R > 0$ such that $\text{supp}(\nu) \subset \Lambda \cap Q(R)$, $\mathcal{G} \in L^1(\mathbb{R}^d)$ and*

$$\int_{\mathbb{R}^d} \mathcal{G}(x) dx = \int_{\Lambda} \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} d\nu(\lambda) < \infty.$$

Then, there exists a constant $\alpha_0 > 0$ such that, if $\alpha := \min\{a_j\} \geq \alpha_0$ and if $\mathcal{F}(x)$ is a real entire minorant of $\mathcal{G}(x)$ of exponential type with respect to $Q(a)$, then

$$\int_{\mathbb{R}^d} \mathcal{F}(x) dx \leq (1 + 5de^{-\pi\alpha^2/R}) \int_{\mathbb{R}^d} \mathcal{L}_a(x) dx.$$

Proof of Corollary 12. By a direct application of Theorem 2 we obtain

$$\int_{\mathbb{R}^d} \mathcal{F}(x) dx \leq \int_{\Lambda_R} \prod_{j=1}^d \Theta\left(\frac{1}{2}; ia_j^2/\lambda_j\right) \lambda_j^{-\frac{1}{2}} d\nu(\lambda).$$

If we choose $\gamma_0 > 0$, as in Theorem 2 and define $\alpha_0 = R\gamma_0$, we can use inequality (1.3) to conclude that

$$\prod_{j=1}^d \Theta\left(\frac{1}{2}; ia_j^2/\lambda_j\right) \lambda_j^{-\frac{1}{2}} \leq (1 + 5de^{-\pi\alpha^2/R}) \int_{\mathbb{R}^d} L_{\lambda,a}(x) dx.$$

for all $\lambda \in \Lambda \cap Q(R)$, if each $a_j \geq \alpha_0$. If we integrate this last inequality with respect to $d\nu(\lambda)$ we obtain the desired result. □

4.1 Proofs of Theorems 9, 10 and 11

Proof of Theorem 9. We follow the proof of Theorem 14 from [12] proving only the majorant case, since the minorant case is nearly identical. Let

$$D(x; \lambda) = F(x; \lambda) - G(x; \lambda) \geq 0.$$

By condition (4.1) and Fubini's theorem the function

$$\mathcal{D}(x) = \int_{\Lambda} D(x; \lambda) d\nu(\lambda) \geq 0,$$

is defined for almost all $x \in \mathbb{R}^d$ and $\mathcal{D} \in L^1(\mathbb{R}^d)$. The Fourier transform of $\mathcal{D}(x)$ is a continuous function given by

$$\widehat{\mathcal{D}}(\xi) = \int_{\Lambda} \widehat{D}(\xi; \lambda) d\nu(\lambda), \quad (4.10)$$

and, due to (4.2), for almost every $\xi \notin K$ we have the alternative representation

$$\widehat{\mathcal{D}}(\xi) = - \int_{\Lambda} \widehat{G}(\xi; \lambda) d\nu(\lambda). \quad (4.11)$$

Let \mathcal{M} be the tempered distribution given by

$$\mathcal{M}(\varphi) = \int_{\mathbb{R}^d} \{\mathcal{D}(x) + \mathcal{G}(x)\} \varphi(x) dx.$$

Now for any $\varphi \in S(\mathbb{R}^d)$ supported in K^c , we have by combining (4.3) and (4.11)

$$\widehat{\mathcal{M}}(\varphi) = \widehat{\mathcal{D}}(\varphi) + \widehat{\mathcal{G}}(\varphi) = 0. \quad (4.12)$$

Hence $\widehat{\mathcal{M}}$ is supported on K , in the distributional sense. By the Theorem 5, it follows that the distribution \mathcal{M} is identified with a analytic function $\mathcal{M} : \mathbb{C}^d \rightarrow \mathbb{C}$ of exponential type with respect to K and that

$$\mathcal{M}(\varphi) = \int_{\mathbb{R}^d} \mathcal{M}(x) \varphi(x) dx \quad (4.13)$$

for every $\varphi \in S(\mathbb{R}^d)$. It then follows from the definition of \mathcal{M} and (4.13) that for almost every $x \in \mathbb{R}^d$

$$\mathcal{M}(x) = \mathcal{D}(x) + \mathcal{G}(x),$$

which implies $\mathcal{M}(x) \geq \mathcal{G}(x)$ for all $x \in \mathbb{R}^d$ since $\mathcal{G}(x)$ is continuous, and

$$\int_{\mathbb{R}^d} \{\mathcal{M}(x) - \mathcal{G}(x)\} dx = \int_{\Lambda} \int_{\mathbb{R}^d} \{F(x; \lambda) - G(x; \lambda)\} dx d\nu(\lambda) < \infty. \quad \square$$

Now we turn to the proof of Theorem 10.

Proof of Theorem 10. We follow the proof of Theorem 14 from [12], skipping some parts but including changes needed for higher dimensions.

By conditions (C1), (C2), (C3) and (C4+) we are at the position of applying Theorem 9 for the functions $M_{\lambda,a}(z)$ defined at the previous section. Let $\mathcal{M}_a(z)$ be the majorant given by Theorem 9 part (i), for $K = Q(a)$, $G_\lambda(x) = G(x; \lambda)$ and $F(z; \lambda) = M_{\lambda,a}(z)$. First, we show that $\mathcal{M}_a(n/a) = \mathcal{G}(n/a)$ for each $n \in \mathbb{Z}^d$ and then we will conclude that $\mathcal{M}_a(z)$ is extremal. Let $\mathcal{D}(x) := \mathcal{M}_a(x) - \mathcal{G}(x)$, by Theorem 9 we know that $\mathcal{D} \in L^1(\mathbb{R}^d)$. Define $\alpha = a_1 a_2 \dots a_d$ and

$$P(x) = \frac{1}{\alpha} \sum_{n \in \mathbb{Z}^d} \mathcal{D}((x+n)/a).$$

It follows from Fubini's theorem that $P(x)$ is defined almost everywhere, is integrable on $\mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$ and $\widehat{P}(k) = \widehat{\mathcal{D}}(ak)$ for all $k \in \mathbb{Z}^d$. Therefore, we have the following identity

$$P * F_R(x) = \sum_{\substack{n \in \mathbb{Z}^d \\ |n_j| \leq R}} \prod_{j=1}^d \left(1 - \frac{|n_j|}{R+1}\right) \widehat{\mathcal{D}}(an) e(x \cdot n) \tag{4.14}$$

for each positive integer R and $x \in \mathbb{R}^d$, where

$$\begin{aligned} F_R(x) &= \prod_{j=1}^d \frac{1}{R+1} \left(\frac{\sin \pi(R+1)x_j}{\sin \pi x_j} \right)^2 \\ &= \sum_{\substack{n \in \mathbb{Z}^d \\ |n_j| \leq R}} \prod_{j=1}^d \left(1 - \frac{|n_j|}{R+1}\right) e(x \cdot n) \end{aligned} \tag{4.15}$$

is the product of one-dimensional Fejér kernels. From (4.11) and (4.14) we have

$$\begin{aligned} P * F_R(0) &= \widehat{\mathcal{D}}(0) - \sum_{\substack{n \neq 0 \\ |n_j| \leq R}} \prod_{j=1}^d \left(1 - \frac{|n_j|}{R+1}\right) \int_{\Lambda} \widehat{G}_\lambda(an) d\nu(\lambda) \\ &= \widehat{\mathcal{D}}(0) - \int_{\Lambda} \left\{ \sum_{\substack{n \neq 0 \\ |n_j| \leq R}} \prod_{j=1}^d \left(1 - \frac{|n_j|}{R+1}\right) \widehat{G}_\lambda(an) \right\} d\nu(\lambda). \end{aligned}$$

Since $P * F_R$ is a non-negative function and the term in the brackets above is positive we may apply Fatou’s lemma to obtain

$$\begin{aligned}
 \widehat{D}(0) &\geq \liminf_{R \rightarrow \infty} P * F_R(0) \\
 &+ \int_{\Lambda} \liminf_{R \rightarrow \infty} \left\{ \sum_{\substack{n \neq 0 \\ |n_j| \leq R}} \prod_{j=1}^d \left(1 - \frac{|n_j|}{R+1}\right) \widehat{G}_{\lambda}(an) \right\} dv(\lambda) \\
 &= \liminf_{R \rightarrow \infty} P * F_R(0) + \int_{\Lambda} \sum_{n \neq 0} \widehat{G}_{\lambda}(an) dv(\lambda). \tag{4.16}
 \end{aligned}$$

From the Poisson Summation formula and the fact that $M_{\lambda,a}(n/a) = G_{\lambda}(n/a)$ for all $n \in \mathbb{Z}^d$, we have

$$\widehat{M}_{\lambda,a}(0) - \widehat{G}_{\lambda}(0) = \sum_{\substack{n \in \mathbb{Z}^d \\ n \neq 0}} \widehat{G}_{\lambda}(an).$$

By (4.10) we conclude that second term on the (RHS) of (4.16) is equal to $\widehat{D}(0)$. This implies that

$$\liminf_{R \rightarrow \infty} P * F_R(0) \leq 0 \implies \liminf_{R \rightarrow \infty} P * F_R(0) = 0.$$

Using this fact with the definition of $P(x)$, we have

$$\begin{aligned}
 \liminf_{R \rightarrow \infty} P * F_R(0) &= \liminf_{R \rightarrow \infty} \int_{\frac{1}{2}Q} P(-y)F_R(y)dy \\
 &= \liminf_{R \rightarrow \infty} \int_{\frac{1}{2}Q} \sum_{n \in \mathbb{Z}^d/a} \frac{1}{\alpha} \mathcal{D}((n - y)/a)F_R(y)dy \\
 &= \liminf_{R \rightarrow \infty} \sum_{n \in \mathbb{Z}^d/a} \int_{\frac{1}{2}Q} \frac{1}{\alpha} \mathcal{D}((n - y)/a)F_R(y)dy \\
 &\geq \sum_{n \in \mathbb{Z}^d} \liminf_{R \rightarrow \infty} \int_{\frac{1}{2}Q} \frac{1}{\alpha} \mathcal{D}((n - y)/a)F_R(y)dy \\
 &= \sum_{n \in \mathbb{Z}^d} \frac{1}{\alpha} \mathcal{D}(n/a),
 \end{aligned}$$

where we have used the positivity of $\mathcal{D}(x)$ and $F_R(x)$, Fubini’s theorem and Fatou’s lemma. The last equality is due to Féjer’s theorem for the continuity of $\mathcal{D}(x)$ at the lattice points \mathbb{Z}^d/a (see Section 3.3 of [23]). Recalling that

$\mathcal{D}(x) \geq 0$ for each $x \in \mathbb{R}^d$, it implies that $\mathcal{D}(n/a) = 0$ which in turn implies $\mathcal{M}_a(n/a) = \mathcal{G}(n/a)$ for each $n \in \mathbb{Z}^d$.

To conclude, note that if $\mathcal{F}(z)$ is an real entire majorant of exponential type with respect to $Q(a)$ such that $\mathcal{F} - \mathcal{G} \in L^1(\mathbb{R}^d)$ then $\mathcal{F} - \mathcal{M}_a \in L^1(\mathbb{R}^d)$. Thus, the following Poisson summation formula holds pointwise

$$\int_{\mathbb{R}^d} \{\mathcal{F}(x) - \mathcal{M}_a(x)\}dx = \frac{1}{\alpha} \sum_{n \in \mathbb{Z}^d} \{\mathcal{F}((n + y)/a) - \mathcal{M}_a((n + y)/a)\}$$

for every $y \in \mathbb{R}^d$. If we take $y = 0$ and use that $\mathcal{M}_a(z)$ interpolates $\mathcal{G}(x)$ at the lattice \mathbb{Z}^d/a we conclude that $\mathcal{M}_a(z)$ is extremal, and this concludes the theorem. □

Before we turn to the proof of Theorem 11, we need a technical lemma, we present the proof of that later.

Lemma 13. *The functions*

$$\phi_a : \lambda \in \Lambda \mapsto \left\{ \sum_{j=1}^d \frac{\Theta(\frac{1}{2}; ia_j^2/\lambda_j)}{\Theta(0; ia_j^2/\lambda_j)} - (d - 1) \right\} \prod_{j=1}^d \Theta(0; ia_j^2/\lambda_j)$$

indexed by $a \in \Lambda$ satisfy

- (i) $\phi_a(\lambda) \leq \phi_b(\lambda)$ if $a_j \leq b_j$ for all $j \in \{1, \dots, d\}$
- (ii) For all $\lambda \in \Lambda$, we have

$$\lim_{a \uparrow \infty} \phi_a(\lambda) = 1$$

where $a \uparrow \infty$ means that $a_j \uparrow \infty$ for all j .

Proof of Theorem 11. First we prove (4.8) and then we conclude with a proof of (4.9).

Let $\mathcal{F}(z)$ be a real entire minorant of $\mathcal{G}(x)$ with exponential type with respect to $Q(a)$. We can assume that $\mathcal{D}(x) = \mathcal{G}(x) - \mathcal{F}(x) \in L^1(\mathbb{R}^d)$ otherwise (4.8) is trivial. Define $\alpha = a_1 \dots a_d$. By Fubini's theorem the function

$$P(x) = \frac{1}{\alpha} \sum_{n \in \mathbb{Z}^d} \mathcal{D}((x + n)/a)$$

is defined almost everywhere, integrable over $\frac{1}{2}Q$ and $\widehat{P}(n) = \widehat{\mathcal{D}}(an)$ for all $n \in \mathbb{Z}^d$. Let $F_R(x)$ be the multidimensional Féjér kernel as defined in (4.15),

hence we have the following equality

$$P * F_R(x) = \sum_{\substack{n \in \mathbb{Z}^d \\ |n_j| \leq R}} \prod_{j=1}^d \left(1 - \frac{|n_j|}{R}\right) \widehat{D}(an) e(x \cdot n). \tag{4.17}$$

By condition (C3) we obtain

$$\begin{aligned} \widehat{D}(0) &= P * F_R(u/2) \\ &\quad - \sum_{\substack{n \neq 0 \\ |n_j| \leq R}} \prod_{j=1}^d \left(1 - \frac{|n_j|}{R+1}\right) \int_{\Lambda} \widehat{G}_{\lambda}(an) d\nu(\lambda) (-1)^{u \cdot n} \\ &= P * F_R(u/2) \\ &\quad + \int_{\Lambda} \left\{ - \sum_{\substack{n \neq 0 \\ |n_j| \leq R}} \prod_{j=1}^d \left(1 - \frac{|n_j|}{R+1}\right) \widehat{G}_{\lambda}(an) (-1)^{u \cdot n} \right\} d\nu(\lambda). \end{aligned}$$

Since $x \mapsto \widehat{G}_{\lambda}(x)$ is a product of radially decreasing functions we easily see that the term in the brackets is positive, thus we can apply Fatou’s lemma to obtain

$$\begin{aligned} \widehat{D}(0) &\geq \liminf_{R \rightarrow \infty} P * F_R(u/2) \\ &\quad + \int_{\Lambda} \liminf_{R \rightarrow \infty} \left\{ - \sum_{\substack{n \neq 0 \\ |n_j| \leq R}} \prod_{j=1}^d \left(1 - \frac{|n_j|}{R+1}\right) \widehat{G}_{\lambda}(an) (-1)^{u \cdot n} \right\} d\nu(\lambda) \\ &= \liminf_{R \rightarrow \infty} P * F_R(u/2) + \int_{\Lambda} - \sum_{n \neq 0} \widehat{G}_{\lambda}(an) (-1)^{u \cdot n} d\nu(\lambda). \tag{4.18} \end{aligned}$$

Using the properties of the Fourier transform of $G_{\lambda}(x)$ and the definition of the Theta Function (2.3), we find that the term inside the integral in (4.18) is equal to

$$\prod_{j=1}^d \lambda_j^{-\frac{1}{2}} - \prod_{j=1}^d \Theta\left(\frac{1}{2}; ia_j^2/\lambda_j\right) \lambda_j^{-\frac{1}{2}}.$$

This proves the lower bound estimate (4.8), since $P * F_R$ is a non-negative function.

Now, we turn to the proof of the L^1 convergence (4.9). By conditions (C1), (C2), (C3) and (C4-) we are at the position of applying Theorem 9 for the functions $L_{\lambda,a}(z)$ defined at the previous section. Let $\mathcal{L}_a(z)$ be the minorant given

by Theorem 9 part (ii), for

$$K = Q(a), \quad G_\lambda(x) = G(x; \lambda) \text{ and } F(z; \lambda) = L_{\lambda,a}(z).$$

We obtain

$$\begin{aligned} \int_{\mathbb{R}^d} G_\lambda(x) - L_a(x) dx &= \int_\Lambda \int_{\mathbb{R}^d} G_\lambda(x) - L_{\lambda,a}(x) dx d\nu(\lambda) \\ &= \int_\Lambda \left\{ 1 - \left\{ \sum_{j=1}^d \frac{\Theta(\frac{1}{2}; ia_j^2/\lambda_j)}{\Theta(0; ia_j^2/\lambda_j)} - (d-1) \right\} \prod_{j=1}^d \Theta(0; ia_j^2/\lambda_j) \right\} \\ &\quad \times \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} d\nu(\lambda). \end{aligned} \tag{4.19}$$

Using Lemma 13 we see that the functions inside the integral form a decreasing sequence (indexed by $a \in \Lambda$) converging to zero as $a \uparrow \infty$. Therefore, by the monotone convergence theorem, the integral goes to 0 and the proof is complete. \square

Proof of Lemma 13. To see that the functions

$$\phi_a : \lambda \in \Lambda \mapsto \left\{ \sum_{j=1}^d \frac{\Theta(\frac{1}{2}; ia_j^2/\lambda_j)}{\Theta(0; ia_j^2/\lambda_j)} - (d-1) \right\} \prod_{j=1}^d \Theta(0; ia_j^2/\lambda_j) \tag{4.20}$$

form an increasing sequence of functions indexed by $a \in \Lambda$ as each $a_j \uparrow \infty$, is enough to prove that for every $\lambda \in \Lambda$ and $k \in \{1, \dots, d\}$

$$\frac{\partial}{\partial a_k} \phi_a(\lambda) \geq 0.$$

For all $t > 0$ denote

$$u_j(t) = \Theta(0; it^2/\lambda_j) \quad \text{and} \quad v_j(t) = \Theta(\frac{1}{2}; it^2/\lambda_j),$$

we can use the Leibniz rule to obtain

$$\frac{\partial}{\partial a_k} \phi_a(\lambda) = \left(v'_k(a_k) + \left\{ \sum_{\substack{j=1 \\ j \neq k}}^d \frac{v_j(a_j)}{u_j(a_j)} - (d-1) \right\} u'_k(a_k) \right) \prod_{\substack{j=1 \\ j \neq k}}^d u_j(a_j). \tag{4.21}$$

Using the summation formula (2.3), we see that the functions $t \mapsto (u_j(t) - 1)$ is a sum of positive decreasing functions that decreases to 0 as $t \uparrow \infty$, thus $u_j(t)$

is a decreasing function that decreases to 1 as $t \uparrow \infty$. Analogously, using the product formula (3.16), each $v_j(t)$ is a product of positive increasing functions that increases to 1 as $t \uparrow \infty$, thus $v_j(t)$ is a positive increasing function that increases to 1 as $t \uparrow \infty$. Therefore, the term inside the parenthesis in (4.21) is positive, which implies that $\frac{\partial}{\partial a_k} \phi_a(\lambda) > 0$, and this proves item (i).

Since

$$\phi_a(\lambda) = \left\{ \sum_{j=1}^d \frac{v_j(a_j)}{u_j(a_j)} - (d - 1) \right\} \prod_{j=1}^d u_j(a_j),$$

we see that $\phi_a(\lambda)$ converges to 1, for every $\lambda \in \Lambda$, as each $a_j \uparrow \infty$ and this proofs item (ii). □

5 Further Results

In this section we will use the machinery of the previous section to construct one-sided approximations by trigonometric polynomials. After, as in Part III of [12], we will give some examples of functions that our method is applicable and then we present some Hilbert-type inequalities that arise from the constructions of section §4.

5.1 Periodic Analogues

In this subsection we find the best approximations by trigonometric polynomials for functions that are, in some sense, subordinate to Theta functions. The proofs of the theorems in this section are almost identical to the proofs of the previous sections, and thus we state the theorems without proof.

Definition 14. Let $a = (a_1, a_2, \dots, a_d) \in \mathbb{Z}_+^d$ (i.e. $a_j \geq 1 \forall j$), we will say that the degree of a trigonometric Polynomial $P(x)$ is less than a (degree $P < a$) if

$$P(x) = \sum_{-a < n < a} \widehat{P}(n)e(n \cdot x).$$

The problems we are interested to solve have the following general form

Periodic Majorization Problem. Fix an $a \in \mathbb{Z}_+^d$ (called the *degree*) and a Lebesgue measurable real periodic function $g : \mathbb{T}^d \rightarrow \mathbb{R}$. Determine the value of

$$\inf \int_{\mathbb{T}^d} |F(x) - g(x)| dx, \tag{5.1}$$

where the infimum is taken over functions $F : \mathbb{T}^d \rightarrow \mathbb{R}$ satisfying

- (i) $F(x)$ is a real trigonometric polynomial.
- (ii) Degree of $F(x)$ is less than a .
- (iii) $F(x) \geq g(x)$ for every $x \in \mathbb{T}^d$.

If the infimum is achieved, then identify the extremal functions $F(z)$. Similarly there exist the minorant problem

Periodic Minorization Problem. Solve the previous problem with condition (iii) replaced by the condition

$$(iv) F(x) \leq g(x) \text{ for every } x \in \mathbb{T}^d.$$

Now we define for every $\lambda \in \Lambda$, the periodization of the Gaussian function $G_\lambda(x)$ by

$$f_\lambda(x) := \sum_{n \in \mathbb{Z}^d} G_\lambda(x + n) = \sum_{n \in \mathbb{Z}^d} \prod_{j=1}^d e^{-\lambda_j \pi (x_j + n_j)^2} = \prod_{j=1}^d \Theta(x_j; i/\lambda_j) \lambda_j^{-\frac{1}{2}}.$$

Theorem 15 (Existence). For a given $a = (a_1, a_2, \dots, a_d) \in \mathbb{Z}_+^d$, let Λ be a measurable space of parameters, and for each $\lambda \in \Lambda$, let $R_\lambda(x)$ be a real trigonometric polynomial with degree less than a . Let ν be a non-negative measure in Λ that satisfies

$$\int_\Lambda \int_{\mathbb{T}^d} |R_\lambda(x) - f_\lambda(x)| dx d\nu(\lambda) < \infty. \tag{5.2}$$

Suppose that $g : \mathbb{T}^d \rightarrow \mathbb{R}$ is a continuous periodic function such that

$$\widehat{g}(k) = \int_\Lambda \widehat{G}_\lambda(k) d\nu(\lambda), \tag{5.3}$$

for all $k \in \mathbb{Z}^d$ such that $|k_j| \geq a_j$ for some $j \in \{1, 2, \dots, d\}$. Then

- (i) if $f_\lambda(x) \leq R_\lambda(x)$ for each $x \in \mathbb{T}^d$ and $\lambda \in \Lambda$, then there exist a trigonometric polynomial $m_a(x)$ with degree $m_a < a$, such that $m_a(x) \geq g(x)$ for all $x \in \mathbb{T}^d$ and

$$\int_{\mathbb{T}^d} m_a(x) - g(x) dx$$

is equal to the (LHS) of (5.2).

(ii) if $R_\lambda(x) \leq f_\lambda(x)$ for each $x \in \mathbb{T}^d$ and $\lambda \in \Lambda$, then there exist a trigonometric polynomial $l_a(x)$ with degree $l_a < a$, such that $l_a(x) \leq g(x)$ in \mathbb{T}^d , and

$$\int_{\mathbb{T}^d} g(x) - l_a(x) dx,$$

is equal to (LHS) of (5.2).

Before we state the main theorems of this section we need some definitions. The functions $M_{\lambda,a}(x)$ and $L_{\lambda,a}(x)$, defined in (3.3) and (3.2), belong to $L^1(\mathbb{R}^d)$, thus, by the Plancharel-Pólya theorem (see [41]) and the periodic Fourier inversion formula, their respective periodizations are trigonometric polynomials of degree less than a , that is

$$m_{\lambda,a}(x) := \sum_{n \in \mathbb{Z}^d} M_{\lambda,a}(x + n) = \sum_{-a < n < a} \widehat{M}_{\lambda,a}(n) e^{2\pi i n \cdot x} \tag{5.4}$$

and

$$l_{\lambda,a}(x) := \sum_{n \in \mathbb{Z}^d} L_{\lambda,a}(x + n) = \sum_{-a < n < a} \widehat{L}_{\lambda,a}(n) e^{2\pi i n \cdot x} \tag{5.5}$$

holds for each $x \in \mathbb{T}^d$. The following theorem offers a resolution to the Majorization Problem for a specific class of functions.

Theorem 16 (Gaussian Subordination – Periodic Majorant). *Let $a \in \mathbb{Z}_+^d$ and ν be non-negative Borel measure on Λ that satisfies*

$$\int_{\Lambda} \prod_{j=1}^d \lambda_k^{-\frac{1}{2}} \left\{ \prod_{j=1}^d \Theta(0; ia_j^2/\lambda_j) - 1 \right\} d\nu(\lambda) < \infty. \tag{5.6}$$

Let $g : \mathbb{T}^d \rightarrow \mathbb{R}$ be a continuous periodic function such that

$$\widehat{g}(k) = \int_{\Lambda} \widehat{G}_\lambda(k) d\nu(\lambda), \tag{5.7}$$

for all $k \in \mathbb{Z}^d$ such that $|k_j| \geq a_j$ for some $j \in \{1, 2, \dots, d\}$. Then for every real trigonometric polynomial $P(x)$, with degree $P < a$ and $P(x) \geq g(x)$ for all $x \in \mathbb{T}^d$, we have

$$\int_{\mathbb{T}^d} P(x) - g(x) dx \geq \int_{\Lambda} \prod_{j=1}^d \lambda_k^{-\frac{1}{2}} \left\{ \prod_{j=1}^d \Theta(0; ia_j^2/\lambda_j) - 1 \right\} d\nu(\lambda). \tag{5.8}$$

Moreover, there exists a real trigonometric polynomial m_a , with degree $m_a < a$, such that $m_a(x)$ is a majorant of $g(x)$ that interpolates $g(x)$ on the lattice \mathbb{Z}^d/a and equality at (5.8) holds.

Theorem 17 (Gaussian Subordination – Periodic Minorant). Let $a \in \mathbb{Z}_+^d$ and ν be non-negative Borel measure on Λ such that

$$\int_{\Lambda} \left\{ 1 - \left\{ \sum_{j=1}^d \frac{\Theta(\frac{1}{2}; ia_j^2/\lambda_j)}{\Theta(0; ia_j^2/\lambda_j)} - (d-1) \right\} \prod_{j=1}^d \Theta(0; ia_j^2/\lambda_j) \right\} \times \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} d\nu(\lambda) < \infty. \tag{5.9}$$

Let $g : \mathbb{T}^d \rightarrow \mathbb{R}$ be continuous periodic function such that

$$\widehat{g}(k) = \int_{\Lambda} \widehat{G}_{\lambda}(k) d\nu(\lambda) \tag{5.10}$$

for all $k \in \mathbb{Z}^d$ such that $|k_j| \geq a_j$ for some $j \in \{1, 2, \dots, d\}$. Then, if $P(z)$ is a real trigonometric polynomial with degree less than a that minorizes $g(x)$, we have

$$\int_{\mathbb{T}^d} g(x) - P(x) dx \geq \int_{\Lambda} \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} \left\{ 1 - \prod_{j=1}^d \Theta(\frac{1}{2}; ia_j^2/\lambda_j) \right\} d\nu(\lambda). \tag{5.11}$$

Furthermore, there exists a family of trigonometric polynomial minorants $\{l_a(x) : a \in \mathbb{Z}_+^d\}$ with degree $l_a < a$, such that the integral

$$\int_{\mathbb{T}^d} g(x) - l_a(x) dx$$

is equal to the quantity in (5.9), and

$$\lim_{a \uparrow \infty} \int_{\mathbb{T}^d} g(x) - l_a(x) dx = 0, \tag{5.12}$$

where $a \uparrow \infty$ means $a_j \uparrow \infty$ for every j .

Corollary 18. Under the hypothesis of Theorem 17, suppose also that there exist an $R > 0$ such that $\text{supp}(\nu) \subset \Lambda \cap Q(R)$, and

$$\int_{\mathbb{T}^d} g(x) dx = \int_{\Lambda} \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} d\nu(\lambda) < \infty.$$

Then, there exist a constant $\alpha_0 > 0$, such that if $\alpha := \min\{a_j\} \geq \alpha_0$ and if $P(x)$ is a trigonometric polynomial with degree $P < a$ that minorizes $g(x)$, we have

$$\int_{\mathbb{T}^d} P(x)dx \leq (1 + 5de^{-\alpha^2/R}) \int_{\mathbb{T}^d} l_a(x)dx.$$

5.2 The Class of Contemplated Functions

We define the class

$$\mathfrak{G}^d = \bigcap_{a \in \Lambda} \mathfrak{G}_-^d(a) \cap \mathfrak{G}_+^d(a).$$

This is the class of pairs such that Theorems 10 and 11 are applicable for every $a \in \Lambda$. In this subsection we present conditions for a pair (\mathcal{G}, ν) belong to this class.

Some interesting properties arise when ν is concentrated in the diagonal. For every $\eta \in [0, 1]$ we define $\Lambda_\eta = \{\lambda \in \Lambda : \eta\lambda_j \leq \lambda_k \forall j, k\}$ and we note that $\Lambda_0 = \Lambda$ and $\Lambda_1 = \{\lambda \in \Lambda : \lambda_j = \lambda_k \forall j, k\}$ is the diagonal.

Proposition 19. *Let (\mathcal{G}, ν) be a pair that satisfies conditions (C1), (C2) and (C3) for every $a \in \Lambda$. Suppose that $\text{supp}(\nu) \subset \Lambda_\eta$ for some $\eta \in (0, 1]$ and $\nu(\Lambda_\eta \setminus \mathcal{Q}(R)) < \infty$ for every $R > 0$. Then $(\mathcal{G}, \nu) \in \mathfrak{G}^d$.*

Proof. We only prove that condition (C4-) holds, the condition (C4+) is analogous. Given an $a \in \Lambda$, define the function

$$\phi_a : \lambda \in \Lambda \mapsto \left(\sum_{j=1}^d \frac{\Theta(\frac{1}{2}; ia_j^2/\lambda_j)}{\Theta(0; ia_j^2/\lambda_j)} - (d - 1) \right) \prod_{j=1}^d \Theta(0; ia_j^2/\lambda_j).$$

By (2.3) and the Poisson summation formula, we have the following estimates

$$1 - \Theta(\frac{1}{2}; i/t) \sim 2e^{-\pi/t} \text{ as } t \rightarrow 0, \tag{5.13}$$

$$\Theta(\frac{1}{2}; i/t) \sim 2t^{\frac{1}{2}}e^{-\pi t/4} \text{ as } t \rightarrow \infty \tag{5.14}$$

and

$$\Theta(0; i/t) - 1 \sim 2e^{-\pi/t} \text{ as } t \rightarrow 0, \tag{5.15}$$

$$\Theta(0; i/t) \sim t^{\frac{1}{2}} \text{ as } t \rightarrow \infty \tag{5.16}$$

where the symbol \sim means that the quotient converges to 1. Using the (LHS) inequality of Lemma 8 we conclude that exists an $R > 0$ and a $C > 0$ such that

$$\phi_a(\lambda) \geq 1 - C \sum_{j=1}^d e^{-\pi a_j^2/\lambda_j}$$

for every $\lambda \in \Lambda \cap Q(R)$. Choose $l \in \{1, \dots, d\}$ such that $a_l \leq a_j$ for every j . If $\lambda \in \Lambda_\eta \cap Q(R)$ we have

$$\begin{aligned} \{1 - \phi_a(\lambda)\} \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} &\leq C \left(\sum_{j=1}^d e^{-\pi a_j^2/\lambda_j} \right) \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} \leq d C e^{-\pi a_l^2 \eta/\lambda_l} \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} \\ &\leq d C \prod_{j=1}^d e^{-\pi a_l^2 \eta^2/(d\lambda_j)} \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} = d C \widehat{G}_\lambda(\beta u), \end{aligned}$$

where $\beta = a_l \eta d^{-\frac{1}{2}}$ and $u = (1, \dots, 1)$.

By estimates (5.13)-(5.16) we see that the functions $\Theta(\frac{1}{2}; i/t)t^{-\frac{1}{2}}$ and $\Theta(0; i/t)t^{-\frac{1}{2}}$ are bounded for $t \in [\eta R, \infty)$, and thus, we conclude that the function

$$\lambda \in \Lambda \mapsto \phi_a(\lambda) \prod_{j=1}^d \lambda_j^{-\frac{1}{2}}$$

is bounded on $\Lambda_\eta \setminus Q(R)$, since it is a finite sum of products of these theta functions.

Since $\eta > 0$, we obtain that the function

$$\lambda \in \Lambda \mapsto \{1 - \phi_a(\lambda)\} \prod_{j=1}^d \lambda_j^{-\frac{1}{2}}$$

is bounded in $\Lambda_\eta \setminus Q(R)$, let say by C' . Therefore, we have

$$\begin{aligned} &\int_{\Lambda_\eta} \{1 - \phi_a(\lambda)\} \prod_{j=1}^d \lambda_j^{-\frac{1}{2}} d\nu(\lambda) \\ &\leq d C \int_{\Lambda_\eta \cap Q(b)} \widehat{G}_\lambda(\beta u) d\nu(\lambda) + C' \nu(\Lambda_\eta \setminus Q(b)) < \infty, \end{aligned}$$

which is finite by condition **(C3)** and the hypotheses of this lemma. Thus ν satisfies condition **(C4-)** and this concludes the proof. \square

Proposition 20. *Let ν be a probability measure on Λ with $\text{supp}(\nu) \subset \Lambda_\eta$ for some $\eta \in (0, 1]$. Define the function*

$$G(x) = \int_{\Lambda} G_\lambda(x) d\nu(\lambda) \tag{5.17}$$

for all $x \in \mathbb{R}^d$. Then $(G, \nu) \in \mathfrak{G}^d$.

Proof. Is easy to see that G is a bounded, continuous and radially decreasing function, thus G satisfies conditions **(C1)**. Note that for every $x \neq 0$, the function

$$\lambda \in \Lambda \mapsto \widehat{G}_\lambda(x)$$

is bounded on Λ_η and this bound can be taken uniform for x outside any neighborhood of the origin. Thus condition **(C2)** holds, and using Fubini’s theorem, condition **(C3)** also holds. We conclude, by Proposition 19, that $(G, \nu) \in \mathfrak{G}^d$. □

Due to a classical result of Schoenberg (see [43]), a radial function $G(x) = G(|x|)$ admits the representation (5.17) for a probability ν supported on the diagonal Λ_1 if and only if the radial extension to \mathbb{R}^n of $G(r)$ is positive definite, for all $n > 0$. And this occurs if and only if the function $G(r^{1/2})$ is completely monotone. As consequence of this fact and Proposition 20 the following multi-dimensional versions of the functions in Section 11 of [12] are contemplated

Example 1.

$$g(x) = e^{-\alpha|x|^r} \in \mathfrak{G}^d, \quad \alpha > 0 \text{ and } 0 < r < 2.$$

Example 2.

$$g(x) = (|x|^2 + \alpha^2)^{-\beta} \in \mathfrak{G}^d, \quad \alpha > 0 \text{ and } \beta > 0.$$

Example 3.

$$g(x) = -\log\left(\frac{|x|^2 + \alpha^2}{|x|^2 + \beta^2}\right) \in \mathfrak{G}^d, \quad \text{for } 0 < \alpha < \beta.$$

The following example is a high dimensional analogue of Corollary 21 [12].

Example 4. *Given a $\sigma \in (0, \infty) \setminus 2\mathbb{Z}_+$ consider the measure ν_σ , that is supported on the diagonal Λ_1 and defined on a Borel set $E \subset \Lambda$ by*

$$\nu_\sigma(E) = C(\sigma) \int_{P(E \cap \Lambda_1)} t^{-\frac{\sigma}{2}-1} dt \tag{5.18}$$

where $P : \Lambda_1 \rightarrow \mathbb{R}$ is the projection $P(tu) = t$ for all $t \geq 0$ and

$$C(\sigma) = \pi^{-\frac{\sigma}{2}} / \Gamma\left(-\frac{\sigma}{2}\right).$$

where $\Gamma(z)$ is the classical Gamma function. Define on \mathbb{R}^d the function

$$G_\sigma(x) = |x|^\sigma \text{ for } \sigma \in (0, \infty) \setminus 2\mathbb{Z}_+$$

we claim that $(G_\sigma, \nu_\sigma) \in \mathfrak{G}^d$. To see this, first we note that Lemma 18 of [12] has a trivial generalization to the several variable setting, that is, for every function $\varphi \in S(\mathbb{R}^d)$ that is zero on a neighborhood of the origin, we have

$$\int_{\mathbb{R}^d} \widehat{\varphi}(x)|x|^\sigma dx = A(d, \sigma) \int_{\mathbb{R}^d} \varphi(x)|x|^{-d-\sigma} dx,$$

where

$$A(d, \sigma) = \pi^{-\sigma-d/2} \Gamma\left(\frac{d+\sigma}{2}\right) / \Gamma\left(-\frac{\sigma}{2}\right).$$

Secondly, note that

$$\frac{A(d, \sigma)}{|x|^{d+\sigma}} = \int_{\Lambda} \widehat{G}_\lambda(x) d\nu_\sigma(\lambda).$$

Hence, by Proposition 19 the pair (G_σ, ν_σ) belongs to \mathfrak{G}^d .

5.2.1 The Periodic Case

Given a pair $(G, \nu) \in \mathfrak{G}^d$, suppose that $G \in L^1(\mathbb{R}^d)$ and the periodization

$$x \in \mathbb{T}^d \mapsto \sum_{n \in \mathbb{Z}^d} G(n+x)$$

is equal almost everywhere to a continuous function $g(x)$. We easily see that the pair (g, ν) is contemplated by the Theorems 16 and 17 for every degree $a \in \mathbb{Z}_+^d$. Thus, the period method contemplates the following functions

Example 5.

$$\begin{aligned} g(x) &= \prod_{j=1}^d \Theta(x_j; i/\lambda_j) \\ &= \prod_{j=1}^d \lambda_j^{\frac{1}{2}} \sum_{n \in \mathbb{Z}^d} G_\lambda(n+x), \text{ for all } \lambda \in \Lambda. \end{aligned} \tag{5.19}$$

Example 6.

$$g(x) = \sum_{n \in \mathbb{Z}^d} e^{-\alpha|n+x|^r}, \text{ for all } \alpha > 0 \text{ and } 0 < r < 2. \tag{5.20}$$

However, we cannot use this construction for the case of the functions $\mathcal{G}_\sigma(x) = |x|^\sigma$ of Example 4. The next proposition tell us that if the Fourier coefficients of \mathcal{G} decay sufficiently fast, then the periodization of \mathcal{G} via Poisson summation formula is contemplated by the periodic method.

Proposition 21. *Let $(\mathcal{G}, \nu) \in \mathfrak{G}^d$. Suppose that exist constants $C > 0$ and $\delta > d/2$ such that*

$$\int_{\Lambda} \widehat{\mathcal{G}}_\lambda(x) d\nu(\lambda) \leq C|x|^{-\delta}$$

if $|x| \geq 1$. Define the function

$$x \mapsto \lim_{N \rightarrow \infty} \sum_{\substack{n \in \mathbb{Z}^d \\ 0 < |n| < N}} \int_{\Lambda} \widehat{\mathcal{G}}_\lambda(n) d\nu(\lambda) e(n \cdot x),$$

where the limit is taken in $L^2(\mathbb{T}^d)$, and suppose this can be identified with a continuous function $g(x)$. Then the pair (g, ν) satisfies all the conditions of Theorems 16 and 17 for every $a \in \mathbb{Z}_+^d$.

With this last proposition we see that the following example is contemplated by the periodic method of subsection 5.1.

Example 7.

$$g_\sigma(x) = \sum_{\substack{n \in \mathbb{Z}^d \\ n \neq 0}} |n|^{-d-\sigma} e(n \cdot x), \text{ for all } \sigma > 0. \tag{5.21}$$

5.3 Hilbert-Type Inequalities

Given an $a \in \Lambda$ and $x \in \mathbb{R}^d$ define the norm

$$|x|_a = \max\{|x_j/a_j| : j = 1, \dots, d\}.$$

Given an $a \in \Lambda$ we say that a sequence $\{\xi_k\}_{k \in \mathbb{Z}}$ of vectors in \mathbb{R}^d is a -separated if $|\xi_k - \xi_l|_a \geq 1$ for all $l \neq k$. We have the following proposition

Proposition 22. *Let $(\mathcal{G}, \nu) \in \mathfrak{G}^d$, $a \in \Lambda$ and $\{\xi_k\}_{k \in \mathbb{Z}}$ an a -separated sequence of vectors. Then for every finite sequence of complex numbers $\{w_{-N}, \dots, w_0,$*

..., w_N }, we have

$$\begin{aligned}
 -A(a, d, \nu) \sum_{n=-N}^N |w_n|^2 &\leq \sum_{\substack{n,m=-N \\ n \neq m}}^N w_n \bar{w}_m \widehat{G}(\xi_n - \xi_m) \\
 &\leq B(a, d, \nu) \sum_{n=-N}^N |w_n|^2
 \end{aligned} \tag{5.22}$$

where $A(a, d, \nu)$ is equal to the quantity (4.6) and $B(a, d, \nu)$ is equal to the quantity (4.5). Furthermore the constant $B(a, d, \nu)$ is sharp.

Remark. For $\xi \neq 0$ we write

$$\widehat{G}(\xi) = \int_{\Lambda} \widehat{G}_\lambda(\xi) d\nu(\lambda).$$

Proof. To prove inequality in (5.22), define the function $D = \mathcal{M}_a - \mathcal{G}$ where \mathcal{M}_a is given by Theorem 10. Since \mathcal{M}_a is of exponential type with respect to $Q(a)$ we obtain

$$\sum_{n,m=-N}^N w_n \bar{w}_m \widehat{D}(\xi_n - \xi_m) = B(a, d, \nu) \sum_{n=-N}^N |w_n|^2 - \sum_{\substack{n,m=-N \\ n \neq m}}^N w_n \bar{w}_m \widehat{G}(\xi_n - \xi_m).$$

But the (LHS) of this last equality is positive since $\mathcal{D}(x)$ is a non-negative function. In an analogous way we can prove the (LHS) inequality of (5.22).

For the sharpness of the (RHS) inequality in (5.22) suppose that we could change $B(a, d, \nu)$ by some other constant B' . Consider the sequence $w_n = 1$ for all $n \in \mathbb{Z}$ and let the sequence $\{\xi_n\}$ be an enumeration of the points in the set $J(R) = Q(Ra) \cap a\mathbb{Z}^d$ where $R \in \mathbb{Z}_+$. We obtain

$$\begin{aligned}
 - \sum_{\xi, \xi' \in J(R)} \widehat{D}(\xi - \xi') &= \sum_{\substack{\xi, \xi' \in J(R) \\ \xi \neq \xi'}} \widehat{G}(\xi - \xi') - (2R + 1)^d \widehat{D}(0) \\
 &\leq (2R + 1)^d (B' - \widehat{D}(0)).
 \end{aligned}$$

On the other hand we have

$$\sum_{\xi, \xi' \in J(R)} \widehat{D}(\xi - \xi') = \sum_{|n_j| \leq 2R} \left\{ \prod_{j=1}^d (2R + 1 - |n_j|) \right\} \widehat{D}(an).$$

Hence, we conclude that

$$B' - \widehat{\mathcal{D}}(0) \geq - \sum_{|n_j| \leq 2R} \prod_{j=1}^d \left(1 - \frac{|n_j|}{2R+1}\right) \widehat{\mathcal{D}}(an) - = F_{2R} * P(0),$$

where $F_{2R}(x)$ is the Féjer’s Kernel defined in (4.15) and

$$P(x) = \frac{1}{\alpha} \sum_{n \in \mathbb{Z}^d} \mathcal{D}((n+x)/a),$$

with $\alpha = a_1 \dots a_d$. Replying the arguments of the Theorem 10 we would conclude that

$$\liminf_{R \rightarrow \infty} F_{2R} * P(0) = 0$$

and this implies that $B' \geq \widehat{\mathcal{D}}(0)$, since $\widehat{\mathcal{D}}(0) = B(a, d, \nu)$ this concludes the proof. □

The next corollary is a generalization of Corollary 22 of [12] in the multidimensional setting and is a direct application of Proposition 22 for Example 4. Below $|\cdot|$ stands for the Euclidean norm in \mathbb{R}^d .

Corollary 23. *Let $\sigma > 0$, $a \in \Lambda$ and $\{\xi_k\}_{k \in \mathbb{Z}}$ an a -separated sequence of vectors. Then for every finite sequence of complex numbers $\{w_{-N}, \dots, w_0, \dots, w_N\}$, we have*

$$\begin{aligned} -A(a, d, \sigma) \sum_{n=-N}^N |w_n|^2 &\leq \sum_{\substack{n,m=-N \\ n \neq m}}^N \frac{w_n \bar{w}_m}{|\xi_n - \xi_m|^{d+\sigma}} \\ &\leq B(a, d, \sigma) \sum_{n=-N}^N |w_n|^2, \end{aligned} \tag{5.23}$$

where

$$A(a, d, \sigma) = - \sum_{j=1}^d \sum_{n \in \mathbb{Z}^d \setminus 0} \frac{(-1)^{n_j}}{|an|^{d+\sigma}} + (d-1) \sum_{n \in \mathbb{Z}^d \setminus 0} \frac{1}{|an|^{d+\sigma}}$$

and

$$B(a, d, \sigma) = \sum_{n \in \mathbb{Z}^d \setminus 0} \frac{1}{|an|^{d+\sigma}}.$$

Furthermore, the constant $B(a, d, \sigma)$ is sharp.

6 Concluding Remarks

We mentioned in the introduction that Selberg generalized his construction of majorizing and minorizing the characteristic function of an interval, to majorizing and minorizing the characteristic function of a box by functions whose Fourier transforms are supported in a (possibly different) box. Another way to generalize Selberg's original construction to the several variables setting is to consider majorizing and minorizing the characteristic function of a ball by functions whose Fourier transforms are supported in a (possibly different) ball. This problem was considered by Holt and Vaaler [28] and their methods were recently extended by Carneiro and Littmann [7].

As far as we know, almost nothing is known about the Beurling-Selberg extremal problem in higher dimensions when the Fourier transform is supported on a fixed symmetric convex body K . The following question (perhaps the simplest Beurling-Selberg extremal problem in higher dimensions) is open:

Let K be a symmetric convex body in \mathbb{R}^d . Determine the value of

$$\eta(K) = \inf \int_K F(x) dx$$

where the infimum is taken over continuous integrable functions $F : \mathbb{R}^d \rightarrow \mathbb{R}$ that satisfy (i) $F(0) \geq 1$, (ii) $F(x) \geq 0$ for all $x \in \mathbb{R}^d$, and (iii) $\hat{F}(\xi) = 0$ if $\xi \notin K$.

In one dimension, the solution to this problem is the Fejer kernel for \mathbb{R} . So the solution to the above problem can be thought of as an analogue of the Fejer kernel for K .

It is conjectured [30] by the third named author and Jeffrey Vaaler, that $\eta(K) = 2^d / \text{vol}_d(K)$. Vaaler [50] has shown that this conjecture is true if K is an *extremal body*. That is, if K achieves equality in Minkowski's convex body theorem, then the conjecture in [30] is true. For example, the regular hexagon in \mathbb{R}^2 is an extremal body. The only other body K for which the conjecture is known to hold is the Euclidean ball.

Acknowledgements. The first named author acknowledges the support from FAPERJ-Brazil scholarship grant. The second named author would like to thank IMPA for hosting his visit in the winter of 2012. He would also like to acknowledge partial support from NSF grants DMS-1045119, DMS-0943832, and DMS-1101326. The third named author acknowledges the support from CAPES-Brazil scholarship grant. Lastly, the authors would like to express their

deepest gratitude to Professor Emanuel Carneiro for all the fruitful discussions and for his guidance throughout the writing of this paper.

References

- [1] Roger C. Baker. Sequences that omit a box (modulo 1). *Adv. Math.*, **227**(5) (2011), 1757–1771.
- [2] Jeffrey T. Barton, Hugh L. Montgomery and Jeffrey D. Vaaler. *Note on a Diophantine inequality in several variables*. *Proc. Amer. Math. Soc.*, **129**(2) (2001), 337–345 (electronic).
- [3] Ralph Philip Boas, Jr. *Entire functions*. Academic Press Inc., New York (1954).
- [4] E Carneiro, V Chandee, F. Littmann and M. Milinovich. *Hilbert spaces and the pair correlation of zeros of the riemann zeta-function*. *J. Reine Angew. Math.* (to appear).
- [5] E. Carneiro and F. Littmann. *Bandlimited approximations to the truncated gaussian and applications*. *Constr. Approx.* (to appear).
- [6] E. Carneiro and F. Littmann. *Entire approximations for a class of truncated and odd functions*. *J. Fourier Anal. Appl.* (to appear).
- [7] E. Carneiro and F. Littmann. *Extremal functions in de branges and euclidean spaces*. *Adv. Math.*, (260) (2014), 281–349.
- [8] Emanuel Carneiro. *Sharp approximations to the Bernoulli periodic functions by trigonometric polynomials*. *J. Approx. Theory*, **154**(2) (2008), 90–104.
- [9] Emanuel Carneiro and Vorrapan Chandee. *Bounding $\zeta(s)$ in the critical strip*. *J. Number Theory*, **131**(3) (2011), 363–384.
- [10] Emanuel Carneiro, Vorrapan Chandee and Micah B. Milinovich. *Bounding $S(t)$ and $S_1(t)$ on the Riemann hypothesis*. *Math. Ann.*, **356**(3) (2013), 939–968.
- [11] Emanuel Carneiro and Felipe Gonçalves. *Extremal problems in de branges spaces: The case of truncated and odd functions*. Preprint (2014).
- [12] Emanuel Carneiro, Friedrich Littmann and Jeffrey D. Vaaler. *Gaussian subordination for the Beurling-Selberg extremal problem*. *Trans. Amer. Math. Soc.*, **365**(7) (2013), 3493–3534.
- [13] Emanuel Carneiro and Jeffrey D. Vaaler. *Some extremal functions in Fourier analysis. II*. *Trans. Amer. Math. Soc.*, **362**(11) (2010), 5803–5843.
- [14] Emanuel Carneiro and Jeffrey D. Vaaler. *Some extremal functions in Fourier analysis. III*. *Constr. Approx.*, **31**(2) (2010), 259–288.
- [15] Vorrapan Chandee and K. Soundararajan. *Bounding $|\zeta(1/2 + it)|$ on the Riemann hypothesis*. *Bull. Lond. Math. Soc.*, **43**(2) (2011), 243–250.
- [16] K. Chandrasekharan. *Elliptic functions, volume 281 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin (1985).

- [17] Todd Cochrane. *Trigonometric approximation and uniform distribution modulo one*. Proc. Amer. Math. Soc., **103**(3) (1988), 695–702.
- [18] Louis de Branges. *Homogeneous and periodic spaces of entire functions*. Duke Math. J., **29** (1962), 203–224.
- [19] Louis de Branges. *Hilbert spaces of entire functions*. Prentice-Hall Inc., Englewood Cliffs, N.J., (1968).
- [20] Michael Drmota and Robert F. Tichy. *Sequences, discrepancies and applications*, volume 1651 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin (1997).
- [21] P.X. Gallagher. *Pair correlation of zeros of the zeta function*. J. Reine Angew. Math., **362** (1985), 72–86.
- [22] D.A. Goldston and S.M. Gonek. *A note on $S(t)$ and the zeros of the Riemann zeta-function*. Bull. Lond. Math. Soc., **39**(3) (2007), 482–486.
- [23] Loukas Grafakos. *Classical Fourier analysis*, volume 249 of *Graduate Texts in Mathematics*. Springer, New York, second edition (2008).
- [24] S.W. Graham and Jeffrey D. Vaaler. *A class of extremal functions for the Fourier transform*. Trans. Amer. Math. Soc., **265**(1) (1981), 283–302.
- [25] Glyn Harman. *Small fractional parts of additive forms*. Philos. Trans. Roy. Soc. London Ser. A, **345**(1676) (1993), 327–338.
- [26] Glyn Harman. *Metric number theory, volume 18 of London Mathematical Society Monographs. New Series*. The Clarendon Press Oxford University Press, New York (1998).
- [27] Alan Haynes, Michael Kelly and Barak Weiss. *Equivalence relations on separated nets arising from linear toral flows*. Proc. London Math. Soc. (to appear), November (2014).
- [28] Jeffrey J. Holt and Jeffrey D. Vaaler. *The Beurling-Selberg extremal functions for a ball in Euclidean space*. Duke Math. J., **83**(1) (1996), 202–248.
- [29] Lars Hörmander. *The analysis of linear partial differential operators. I*, volume 256 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1983. Distribution theory and Fourier analysis.
- [30] Michael Kelly and Gabriele Bianchi. *The Blaschke-Santaló Inequality*. Preprint (2014).
- [31] Michael Kelly and Thái Hoàng Lê. *Uniform dilations in higher dimensions*. Journal of the London Mathematical Society, **88**(3) (2013), 925–940.
- [32] M.G. Krein. *On the best approximation of continuous differentiable functions on the whole real axis*. Dokl. Akad. Nauk SSSR (Russian), **18** (1938), 615–624.
- [33] Xian-Jin Li and Jeffrey D. Vaaler. *Some trigonometric extremal functions and the Erdős-Turán type inequalities*. Indiana Univ. Math. J., **48**(1) (1999), 183–236.
- [34] Friedrich Littmann. *Entire approximations to the truncated powers*. Constr. Approx., **22**(2) (2005), 273–295.

- [35] Friedrich Littmann. *One-sided approximation by entire functions*. J. Approx. Theory, **141**(1) (2006), 1–7.
- [36] Friedrich Littmann. *Quadrature and extremal bandlimited functions*. SIAM J. Math. Anal., **45**(2) (2013), 732–747.
- [37] B.F. Logan. *Bandlimited functions bounded below over an interval*. Notices of the Amer. Math. Soc., **24** (1977).
- [38] H.L. Montgomery and R.C. Vaughan. *Hilbert's inequality*. J. London Math. Soc. (2), **8** (1974), 73–82.
- [39] Hugh L. Montgomery. *The analytic principle of the large sieve*. Bull. Amer. Math. Soc., **84**(4) (1978), 547–567.
- [40] Hugh L. Montgomery. *Ten lectures on the interface between analytic number theory and harmonic analysis*, volume 84 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI (1994).
- [41] M. Plancherel and G. Pólya. *Fonctions entières et intégrales de fourier multiples*. Comment. Math. Helv., **10**(1) (1937), 110–163.
- [42] Marvin Rosenblum and James Rovnyak. *Topics in Hardy classes and univalent functions*. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser Verlag, Basel (1994).
- [43] I.J. Schoenberg. *Metric spaces and completely monotone functions*. Ann. of Math. (2), **39**(4) (1938), 811–841.
- [44] Atle Selberg. *Remarks on sieves*. In *Proceedings of the Number Theory Conference (Univ. Colorado, Boulder, Colo., 1972)*, pages 205–216. Univ. Colorado, Boulder, Colo., (1972).
- [45] Atle Selberg. *Collected papers*. Vol. II. Springer-Verlag, Berlin, 1991. With a foreword by K. Chandrasekharan.
- [46] E.M. Stein. *Functions of exponential type*. Ann. of Math. (2), **65** (1957), 582–592.
- [47] Elias M. Stein. *Singular integrals and differentiability properties of functions*. Princeton Mathematical Series, No. 30. Princeton University Press, Princeton, N.J., (1970).
- [48] Elias M. Stein and Rami Shakarchi. *Complex analysis*. Princeton Lectures in Analysis, II. Princeton University Press, Princeton, N.J., (2003).
- [49] Elias M. Stein and Guido Weiss. *Introduction to Fourier analysis on Euclidean spaces*. Princeton University Press, Princeton, N.J., 1971. Princeton Mathematical Series, No. 32.
- [50] Jeffrey Vaaler. Personal Communication. (2012).
- [51] Jeffrey D. Vaaler. *Some extremal functions in Fourier analysis*. Bull. Amer. Math. Soc. (N.S.), **12**(2) (1985), 183–216.
- [52] A. Zygmund. *Trigonometric series. 2nd ed. Vols. I, II*. Cambridge University Press, New York (1959).

Felipe Gonçalves

IMPA – Instituto de Matemática Pura e Aplicada
Estrada Dona Castorina, 110
22460-320 Rio de Janeiro, RJ
BRAZIL

E-mail: ffgoncalves@impa.br

Michael Kelly

Department of Mathematics
Univ. of Texas
1 University Sta. C1200 Austin
Texas 78712
U.S.A.

E-mail: mkelly@math.utexas.edu

Jose Madrid

IMPA – Instituto de Matemática Pura e Aplicada
Estrada Dona Castorina, 110
22460-320 Rio de Janeiro, RJ
BRAZIL

E-mail: josermp@impa.br