



Rehabilitation to improve outcomes of lumbar fusion surgery: a systematic review with meta-analysis

Liedewij Bogaert^{1,2} · Tinne Thys¹ · Bart Depreitere³ · Wim Dankaerts⁴ · Charlotte Amerijckx² · Peter Van Wambeke¹ · Karel Jacobs^{5,7} · Helena Boonen⁴ · Simon Brumagne^{1,4} · Lieven Moke^{5,6} · Sebastiaan Schelfaut^{5,6} · Ann Spriet¹ · Koen Peers¹ · Thijs Willem Swinnen⁴ · Lotte Janssens²

Received: 23 August 2021 / Accepted: 16 February 2022 / Published online: 8 March 2022
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2022

Abstract

Purpose To evaluate the effectiveness of rehabilitation strategies on disability, pain, pain-related fear, and return-to-work in patients undergoing lumbar fusion surgery for degenerative conditions or adult isthmic spondylolisthesis.

Methods Six electronic databases were systematically searched for randomized controlled trials (RCTs) evaluating the effect of rehabilitation (unimodal or multimodal). The estimated effect size was calculated for interventions with homogeneous content using a random-effects model. Certainty of evidence was assessed by GRADE.

Results In total, 18 RCTs, including 1402 unique patients, compared specific rehabilitation to other rehabilitation strategies or usual care. Most described indications were degenerative disc disease and spondylolisthesis. All rehabilitation interventions were delivered in the postoperative period, and six of them also included a preoperative component. Intervention dose and intensity varied between studies (ranging from one session to daily sessions for one month). Usual care consisted mostly of information and postoperative mobilization. At short term, low quality of evidence shows that exercise therapy was more effective for reducing disability and pain than usual care (standardized mean difference [95% CI]: -0.41 [-0.71; -0.10] and -0.36 [-0.65; -0.08], four and five studies, respectively). Multimodal rehabilitation consisted mostly of exercise therapy combined with cognitive behavioral training, and was more effective in reducing disability and pain-related fear than exercise therapy alone (-0.31 [-0.49; -0.13] and -0.64 [-1.11; -0.17], six and four studies, respectively). Effects disappeared beyond one year. Rehabilitation showed a positive tendency towards a higher return-to-work rate (pooled relative risk [95% CI]: 1.30 [0.99; 1.69], four studies).

Conclusion There is low-quality evidence showing that both exercise therapy and multimodal rehabilitation are effective for improving outcomes up to six months after lumbar fusion, with multimodal rehabilitation providing additional benefits over exercise alone in reducing disability and pain-related fear. Additional high-quality studies are needed to demonstrate the effectiveness of rehabilitation strategies in the long term and for work-related outcomes.

Keywords Rehabilitation · Spinal fusion · Exercise · Physical therapy · Cognitive behavioral therapy · Return to work

Thijs Willem Swinnen and Lotte Janssens are shared last authors.

✉ Liedewij Bogaert
Liedewij.bogaert@uzleuven.be

¹ Department of Physical and Rehabilitation Medicine, University Hospitals Leuven, Leuven, Belgium

² Reval Rehabilitation Research Center, Hasselt University, Diepenbeek, Belgium

³ Department of Neurosurgery, University Hospitals Leuven, Leuven, Belgium

⁴ Department of Rehabilitation Sciences, KU Leuven, Leuven, Belgium

⁵ Institute for Orthopaedic Research and Training (IORT), Department of Development and Regeneration, KU Leuven, Leuven, Belgium

⁶ Department of Orthopaedics, University Hospitals Leuven, Leuven, Belgium

⁷ Department of Public Health and Primary Care, Leuven Institute for Health Care Policy, KU Leuven, Leuven, Belgium

Introduction

Internationally, a rising trend in the number of spinal fusion procedures is observed [1, 2]. Over the years, technical advancements have been translated into higher *radiographic* success rates of bony fusion and sagittal alignment [3, 4]. In contrast, the *clinical* success rate remains only modest with up to 40% of patients reporting persistent pain, sub-optimal functional improvement and dissatisfaction [5–8], and a work resumption in only half of the patients below normal retirement age [7]. Therefore, an urgent need exists to optimize clinical outcomes after lumbar fusion.

Rehabilitation has been put forward as a window of opportunity to enhance the value of spine care [9, 10]. However, the golden standard of rehabilitation for lumbar fusion remains largely unclear. This is reflected by extensive variation in everyday practice. For example, no consensus regarding timing and content of rehabilitation was found between surgeons in the Netherlands and Sweden [11]. This considerable variability in physiotherapy practice was also demonstrated in Australia and the United Kingdom [12, 13].

The shortcomings of previous reviews in this field are summarized below: *firstly*, previous reviews were focused on either the pre- or postoperative period but not on the entire care continuum [9, 10, 14, 15]; *secondly*, extrapolated or included evidence from other types of lumbar surgery [10, 15]; and/or *thirdly*, were out-of-dated [9, 14]. Hence, an updated review and meta-analysis of the effectiveness of rehabilitation strategies for lumbar fusion across the entire care continuum was warranted.

Therefore, the primary aim of this systematic review and meta-analysis was to assess and compare the effectiveness of unimodal and multimodal rehabilitation strategies on disability, pain, and pain-related fear in patients undergoing lumbar fusion surgery for degenerative conditions and (adult) isthmic spondylolisthesis. The secondary aim was to assess the effectiveness on return-to-work.

Methods

This systematic review followed the methods of the Cochrane Handbook for Systematic Reviews of Interventions [16], and is reported in line with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement [17]. The protocol has been prospectively registered in PROSPERO (CRD42018083422).

Search strategy

Our search strategy included *lumbar, fusion, rehabilitation, randomized controlled trial* and related terms. To optimize

the sensitivity of the search, no terms related to relevant outcomes were applied. This search strategy was developed in conjunction with a research librarian, peer reviewed for completeness within our team (LB, TT, TWS, LJ), and validated by testing whether it could identify eight relevant studies in PubMed and Embase. The full search strategy is outlined in Appendix A. We searched PubMed, Embase, Web of Science, Pedro, Cinahl and Cochrane Library from inception until April 28, 2021. To identify ongoing research, Clinicaltrials.gov was additionally searched. Thereafter, we scanned references of identified articles and relevant reviews. Our search output was managed in EndNote X9, which facilitated removal of duplicates in a stepwise manner [18]. After deduplication, two reviewers with complementary methodological and clinical expertise (LB, TT) independently screened titles and abstracts (phase 1) and full texts (phase 2) using blinded Rayyan software [19]. In case of disagreement, consensus was obtained after each phase by discussion and, if necessary, mediation by a third reviewer (LJ).

Eligibility criteria

RCTs investigating the effect of specified rehabilitation in the pre-, peri- and/or postoperative period of lumbar fusion on disability, pain and/or pain-related fear were eligible for inclusion (Table 1). Outcomes were narrowed from our registered protocol, representing most of the components of the International Classification of Functioning, Disability and Health (ICF) framework: pain (function), disability (activities), return-to-work (participation) and pain-related fear (personal factors), except for environmental factors as an a priori exploratory search indicated that these were not reported in this context. A pilot test was used to ensure that the eligibility criteria were applied consistently between the reviewers.

Risk of bias

The quality of the included RCTs was independently assessed as ‘low’, ‘uncertain’ or ‘high’ risk of bias by two reviewers (LB, TT), using the Cochrane Collaboration Revised Risk of Bias tool for RCTs (RoB 2.0, version 22 August 2019, facilitated by Cochrane RoB 2: Learning Live series) [20, 21]. Given the nature of rehabilitation interventions, blinding of participants was not feasible. Therefore, this domain was not considered in the overall summary risk of bias judgment, which is in line with previous reviews of rehabilitation interventions [22].

Data extraction and synthesis

Data extraction was completed by two reviewers (LB, CA), using a predefined extraction form based on the TIDieR

Table 1 Eligibility criteria for inclusion

Patients	Aged over 18 years, undergoing lumbar fusion surgery for degenerative conditions and (adult) isthmic spondylolisthesis (without restrictions on fusion technique, limited to single-, double-, and three-level fusion)
Intervention	Rehabilitation as defined by the World Health Organization. This encompasses interventions with a physical, psychological, social, or occupational dimension (i.e., unimodal rehabilitation interventions); and multimodal rehabilitation (i.e., simultaneous or sequential application of different dimensions). Both preoperative (as from the decision to perform lumbar fusion surgery) and postoperative (until six months after surgery) starting points of rehabilitation were eligible for inclusion
Comparator	No treatment (or placebo), usual care or other rehabilitation strategies
Outcomes	Studies concerning disability, pain, pain-related fear or working rate. A validated outcome measure of disability and pain was eligible. Pain-related fear was defined for this review as fear-avoidance behavior and beliefs related to low back pain, targeted to Tampa Scale of Kinesiophobia (TSK) or Fear-Avoidance Beliefs Questionnaire (FABQ)
Other	Studies were excluded if only an abstract was available, and further data could not be obtained on request (after two attempts by mail to contact the corresponding author). Language was restricted to English and Dutch. No restriction of publication date was applied

checklist (for details, see Table 2) [23]. Consistent data extraction by the two extracting authors was ensured by piloting the extraction form (on two articles).

Primary outcomes were patient-reported disability, pain, and pain-related fear at short term (≤ 6 months postoperatively) and/or long term (≥ 1 year postoperatively). *Secondary outcome* was return-to-work at short- and/or long term. If studies reported multiple follow-up moments, data closest to three months and one year postoperatively were used for meta-analyses for short term and long term, respectively.

Across all outcomes, random-effects meta-analyses were conducted of studies that were sufficiently homogeneous in terms of the rehabilitation procedure, procedure of the comparator and outcome measurement (by LB, TWS, LJ). Effect estimates were reported as relative risks (RR) and 95% confidence interval (95% CI) for dichotomous outcomes and standardized mean difference (SMD) with 95% CI for continuous outcomes. A SMD was applied, since different valid measurement scales of the same continuous outcomes were used across studies (e.g., for pain). Based on Cohen's interpretation of effect size, a SMD of ≥ 0.2 , ≥ 0.5 and ≥ 0.8 represents a small, moderate, and large effect, respectively. Post-rehabilitation measurements were used for effect size estimation as these yields more precise analysis for the included trials than change from baseline measurements (i.e., correlation coefficient of change scores was less than 0.5) [16]. Inverse variance weighting was used for pooling, which gives studies with more precise results (narrower confidence intervals) more weight. If sample mean and standard deviation could not be retrieved upon request from the corresponding authors, sample mean and standard deviation were estimated from reported CI; or from median and range. If multiple randomized arms were included in one RCT, each comparison was separately included but with the shared control group divided evenly among the comparisons [16]. Outliers were defined as studies in which the 95% CI of the studies effect size was outside the 95% CI of the pooled effect size. In case an outlier was detected, a sensitivity

analysis by pooling the effect size again, this time excluding the identified outlier, was conducted. Statistical heterogeneity among the included studies was considered by calculation of I^2 statistics, with 75% as boundary for high heterogeneity. High statistical heterogeneity did not preclude meta-analysis, but it downgraded ratings of the quality of evidence. Exploration of publication bias could not be visualized in funnel plots, since less than ten studies were included in our meta-analyses. All statistical analyses and visualizations of data were performed in R software (version 4.0.3), using meta package [24–26].

Certainty of evidence

The certainty of evidence was evaluated for each pooled estimate according to the GRADE system, as high, moderate, low, or very low [27]. The GRADE profile was downrated from high quality by one level for each of the following limitations: low methodological quality, inconsistency, indirectness, imprecision, or publication bias (operational rules are outlined in Table 3).

Results

A total of 4425 records were identified through electronic database searching (Fig. 1). After removal of duplicates, 2085 titles and abstracts were screened; and subsequently 86 full-text articles were reviewed for eligibility. Finally, 21 articles, reporting data from 18 different RCTs were included, with a total of 1402 participants (mean age 43–61 years, 57% female). Indications and fusion techniques varied across and within studies. Most described indications for lumbar fusion surgery were degenerative disc disease (39%) and spondylolisthesis (25%) (Appendix B). All articles were published in 2003 or later, and the trials were conducted in Europe ($n = 15$), Asia ($n = 2$) or Africa ($n = 1$).

Table 2 provides an overview of the data extraction. The 18 included trials investigated 21 different rehabilitation interventions in total.

Table 2 Description of study characteristics and key intervention items according to TIDieR guide

Author, year (Country)	No. of participants (% female, Mean age ± SD)	Fusion technique (primary diagnosis)	Intervention condition (TIDieR items)			When and how much	Compliance (% of patients)	Control condition TIDieR items restrictions†	Outcome assessment Follow-up points and outcome measurement of interest	
			Procedure	Who provided	Mode of delivery; individual/ in group					Where
<i>Multimodal rehabilitation intervention vs exercise therapy</i>										
Rolving et al. [28], 2015 (Denmark)	90 (56,7% female; 50.1 ± 9.2y)	PLF, TLIF, uninstrumented fusion (DDD, spondylolisthesis grades I to II)	CBT covering interaction of cognition and pain perception, coping strategies, pacing principles, ergonomic directions, return to work and details about the surgical procedure (in addition to control condition)	Multi-disciplinary	Face to face, group	Hospital	4 pre- and 2 postoperative sessions (3 h)	83%	Preoperative information, postoperative physiotherapist-supervised exercises in group or individual, starting 3 m after surgery †Information contains physical restrictions after surgery NOS	1w: NRS 12 and 52w: ODI, LBPRS (leg, back), FABQ-PA, work status
Abbott et al. [29], 2010 (Sweden)	107 (61,7% female; 50.7 ± 10.4y)	PLF, TLIF, uninstrumented fusion (DDD, spinal stenosis, degenerative / isthmic spondylolisthesis, spondylolysis)	Psychomotor therapy including CBT and graded motor relearning therapy (lumbopelvic stabilization) (in addition to control condition)	Physiotherapist (trained in behavioral medicine)	Face to face, individual	Outpatient, home	Three sessions (90 min) between 0-3 m postoperative	94%	Home-based exercise program (dynamic exercises, stretches, cardiovascular), 0-3 m postoperative †No running, contact sports, heavy lifting or outer-range lumbar spine movements until 6 m	3 m, 6 m, 1y, 2y, 3y: ODI, VAS, TSK, work status

Table 2 (continued)

Author, year (Country)	No. of participants (% female, Mean age ± SD)	Fusion technique (primary diagnosis)	Intervention condition (TIDieR items)				Control condition TIDieR items restrictions [†]	Outcome assessment Follow-up points and outcome measurement of interest		
			Procedure	Who provided	Mode of delivery; individual/ in group	Where			When and how much	Compliance (% of patients)
Ilves et al. [30], 2016 (Finland)	98 (73.5% female; 58.9 ± 9.5y)	PLF, PLIF, TLIF (degenerative/ isthmia spondylolisthesis)	Progressive back-specific exercise (control, coordination, strength, and endurance of back, abdominal, gluteal and thigh muscles) and aerobic training (walking sessions) and fear-avoidance counseling (identifying barriers to physical activity, correction of harmful, irrational beliefs and fears towards activity, goal-setting)	Physiotherapist	Face to face, individual	Unknown	6 sessions between 3–15 m postoperative	Unknown	Home-based exercises for endurance (abdominal, back and hip muscles), stretching and balance with one physiotherapist guiding session 3 m postoperative	3, 15, 27 m: ODI, VAS (leg, back), TSK,
Greenwood et al. [31], 2019 (United Kingdom)	52 (67.3% female; 54.2 ± 13.5y)	Unknown (DDD, spinal stenosis, isthmia spondylolisthesis)	Peer support, education, exercise training (cardiovascular, limb and spine strengthening)	Physiotherapist	Face to face, group	Unknown	10 sessions, starting 3 m postoperative	95%	Referral to external physiotherapy [†] No lifting heavier than a full kettle until 3 m	3, 6, 12 m: ODI

Table 2 (continued)

Author, year (Country)	No. of participants (% female, Mean age ± SD)	Fusion technique (primary diagnosis)	Intervention condition (TIDieR items)				When and how much	Compliance (% of patients)	Control condition TIDieR items restrictions [†]	Outcome assessment Follow-up points and outcome measurement of interest
			Procedure	Who provided	Mode of delivery; individual/ in group	Where				
Monticone et al. [32], 2013 (Italy)	130 (60.8% female; 57.3 ± 13.1y)	Unknown (spinal stenosis, degenerative/isthmic spondylolisthesis)	CBT (controlling catastrophizing, kinesiophobia and maladaptive behavior) (in addition to control condition)	Multi-disciplinary	Face to face, individual	Hospital	8 postoperative sessions (1 h) in 4w	Unknown	Hospital-based, physiotherapist-supervised exercise program, 5 times/w during 4w with active spinal mobilization and exercise training (strengthening deep spinal muscles, postural control, stretching, walking, changing of positions)	1 m, 1y post-treatment: ODI, TSK, NRS
Lotzke et al. [33], 2019 (Sweden)	118 (53.4% female; 45.7 ± 8.3y)	PLF, ALIF, PLIF (disc herniation, foraminal stenosis, isthmic spondylolisthesis)	CBT-based prehabilitation (targeting psychological risk factors and promoting physical activity)	Physiotherapist (experienced and graduated in CBT)	Face to face (pre-), telephone (postoperative), individual	Spine clinic, telephone	4 pre- (1 h) and 1 postoperative (30 min) session	Unknown	Referral to local physiotherapist (i.e., one preoperative session with core exercise program, information and advice to stay active)	1w pre- and 3, 8, 12, 26w post-operative: ODI, VAS (leg, back), TSK
Strom et al. [34], 2019 (Denmark)	114 (65% female; 54y (range 29–79y))	PLF, TLIF (spondylolisthesis, degenerative disease)	Preoperative access to w-SPIINA (web-based information/animation/peer support, in line with CBT principles) (in addition to control condition)	Web-based (including peer support of 6 former patients)	Online, individual and group features	Online platform	All time access	100% (48% active users)	Preoperative information session (2 h). Physiotherapist-supervised exercise program, starting 3 m postoperative	2d, 3 m, 6 m: HADS, ODI, EQ5D, LBPRS

Table 2 (continued)

Author, year (Country)	No. of participants (% female, Mean age ± SD)	Fusion technique (primary diagnosis)	Intervention condition (TIDieR items)				When and how much	Compliance (% of patients)	Control condition TIDieR items restrictions [†]	Outcome assessment Follow-up points and outcome measurement of interest
			Procedure	Who provided	Mode of delivery; individual/ in group	Where				
Oestergaard et al. [35], 2020 (Denmark)	82 (50% female; 46.8 ± 8.8y)	PLF, TLIF (DDD, isthmic spondylolisthesis grades I to II)	Case-manager assisted rehabilitation program (preoperative meeting to determine rehabilitation program, postoperative (phone-)meeting and possibility for roundtable-meeting and workplace visit) (in addition to control condition)	Multi-disciplinary (with trained case managers)	Face to face, telephone, individual	Hospital, rehabilitation unit, possible workplace or municipality visit, telephone	1 pre- and voluntary multiple postoperative sessions	100% (59% did sent a summary to the municipality)	Rehabilitation-unit-based, physiotherapist-supervised, exercise program, 1–2 times/w during 8–10w, starting at 8w after surgery	3, 6, 9, 12, 24 m: ODI, VAS (leg, back), work status
<i>Multimodal rehabilitation intervention: timing</i>										
Oestergaard et al. [36], 2012 (Denmark)	82 (53,7% female; 51.7 ± 9.2y)	PLF, TLIF (DDD, isthmic spondylolisthesis grades I to II)	Peer and psychological support, home exercise program (stability of trunci and large muscle groups), occupational advice	Multi-disciplinary	Face to face, group	rehabilitation unit	4 sessions (2 h), starting 6w postoperative	Unknown	Rehabilitation starting at 12w instead of 6w (same procedure, provider, mode of delivery and place as the intervention group)	6w, 3, 6, 12 m: ODI, LBPRS (leg, back), sick leave

Table 2 (continued)

Author, year (Country)	No. of participants (% female, Mean age \pm SD)	Fusion technique (primary diagnosis)	Intervention condition (TIDieR items)				Control condition TIDieR items restrictions [†]	Outcome assessment Follow-up points and outcome measurement of interest		
			Procedure	Who provided	Mode of delivery; individual/ in group	Where			When and how much	Compliance (% of patients)
<i>Exercise therapy vs usual care</i>										
Christensen et al. [37], 2003 (Denmark)	90 (66.7% female; median 47y)	PLF, Circumferential fusion (isthmic spondylolisthesis grades I to II, degeneration)	Group II: Exercise training (conditioning training, dynamic muscular endurance training (back/abdominal/leg muscles) and stretching)	Physiotherapist	Face to face, group	Hospital's physical therapy section	16 postoperative sessions (90 min) during 8w	Unknown	Video-recorded demonstration, with one-time oral instruction, of exercises (dynamic muscular endurance training of back/abdominal/leg muscles) [†] No contact sports, training on machines at a fitness center, jogging/running until 3 m	3, 6, 12, 24 m: LBPRS (leg, back, physical and emotional components), work status
Kang et al. [38], 2012 (South-Korea)	60 (48.3% female; 60.6 \pm 9.7y)	PLIF (unknown)	Group I: Extension exercise group: exercises using MedX device, McKenzie extension exercises Group II: Lumbar stability exercise group: transverse abdominis and multifidus co-contraction method	Physiotherapist	Face to face, group	Unknown	24 sessions (30 min) during 8w, starting at 3 m postoperative	Unknown	William and McKenzie exercise program	Posttreatment: ODI, VAS

Table 2 (continued)

Author, year (Country)	No. of participants (% female, Mean age ± SD)	Fusion technique (primary diagnosis)	Intervention condition (TIDieR items)				When and how much	Compliance (% of patients)	Control condition TIDieR items restrictions [†]	Outcome assessment Follow-up points and outcome measurement of interest
			Procedure	Who provided	Mode of delivery; individual/ in group	Where				
Nielsen et al. [39], 2010 (Denmark)	60 (58.9% female; 50.1y)	PLF, uninstrumented fusion (DDD)	Home-based exercise program (strengthening back and abdomen muscle, cardiovascular), information and advice smoking and alcohol cessation. Post-operative early mobilization (aim discharge on day 5) and exercise program	Physiotherapist	Face to face, telephone, individual	Hospital, home	Unknown	Preoperative information. Postoperative mobilization once/day (aim discharge on day 8)	2 m pre-, admission, discharge, 1, 3, 6 m post-operative: RMQ, VAS (leg, back)	
Kernc et al. [40], 2018 (Slovenia)	27 (48.1% female; 60.7 ± 7.9y)	TLIF (DDD, degenerative/ ischmic spondylolisthesis)	Strength training (focus on lumbopelvic stabilization muscles, IAP utilization)	Physiotherapist	Face to face, unknown	Unknown	Unknown	No exercises prior to 3 m postoperative	3w, 3, 18 m: ODI, VAS, IAP activation, performance	
Sengul et al. [41], 2020 (Turkey)	37 (50.2% female; 53.9y)	PLF (spinal stenosis, degenerative/ isthmic spondylolisthesis, DDD with instability, spondylosis)	Home-based exercise program, each time preceded by motor imagery exercises, facilitated by voice commands	Physiotherapist	Sound records, telephone individual	Home, Follow-up by telephone	100%	Home-based exercise program (education, neutral spine control, maximal voluntary isometric contraction), starting 2d postoperative [†] Education on how to use a corset	3, 6w: VAS (back), ODI, TSK,	

Table 2 (continued)

Author, year (Country)	No. of participants (% female, Mean age ± SD)	Fusion technique (primary diagnosis)	Intervention condition (TIDieR items)			When and how much	Compliance (%) of patients	Control condition TIDieR items restrictions†	Outcome assessment Follow-up points and outcome measurement of interest	
			Procedure	Who provided	Mode of delivery; individual/ in group					Where
<i>Other physiotherapeutic interventions vs usual care</i>										
Elsayyad et al. [42], 2020 (Egypt)	60 (41.6% female; 42.9 ± 4.1y)	Unknown (DDD ± spinal stenosis)	Group I: Neural mobilization (longitudinal traction along the sciatic nerve; 12 to 15 min per session) (in addition to control condition) Group II: Myofascial release (thoracolumbar fascia, quadratus lumborum, piriformis, hamstring muscles) (in addition to control condition)	Physiotherapist	Face to face, individual	Hospital	12 sessions for 4w	Unknown	Hospital-based, physiotherapist-supervised, stabilization exercise program, 12 sessions for 4w	1, 2 m: ODI, VAS
Zhao et al. [43], 2008 (China)	69 (43.5% female; 52.4 ± 8.2y)	Unknown (disk prolapse)	Electroacupuncture and cupping	Unknown	Face to face, individual	Unknown	Daily session (30 min) electroacupuncture, every other day cupping. Min 6 courses of 15d with 5d interval	Unknown	Deaquation and nerve nutrition medication 6w †bedrest	3, 6, 12 m: JOA
<i>Occupational therapy vs usual care</i>										
Oestergaard et al. [44], 2012 (Denmark)	87 (65.5% female; 54.6y)	PLF, ALIF, TLIF, uninstrumented fusion (DDD, spondylosis, instability)	Semi-structured interview using COPM during inpatient rehabilitation to identify ADL-problems and set rehabilitation goals	Occupational therapist	Face to face, individual	Hospital	1 session during hospitalization	Unknown	Usual instructions for ADL	1, 4, 12w and 3y: DPQ

Table 2 (continued)

Author, year (Country)	No. of participants (% female, Mean age ± SD)	Fusion technique (primary diagnosis)	Intervention condition (TIDieR items)			When and how much	Compliance (% of patients)	Control condition TIDieR items restrictions [†]	Outcome assessment Follow-up points and outcome measurement of interest
			Procedure	Who provided	Mode of delivery; individual/ in group				
<i>Psychological intervention vs usual care</i>									
Reichart et al. [45], 2011 (Germany)	39 (56.4% female; 59.1y)	PLIF (spinal stenosis and instability)	Shortpsychological intervention focusing on reducing fear-avoidance and motivational strategies	Psychotherapist (trained in behavioral therapy)	Face to face, individual	Unknown	1 pre- and 1 post-operative session (30 min)	Unknown	1d, 6w: VAS, FABQ
<i>Peer support vs usual care</i>									
Christensen et al. [37], 2003 (Denmark)	— (details provided above)	—	Group 1: Back-café with peer support of other patients (exchange of experiences, doubts, tips) (in addition to control condition)	Physiotherapist	Face to face, group	Hospital's physical therapy section	8 post-operative sessions (90 min) during 8w	Unknown	—

PLF posterior/posterolateral lumbar fusion; *PLIF* posterior lumbar interbody fusion; *TLIF* transforaminal lumbar interbody fusion; *ALIF* anterior lumbar interbody fusion and lateral lumbar interbody fusion; *DDD* degenerative disc disease; *min* minutes; *h* hours; *w* weeks; *m* months; *y* years; *CBT* cognitive behavioral therapy; *ODI* Oswestri Disability Index; *NRS* Numeric Rating Scale; *VAS* Visual Analog Scale; *LBPRS* Low Back Pain Rating Scale; *DPQ* Dallas Pain Questionnaire; *FABQ* Fear-avoidance beliefs questionnaire; *RMQ* Rolland-Morris Questionnaire; *JOA* Japanese Orthopedics Association

As Fig. 2 shows, rehabilitation was either initiated preoperatively ($n = 6$); postoperatively within three months ($n = 8$), from three months ($n = 6$), or unspecified ($n = 1$), yet all rehabilitation interventions included a postoperative rehabilitation component. Ten trials provided follow-up beyond one year.

Nine interventions consisted of multimodal rehabilitation [28–36]. Of these, eight compared this multimodal rehabilitation to exercise therapy alone, and were included for meta-analyses [28–35]. All multimodal rehabilitation interventions featured exercise training, most often combined with CBT ($n = 5$), with fear-avoidance counseling ($n = 1$), case manager guidance ($n = 1$) or education and peer support ($n = 1$). Despite the multimodal nature of those interventions, these interventions were mostly provided by physiotherapists ($n = 5$) rather than by a multidisciplinary rehabilitation team ($n = 3$).

On the other hand, six interventions including exercise therapy alone were compared to usual care, and were included for meta-analyses [37–41]. Although they shared similar durations of at least six weeks, the exercise methods varied. All exercise interventions used strength training, and

in two studies this was combined with cardiovascular conditioning [37, 39].

Due to heterogeneity in the content of the remaining six unimodal interventions (i.e., occupational therapy, psychological therapy, peer support and three different types of passive physiotherapy), no inclusion for meta-analysis was possible [37, 42–45].

Risk of bias

As shown in Fig. 3, the overall bias was scored unclear ($n = 13$; 72%) or high ($n = 5$; 28%). A high proportion of studies had an unclear or high risk for selective outcome reporting. This was mainly explained by a lack of registered protocols in the majority of RCTs ($n = 11$; 61%).

Certainty of evidence and sensitivity analysis

A summary of pooled effect sizes and GRADE quality ratings are provided in Table 3. A sensitivity analysis showed that one outlier in the meta-analysis of the effect of multimodal rehabilitation on disability and pain, Monticone

Table 3 Overview of estimated effect of rehabilitation interventions according to their content and GRADE assessment

Outcomes	No. of participants (No. of RCTs)	Certainty of evidence (GRADE)	SMD* (95% CI)	SMD with exclusion of outlier**
<i>Multimodal rehabilitation vs exercise</i>				
At short-term follow-up				
Disability	620 (7)	Low ^{a,b}	−0.78 [−1.55; −0.01]	−0.31 [−0.49; −0.13]
Pain	580 (6)	Low ^{a,b}	−0.43 [−0.89; 0.03]	−0.23 [−0.51; 0.04]
Fear-avoidance	412 (4)	Low ^{a,b}	−0.64 [−1.11; −0.17]	
At long-term follow-up				
Disability	524 (6)	Low ^{a,b}	−0.49 [−1.16; 0.18]	−0.18 [−0.49; 0.14]
Pain	480 (5)	Low ^{a,b}	−0.61 [−1.49; 0.26]	−0.16 [−0.37; 0.05]
Fear-avoidance	409 (4)	Low ^{a,b}	−0.85 [−1.74; 0.04]	
<i>Exercise vs usual care</i>				
At short-term follow-up				
Disability	180 (4)	Low ^{a,d}	−0.41 [−0.71; −0.10]	
Pain	235 (5)	Low ^{a,d}	−0.36 [−0.65; −0.08]	
At long-term follow-up				
Pain	82 (2)	Low ^{a,d}	−0.10 [−0.53; 0.34]	

RCT randomized controlled trial; No. number; GRADE Grading of Recommendations, Assessment, Development and Evaluations; SMD standardized mean difference; CI confidence interval

*SMD with 95%CI is used to express the estimated effect since different scales are used to measure the same outcome. A SMD of 0.2, 0.5 and 0.8 represents a small, moderate, and large effect respectively

**Monticone et al. [32] was identified as outlier and contributed to high statistical heterogeneity (I^2 without exclusion: 95% for disability at short term, 86% for pain at short term, 93% for disability at long term and 95% for pain at long-term follow-up)

^aLow methodological quality: more than 75% of patients out of studies with unclear or high risk of bias

^bInconsistency: statistical heterogeneity measured as I^2 is more than 75%, or outlier detected in sensitivity analysis

^cInconsistency: wide variation in the effect estimates across studies (i.e. minimal or no overlap of CI)

^dImprecision: magnitude of the sample is less than 400 (recommended informative sample size for continuous outcomes). No downgrade was done for publication bias (since the comprehensive search and no industry influence), neither for indirectness (since inclusion of a specific population, relevant outcomes, and predefined comparisons)

et al., highly contributed to heterogeneity and possibly lead to an overestimation of the effect size. This could partly be explained by clinical variation between the intervention of Monticone et al. and other multimodal rehabilitation interventions (i.e. more dose-intense rehabilitation program, less well described population). Therefore, this outlier was excluded [32], leading to a decrease in pooled effect size and a reduction from high to low heterogeneity.

Effects on disability and pain (primary outcomes)

Effects on disability were reported for 13 interventions (five exercise and eight multimodal interventions), using the Oswestry Disability Index (ODI) [28–30, 32–36, 38, 40–42] or the Roland Morris Disability Questionnaire (RMDQ) [39]. Effectiveness on pain was measured with Visual Analog Scale (VAS) [29, 30, 33, 35, 38–41, 45], Numeric Rating Scale (NRS) [32] or Low Back Pain Rating Scale (LBPRS) [28, 34, 36, 37], for six exercise and eight multimodal interventions.

Exercise vs usual care

There is low-quality evidence that an exercise intervention was more effective than usual care for reducing disability at short term (four trials with a total of five interventions and 180 participants, SMD with 95%CI: -0.41 [-0.71 ; -0.10]) (Fig. 4). Only one study with a high overall risk of bias investigated the long-term effect of exercise treatment on disability, and reported no significant differences between exercise and usual care (SMD with 95%CI: -0.10 [-0.85 ; 0.66]) [40].

Low-quality evidence from five studies (235 participants) indicated significantly more pain reduction after rehabilitation with an exercise component (SMD with 95%CI: -0.36 [-0.65 ; -0.08]). The pooled results of two studies (82 participants) provided low-quality evidence for no difference on the long term (SMD with 95%CI: -0.10 [-0.53 ; 0.34]).

Multimodal rehabilitation vs exercise

Participants who received a multimodal rehabilitation intervention ($n = 255$), which was in more than half of the patients initiated preoperatively, showed less disability at short-term follow-up than those who received only exercise therapy ($n = 235$) (SMD with 95%CI: -0.31 [-0.49 ; -0.13], low-quality evidence, six trials) (Fig. 5).

In the long term, the pooled result of five trials (including 394 participants) provided low-quality evidence for no significant effect on disability (SMD with 95%CI: -0.18 [-0.49 ; 0.14]).

For pain, low-quality evidence suggests no significant effect of multimodal rehabilitation compared to exercise

alone at both short term (SMD with 95%CI: -0.23 [-0.51 ; 0.04], five trials with 450 participants) and long-term follow-up (SMD with 95%CI: -0.16 [-0.37 ; 0.05], four trials with 350 participants) (Fig. 5).

Peer support, occupational therapy, psychological intervention, or passive physiotherapy vs usual care

Christensen et al. compared a postoperative ‘back café’ to usual care. There was no group difference in back pain at two-year follow-up, and whereas peer support improved the ability to raise a chair, carry a bag and take stairs, no superiority was reported for the other daily functions. [37] Also, occupational therapy guided by a questionnaire in the immediate postoperative period was not associated with better daily functioning performance [44]. In contrast, Reichart et al. demonstrated that participants receiving a short perioperative psychological intervention to increase their self-efficacy reported less pain and better functionality than those receiving usual care [45].

Two trials investigated the effectiveness of passive, postoperative physiotherapeutic interventions. More specifically, Elsayyad et al. [42] reported less disability and pain when myofascial release or neural mobilization (under the form of longitudinal traction) were added to stabilization exercises compared to stabilization exercises only. On the other hand, Zhao et al. [43] favored acupuncture to improve functioning over complete bedrest for six weeks, however not reaching the minimal clinically important difference (MCID) for the Japanese Orthopaedics Association (JOA) score. Due to this striking contrast in comparator between both RCTs, those interventions were excluded for meta-analysis.

Effects on pain-related fear (primary outcome)

The effects on pain-related fear were reported in seven studies including five multimodal, one psychological and one exercise alone intervention, using the Tampa Scale of Kinesiophobia (TSK) [29, 30, 32, 33, 41] or Fear-Avoidance Beliefs Questionnaire (FABQ) [28, 45].

Exercise vs usual care

One study of uncertain quality including 37 participants showed no significant difference in pain-related fear between exercise and usual care at six weeks postoperative (SMD with 95%CI: -0.25 [-0.90 ; 0.40]) and attributed this partly to the absence of a longer follow-up [41].

Multimodal rehabilitation vs exercise

Participants who received a multimodal intervention showed less pain-related fear at short term, compared to those who




Fig. 1 Study selection flowchart, according to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) diagram. *RCT* randomized controlled trial. *When multiple publications

reported data from the same RCT, the first publication was referred to as primary publication and any additional publications as companion reports. Companion reports without relevant outcomes were excluded

received exercise therapy alone (four RCTs with 412 participants; observed SMD with 95%CI ranging from -0.02 [$-0.40; 0.37$] to -1.10 [$-1.47; -0.73$], low-quality evidence). At long-term follow-up, however, no significant difference in pain-related fear was present between participants of the multimodal intervention or those of the exercise intervention (four RCTs, including 409 patients; observed SMD with 95%CI ranging from 0.00 [$-0.40; 0.40$] to -1.91 [$-2.33; -1.50$], low-quality evidence) (Fig. 6). Both estimates were imprecise owing to the low absolute sample sizes, as indicated by the width of the confidence interval. High statistical heterogeneity across trials was present, yet no outlier was detected, and an additional sensitivity analysis was not performed because of the low number of trials.

Psychological intervention vs usual care

At short-term follow-up, Reichart et al. described a trend towards an increase in fear-avoidance beliefs after usual care and a decrease after a psychological intervention ($p=0.11$).

This study was limited by an uncertain risk of bias, sample of 39 participants and a follow-up of only 6 weeks [45].

Effects on return-to-work (secondary outcome)

Four studies evaluated the efficacy of specific rehabilitation on return-to-work at long-term follow-up [28, 29, 35, 37]. Taken together, the estimated relative risk for return-to-work tends to favor rehabilitation modes of various content (i.e., peer support, occupational therapy, exercise, multimodal rehabilitation). However, this difference was not statistically significant (pooled RR with 95%CI: 1.30 [$0.99-1.69$]) (Fig. 7).

Discussion

The results of this systematic review and meta-analysis indicate that exercise is likely to reduce disability and pain up to six months after lumbar fusion. Moreover, multimodal rehabilitation combining exercise training with CBT, peer




Fig. 2 Overview of the timing, duration, intensity and outcomes of the included studies. Timing and duration are visualized by the number of weeks, intensity is indicated by the number of sessions (and duration per session in minutes)

support or counseling, is associated with a greater reduction in disability and pain-related fear than exercise alone. It is uncertain, however, which effects of exercise and multimodal rehabilitation persist in the long term and to what extent they remain beneficial. Also, since multimodal rehabilitation was compared to exercise, the magnitude of effect of multimodal rehabilitation compared to no rehabilitation remains unclear.

Exercise therapy reduces pain up to six months after lumbar fusion, when compared to usual care. ‘Usual care’ varied between studies but mostly consisted of providing information and postoperative mobilization. It is unclear if

exercise-induced hypoalgesia is the mechanism to explain the pain reduction. While in healthy persons, pain and pain sensitivity decreases during and shortly after exercise, the evidence of exercise-induced hypoalgesia in patients with chronic pain is less substantiated [46]. Multimodal rehabilitation has no additional effect on pain when compared to exercise in isolation. For disability, multimodal rehabilitation seems to be more effective than exercise alone at short-term follow-up.

Greenwood et al. included two RCTs in their meta-analysis and concluded that multimodal rehabilitation reduces

Author	Randomisation process	Deviations from the intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported result	Overall bias
Rolving et al, 2015	!	+	+	-	+	!
Abbott et al, 2010	+	+	!	-	!	!
Ilves et al, 2016	!	+	+	-	!	!
Greenwood et al, 2019	+	!	+	-	+	!
Monticone et al, 2013	+	+	+	-	!	!
Lotzke et al, 2019	!	+	+	-	+	!
Strom et al, 2019	+	-	-	-	+	-
Oestergaard et al, 2020	+	!	!	-	+	!
Oestergaard et al, 2012	+	-	!	-	!	-
Christensen et al, 2003	!	-	!	-	-	-
Kang et al, 2013	!	+	+	-	!	!
Nielsen et al, 2010	+	+	!	-	+	!
Kernc et al, 2018	!	-	-	-	!	-
Sengul et al, 2020	!	!	+	-	!	!
Elsayyad et al, 2020	+	+	+	-	!	!
Zhao et al, 2008	!	-	-	-	-	-
Oestergaard et al, 2013	+	+	!	-	!	!
Reichart et al, 2011	+	!	+	-	!	!

+ Low risk
! Some concerns
- High risk

Fig. 3 Risk of Bias assessment using the ROB2.0. Since blinding of participants is not feasible in rehabilitation interventions, thereby leading to high risk of outcome measurement, this was not considered in overall risk of bias assessment, as is outlined in the method section




Fig. 4 Forest plot for the meta-analysis of the effectiveness of exercise versus usual care for reducing disability and pain. All studies are ordered from most to least effective. Random-effects model was used. Negative effect sizes favor exercise therapy

disability and pain-related fear in both short and long-term follow-up. The current findings confirmed this beneficial effect of multimodal rehabilitation at short term [9]. In contrast, no significant benefit of multimodal rehabilitation in the long term was detected in our meta-analysis. Greenwood’s conclusion was skewed by inclusion of Monticone

et al., while the current review clearly identified this study as an outlier.

In patients undergoing lumbar surgery, greater fear of movement is associated with higher levels of pain, more disability and poorer quality of live [47–49]. Several authors, therefore, have pointed to fear-avoidance as a potential treatment target in rehabilitation of lumbar surgery [47,




Fig. 5 Forest plot for the meta-analysis of the effectiveness of multimodal rehabilitation versus exercise alone for reducing disability and pain. All studies are ordered from most to least effective. Random-effects model was used. Negative effect sizes favor multimodal rehabilitation

50]. Recently, Hanel et al. [22] demonstrated in their meta-analysis that exercise training effectively reduces fear-avoidance in a population with chronic low back pain. A single study included in our review could not confirm a fear-reducing effect of exercise alone in patients undergoing lumbar fusion [41]. However, the combination of exercise with psychosocial modalities was associated with less fear-avoidance up to six months after lumbar fusion. Given the high prevalence of fear-avoidance in patients with chronic musculoskeletal pain (56%) [51], a multimodal framework should be considered for patients undergoing lumbar fusion. In particular, patients with pain-related fear and in extent other interfering psychological components as outlined in the fear-avoidance model of Vlaeyen and Linton (e.g., anxiety and depression) [52], could benefit from multimodal rehabilitation tailored to their patient-specific characteristics and needs. Besides avoidance of activities, persistence of pain-provoking activities or a combination of pain persistence and avoidance, are also well-known maladaptive coping strategies, that may guide therapeutic approaches. It

should be pointed out, however, that none of the included multimodal interventions preselected patients based on their psychological profile or coping strategy.

Compared to *prehabilitation* in other orthopedic interventions such as hip and knee replacement, prehabilitation of lumbar fusion is still in its infancy. The fact that the majority (71%) of RCTs in this review skipped the preoperative period and only started rehabilitation postoperatively, may partly be an expression of prehabilitation being “unknown, unloved”. Four RCTs started preoperatively with CBT, but could not demonstrate less disability at last follow-up, which is in line with a recent meta-analysis that provided very low to low-certainty evidence that preoperative CBT is not effective for disability in patients undergoing lumbar surgery [15]. Nevertheless, preoperative physiotherapy and psychological therapy, improved pain after lumbar fusion surgery, in the study of Nielsen et al. and Reichart et al., respectively [39, 45]. Overall, we hope to set the scene for new (needed) studies rethinking rehabilitation across the entire care





Fig. 6 Forest plot for the meta-analysis of the effectiveness of multimodal rehabilitation versus exercise alone for reducing fear-avoidance. All studies are ordered from most to least effective. Random-effects model was used. Negative effect sizes favor multimodal rehabilitation

Fig. 7 Relative risk (RR) of return to work at long-term follow-up (Rolving et al. at 1 year, the remaining studies at 2 years postoperative) of rehabilitation interventions versus control group



continuum of lumbar fusion to unravel opportunities for value improvement. Given that all interventions that started preoperatively also continued postoperatively, we were not able to distinguish prehabilitation and postoperative rehabilitation. Consequently, the value of the optimal rehabilitation period (preoperatively, postoperatively or both) remains still unclear and in need of further investigation.

One *unexpected finding* is the variability of reported restrictions in the included trials, reflecting uncertainty among authors in whether and which restrictions are necessary following lumbar fusion. Restrictions ranged from prohibition of sports for three or six months [29, 37], or postoperative bracing [41], to six weeks of complete bedrest [43]. Noteworthy, overgeneralizing (unnecessary) restrictions may fuel iatrogenic pain-related fear and fear of movement, which are reported barriers for physical activity [53]. Restrictions not tailored to patient- and technique specific factors may thereby jeopardize the effects of rehabilitation interventions and a timely return-to-work. Hence, a call for evidence on the impact of postoperative restrictions emerges, requiring future research to clearly report on implied restrictions.

Our results suggest a tendency towards a higher *return-to-work* ratio after participation in a rehabilitation intervention compared to control condition in the long run. It would be interesting to also map out the time to return to work, however this was precluded due to underreporting of return-to-work at short-term follow-up in the included studies. Even small improvements in the return-to-work timeframes may have large impact on patients and our society. In this light, future rehabilitation trials should consistently measure return-to-work, and this already shortly after lumbar fusion surgery.

Based on our meta-analysis, exercise as a centerpiece of a multimodal framework is suggested. To translate this framework into a more detailed blueprint ready for clinical use, perspectives from the important stakeholders, such as patients, their caregivers, and policy makers, need to be included.

Study limitations

This study has several limitations. First, a small number of eligible trials with an unclear (72%) or high (28%) risk of bias, limited the level of evidence to low. Nonetheless, 15 additional RCTs were identified since the previous meta-analysis of Greenwood et al. [9]. Second, due to language other than English or Dutch, one record could not be retrieved, and one full-text article was excluded. Third, most trials were conducted in European countries (83%). Six author groups were affiliated to the same university in Denmark [28, 34–37, 44], thereby potentially limiting generalizability to other settings.

Fourth, rehabilitation interventions and comparisons were often insufficiently described. To enhance transparency and enable replication of exercises and other modalities, future studies should follow description guidelines. [23, 54, 55]. Moreover, transparency of trials also requires prospective protocol registration, which was only present in a minority of included trials.

Finally, the comparison of multimodal rehabilitation with exercise had a large degree of statistical heterogeneity, as indicated by an outlier and large I^2 statistics. Inclusion in the meta-analysis was based on sufficient clinical homogeneity in terms of rehabilitation modality. Remaining clinical heterogeneity could be related to differences in timing, duration, intensity and setting of the rehabilitation. Additionally, it is possible that non-reaching of surgical goals (e.g., unsuccessful fusion, alignment or decompression) interferes with the long-term effects of rehabilitation. The inclusion of different fusion techniques and indications across RCTs, may imply variable structural success rates. Surprisingly, four included RCTs reported non-instrumented fusion [28, 29, 39, 44], which increases the risk for pseudarthrosis. Given paucity of surgical success data in included studies, we could not correct for this variability. One study with uncertain risk of bias and no description of used fusion technique reported an effect size on disability and pain much larger than any of the other included studies. This result is presumably attributed to the very high intensity of the rehabilitation program [32]. Exclusion of

this outlier from the meta-analyses substantially reduces heterogeneity and the magnitude of the summary effect sizes. This observation may raise the question whether rehabilitation shows a dose–response effect, which should be investigated by future research.

Conclusions

The results of this systematic review with meta-analysis encourage exercise for all patients undergoing lumbar fusion given the positive impact on disability and pain up to six months postoperative. Embedding exercise in a multimodal rehabilitation context is suggested given the additional positive effect on disability and pain-related fear, compared to exercise alone. It remains uncertain if these beneficial effects of exercise and multimodal rehabilitation persist in the long term. Additional high-quality research is needed to evaluate these long-term functional and work-related outcomes and to establish the optimal period (pre-, postoperative or both) and dose of rehabilitation.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00586-022-07158-2>.

Acknowledgements The authors wish to thank Thomas Vandendriessche, Kristel Paque and Krizia Tuand, the biomedical reference librarians of the KU Leuven Libraries, 2Bergen Learning Centre Désiré Collen (Leuven, Belgium), for their help in conducting the systematic literature search.

Author contributions All authors contributed to the study conception and design. The articles screening and selection was performed by LB and TT. Studies' quality assessment and data extraction was performed by LB, TT and CA. Analysis and interpretation of results was performed by LB, LJ and TWS. The first draft of the manuscript was written by LB and all authors revised it critically for important intellectual content. All authors read and approved the final manuscript.

Funding This work was supported by the Research Foundation—Flanders (FWO, grant 1181521N) and Competence Centre Physiotherapy UZ/KU Leuven.

Declarations

Conflict of interest The authors have no conflicts of interest to declare that are relevant to the content of this article.

Availability of data and material NA.

Code availability NA.

References

- Rajae SS, Bae HW, Kanim LE, Delamarter RB (2012) Spinal fusion in the United States: analysis of trends from 1998 to 2008. *Spine (Phila Pa 1976)* 37:67–76. <https://doi.org/10.1097/BRS.0b013e31820cccfb>
- Al Jammal OM, Delavar A, Maguire KR, Hirshman BR, Wali AR, Kazzaz M et al (2019) National trends in the surgical management of lumbar spinal stenosis in adult spinal deformity patients. *Spine (Phila Pa 1976)* 44:E1369–E1378. <https://doi.org/10.1097/brs.0000000000003155>
- Van Grinsven V (2018) Lumbar fusion surgery: are fusion success and clinical outcome associated? <https://www.scriptiebank.be/scriptie/2018/lumbar-fusion-surgery-are-fusion-success-and-clinical-outcome-associated>. Accessed 28 Apr 2021
- Reisener M-J, Pumberger M, Shue J, Girardi FP, Hughes AP (2020) Trends in lumbar spinal fusion—a literature review. *J Spine Surg* 6:752–761. <https://doi.org/10.21037/jss-20-492>
- Khor S, Lavalley DC, Cizik AM, Bellabarba C, Dagal A, Hart RA et al (2020) Hospital and surgeon variation in patient-reported functional outcomes after lumbar spine fusion: a statewide evaluation. *Spine (Phila Pa 1976)* 45:465–472. <https://doi.org/10.1097/BRS.0000000000003299>
- Strömqvist B, Fritzell P, Hägg O, Jönsson B, Sandén B (2013) Swespine: the Swedish spine register: the 2012 report. *Eur Spine J* 22:953–974. <https://doi.org/10.1007/s00586-013-2758-9>
- Hedlund R, Johansson C, Hägg O, Fritzell P, Tullberg T (2016) The long-term outcome of lumbar fusion in the Swedish lumbar spine study. *Spine J* 16:579–587. <https://doi.org/10.1016/j.spinee.2015.08.065>
- Mannion AF, Brox JI, Fairbank JC (2013) Comparison of spinal fusion and nonoperative treatment in patients with chronic low back pain: long-term follow-up of three randomized controlled trials. *Spine J* 13:1438–1448
- Greenwood J, McGregor A, Jones F, Mullane J, Hurley M (2016) Rehabilitation following lumbar fusion surgery: a systematic review and meta-analysis. *Spine (Phila Pa 1976)* 41:E28–E36. <https://doi.org/10.1097/brs.0000000000001132>
- Madera M, Brady J, Deily S, McGinty T, Moroz L, Singh D et al (2017) The role of physical therapy and rehabilitation after lumbar fusion surgery for degenerative disease: a systematic review. *J Neurosurg Spine* 26:694–704. <https://doi.org/10.3171/2016.10.Spine16627>
- van Erp RMA, Jelsma J, Huijnen IPJ, Lundberg M, Willems PC, Smeets R (2018) Spinal surgeons' opinions on pre- and postoperative rehabilitation in patients undergoing lumbar spinal fusion surgery: a survey-based study in the Netherlands and Sweden. *Spine (Phila Pa 1976)* 43:713–719. <https://doi.org/10.1097/brs.0000000000002406>
- Gilmore SJ, McClelland JA, Davidson M (2016) Physiotherapy management of patients undergoing lumbar spinal surgery: a survey of Australian physiotherapists. *N Z J Physiother* 44:105–112
- Rushton A, Wright C, Heap A, White L, Eveleigh G, Heneghan N (2014) Survey of current physiotherapy practice for patients undergoing lumbar spinal fusion in the United Kingdom. *Spine (Phila Pa 1976)* 39:E1380–E1387. <https://doi.org/10.1097/brs.0000000000000573>
- Rushton A, Eveleigh G, Petherick EJ, Heneghan N, Bennett R, James G et al (2012) Physiotherapy rehabilitation following lumbar spinal fusion: a systematic review and meta-analysis of randomised controlled trials. *BMJ Open*. <https://doi.org/10.1136/bmjopen-2012-000829>
- Janssen ERC, Punt IM, Clemens MJ, Staal JB, Hoogeboom TJ, Willems PC (2021) Current prehabilitation programs do not improve the postoperative outcomes of patients scheduled for lumbar spine surgery: a systematic review with meta-analysis. *J Orthop Sports Phys Ther* 51:103–114. <https://doi.org/10.2519/jospt.2021.9748>
- HHiggins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (2022) *Cochrane handbook for systematic reviews*

- of interventions version 6.3. <https://www.training.cochrane.org/handbook>
17. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD et al (2021) The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 372:n71. <https://doi.org/10.1136/bmj.n71>
 18. Jane F (2018) Removing duplicates from an EndNote library. <https://blogs.lshrm.ac.uk/library/2018/12/07/removing-duplicates-from-an-endnote-library/>. Accessed 28 Apr 2021
 19. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A (2016) Rayyan—a web and mobile app for systematic reviews. *Syst Rev* 5:210. <https://doi.org/10.1186/s13643-016-0384-4>
 20. Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I et al (2019) RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 366:l4898. <https://doi.org/10.1136/bmj.l4898>
 21. Cochrane Training (2021) RoB 2: learning live webinar series. <https://training.cochrane.org/rob-2-learning-live-webinar-series>
 22. Hanel J, Owen PJ, Held S, Tagliaferri SD, Miller CT, Donath L et al (2020) Effects of exercise training on fear-avoidance in pain and pain-free populations: systematic review and meta-analysis. *Sports Med* 50:2193–2207. <https://doi.org/10.1007/s40279-020-01345-1>
 23. Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D et al (2014) Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 348:g1687. <https://doi.org/10.1136/bmj.g1687>
 24. Balduzzi S, Schwarzer G (2019) How to perform a meta-analysis with R: a practical tutorial. *Evid Based Ment Health* 22:153–160
 25. Harrer M, Cuijpers P, Furukawa TA, Ebert DD (2019) Doing meta-analysis in R: a hands-on guide. PROTECT Lab, Erlangen
 26. R Core Team (2020) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Australia
 27. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P et al (2008) GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 336:924–926. <https://doi.org/10.1136/bmj.39489.470347.AD>
 28. Rolving N, Nielsen CV, Christensen FB, Holm R, Bünger CE, Oestergaard LG (2015) Does a preoperative cognitive-behavioral intervention affect disability, pain behavior, pain, and return to work the first year after lumbar spinal fusion surgery? *Spine (Phila Pa 1976)* 40:593–600. <https://doi.org/10.1097/brs.0000000000000843>
 29. Abbott AD, Tyni-Lenné R, Hedlund R (2010) Early rehabilitation targeting cognition, behavior, and motor function after lumbar fusion: a randomized controlled trial. *Spine (Phila Pa 1976)* 35:848–857. <https://doi.org/10.1097/BRS.0b013e3181d1049f>
 30. Ilves O, Häkkinen A, Dekker J, Wahlman M, Tarnanen S, Pekkanen L et al (2017) Effectiveness of postoperative home-exercise compared with usual care on kinesiophobia and physical activity in spondylolisthesis: a randomized controlled trial. *J Rehabil Med* 49:751–757. <https://doi.org/10.2340/16501977-2268>
 31. Greenwood J, McGregor A, Jones F, Hurley M (2019) Rehabilitation following lumbar fusion surgery (REFS) a randomised controlled feasibility study. *Eur Spine J* 28:735–744. <https://doi.org/10.1007/s00586-019-05913-6>
 32. Monticone M, Ferrante S, Teli M, Rocca B, Foti C, Lovi A et al (2014) Management of catastrophising and kinesiophobia improves rehabilitation after fusion for lumbar spondylolisthesis and stenosis. A randomised controlled trial. *Eur Spine J* 23:87–95. <https://doi.org/10.1007/s00586-013-2889-z>
 33. Lotzke H, Brisby H, Gutke A, Hägg O, Jakobsson M, Smeets R et al (2019) A person-centered prehabilitation program based on cognitive-behavioral physical therapy for patients scheduled for lumbar fusion surgery: a randomized controlled trial. *Phys Ther* 99:1069–1088. <https://doi.org/10.1093/ptj/pzz020>
 34. Strøm J, Nielsen CV, Jørgensen LB, Andersen NT, Laursen M (2019) A web-based platform to accommodate symptoms of anxiety and depression by featuring social interaction and animated information in patients undergoing lumbar spine fusion: a randomized clinical trial. *Spine J* 19:827–839. <https://doi.org/10.1016/j.spinee.2018.11.011>
 35. Oestergaard LG, Christensen FB, Bünger CE, Sogaard R, Holm R, Helmig P et al (2020) Does adding case management to standard rehabilitation affect functional ability, pain, or the rate of return to work after lumbar spinal fusion? A randomized controlled trial with two-year follow-up. *Clin Rehabil* 34:357–368. <https://doi.org/10.1177/0269215519897106>
 36. Oestergaard LG, Nielsen CV, Bünger CE, Sogaard R, Fruensgaard S, Helmig P et al (2012) The effect of early initiation of rehabilitation after lumbar spinal fusion: a randomized clinical study. *Spine (Phila Pa 1976)* 37:1803–1809. <https://doi.org/10.1097/BRS.0b013e31825a17ab>
 37. Christensen FB, Laurberg I, Bünger CE (2003) Importance of the back-café concept to rehabilitation after lumbar spinal fusion: a randomized clinical study with a 2-year follow-up. *Spine (Phila Pa 1976)* 28:2561–2569. <https://doi.org/10.1097/01.Brs.0000097890.96524.A1>
 38. Kang H, Cho K, Shim S, Yu J, Jung J (2012) Effects of exercise rehabilitation on pain, disability, and muscle strength after posterior lumbar interbody fusion surgery: a randomized controlled trial. *J Phys Ther Sci* 24:1037–1040. <https://doi.org/10.1589/jpts.24.1037>
 39. Nielsen PR, Jørgensen LD, Dahl B, Pedersen T, Tønnesen H (2010) Prehabilitation and early rehabilitation after spinal surgery: randomized clinical trial. *Clin Rehabil* 24:137–148. <https://doi.org/10.1177/0269215509347432>
 40. Kernc D, Strojnik V, Vengust R (2018) Early initiation of a strength training based rehabilitation after lumbar spine fusion improves core muscle strength: a randomized controlled trial. *J Orthop Surg Res* 13:151. <https://doi.org/10.1186/s13018-018-0853-7>
 41. Salik Sengul Y, Kaya N, Yalcinkaya G, Kirmizi M, Kalemci O (2020) The effects of the addition of motor imagery to home exercises on pain, disability and psychosocial parameters in patients undergoing lumbar spinal surgery: a randomized controlled trial. *Explore (NY)*. <https://doi.org/10.1016/j.explore.2020.02.001>
 42. Elsayyad MM, Abdel-Aal NM, Helal ME (2021) Effect of adding neural mobilization versus myofascial release to stabilization exercises after lumbar spine fusion: a randomized controlled trial. *Arch Phys Med Rehabil* 102:251–260. <https://doi.org/10.1016/j.apmr.2020.07.009>
 43. Zhao BX, Wang KZ, Zhao JX, Wang CS, Huang XH, Shu-qiang M et al (2008) Clinical effects of acupuncture after surgical operation in patients with prolapse of the lumbar intervertebral disc. *J Tradit Chin Med* 28:250–254. [https://doi.org/10.1016/s0254-6272\(09\)60003-0](https://doi.org/10.1016/s0254-6272(09)60003-0)
 44. Oestergaard LG, Maribo T, Bünger CE, Christensen FB (2012) The Canadian Occupational Performance Measure’s semi-structured interview: its applicability to lumbar spinal fusion patients. A prospective randomized clinical study. *Eur Spine J* 21:115–121. <https://doi.org/10.1007/s00586-011-1957-5>
 45. Reichart R, Vogel I, Weiss T, Hennig S, Walter J, Kalf R (2011) Short Psychological intervention as a perioperative pain reduction treatment in spinal neurosurgery. *J Neurol Surg A Cent Eur Neurosurg* 73:387–396. <https://doi.org/10.1055/s-0032-1313642>
 46. Wewege MA, Jones MD (2021) Exercise-induced hypoalgesia in healthy individuals and people with chronic musculoskeletal pain: a systematic review and meta-analysis. *J Pain* 22:21–31. <https://doi.org/10.1016/j.jpain.2020.04.003>

47. Kemani MK, Hägg O, Jakobsson M, Lundberg M (2020) Fear of movement is related to low back disability during a two-year period in patients who have undergone elective lumbar spine surgery. *World Neurosurg* 137:e416–e424. <https://doi.org/10.1016/j.wneu.2020.01.218>
48. Johansson A-C, Linton SJ, Rosenblad A, Bergkvist L, Nilsson O (2010) A prospective study of cognitive behavioural factors as predictors of pain, disability and quality of life one year after lumbar disc surgery. *Disabil Rehabil* 32:521–529. <https://doi.org/10.3109/09638280903177243>
49. Abbott AD, Tyni-Lenné R, Hedlund R (2011) Leg pain and psychological variables predict outcome 2–3 years after lumbar fusion surgery. *Eur Spine J* 20:1626–1634. <https://doi.org/10.1007/s00586-011-1709-6>
50. Leeuw M, Goossens ME, Linton SJ, Crombez G, Boersma K, Vlaeyen JW (2007) The fear-avoidance model of musculoskeletal pain: current state of scientific evidence. *J Behav Med* 30:77–94. <https://doi.org/10.1007/s10865-006-9085-0>
51. Bränström H, Fahlström M (2008) Kinesiophobia in patients with chronic musculoskeletal pain: differences between men and women. *J Rehabil Med* 40:375–380. <https://doi.org/10.2340/16501977-0186>
52. Vlaeyen JWS, Linton SJ (2000) Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain* 85:317–332. [https://doi.org/10.1016/s0304-3959\(99\)00242-0](https://doi.org/10.1016/s0304-3959(99)00242-0)
53. Boutevillain L, Dupeyron A, Rouch C, Richard E, Coudeyre E (2017) Facilitators and barriers to physical activity in people with chronic low back pain: a qualitative study. *PLoS ONE* 12:e0179826. <https://doi.org/10.1371/journal.pone.0179826>
54. Slade SC, Dionne CE, Underwood M, Buchbinder R (2016) Consensus on exercise reporting template (CERT): explanation and elaboration statement. *Br J Sports Med* 50:1428–1437. <https://doi.org/10.1136/bjsports-2016-096651>
55. Whyte J, Dijkers MP, Fasoli SE, Ferraro M, Katz LW, Norton S et al (2021) Recommendations for reporting on rehabilitation interventions. *Am J Phys Med Rehabil* 100:5–16. <https://doi.org/10.1097/phm.0000000000001581>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.