

A Lower Bound for the Height of a Rational Function at S -unit Points

By

Pietro Corvaja¹ and Umberto Zannier²

¹ University of Udine, Italy

² Scuola Normale Superiore, Pisa, Italy

Received September 29, 2003; accepted in revised form March 30, 2004

Published online December 13, 2004 © Springer-Verlag 2004

Abstract. Let a, b be given, multiplicatively independent positive integers and let $\epsilon > 0$. In a recent paper jointly with Y. Bugeaud we proved the upper bound $\exp(\epsilon n)$ for $\text{g.c.d.}(a^n - 1, b^n - 1)$; shortly afterwards we generalized this to the estimate $\text{g.c.d.}(u - 1, v - 1) < \max(|u|, |v|)^\epsilon$ for multiplicatively independent S -units $u, v \in \mathbf{Z}$. In a subsequent analysis of those results it turned out that a perhaps better formulation of them may be obtained in terms of the language of heights of algebraic numbers. In fact, the purposes of the present paper are: to generalize the upper bound for the g.c.d. to pairs of rational functions other than $\{u - 1, v - 1\}$ and to extend the results to the realm of algebraic numbers, giving at the same time a new formulation of the bounds in terms of height functions and algebraic subgroups of \mathbf{G}_m^2 .

2000 Mathematics Subject Classification: 11J25

Key words: Lower bounds for the height, Subspace Theorem, Linear tori

1. Introduction

Let a, b be given, multiplicatively independent positive integers and let $\epsilon > 0$. In the paper [2] a nearly best-possible upper bound $\exp(\epsilon n)$ for $\text{g.c.d.}(a^n - 1, b^n - 1)$ was proved, as the integer n tends to infinity. In the subsequent paper [7] we observed (see Remark 1) that the combination of those arguments with [2] yields a generalization, where a^n, b^n are replaced by multiplicatively independent S -units $u, v \in \mathbf{Z}$. Namely the upper bound $\text{g.c.d.}(u - 1, v - 1) < \max(|u|, |v|)^\epsilon$ holds with finitely many exceptions. (Full details for this can be obtained in a straightforward way from [2] and [7], as is done in [14] and [5].)

In a subsequent analysis of those results it turned out that a perhaps better formulation of them may be obtained in terms of the language of heights of algebraic numbers. This also suggested an extension to general number fields. Finally, in view of possible applications (like, e.g., the one in [7], where a conjecture about the greatest prime factor of $(ab + 1)(ac + 1)$ was settled) we have tried to replace $u - 1, v - 1$ with more general functions.

In fact, the purposes of the present paper are:

(i) to generalize the upper bound for the g.c.d. to pairs of rational functions other than $\{u - 1, v - 1\}$ and

(ii) to extend the results to the realm of algebraic numbers, giving at the same time a new formulation of the bounds in terms of height functions and algebraic subgroups of \mathbf{G}_m^2 .

More precisely, note that a “good” upper bound for the g.c.d. of two integers amounts to a lower bound for the height of their ratio, essentially by their maximum height. With this in mind, we seek lower bounds for the height $h(\varphi(u, v))$, where φ is a rational function and u, v run along S -units in a number field, for a finite set S . Equivalently, we may let (u, v) run through a finitely generated group $\Gamma \subset \mathbf{G}_m^2(\overline{\mathbf{Q}})$. Our main results will give, under suitable assumptions, the expected lower bounds, with the possible exceptions of the pairs (u, v) contained in a finite union of proper algebraic subgroups (or translates) of \mathbf{G}_m^2 : these kind of subvarieties are certainly relevant in the context; for instance, a pair (u, v) lies in some proper algebraic subgroup of \mathbf{G}_m^2 if and only if u, v are multiplicatively dependent.

Before stating the results we observe that analogous investigations appear in [1] in the function field case, and from the quantitative point of view appear in [3] (for the greatest prime factor of $(ab + 1)(ac + 1)(bc + 1)$) and [4] (where some results of [8] are in part quantified).

We immediately proceed to give the formal statements.

Theorem 1. *Let $\Gamma \subset \mathbf{G}_m^2(\overline{\mathbf{Q}})$ be a finitely generated group, $p(X, Y), q(X, Y) \in \overline{\mathbf{Q}}[X, Y]$ be non constant coprime polynomials and suppose that not both of them vanish at $(0, 0)$. For every positive ϵ , the Zariski closure of the set of solutions $(u, v) \in \Gamma$ to the inequality*

$$h\left(\frac{p(u, v)}{q(u, v)}\right) < h(p(u, v) : q(u, v) : 1) - \epsilon \cdot \max\{h(u), h(v)\} \tag{1.1}$$

is the union of finitely many translates of 1-dimensional subtori of \mathbf{G}_m^2 which can be effectively determined and a finite set.

Note that, since $h(x : y : 1) \geq \max\{h(x), h(y)\}$, in (1.1) we can replace $h(p(u, v) : q(u, v) : 1)$ with $\max\{h(p(u, v)), h(q(u, v))\}$ (but this would lead to a weaker statement). Also, we could have replaced (1.1) with the equivalent inequality

$$\sum_{\mu \in M_K} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\} < -\epsilon \max\{h(u), h(v)\}, \tag{1.2}$$

where $\log^-(\cdot)$ stands for $\min\{0, \log(\cdot)\}$. (See the beginning of next section for conventions about valuations.)

Actually, the proof will go through (1.2) and we shall show later the equivalence. Note that the left hand side of (1.2) is an analogue of the g.c.d. (in a sense which considers also archimedean valuations), and for rational integers it is precisely its logarithm.

Theorem 1 easily implies the following corollary:

Corollary 1. *Let Γ be as before. Then for multiplicatively independent pairs $(u, v) \in \Gamma$ the height of the ratio $(u - 1)/(v - 1)$ verifies the asymptotic equivalence,*

as $\max\{h(u), h(v)\} \rightarrow \infty$:

$$h((u - 1)/(v - 1)) \sim h(1 : u : v).$$

Note that, as before, this is (essentially) equivalent to the bound

$$\sum_{\mu \in M_K} \log^- \max\{|u - 1|_\mu, |v - 1|_\mu\} > -\epsilon \cdot \max\{h(u), h(v)\}, \tag{1.3}$$

valid for all ϵ and all multiplicatively independent pairs $(u, v) \in \Gamma$ apart from a finite set depending on ϵ and Γ .

Also, the main results of [2] and [7] are immediate consequences of this Corollary.

Our most general result is the following

Main Theorem. *Let $f(X, Y) \in \overline{\mathbf{Q}}(X, Y)$ be a rational function and $\Gamma \subset \mathbf{G}_m^2(\overline{\mathbf{Q}})$ be a finitely generated subgroup. Denote by $T_1(X, Y), \dots, T_N(X, Y)$ the monomials appearing in the numerator and denominator of $f(X, Y)$ and suppose that $1 \in \{T_1, \dots, T_N\}$. Then for every $\epsilon > 0$ the Zariski closure of the set of solutions $(u, v) \in \Gamma$ of the inequality*

$$h(f(u, v)) < (1 - \epsilon) \max\{h(T_1(u, v)), \dots, h(T_N(u, v))\} \tag{1.4}$$

is a finite union of translates of proper subtori of \mathbf{G}_m^2 . Also, outside a finite union of translates of proper subtori of \mathbf{G}_m^2 , one has

$$h(f(u, v)) > (1 - \epsilon) \max\left\{\frac{h(u)}{2 \deg_Y f}, \frac{h(v)}{2 \deg_X f}\right\}. \tag{1.5}$$

We shall now explain the strategy behind our proofs. For instance, to prove our Main Theorem above, observe first that we can reduce to the case when f is defined over a number field K and Γ is the group $(\mathcal{O}_S^\times)^2$ of points whose coordinates are S -units for a certain fixed finite set of places $S \subset M_K$. Now, put $f(X, Y) = \frac{p(X, Y)}{q(X, Y)}$, where $p, q \in K[X, Y]$ are coprime polynomials; for every point $(u, v) \in K^2$, where f is defined, the height of $f(u, v)$ is $h(f(u, v)) = h(p(u, v) : q(u, v))$. Hence it verifies

$$\begin{aligned} h(f(u, v)) &= \sum_{\mu \in M_K} \log \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\} \\ &= \sum_{\mu \in M_K} \log^+ \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\} \\ &\quad + \sum_{\mu \in M_K} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\} \end{aligned}$$

where $\log^+(\cdot) = \max\{0, \log(\cdot)\}$ and $\log^-(\cdot) = \min\{0, \log(\cdot)\}$. Hence

$$h(f(u, v)) \geq \max\{h(p(u, v)), h(q(u, v))\} + \sum_{\mu \in M_K} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\}.$$

We then need lower bounds for $\max\{h(p(u, v)), h(q(u, v))\}$ and for $\sum_{\mu \in M_K} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\}$. These estimates are the object of Propositions 1 and 4 and will be proved by using in an essential way the

Subspace Theorem. While the techniques to prove lower bounds like in Proposition 1 are well known, to prove upper bounds as in Proposition 4 we need a new method, introduced in [6] and developed in [2] and [7], which consists in applying the Subspace Theorem to suitable linear combinations of S -units and S -integers.

2. Proofs

We start by recalling the definitions:

$$\log^+ x := \max\{0, \log x\}, \quad \log^- x := \min\{0, \log x\} \quad x > 0.$$

Note that these functions are nondecreasing.

In the sequel, K will denote a number field, M_K the set of places of K , M_0 the subset of finite places; for a finite subset $S \subset M_K$ containing all archimedean places, let us denote by \mathcal{O}_S the ring of S -integers and by \mathcal{O}_S^\times its group of units (S -units). For a place $\mu \in M_K$, let us denote by $|\cdot|_\mu$ the corresponding absolute value, normalized *with respect to* K , i.e. in such a way that the product formula holds, and the absolute Weil height reads $H(x) = \exp(h(x)) = \prod_\mu \max\{1, |x|_\mu\}$. For a point $\mathbf{x} = (x_1, \dots, x_n) \in K^n$ and a place $\mu \in M_K$, put $|\mathbf{x}|_\mu = \max_i |x_i|_\mu$. For a point $\mathbf{x} = (x_1 : \dots : x_n) \in \mathbf{P}_{n-1}(K)$, put $H(\mathbf{x}) = \prod_\mu \max\{|x_1|_\mu, \dots, |x_n|_\mu\} = \prod_\mu |\mathbf{x}|_\mu$ (where in the last term the point \mathbf{x} is considered in K^n), and call it the height of the point \mathbf{x} ; note that it is well defined in view of the product formula.

We begin by stating the so called Subspace Theorem, which represents the main technical tool in the present proofs.

Subspace Theorem. *Let S be a finite set of absolute values of a number field K , including the archimedean ones. For $\mu \in S$, let $L_{1,\mu}, \dots, L_{N,\mu}$ be linearly independent linear forms in N variables, defined over K ; let $\delta > 0$. Then the solutions $P = (x_1, \dots, x_N) \in K^N \setminus \{0\}$ to the inequality*

$$\prod_{\mu \in S} \prod_{i=1}^N \frac{|L_{i,\mu}(P)|_\mu}{|P|_\mu} < H(P)^{-N-\delta}$$

are contained in finitely many proper subspaces of K^N .

For a proof, see [12, 13].

In the sequel, we shall also make use of the following two lemmas:

Lemma 1. *Let \tilde{K} be a number field, $\varphi(T) \in \tilde{K}(T)$ be a rational function of degree d . Then for $t \in \tilde{K}$,*

$$H(\varphi(t)) \gg H(t)^d,$$

where the constant appearing implicitly in the symbol \gg can be effectively determined.

The proof of Lemma 1 can be found for instance in [10]: it is a very particular case of Theorem B.2.5 of [10].

Lemma 2. *Let $\{T_1(X, Y), \dots, T_M(X, Y)\}$ be a set of non constant monomials in X, Y and suppose that $\deg_X T_i \leq d_1, \deg_Y T_i \leq d_2$ for $i = 1, \dots, M$. Suppose also*

that the intersection of the M subgroups defined by the equations $T_i(X, Y) = 1$ is zero-dimensional. Let $(u, v) \in (K^\times)^2$ be an algebraic point of \mathbf{G}_m^2 . Then

$$\max\{h(T_1(u, v)), \dots, h(T_M(u, v))\} \geq \frac{1}{2} \max\left\{\frac{h(u)}{d_2}, \frac{h(v)}{d_1}\right\}.$$

Proof. Denote by $T \subset \mathbf{Z}^2$ the set of exponents (i, j) appearing in the monomials T_1, \dots, T_M . The group generated by T in \mathbf{Z}^2 is of rank two by our assumption. Hence the set T contains two linearly independent points $(i_1, i_2), (j_1, j_2)$. Hence $X^d = (X^{i_1} Y^{i_2})^{j_2} (X^{j_1} Y^{j_2})^{-i_2}$ where $d = i_1 j_2 - i_2 j_1 \neq 0$. By substituting X, Y by u, v and considering the corresponding height one obtains

$$|d| \cdot h(u) \leq |j_2| \cdot h(u^{i_1} v^{i_2}) + |i_2| \cdot h(u^{j_1} v^{j_2}) \leq 2d_2 \max\{h(u^{i_1} v^{i_2}), h(u^{j_1} v^{j_2})\}.$$

Hence the maximum of the heights of the monomials is bounded from below by $\frac{1}{2} \frac{h(u)}{d_2}$. Of course the analogous inequality holds for $h(v)$, thus proving the lemma. The following proposition is essentially a corollary of Evertse’s paper [9]. For the readers’ convenience, we give here a complete proof.

Proposition 1. *Let $S \subset M_K$ and $\Gamma \subset \mathbf{G}_m^2(\overline{\mathbf{Q}})$ be as above, $f(X, Y) \in K[X, Y, X^{-1}, Y^{-1}]$ be a regular function on \mathbf{G}_m^2 . Denote by $T_1(X, Y), \dots, T_N(X, Y)$ the monomials in X, X^{-1}, Y, Y^{-1} which appear in f and suppose that $1 \in \{T_1, \dots, T_N\}$. Let $\epsilon > 0$ be fixed. Then the solutions $(u, v) \in \Gamma$ to the inequality*

$$h(f(u, v)) < (1 - \epsilon) \max\{h(T_1(u, v)), \dots, h(T_N(u, v))\} \tag{2.1}$$

lie in the union of finitely many translates of 1-dimensional subtori of \mathbf{G}_m^2 , which can be effectively determined, and a finite set. The same conclusion holds for the solutions to the inequality

$$\sum_{\mu \in S} \log^- |f(u, v)|_\mu < -\epsilon \max\{h(u), h(v)\}. \tag{2.2}$$

The idea of the proof is two-fold: first, by an application of the Subspace Theorem, we obtain that the solutions to the above inequality belong to the union of finitely many proper subgroups. Then, a Liouville-type argument enables to bound the degree of the minimal equations satisfied by any solution.

Proof of Proposition 1. We shall first prove our contention for the solutions to the inequality (2.1). Let us write

$$f(X, Y) = \sum_{(i,j) \in T} a_{i,j} X^i Y^j = b_1 T_1 + \dots + b_N T_N$$

where $T \subset \mathbf{Z}^2$ is a finite set of cardinality N and, for $(i, j) \in T$, $a_{i,j} \in K^\times$ are nonzero algebraic numbers in K ordered as b_1, \dots, b_N .

We first treat the (easier) case where T is contained in a line of \mathbf{Z}^2 . In this case, there exists a rational function $\varphi(T) \in K[T, T^{-1}]$ and a pair (i_0, j_0) such that identically

$$f(u, v) = \varphi(u^{i_0} v^{j_0}).$$

Also, for $(u, v) \in \mathbf{G}_m^2(K)$, $\deg(\varphi) \cdot h(u^{i_0}v^{j_0}) \geq \max\{h(T_1(u, v)), \dots, h(T_N(u, v))\}$. So, by Lemma 1, there exists a computable constant c such that for all $(u, v) \in \mathbf{G}_m^2(K)$

$$h(f(u, v)) \geq \max\{h(T_1(u, v)), \dots, h(T_N(u, v))\} - c.$$

Then, if inequality (2.1) holds, the above lower bound gives

$$\max\{h(T_1(u, v)), \dots, h(T_N(u, v))\} \leq c/\epsilon.$$

This implies that the pair (u, v) is contained in a finite translate of a subtorus of \mathbf{G}_m^2 defined by an equation of the form

$$u^{i_0}v^{j_0} = \lambda$$

where $\lambda \in K$ is an algebraic number with $h(\lambda) \leq c/\epsilon$. This implies our contention in that particular case.

We now consider the generic case where the subset $T \subset \mathbf{Z}^2$ is not contained in a line.

As we have already remarked, we can choose a finite set S of places of K containing the archimedean ones such that the subgroup Γ is contained in $(\mathcal{O}_S^\times)^2$. Now, for each solution $(u, v) \in (\mathcal{O}_S^\times)^2$ of the inequality (2.1) and each $\mu \in S$, let $k(u, v; \mu)$ be an index in $\{1, \dots, N\}$ for which the quantity $|T_k(u, v)|_\mu$ is maximal.

By partitioning the set of solutions of inequality (2.1) in at most N subsets, and working separately with each subset, we can suppose that for each $\mu \in S$, $k(u, v; \mu) =: k_\mu$ does not depend on (u, v) .

Let us define, for each $\mu \in S$, N linearly independent linear forms in T_1, \dots, T_N as follows: for $i \neq k_\mu$, put $L_{i,\mu} = T_i$, while for $i = k_\mu$, put

$$L_{k_\mu,\mu} = b_1T_1 + \dots + b_N T_N.$$

Note that for each $\mu \in S$, the above defined linear forms are indeed independent, since all coefficients b_i are nonzero.

For a solution $(u, v) \in (\mathcal{O}_S^\times)^2$ of the inequality (2.1) we put $P = P(u, v) = (T_1(u, v), \dots, T_N(u, v)) \in \mathcal{O}_S^N$, so that for each $\mu \in S$ we have $L_{k_\mu,\mu}(P) = f(u, v)$. Note that, by Lemma 2 and the assumption that T is not contained in a line, the height of P tends to infinity on every infinite sequence of points $(u, v) \in K^2$. We now consider the double product

$$\prod_{\mu \in S} \prod_{i=1}^N \frac{|L_{i,\mu}(P)|_\mu}{|P|_\mu}; \tag{2.3}$$

an application of the Subspace Theorem to the above double product will give what we want. We note that, since the coordinates of P are S -integers (in fact even S -units), the quantity $\prod_{\mu \in S} |P|_\mu$ equals the height $H(P)$ of P . We can then rewrite the above double product as

$$\begin{aligned} \prod_{\mu \in S} \prod_{i=1}^N \frac{|L_{i,\mu}(P)|_\mu}{|P|_\mu} &= H(P)^{-N} \cdot \left(\prod_{\mu \in S} |f(u, v)|_\mu \right) \\ &\cdot \left(\prod_{\mu \in S} \prod_{i=1}^N |T_i(u, v)|_\mu \right) \cdot \prod_{\mu \in S} |T_{k_\mu,\mu}(u, v)^{-1}|_\mu. \end{aligned} \tag{2.4}$$

Clearly, the inequality $\prod_{\mu \in S} |f(u, v)|_{\mu} \leq H(f(u, v))$ holds and by assumption (2.1) we also have $H(f(u, v)) < \max\{H(T_1(u, v)), \dots, H(T_N(u, v))\}^{1-\epsilon}$. In turn, since by assumption $1 \in \{T_1, \dots, T_N\}$, the height of P is larger than the maximum of the heights of its coordinates. Then from (2.1) it follows that $H(f(u, v)) < H(P)^{1-\epsilon}$.

Also the third factor in (2.4) equals 1 since all the $T_i(u, v)$ are S -units. The last factor $\prod_{\mu \in S} |T_{k_{\mu}, \mu}(u, v)|_{\mu}$ is equal to the height $H(P)$, by the choice of the k_{μ} . Hence the double product (2.3) can be bounded as

$$\prod_{\mu \in S} \prod_{i=1}^N \frac{|L_{i, \mu}(P)|_{\mu}}{|P|_{\mu}} \leq H(P)^{-N-\epsilon}.$$

The Subspace Theorem, applied with $\delta = \epsilon$, asserts that the points P arising from the solutions (u, v) to the inequality (2.1) lie in finitely many hyperplanes. Then we obtain finitely many equations of the form $\psi_j(u, v) = 0$ ($j = 1, \dots, M$), such that: (i) $\psi_j(X, Y)$ ($j = 1, \dots, M$) are irreducible polynomials in $\bar{K}[X, Y]$; (ii) all but finitely many solutions of (2.1) satisfy one of these equations and (iii) each such equation has infinitely many solutions in S -units $(u, v) \in (\mathcal{O}_S^{\times})^2$. By a Theorem of Lang [10, Theorem 7.3, p. 207], each equation $\psi_j(X, Y) = 0$ defines a translate of a proper subtorus. Every translate of a proper subtorus is either a point or a curve defined by an equation of the form

$$U^p V^q = w \tag{2.5}$$

for coprime p, q and nonzero $w \in K^{\times}$. Our next goal is to show that the pairs (p, q) such that infinitely many solutions of (2.1) satisfy the above equation (2.5) can be effectively determined.

Fix such a translate, given by (2.5), and suppose it contains infinitely many solutions $(u, v) \in (\mathcal{O}_S^{\times})^2$ to the inequality (2.1). For each $(u, v) \in (\mathcal{O}_S^{\times})^2$, write

$$u = t^q, \quad v = s^{-p}$$

where t, s lie in a fixed number field \tilde{K} . The relation $u^p v^q = w$ can be rewritten as $(t/s)^{pq} = w$; so that $(t/s)^p$ is a q -th root of w . Select one such root \bar{w} . We obtain the parametrization

$$\begin{cases} u = t^q \\ v = \bar{w}t^{-p} \end{cases} \tag{2.6}$$

Then, using the above parametrization, we can write

$$f(u, v) = f(t^q, \bar{w}t^{-p}) =: \varphi(t), \tag{2.7}$$

for a rational function $\varphi(t) \in \tilde{K}(t)$.

Note that in our case the function $\varphi(t)$ in (2.7) can be written as

$$\varphi(t) = \sum_{(i,j) \in T} a_{i,j} \bar{w}^j t^{qi-pj} = \sum_{l \in \mathbf{Z}} c_l t^l$$

where, for every $l \in \mathbf{Z}$,

$$c_l = \sum_{(i,j) \in T : qi-pj=l} a_{i,j} \bar{w}^j,$$

the last sum running over those pairs $(i, j) \in T$ with $qi - pj = l$. Since the degree d of $\varphi(t)$ is at least $\max\{|l| : c_l \neq 0\}$, Lemma 1 implies that

$$h(f(u, v)) = h(\varphi(t)) \geq \max\{|l| : c_l \neq 0\}h(t) + O(1)$$

Now, if $l = qi - pj$, for a pair $(i, j) \in T$, the height of the corresponding monomial $u^i v^j$ is $|l|h(t) + O(1)$. Hence, if all the coefficients c_l , for l of the form $l = qi - pj$ for at least one $(i, j) \in T$, are nonzero, we have the inequality

$$h(f(u, v)) \geq \max\{h(T_1(u, v)), \dots, h(T_N(u, v))\} + O(1)$$

which contradicts (2.1) for large values of $\max\{h(T_1(u, v)), \dots, h(T_N(u, v))\}$ (recall that by Lemma 2 such a maximum does tend to infinity, so we indeed obtain a contradiction with the hypothesis that the given subgroups contains infinitely many solutions of (2.1)). Hence some c_l must vanish, which means that, after the substitution (2.6), some cancellation occurs among the monomials T_1, \dots, T_N . This fact implies that there exist two distinct pairs of exponents $(i, j), (i', j') \in T$ such that $qi - pj = qi' - pj'$, so (p, q) verifies a (non trivial) equation of the form

$$q(i - i') - p(j - j') = 0.$$

Each such equation has exactly two solutions in coprime integers $p, q \in \mathbf{Z}$, which can be effectively determined. This proves the first contention in Proposition 1.

We shall now obtain the same conclusion under assumption (2.2). Chosing linear forms as before, we arrive again at equality (2.4). As we have already remarked, the third factor equals 1, since $T_i(u, v)$ are S -units for all $i = 1, \dots, N$. The product of the second and fourth factors in (2.4) can be rewritten as

$$\begin{aligned} & \left(\prod_{\mu \in S} |f(u, v)|_{\mu} \right) \prod_{\mu \in S} |T_{k_{\mu}, \mu}(u, v)^{-1}|_{\mu} \\ &= \left(\prod_{\mu \in S} \min\{1, |f(u, v)|_{\mu}\} \right) \cdot \left(\prod_{\mu \in S} \max\{1, |f(u, v)|_{\mu}\} \right) \cdot \prod_{\mu \in S} |T_{k_{\mu}, \mu}(u, v)^{-1}|_{\mu}. \end{aligned}$$

Clearly, for every $\mu \in S$,

$$\max\{1, |f(u, v)|_{\mu}\} \leq \max\{1, |N|_{\mu}\} \cdot \max\{|b_1|_{\mu}, \dots, |b_N|_{\mu}\} \cdot |T_{k_{\mu}, \mu}(u, v)|_{\mu},$$

so we have

$$\left(\prod_{\mu \in S} \max\{1, |f(u, v)|_{\mu}\} \right) \prod_{\mu \in S} |T_{k_{\mu}, \mu}(u, v)^{-1}|_{\mu} = O(1).$$

Now from (2.4) we obtain (recall that the third factor in (2.4) equals 1)

$$\prod_{\mu \in S} \prod_{i=1}^N \frac{|L_{i, \mu}(P)|_{\mu}}{|P|_{\mu}} \ll H(P)^{-N} \cdot \prod_{\mu \in S} \min\{1, |f(u, v)|_{\mu}\}$$

where the constant implicit in the symbol \ll does not depend on (u, v) . The inequality (2.2) reads

$$\prod_{\mu \in S} \min\{1, |f(u, v)|_{\mu}\} < \max\{H(u), H(v)\}^{-\epsilon}.$$

Clearly, $H(P) \leq \max\{H(u), H(v)\}^D$ for a positive constant D (which can be taken to be $2 \max\{|i| + |j| : (i, j) \in T\}$). Then from the last displayed inequality we obtain

$$\prod_{\mu \in S} \prod_{i=1}^N \frac{|L_{i,\mu}(P)|_\mu}{|P|_\mu} \ll H(P)^{-N-\epsilon/D}.$$

The Subspace Theorem, applied with $\delta = \epsilon/(2D)$, say, asserts that all points $P(u, v)$ are contained in the union of finitely many hyperplanes. The final argument to obtain the conclusion is exactly the same as in the previous case, so we do not repeat it.

Our next goal is to prove an “explicit” version of Corollary 1, which will be used in the subsequent proofs of Theorem 1 and the Main Theorem. Corollary 1 can be rephrased by saying that if the ratio $(u - 1)/(v - 1)$ has “small” height, where u and v belong to a given finitely generated multiplicative group, then, apart from finitely many exceptions, the pair (u, v) verifies a multiplicative dependence relation, i.e. a relation of the form

$$u^p \cdot v^q = 1$$

for a vector $(p, q) \in \mathbf{Z}^2 \setminus \{(0, 0)\}$. In other words, the point $(u, v) \in \mathbf{G}_m^2$ belongs to a one-dimensional subgroup. Note that if u and v are not both roots of unity, then vectors $(p, q) \in \mathbf{Z}^2$ such that the above relation holds form a rank-one subgroup. The following Proposition 2 quantifies the above mentioned Corollary, in the sense that it permits to explicitly obtain the dependence relations satisfied by all but finitely many exceptions to the inequality (1.3).

Proposition 2. *Let K be a number field, S a finite set of places as before and $\epsilon > 0$ a real number. All but finitely many solutions $(u, v) \in (\mathcal{O}_S^\times)^2$ to the inequality*

$$\sum_{\mu \in M_K} \log^- \max\{|u - 1|_\mu, |v - 1|_\mu\} < -\epsilon \cdot \max\{h(u), h(v)\} \tag{2.8}$$

are contained in finitely many 1-dimensional subgroups of \mathbf{G}_m^2 which can be effectively determined. Namely they are defined by an equation of the form $u^p = v^q$ for coprime integers p, q with $\max\{|p|, |q|\} \leq \epsilon^{-1}$.

Proof of Proposition 2. Let us suppose that (u_i, v_i) is an infinite sequence in $(\mathcal{O}_S^\times)^2$ verifying the above inequality (2.8). We shall prove the existence of finitely many pairs $(p, q) \in \mathbf{Z}^2 \setminus \{(0, 0)\}$ such that for all but finitely many indices, u_i and v_i verify one dependence relation of the form $u^p v^q = 1$ (in particular for all but finitely many indices, u_i, v_i are multiplicatively dependent). We shall later show how to bound the size of p and q . In the sequel, we will drop the index i for convenience of notation.

First of all, notice that by Ridout’s theorem (the case $N = 2$ of the above Subspace Theorem) for all but finitely many $(u, v) \in \Gamma$ we have

$$\prod_{\mu \in S} \min\{1, |u - 1|_\mu\} > H(u)^{-\epsilon}, \quad \text{and} \quad \prod_{\mu \in S} \min\{1, |v - 1|_\mu\} > H(v)^{-\epsilon}.$$

Hence we clearly have that except for finitely many pairs $(u, v) \in (\mathcal{O}_S^\times)^2$

$$\sum_{\mu \in S} \log^- \max\{|u - 1|_\mu, |v - 1|_\mu\} > -\frac{\epsilon}{2} \cdot \max\{h(u), h(v)\};$$

so all but finitely many solutions of (2.8) also verify

$$\begin{aligned} \sum_{\mu \in M_K \setminus S} \log^- \max\{|u - 1|_\mu, |v - 1|_\mu\} &= \sum_{\mu \in M_K \setminus S} \log \max\{|u - 1|_\mu, |v - 1|_\mu\} \\ &< -\frac{\epsilon}{2} \cdot \max\{h(u), h(v)\}. \end{aligned} \tag{2.8'}$$

We can suppose (since the expressions (2.8), (2.8') are symmetric in u, v) that $H(v) \geq H(u)$. Let us denote by S^+ (resp. S^-), the subset of S made up of those absolute values μ such that $|v|_\mu > 1$ (resp. $|v|_\mu \leq 1$). Since there are only finitely many choices for the pair S^+, S^- , we can partition the solutions in a finite number of subsets for which S^+ and S^- are constant. From now on, we shall work separately with each subset.

Now the proof follows the same lines as [7]. Let (u, v) be a pair in $(\mathcal{O}_S^\times)^2$, satisfying inequality (2.8'). We write, for a positive integer j ,

$$z_j(u, v) = z_j := \frac{u^j - 1}{v - 1}. \tag{2.9}$$

Note that

$$z_j = z_1 \cdot (u^{j-1} + \dots + u + 1). \tag{2.10}$$

Fix an integer $h > 0$ and consider the identity

$$\frac{1}{v - 1} = \frac{1}{v} \cdot \frac{1}{1 - v^{-1}} = \frac{1}{v} \left(1 + v^{-1} + \dots + v^{-h+1} + \frac{v^{-h}}{1 - v^{-1}} \right).$$

Let us fix a second integer $k > 0$. Then for $j \in \{1, \dots, k\}$ we obtain, on multiplying by $u^j - 1$ in the above identity,

$$z_j = (u^j - 1) \cdot \left(v^{-1} + v^{-2} + \dots + v^{-h} + \frac{v^{-h-1}}{v - 1} \right).$$

For $\mu \in S^+$, we then derive the inequality (in fact an equality)

$$|z_j - u^j v^{-1} - \dots - u^j v^{-h} + v^{-1} + \dots + v^{-h}|_\mu \leq |u^j - 1|_\mu \cdot \frac{|v|_\mu^{-h-1}}{|1 - v|_\mu}. \tag{2.11}$$

We put $N = hk + h + k$; for convenience we shall write vectors in K^N as

$$\mathbf{x} = (x_1, \dots, x_N) = (z_1, \dots, z_k, y_{0,1}, \dots, y_{0,h}, \dots, y_{k,1}, \dots, y_{k,h}).$$

In this notation we choose linear forms with integral coefficients as follows. For $j = 1, \dots, k$, and $\mu \in S^+$, we put

$$L_{j,\mu} = z_j + y_{0,1} + \dots + y_{0,h} - y_{j,1} - \dots - y_{j,h},$$

while, for $(j, \mu) \notin \{1, \dots, k\} \times S^+$ we put

$$L_{j,\mu} = x_j.$$

Observe that for each $\mu \in S$, the linear forms are indeed linearly independent. For a given pair (u, v) we also set

$$P = P(u, v) = (z_1, \dots, z_k, v^{-1}, \dots, v^{-h}, uv^{-1}, \dots, uv^{-h}, \dots, u^k v^{-1}, \dots, u^k v^{-h}).$$

Note that all the coordinates of P apart from the first k of them are S -units.

To apply the Subspace Theorem we have to estimate the double product $\prod_{j=1}^N \prod_{\mu \in S} \frac{|L_{j,\mu}(P)|_\mu}{|P|_\mu}$ at points $P = P(u, v)$. First note that for $j > k$,

$$\prod_{\mu \in S} |L_{j,\mu}(P)|_\mu = 1$$

since all the involved linear forms are projections on the coordinates, which are S -units. Then

$$\prod_{j=k+1}^N \prod_{\mu \in S} \frac{|L_{j,\mu}(P)|_\mu}{|P|_\mu} = \prod_{\mu \in S} |P|_\mu^{-N+k}. \tag{2.12}$$

Let now j be an index in $\{1, \dots, k\}$. For $\mu \in S^-$, we recall that $L_{j,\mu}(P) = z_j = (u^j - 1)/(v - 1)$, so in particular

$$\frac{|L_{j,\mu}(P)|_\mu}{|P|_\mu} \leq \frac{|u^j - 1|_\mu}{|v - 1|_\mu} \cdot |P|_\mu^{-1}. \tag{2.13}$$

For $\mu \in S^+$ we use (2.11) which gives, since $L_{j,\mu}(P)$ coincides with the left hand side term of (2.11),

$$|L_{j,\mu}(P)|_\mu \leq |u^j - 1|_\mu \cdot \frac{|v|_\mu^{-h}}{|1 - v|_\mu}$$

so, since $|u^j - 1|_\mu \leq \max\{1, |2|_\mu\} \cdot \max\{1, |u|_\mu\}^j$,

$$\frac{|L_{j,\mu}(P)|_\mu}{|P|_\mu} \leq \max\{1, |2|_\mu\} \cdot \max\{1, |u|_\mu\}^j \cdot \frac{1}{|1 - v|_\mu} \cdot |v|_\mu^{-h} \cdot |P|_\mu^{-1}. \tag{2.14}$$

Then, from (2.13), (2.14), we have

$$\prod_{j=1}^k \prod_{\mu \in S} \frac{|L_{j,\mu}(P)|_\mu}{|P|_\mu} \leq \left(\prod_{j=1}^k \prod_{\mu \in S} \max\{1, |2|_\mu\} \cdot \max\{1, |u|_\mu\}^j \cdot \frac{1}{|1 - v|_\mu} \cdot |P|_\mu^{-1} \right) \cdot \left(\prod_{\mu \in S^+} |v|_\mu^{-h} \right)^k.$$

Now, using the fact that $\prod_{\mu \in S} \max\{1, |2|_\mu\} \leq H(2) = 2$ (and the analogue estimates for the products of $\max\{1, |u|_\mu\}$ and $\frac{1}{|1-v|_\mu}$), and using also the equality $H(v) = \prod_{\mu \in S^+} |v|_\mu$, we obtain

$$\prod_{j=1}^k \prod_{\mu \in S} \frac{|L_{j,\mu}(P)|_\mu}{|P|_\mu} \leq H(v)^{-hk} (2H(u))^{1+\dots+k} H(1-v)^k \prod_{\mu \in S} |P|_\mu^{-k}. \tag{2.15}$$

Finally, from (2.12), (2.15) we obtain

$$\prod_{j=1}^N \prod_{\mu \in S} \frac{|L_{j,\mu}(P)|_\mu}{|P|_\mu} \leq \left(\prod_{\mu \in S} |P|_\mu^{-N+k} \right) H(v)^{-hk} (2H(u))^{k^2} H(1-v)^k \prod_{\mu \in S} |P|_\mu^{-k}.$$

Observe that the product of the first and last factors can be written as

$$\left(\prod_{\mu \in S} |P|_\mu \right)^{-N} = H(P)^{-N} \cdot \left(\prod_{\mu \notin S} |P|_\mu \right)^N.$$

Hence we can rewrite the above inequality as

$$\prod_{j=1}^N \prod_{\mu \in S} \frac{|L_{j,\mu}(P)|_\mu}{|P|_\mu} \leq H(P)^{-N} \left(\prod_{\mu \notin S} |P|_\mu \right)^N H(v)^{-hk} (2H(u))^{k^2} H(1-v)^k. \tag{2.16}$$

Our next goal will be to estimate the quantity $\prod_{\mu \notin S} |P|_\mu$. Now, observe that the only coordinates which are not S -integers are the first k , i.e. z_1, \dots, z_k , and, for $j = 1, \dots, k$,

$$z_j = \frac{u^j - 1}{v - 1} = z_1 \cdot (u^{j-1} + u^{j-2} + \dots + u + 1);$$

also, the factor $u^{j-1} + \dots + u + 1$ is an S -integer, so the only contribution comes from the first factor $z_1 = (u - 1)/(v - 1)$; hence we have the bound

$$\prod_{\mu \notin S} |P|_\mu \leq \prod_{\mu \notin S} \max \left\{ 1, \left| \frac{u - 1}{v - 1} \right|_\mu \right\}.$$

Now, we have

$$\begin{aligned} \prod_{\mu \notin S} \max \left\{ 1, \left| \frac{u - 1}{v - 1} \right|_\mu \right\} &= \prod_{\mu \notin S} \frac{1}{|v - 1|_\mu} \cdot \max\{|u - 1|_\mu, |v - 1|_\mu\} \\ &\leq H(v - 1) \prod_{\mu \notin S} \max\{|u - 1|_\mu, |v - 1|_\mu\}; \end{aligned}$$

we now use the hypothesis that our pairs (u, v) satisfy (2.8') so that

$$\prod_{\mu \notin S} \max\{|u - 1|_\mu, |v - 1|_\mu\} < H(v)^{-\epsilon/2},$$

so finally

$$\prod_{\mu \notin S} |P|_\mu \leq H(v - 1) \cdot H(v)^{-\epsilon/2}.$$

Using the above estimate and the fact that $H(1 - v) \leq 2H(v)$, we can rewrite (2.16) as

$$\begin{aligned} \prod_{j=1}^N \prod_{\mu \in S} \frac{|L_{j,\mu}(P)|_\mu}{|P|_\mu} &\leq H(v)^{-hk} H(P)^{-N} H(v)^{(1-\epsilon/2)N} 2^N (2H(u))^{k^2} (2H(v))^k \\ &\leq H(v)^{-hk} H(P)^{-N} H(v)^{(1-\epsilon/2)N} 2^N (2H(v))^{k^2+k}. \end{aligned} \tag{2.17}$$

Since $N = hk + h + k$, the exponent of $H(v)$ can be estimated (at least when $k \geq 2$), by

$$\left(1 - \frac{\epsilon}{2}\right)N + k^2 + k - hk < -\frac{\epsilon}{2}hk + h + k + k^2 + k \leq -\frac{\epsilon}{2}hk + h + 2k^2.$$

By choosing $k > \frac{4}{\epsilon}$ we obtain that for large enough h (in particular $h > 2k^2 + 1$), $\epsilon hk/2 - h - 2k^2 =: \epsilon_0 > 0$. Choose k, h in such a way. From (2.17) we then obtain that the double product is bounded as

$$\prod_{j=1}^N \prod_{\mu \in S} \frac{|L_{j,\mu}(P)|_\mu}{|P|_\mu} \leq H(P)^{-N} H(v)^{-\epsilon_0} \cdot 2^{N+k+k^2}. \tag{2.18}$$

Our next goal, in view of the application of the Subspace Theorem, will be to compare the height of P with the height of v . A rough estimate gives

$$H(P) \leq H(u)^k H(v)^h H(1-v) \leq H(u)^k 2H(v)^{h+1} \leq H(v)^{h+k+2}$$

at least for $H(v) \geq 2$, which we may suppose. Hence

$$H(v)^{-\epsilon_0} \leq H(P)^{\frac{-\epsilon_0}{h+k+2}}$$

so that, for $\delta := \frac{\epsilon_0}{h+k+3}$ and large values of $H(v)$, we obtain from (2.18)

$$\prod_{j=1}^N \prod_{\mu \in S} \frac{|L_{j,\mu}(P)|_\mu}{|P|_\mu} \leq H(P)^{-N-\delta}.$$

Now we are able to apply the Subspace Theorem, which implies that there exist finitely many hyperplanes containing all the points P . Choose one such hyperplane, containing infinitely many points P . Then we obtain an equation of the kind

$$\eta_k \frac{u^k - 1}{v - 1} + \dots + \eta_1 \frac{u - 1}{v - 1} + \sum_{i,j} \rho_{i,j} u^j v^i = 0, \tag{2.19}$$

where the sum runs over the pairs $(i, j) \in \{-1, \dots, -h\} \times \{0, \dots, k\}$. Here η_1, \dots, η_k and the $\rho_{i,j}$ are elements of the number field K , not all zero. By multiplying by $v^h(v - 1)$ we obtain that the point (u, v) lies in the affine curve given by the equation

$$\eta_k V^h (U^k - 1) + \dots + \eta_1 V^h (U - 1) + (V - 1) \sum_{i,j} \rho_{i,j} U^j V^{h+i} = 0.$$

We pause to show that this equation is non trivial, so it defines in fact a curve. If the left hand side term vanishes identically, then the binomial $(V - 1)$ would divide $\eta_k V^h (U^k - 1) + \dots + \eta_1 V^h (U - 1)$, which implies that $\eta_1 = \dots = \eta_k = 0$. But then all coefficients $\rho_{i,j}$ would vanish, contrary to the assumption.

In view of (2.19), infinitely many points (u, v) lie in the curve defined by the above equation. Then, by the mentioned theorem of Lang [11, Thm. 7.3, p. 207], all such points lie in the union of finitely many translates of subtori of \mathbf{G}_m^2 ; every translate of a proper subtorus is either a point or a curve defined by an equation of the form (2.5) for coprime p, q . The remaining part of the proof is similar to

the last part of the proof of Proposition 1: Fix such a translate and suppose it contains infinitely many solutions $(u, v) \in (\mathcal{O}_S^\times)^2$ to the inequality (2.8). For each $(u, v) \in (\mathcal{O}_S^\times)^2$, write

$$u = t^q, \quad v = s^{-p}$$

where t, s lie in a fixed number field \tilde{K} . As in the proof of Proposition 1, we arrive at the parametrization (2.6). Then

$$\frac{u - 1}{v - 1} = \frac{t^q - 1}{\bar{w}t^{-p} - 1} =: \varphi(t), \tag{2.20}$$

say.

Now, we distinguish four cases:

First Case: $pq > 0$ and $\bar{w}^q \neq 1$. In this case the rational function φ in (2.20), which can be written as $\varphi(t) = -t^p \cdot \frac{t^q - 1}{t^p - \bar{w}}$, has degree $|p| + |q|$ (note that the numerator and denominator are coprime polynomials in t , or in t^{-1}). So, in this case, we have $H((u - 1)/(v - 1)) \gg H(u)H(v)$, i.e.

$$h((u - 1)/(v - 1)) \geq h(u) + h(v) + O(1)$$

where the implied constant in the $O(1)$ term does not depend on u, v . Since

$$\begin{aligned} h((u - 1)/(v - 1)) &= h(u - 1 : v - 1) \\ &\leq h(u) + h(v) + \sum_{\mu \in M_K} \log^- \max\{|u - 1|_\mu, |v - 1|_\mu\} + O(1), \end{aligned}$$

we obtain from the above inequalities and from (2.8) a uniform bound for the $\max\{h(u), h(v)\}$, contrary to the assumption that the given subgroup contains infinitely many solutions to the inequality (2.8).

Second Case: $pq > 0$ and $\bar{w}^q = 1$. First of all, note that in this case $w = \bar{w}^q = 1$, so that the dependence relation (2.5) takes the form $u^p v^q = 1$. Also, the rational function $\varphi(t)$ has degree $|p| + |q| - 1$, since the polynomials $t^p(t^q - 1)$, and $t^p - \bar{w}$ have a greatest common divisor of degree one. Then from the parametrization (2.6), we clearly obtain

$$h(1 : u - 1 : v - 1) \leq (|p| + |q|)h(t) + O(1)$$

and from Lemma 1

$$h((u - 1)/(v - 1)) \geq (|p| + |q| - 1)h(t).$$

Since we always have

$$\begin{aligned} h((u - 1)/(v - 1)) &= h(u - 1 : v - 1) \\ &\leq h(1 : u - 1 : v - 1) + \sum_{\mu \in M_K} \log^- \max\{|u - 1|_\mu, |v - 1|_\mu\} \end{aligned}$$

we deduce

$$(|p| + |q| - 1)h(t) \leq (|p| + |q|)h(t) + \sum_{\mu \in M_K} \log^- \max\{|u - 1|_\mu, |v - 1|_\mu\} + O(1)$$

from which

$$- \sum_{\mu \in M_K} \log^- \max\{|u - 1|_\mu, |v - 1|_\mu\} \leq h(t) + O(1).$$

Since $h(t) = \max\{h(u), h(v)\}(\max\{|p|, |q|\})^{-1}$, the above inequality can be satisfied by infinitely many solutions of (2.8) only if $\max\{|p|, |q|\} \leq \epsilon^{-1}$, as wanted.

Third Case: $pq < 0$ and $\bar{w}^q \neq 1$. Now, the rational function φ has degree $\max\{|p|, |q|\}$, so, by Lemma 1, we have $H((u - 1)/(v - 1)) \gg \max\{H(u), H(v)\}$, i.e.

$$h(u - 1 : v - 1) \geq \max\{h(u), h(v)\} + O(1). \tag{2.21}$$

Note that in this case it follows from the parametrization (2.6) that u and v are, apart from a constant \bar{w} , positive powers of a same element t (or t^{-1}). From this fact it easily follows that

$$h(1 : u : v) = \max\{h(u), h(v)\} + O(1). \tag{2.22}$$

Using again the inequality

$$h(u - 1 : v - 1) \leq h(1 : u : v) + \sum_{\mu \in M_K} \log^- \max\{|u - 1|_\mu, |v - 1|_\mu\}$$

we obtain from (2.21) and (2.22)

$$\begin{aligned} \max\{h(u), h(v)\} + O(1) &\leq h(u - 1 : v - 1) \\ &\leq \max\{h(u), h(v)\} + \sum_{\mu \in M_K} \log^- \max\{|u - 1|_\mu, |v - 1|_\mu\}. \end{aligned}$$

Then from (2.8) we obtain, as before, a uniform upper bound for the $\max\{h(u), h(v)\}$, contrary to the assumption that our subgroup contains infinitely many solutions of (2.8).

Fourth Case: $pq < 0$ and $\bar{w}^q = 1$. Then from the parametrization (2.6) we obtain, as in the Second Case, $u^p v^q = 1$, i.e. a multiplicative dependence relation as wanted.

Now, since p, q are coprime and of opposite sign, the rational function $\varphi(t) = (t^q - 1)/(t^{-p} - 1)$ has degree $d = \max\{|q| - 1, |p| - 1\}$. Then the Lemma 1 gives

$$\begin{aligned} h((u - 1)/(v - 1)) &\geq \max\{|q| - 1, |p| - 1\}h(t) + O(1) \\ &= (\max\{|q|, |p|\} - 1)h(t) + O(1) \end{aligned}$$

On the other hand, the above estimates and (2.8) give

$$h(u - 1 : v - 1) \leq \max\{|q|, |p|\}h(t) - \epsilon \max\{|q|, |p|\}h(t),$$

so that we finally obtain

$$(\max\{|q|, |p|\} - 1)h(t) + O(1) \leq \max\{|q|, |p|\}h(t) - \epsilon \max\{|q|, |p|\}h(t)$$

which gives $\max\{|q|, |p|\} \leq \epsilon^{-1}$, as wanted.

Notice that we have also proved that, in the only cases when infinitely many solutions can occur (i.e. the second and fourth cases), the constant w in (2.5) equals 1. In other terms, the minimal dependence relation has coprime exponents, i.e. the infinite families of solutions of (2.8) are contained in finitely many *connected* one-dimensional subgroups.

From Proposition 2 we easily obtain the following statement:

Proposition 3. *Let S , as before, be a finite set of places of a number field K containing the archimedean ones, θ, η be nonzero elements of K , $\epsilon > 0$. The solutions $(u, v) \in (\mathcal{O}_S^\times)^2$ to the inequality*

$$\sum_{\mu \in M_K} \log^- \max\{|u - \theta|_\mu, |v - \eta|_\mu\} < -\epsilon \cdot \max\{h(u), h(v)\}$$

are contained in finitely many translates of proper subgroups of \mathbf{G}_m^2 . Those of dimension one can be effectively determined.

Proof. Fix $\theta, \eta \in K^\times$. By enlarging S , we can suppose that they both are S -units. Now, note that

$$\begin{aligned} \sum_{\mu \in M_K} \log^- \max\{|u - \theta|_\mu, |v - \eta|_\mu\} \\ = \sum_{\mu \in M_K} \log^- \max\{|u\theta^{-1} - 1|_\mu, |v\eta^{-1} - 1|_\mu\} + O(1). \end{aligned}$$

Then Proposition 2 implies that apart from finitely many exceptions, the pairs $(u\theta^{-1}, v\eta^{-1})$ are contained in finitely many one-dimensional subgroups, which can be effectively determined; this in turn implies that the pairs (u, v) , apart from finitely many of them, are contained in finitely many translates of one-dimensional subgroups, which can be effectively determined.

The following Proposition represents a particular but crucial case of Theorem 1.

Proposition 4. *Let $r(X), s(X) \in \bar{\mathbf{Q}}[X]$ be two non zero polynomials. Then for every $\epsilon > 0$, all but finitely many solutions $(u, v) \in (\mathcal{O}_S^\times)^2$ to the inequality*

$$\sum_{\mu \in M_K \setminus S} \log^- \max\{|r(u)|_\mu, |s(v)|_\mu\} < -\epsilon \cdot (\max\{h(u), h(v)\}) \tag{2.23}$$

are contained in finitely many translates of one-dimensional subgroups of \mathbf{G}_m^2 , which can be effectively determined. Moreover, if $r(X)$ and $s(X)$ do not both vanish at 0, the same conclusion holds for the solutions to the inequality

$$\sum_{\mu \in M_K} \log^- \max\{|r(u)|_\mu, |s(v)|_\mu\} < -\epsilon \cdot (\max\{h(u), h(v)\}) \tag{2.24}$$

Proof. We begin by observing that the points (u, v) such that $r(u) \cdot s(v) = 0$ are contained in a finite union of translates of proper subtori; hence we will tacitely disregard these pairs. Also, for each constant C , the pairs $(u, v) \in (\mathcal{O}_S^\times)^2$ such that

$\max\{h(u), h(v)\} \leq C$ are finite in number, so they certainly lie in finitely many translates of proper subtori. For this reason, we can restrict our attention to “large” solutions of the inequality (2.23) or (2.24). Let us write

$$\begin{aligned} r(X) &= r_0 X^a (X - \theta_1)^{a_1} \cdots (X - \theta_m)^{a_m}, \\ s(X) &= s_0 X^b (X - \eta_1)^{b_1} \cdots (X - \eta_n)^{b_n}; \end{aligned}$$

here a and b are non-negative integers with $ab = 0$, $a_1, \dots, a_m, b_1, \dots, b_n$ are positive integers and $\theta_1, \dots, \theta_m$, (*resp.* η_1, \dots, η_n) are pairwise distinct nonzero algebraic numbers in a number field K . We shall first prove the first part of the Proposition, so we shall estimate the relevant sum for μ running in the complement of S . Let, for each place $\mu \in M_K \setminus S$, δ_μ be the minimum of the set

$$\begin{aligned} &\{|\theta_i - \theta_j|_\mu, \text{ for } (i, j) \in \{1, \dots, m\} \times \{1, \dots, m\}, (i \neq j)\} \\ &\cup \{|\eta_i - \eta_j|_\mu, \text{ for } (i, j) \in \{1, \dots, n\} \times \{1, \dots, n\}, (i \neq j)\} \\ &\cup \{|\theta_1|_\mu, \dots, |\theta_m|_\mu, |\eta_1|_\mu, \dots, |\eta_n|_\mu\}. \end{aligned}$$

Clearly, $\delta_\mu > 0$ for all μ and $\delta_\mu = 1$ for all but finitely many places. Let us divide the places of $M_K \setminus S$ in two classes \mathcal{A} and \mathcal{B} : namely,

$$\begin{aligned} \mathcal{A} := &\left\{ \mu \in M_K \mid \text{there exist } i \in \{1, \dots, m\} \text{ and } j \in \{1, \dots, n\} \right. \\ &\left. \text{such that } |u - \theta_i|_\mu < \frac{\delta_\mu}{2} \text{ and } |v - \eta_j|_\mu < \frac{\delta_\mu}{2} \right\} \end{aligned}$$

while \mathcal{B} is its complement in the subset of $M_K \setminus S$ composed by the places μ such that $\max\{|r(u)|_\mu, |s(v)|_\mu\} < 1$. Then the left hand side of (2.23) becomes

$$\begin{aligned} \sum_{\mu \in M_K \setminus S} \log^-(\max\{|r(u)|_\mu, |s(v)|_\mu\}) &= \sum_{\mu \in \mathcal{A}} \log^-(\max\{|r(u)|_\mu, |s(v)|_\mu\}) \\ &+ \sum_{\mu \in \mathcal{B}} \log^-(\max\{|r(u)|_\mu, |s(v)|_\mu\}). \end{aligned}$$

The second term is simply bounded independently of u, v by

$$\begin{aligned} \sum_{\mu \in \mathcal{B}} \log^-(\max\{|r(u)|_\mu, |s(v)|_\mu\}) &\geq -h(r_0) - h(s_0) \\ &+ \max\{\deg r, \deg s\} \sum_{\mu \in \mathcal{B}} \log^-\frac{\delta_\mu}{2}. \end{aligned} \tag{2.25}$$

To estimate the first term, let us notice that for each $\mu \in \mathcal{A}$ there exists exactly one pair $(i_\mu, j_\mu) \in \{1, \dots, m\} \times \{1, \dots, n\}$ such that $|u - \theta_{i_\mu}|_\mu < \frac{\delta_\mu}{2}$, and $|v - \eta_{j_\mu}|_\mu < \frac{\delta_\mu}{2}$. Then, for each such place μ , we can bound

$$\begin{aligned} \log^-\max\{|r(u)|_\mu, |s(v)|_\mu\} &\geq \log^-|r_0|_\mu + \log^-|s_0|_\mu + \max\{\deg r, \deg s\} \log^-\frac{\delta_\mu}{2} \\ &+ \log^-\max\{|u - \theta_{i_\mu}|_\mu^{a_{i_\mu}}, |v - \eta_{j_\mu}|_\mu^{b_{j_\mu}}\}, \end{aligned} \tag{2.26}$$

where we have used the fact that $|u|_\mu^a = |v|_\mu^b = 1$ for $\mu \in M_K \setminus S$. Again the quantity

$$-\sum_{\mu \in \mathcal{A}} \left(\log^- |r_0|_\mu + \log^- |s_0|_\mu + \max\{\deg r, \deg s\} \log^- \frac{\delta_\mu}{2} \right)$$

is bounded independently of u, v simply by $h(r_0) + h(s_0) + \max\{\deg(r), \deg(s)\} \sum_{\mu \in \mathcal{A}} \log^- \frac{\delta_\mu}{2}$. It remains to estimate the term

$$\sum_{\mu \in \mathcal{A}} \log^- \max\{|u - \theta_{i_\mu} |_\mu^{a_{i_\mu}}, |v - \eta_{j_\mu} |_\mu^{b_{j_\mu}}\}$$

in the quantity appearing in (2.26). This can be rewritten as

$$\sum_{(i,j) \in \{1, \dots, m\} \times \{1, \dots, n\}} \sum_{\mu \in \mathcal{A}_{i,j}} \log^- \max\{|u - \theta_i |_\mu^{a_{i_\mu}}, |v - \eta_j |_\mu^{b_{j_\mu}}\}$$

where we define $\mathcal{A}_{i,j}$ to be the subset of \mathcal{A} composed of the places μ for which $(i_\mu, j_\mu) = (i, j)$. Proposition 3, applied with $\epsilon/(2(\deg r)(\deg s))$ instead of ϵ , gives for each pair (i, j) the upper bound

$$\sum_{\mu \in \mathcal{A}_{i,j}} \log^- \max\{|u - \theta_i |_\mu, |v - \eta_j |_\mu\} \geq -\frac{\epsilon}{2(\deg r)(\deg s)} \cdot \max\{h(u), h(v)\} \quad (2.27)$$

for $(u, v) \in (\mathcal{O}_S^\times)^2$ outside a finite union of translates of proper subgroups, of which those of dimension one can be effectively determined. Then, outside the union of such exceptional subvarieties, we get from (2.25), (2.26) and (2.27),

$$\sum_{\mu \in M_K \setminus S} \log^- \max\{|r(u)|_\mu, |s(v)|_\mu\} \geq -\frac{\epsilon}{2} \max\{h(u), h(v)\} + O(1). \quad (2.28)$$

This clearly implies the first part of Proposition 4. We shall now estimate the relevant sum for μ running over the places in S . Recall that by hypothesis one at least between the exponents a and b vanishes. Suppose that $b = 0$, say. We shall apply Ridout’s theorem (i.e. the case $N = 2$ of the Subspace Theorem) to give a lower bound for $\log^- |s(v)|_\mu$, valid for all but finitely many v (of course, in our situation, we could also apply the stronger Baker’s bound for linear forms in logarithms). For each $\mu \in S$, we denote by $j_\mu \in \{1, \dots, n\}$ an index such that for all $j = 1, \dots, n$

$$|v - \eta_{j_\mu} |_\mu \leq |v - \eta_j |_\mu.$$

Then

$$|s(v)|_\mu \geq |s_0|_\mu \cdot |v - \eta_{j_\mu} |_\mu^{\deg s}.$$

By Ridout’s theorem we have that for all but finitely many S -units $v \in \mathcal{O}_S^\times$,

$$|v - \eta_{j_\mu} |_\mu > H(v)^{-\frac{\epsilon}{3(\deg s)|S|}} \geq \max\{H(u), H(v)\}^{-\frac{\epsilon}{3(\deg s)|S|}}$$

so that, for all but finitely many $v \in \mathcal{O}_S^\times$,

$$\begin{aligned} \sum_{\mu \in S} \log^- \max\{|r(u)|_\mu, |s(v)|_\mu\} &\geq \sum_{\mu \in S} \log^- |s(v)|_\mu \\ &\geq -\frac{\epsilon}{3} \max\{h(u), h(v)\} + O(1). \end{aligned} \tag{2.29}$$

Putting together (2.28) and (2.29) we obtain that outside the union of a finite set and a finite number of effectively computable translates of one dimensional subgroups we have

$$\sum_{\mu \in S} \log^- \max\{|r(u)|_\mu, |s(v)|_\mu\} \geq -\frac{5}{6}\epsilon \cdot \max\{h(u), h(v)\} + O(1)$$

from which the Proposition follows immediately.

Proof of Theorem 1. We start by remarking that it suffices to prove that the relevant set is contained in a finite union of translates of proper subtori of \mathbf{G}_m^2 . This is very easy to see directly, since we are working in dimension 2; in any case, this equivalence follows from a well-known theorem of M. Laurent (see e.g. [14]). This remark applies also to the Main Theorem.

To go on with the proof, we may clearly assume that the polynomials $p(X, Y), q(X, Y)$ are defined over a number field K and that Γ is the group $(\mathcal{O}_S^\times)^2$ for a suitable finite subset S of M_K .

We first slightly transform the relevant inequality. Observe that, by definition,

$$\begin{aligned} h(p(u, v) : q(u, v) : 1) &= \sum_{\mu \in M_K} \log \max\{|p(u, v)|_\mu, |q(u, v)|_\mu, 1\} \\ &= \sum_{\mu \in M_K} \max\{0, \log \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\}\}. \end{aligned}$$

Then, using the formula $\max\{0, \log(\cdot)\} = \log^+(\cdot) = \log(\cdot) - \log^-(\cdot)$, we can write

$$\begin{aligned} h(p(u, v) : q(u, v) : 1) &= \sum_{\mu \in M_K} \log \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\} \\ &\quad - \sum_{\mu \in M_K} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\} \\ &= h(p(u, v) : q(u, v)) \\ &\quad - \sum_{\mu \in M_K} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\}. \end{aligned}$$

Note that $h(p(u, v)/q(u, v)) = h(p(u, v) : q(u, v))$ so the Theorem is reduced to proving that the solutions to the inequality

$$\sum_{\mu \in M_K} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\} < -\epsilon \max\{h(u), h(v)\} \tag{2.30}$$

are contained in finitely many translates of proper subtori of \mathbf{G}_m^2 , of which those of dimension one can be effectively determined. We shall prove this contention, by

estimating separately the two subsums $\sum_{\mu \in M_K \setminus S} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\}$ and $\sum_{\mu \in S} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\}$.

Let us begin with the second. By hypothesis not both $p(X, Y)$ and $q(X, Y)$ vanish at the origin; suppose, for instance, that $p(0, 0) \neq 0$. We apply Proposition 1, with $f(X, Y) = p(X, Y)$ and $\epsilon/2$ instead of ϵ . Then (the second part of) Proposition 1 states that, outside the union of a finite set and finitely many effectively computable translates of one dimensional subgroups, one has

$$\sum_{\mu \in S} \log^- |p(u, v)|_\mu > -\frac{\epsilon}{2} \max\{h(u), h(v)\}$$

so clearly

$$\sum_{\mu \in S} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\} > -\frac{\epsilon}{2} \max\{h(u), h(v)\}. \tag{2.31}$$

To estimate the sum for μ running over the complement of S , we argue as follows: Since $p(X, Y), q(X, Y)$ are coprime polynomials, there exist polynomials $A(X, Y), B(X, Y), C(X, Y), D(X, Y) \in \mathcal{O}_S[X, Y]$ and nonzero polynomials $r(X), s(Y)$ such that

$$\begin{aligned} A(X, Y)p(X, Y) + B(X, Y)q(X, Y) &= r(X) \\ C(X, Y)p(X, Y) + D(X, Y)q(X, Y) &= s(Y) \end{aligned}$$

Clearly, for a place $\mu \notin S$ and S -unit point (u, v) , we can bound by 1 the quantities $|A(u, v)|_\mu, |B(u, v)|_\mu, |C(u, v)|_\mu, |D(u, v)|_\mu$. Then

$$\sum_{\mu \in M_K \setminus S} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\} \geq \sum_{\mu \in M_K \setminus S} \log^- \max\{|r(u)|_\mu, |s(v)|_\mu\}.$$

The last quantity can be bounded, using Proposition 4, by $-(\epsilon/2) \max\{h(u), h(v)\}$ on the S -unit points (u, v) outside a finite union of translates of proper subgroups of \mathbf{G}_m^2 . This inequality, combined with (2.31), concludes the proof of Theorem 1.

Proof of Corollary 1. We could derive Corollary 1 both from Theorem 1 and from Proposition 2. We choose the latter possibility. We always have

$$\begin{aligned} h((u-1)/(v-1)) &= h(u-1 : v-1) \\ &= \sum_{\mu \in M_K} \log \max\{|u-1|_\mu, |v-1|_\mu\} \\ &\leq \sum_{\mu \in M_K} \log \max\{1, |u-1|_\mu, |v-1|_\mu\} = h(1 : u-1 : v-1). \end{aligned}$$

Also, $h(1 : u-1 : v-1) \leq h(1 : u : v) + O(1)$. So in particular

$$\limsup_{(u,v)} \frac{h((u-1)/(v-1))}{h(1 : u : v)} \leq 1.$$

We shall prove the non trivial inequality

$$\liminf_{(u,v) \in \Gamma} \frac{h((u-1)/(v-1))}{h(1 : u : v)} \geq 1,$$

where now the limit is taken only over the multiplicative independent pairs $(u, v) \in \Gamma$. This means that given a positive ϵ , only finitely many such pairs should verify the inequality

$$h((u - 1)/(v - 1)) < (1 - \epsilon)h(1 : u : v).$$

As we have already remarked,

$$h((u - 1)/(v - 1)) \geq h(1 : u - 1 : v - 1) + \sum_{\mu \in M_K} \log^- \max\{|u - 1|_\mu, |v - 1|_\mu\}$$

and by Proposition 2 all but finitely many multiplicative independent pairs $(u, v) \in \Gamma$ satisfy

$$\sum_{\mu \in M_K} \log^- \max\{|u - 1|_\mu, |v - 1|_\mu\} \geq -\epsilon \max\{h(u), h(v)\} \geq -\epsilon h(1 : u : v).$$

Proof of Main Theorem. Let us write $f(X, Y) = p(X, Y)/q(X, Y)$ for polynomials $p(X, Y), q(X, Y)$ defined over a number field K . Note that the hypothesis that $1 \in \{T_1(X, Y), \dots, T_N(X, Y)\}$ appearing on the Main Theorem is equivalent to the hypothesis in Theorem 1 that $p(X, Y), q(X, Y)$ do not both vanish at $(0, 0)$. The Main Theorem then follows essentially from Proposition 1 and Theorem 1. We have just to compare the maximum of the heights of the monomials appearing in the development of $p(X, Y), q(X, Y)$ with the maximum of the heights of u and v . This is the object of Lemma 2.

First of all, note that (the elementary) Lemma 1 immediately implies the result in a stronger form if all the non constant monomials $T_i(X, Y)$ vanish on a same one-dimensional subtorus. In fact, in such a case, the function $f(X, Y)$ could be written as $f(X, Y) = \varphi(X^a Y^b)$ for a suitable one-variable rational function $\varphi(T) \in K(T)$ and a pair $(a, b) \in \mathbf{Z}^2$; in that case the proof is exactly the same as in the corresponding case of the proof of Proposition 1. So we suppose that the set of monomials $\{T_1, \dots, T_N\} \setminus \{1\}$ satisfies the hypothesis of Lemma 2.

Fix a positive ϵ . Since $h(f(u, v)) = h(p(u, v) : q(u, v))$, we have

$$h(f(u, v)) \geq \max\{h(p(u, v)), h(q(u, v))\} + \sum_{\mu \in M_K} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\}; \tag{2.32}$$

by Proposition 1 the first summand is bounded from below as

$$\max\{h(p(u, v)), h(q(u, v))\} \geq (1 - \epsilon/2) \max\{h(T_1(u, v)), \dots, h(T_N(u, v))\} \tag{2.33}$$

outside a finite union of translates of proper subgroups. From the proof of Theorem 1 (see inequality (2.31)), the second summand in (2.32) is bounded, outside a finite union of translates of proper subgroups of \mathbf{G}_m^2 , as

$$\sum_{\mu \in M_K} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\} > -\frac{\epsilon}{4 \max\{\deg_X p, \deg_Y p, \deg_X q, \deg_Y q\}} \max\{h(u), h(v)\}$$

so by Lemma 2 we have

$$\sum_{\mu \in M_K} \log^- \max\{|p(u, v)|_\mu, |q(u, v)|_\mu\} > -\frac{\epsilon}{2} \max\{h(T_1(u, v)), \dots, h(T_N(u, v))\}. \quad (2.34)$$

Putting together (2.32), (2.33) and (2.34) we obtain that the solutions to (1.4) are contained in a finite union of proper translates, as wanted. Using this fact and Lemma 2 we obtain (1.5).

Acknowledgements. After the present paper was written, Prof. J. Silverman kindly sent us a preprint exploring related questions. Concerning the formulations in terms of heights, he goes further than we, relating with certain cases of Vojta's conjecture: for instance he proves that Corollary 1 follows from the Vojta conjecture suitably applied to the blow-up of \mathbf{P}_1^2 at one point. A paper of Silverman in this respect is expected soon. The authors are grateful to the referee for his comments.

References

- [1] Ailon N, Rudnick Z (2004) Torsion points on curves and common divisors of $a^k - 1$ and $b^k - 1$. *Acta Arith* **113**: 31–38
- [2] Bugeaud Y, Corvaja P, Zannier U (2003) An upper bound for the g.c.d. of $a^n - 1$ and $b^n - 1$. *Math Z* **243**: 79–84
- [3] Bugeaud Y, Luca F (2004) A quantitative upper bound for the greatest prime factor of $(ab + 1)(ac + 1)(bc + 1)$. *Acta Arith* (to appear)
- [4] Fuchs C (2003) An upper bound for the g.c.d. of two linear recurring sequences. *Math Slovaca* **53**: 21–42
- [5] Hernandez S, Luca F (2004) On the largest prime factor of $(ab + 1)(ac + 1)(bc + 1)$. *Bol Soc Mat Mexicana* (to appear)
- [6] Corvaja P, Zannier U (1998) Diophantine equations with power sums and universal Hilbert sets. *Indag Math NS* **9**: 317–332
- [7] Corvaja P, Zannier U (2003) On the greatest prime factor of $(ab + 1)(ac + 1)$. *Proc Amer Math Soc* **131**: 1705–1709
- [8] Corvaja P, Zannier U (2002) Finiteness of integral values for the ratio of two linear recurrences. *Invent Math* **149**: 431–451
- [9] Evertse JH (1984) On sums of S -units and linear recurrences. *Compositio Math* **53**: 225–244
- [10] Hindry M, Silverman JH (2000) *Diophantine Geometry*. Berlin Heidelberg New York: Springer
- [11] Lang S (1983) *Fundamentals of Diophantine Geometry*. Berlin Heidelberg New York: Springer
- [12] Schmidt WM (1980) *Diophantine Approximation*. *Lect Notes Math* **785**. Berlin Heidelberg New York: Springer
- [13] Schmidt WM (1991) *Diophantine Approximations and Diophantine Equations*. *Lect Notes Math* **1467**. Berlin Heidelberg New York: Springer
- [14] Zannier U (2003) *Some applications of Diophantine Approximation to Diophantine Equations*. Udine: Forum Editrice

Authors' addresses: P. Corvaja, Dip. di Matematica e Informatica, Via delle Scienze, 33100 Udine, Italy, e-mail: corvaja@dimi.uniud.it; U. Zannier, Scuola Normale Superiore, Piazza dei Cavalieri 7, 56100 Pisa, Italy, e-mail: zannier@dimi.uniud.it