

Unique strong and strict solutions for inhomogeneous hyperbolic differential equations in Banach space

Salih Jawad¹

Received: 28 October 2016 / Accepted: 26 February 2018 / Published online: 6 March 2018
© Springer-Verlag GmbH Austria, part of Springer Nature 2018

Abstract The purpose of the present paper is the improvement of results for solutions of the inhomogeneous differential equation:

$$\begin{aligned}u'(t) + A(t)u(t) + f(t) &= 0, \quad t \in (t_1, t_2) \\ u(t_1) &= \varphi\end{aligned}$$

in reflexive Banach space X . For $f(s) \in C^1([t_1, t_2], X)$, Kato obtained a unique strict solution under some conditions on the operator family $\{A(t)\}_{t_1 \leq t \leq t_2}$ to ensure the hyperbolicity of the problem. In a previous paper, the author obtained in abstract Hilbert space H a unique strong solution $u(t) \in C^{0,1}([t_1, t_2], H) \cap D$ if $f \in BV([t_1, t_2], H)$ and a strict solution if additionally $f \in C^0([t_1, t_2], H)$. Here is $D = D((A(t)))$ independent of t . In the present paper we extend these results to reflexive Banach spaces X .

Keywords Evolution equations · Regularity of integrals · Abstract hyperbolic differential equations · Strong solutions · Strict solutions

Mathematics Subject Classification Primary 47D06, 47D03, 35L90 · Secondary 64G05, 58D25, 34K30, 34G20

Communicated by A. Constantin.

✉ Salih Jawad
jassir83@yahoo.de

¹ Elisenstr. 17, 30451 Hannover, Germany

1 Introduction

For the treatment of the inhomogeneous differential equation:

$$\begin{cases} u'(t) + A(t)u(t) + f(t) = 0, & t_1 < t < t_2 \\ u(t_1) = \varphi \end{cases} \quad (1.1)$$

in reflexive Banach space X , we use as usual the associated integral equation:

$$u(t) = U(t, t_1)\varphi - \int_{t_1}^t U(t, s)f(s)ds. \quad (1.2)$$

Here $U(t, s)$ is the evolution operator generated by the family of closed operators $\{A(t)\}_{t_1 \leq t \leq t_2}$ according to Kato [7], page 246 (see precisely the assumptions on $A(t)$ in Sect. 2).

The classic of Kato [6] from 1953 represents the beginning. After his introduction of the stability concept in his revision [7] from 1970 intense activities arose with respect to the homogeneous part of (1.1) while the inhomogeneous problem was remaining disregarded. So, this paper is devoted to this topic. For details on these activities around the homogeneous equation, we refer to the paper of Bárta [1].

The stability concept due to Kato means essentially the admission that the operators $A(t)$ generate C_0 -semigroups instead of the merely C_0 -semigroups of contractions at Kato [6].

As is to be observed, the integral term in (1.2) plays the leading role. In this context, we show in Theorem 3.1(I) for $f \in BV([t_1, t_2], X)$ first of all the relation:

$$v(t) := \int_{t_1}^t U(t, s)f(s)ds \in D, \quad t \in [t_1, t_2]. \quad (1.3)$$

This relation means nearness to the a.e. differentiability of $v(t)$. For comparison see Kato [7], page 255 and Pazy [8], page 148.

Further we also obtain in Theorem 3.1 a similar estimate to the estimate (3.2) for $\|A(t)v(t)\|$ in $[t_1, t_2]$ from Jawad [5].

The chapter after deals with the investigation of the differentiability of $v(t)$ by showing of the Lipschitz continuity of $v(t)$ so that $u(t) := U(t, t_1)\varphi - v(t)$ fullfils (1.1) strongly and uniquely for $f \in BV([t_1, t_2], X)$ and strictly for additionally $f \in C^0([t_1, t_2], X)$. Regarding the strict solution, that means the renunciation of the usually requirement of the continuously differentiability of $f(t)$ (see Kato 1953 [6], Theorem 5, and Kato 1970 [7], Theorem 7.2). Further we show the uniqueness of $u(t)$ even for absolutely instead of Lipschitz continuous $u(t)$.

Regarding the classification of the solution, we note :

1. Mild Solution: The solution $u(t)$ satisfies the integral equation (1.2). For that it is sufficient that $f \in L^1((t_1, t_2), X)$.
2. Strong Solution: $u(t)$ fullfils the differential equation (1.1) a.e. in $[t_1, t_2]$ with $u'(t) \in L^1((t_1, t_2), X)$ and $u(t) \in D, t \in [t_1, t_2]$. Pazy [8], page 109.

3. Classical Solution: $u(t) \in C^0([t_1, t_2], X) \cap C^1((t_1, t_2), X)$, $u(t) \in D$, $t \in [t_1, t_2]$. The differential equation (1.1) is satisfied on $(t_1, t_2]$, see Pazy [8], page 139.
4. Strict Solution: $u(t) \in C^0([t_1, t_2], D) \cap C^1([t_1, t_2], X)$ and $u(t)$ satisfies the differential equation on the whole $[t_1, t_2]$.

2 Preliminaries

In this chapter we state the conditions on the operator family $\{A(t)\}_{t_1 \leq t \leq t_2}$ as well as the resulting conclusions and tools:

- (1) The closed operators $A(t)$ are infinitesimal generators of C_0 semigroups on the reflexive Banach space X .
- (2) The operator family $\{A(t)\}_{t_1 \leq t \leq t_2}$ is stable, i.e. there exist constants $M \geq 1$ and ω such that:

$$\left\| \prod_{j=1}^k (\lambda + A(\tau_j))^{-1} \right\| \leq \frac{M}{(\lambda - \omega)^k} \tag{2.1}$$

for $\lambda > \omega$ and every finite sequence $t_1 \leq \tau_1 \leq \tau_2 \leq \dots \leq \tau_k \leq t_2$, $k = 1, 2, \dots$. The constants M, ω are called the stability constants.

- (3) The domains of definition $D(A(t))$ are independent of t , i.e. $D(A(t)) = D$, $t \in [t_1, t_2]$.
- (4) For every $x \in D$, $A(t)x$ is continuously differentiable in $[t_1, t_2]$.

These conditions are the basis for Theorem 7.2 of Kato [7]. They imply the following propositions:

- (a) For the operator family $\{A(t)\}_{t_1 \leq t \leq t_2}$ there exists the unique evolution operator $U(t, s) \in B(X)$, $t_1 \leq s \leq t \leq t_2$, with the following properties according to Kato [7] (Theorem 6.1, page 252):
- (b) For every $x \in X$, $U(t, s)x$ is jointly continuous in $t, s \in [t_1, t_2]$ with $U(s, s) = I$ and $\|U(t, s)\| \leq Me^{\omega(t-s)}$.
- (c) $U(t, s) = U(t, r)U(r, s)$, $s \leq r \leq t$.
- (d) $U(t, s)D \subset D$.
- (e) For $x \in D$ it is in $[t_1, t_2]$:

$$\frac{\partial}{\partial t} U(t, s)x = -A(t)U(t, s)x \tag{2.2}$$

$$\frac{\partial}{\partial s} U(t, s)x = U(t, s)A(s)x. \tag{2.3}$$

- (f) For every $y \in D$, $A(t)U(t, s)y$ and $U(t, s)A(s)y$ are continuous in t, s . As it is usual in the sources (see Kato [7], page 254, 255) we may also assume for simplicity and without loss of generality the existence of the inverses $A^{-1}(t)$, $t \in [t_1, t_2]$, on X .

By setting $S(t) := A(t)$ on page 253 from Kato [7], we have further:

(g) For every $x \in X$, the expression $A(t)U(t, s)A^{-1}(s)x$ is continuous in t, s (separately) and there is a constant k_1 such that:

$$\|A(t)U(t, s)A^{-1}(s)\| \leq k_1, \quad t, s \in [t_1, t_2]. \tag{2.4}$$

(h) For every $x \in X$, $A'(s)A^{-1}(s)x$ is continuous in s with:

$$\|A'(s)A^{-1}(s)\| \leq k_2 \quad \text{on } [t_1, t_2]. \tag{2.5}$$

For the proof of (h) see the author [5], Hilfssatz 2.3.

3 The regularity of v

Theorem 3.1 *Let the Banach space X be reflexive and the conditions (1)–(4) from the Preliminaries be satisfied. Then it holds:*

(I) *For $f \in BV([t_1, t_2], X)$ and every $t \in [t_1, t_2]$ it is:*

$$v(t) = \int_{t_1}^t U(t, s)f(s)ds \in D \tag{3.1}$$

with

$$\begin{aligned} \|A(t)v(t)\| &\leq \|[I - A(t)U(t, t_1)A^{-1}(t_1)]f(t_1)\| \\ &\quad + k_1k_2\left(\sup_{s \in [t_1, t_2]} \|f(s)\|\right)(t - t_1) + (2k_1 + 1) \int_{t_1}^t \|df(s)\| \end{aligned} \tag{3.2}$$

in $[t_1, t_2]$, where k_1, k_2 are from (2.4) and (2.5).

- (II) $v(t) \in C^{0,1}([t_1, t_2], X)$.
- (III) *If additionally $f \in C^0([t_1, t_2], X)$ then:*

$$A(t)v(t) \in C^0([t_1, t_2], X).$$

Proof : (I) For a fixed $t \in [t_1, t_2]$, our proceeding consists of the approximation of the integral $v(t)$ by a sequence of elements $v_n \in D, n = 1, 2, \dots$. Afterwards we show the boundedness of $A(t)v_n$ so that finally $v(t) \in D$ because X is reflexive. The estimate (3.2) is then a consequence of the weak convergence:

$$A(t)v_n \rightharpoonup A(t)v(t), \quad (n \rightarrow \infty).$$

So, let $t \in [t_1, t_2]$ be fixed. Since $f \in BV([t_1, t_2]X)$, there exists a sequence of step functions $\{f_n(s)\}_{n=0}^\infty$ which converges uniformly to $f(s)$ on $[t_1, t_2]$ (see Dieudonné [2], page 139). For every $f_n(s), n = 0, 1, 2, \dots$, there exists a partition of the interval $[t_1, t]$:

$$Z : t_1 = s_0 < s_1 < s_2 < \dots < s_m < s_{m+1} = t$$

so that it holds with $f_n = c_i$ on (s_{i-1}, s_i) for $i = 1, \dots, m$:

$$\begin{aligned}
 v_n(t) &:= \int_{t_1}^t U(t, s) f_n(s) ds \\
 &= \int_{t_1=s_0}^{s_1} U(t, s) c_0 ds + \int_{s_1}^{s_2} U(t, s) c_1 ds + \int_{s_2}^{s_3} U(t, s) c_2 ds \\
 &\quad + \int_{s_3}^{s_4} U(t, s) c_3 ds + \int_{s_4}^{s_5} U(t, s) c_4 ds + \int_{s_5}^{s_6} U(t, s) c_5 ds \\
 &\quad + \dots + \int_{s_{m-1}}^{s_m} U(t, s) c_{m-1} ds + \int_{s_m}^{s_{m+1}=t} U(t, s) c_m ds, \tag{3.3}
 \end{aligned}$$

where $c_i = f(s_i^*)$, $s_i < s_i^* < s_{i+1}$, $i = 0, 1, \dots, m$ and $c_i \in X$ are constants.

We show now that every one of these integrals belongs to D and that $A(t)$ applied on these integrals (i.e. on v_n) implies the boundedness of the sequence $A(t)v_n(t)$.

Since $f(s)$ is of bounded variation on $[t_1, t_2]$, $f(s)$ is bounded on $[t_1, t_2]$. The uniform boundedness of the operator $U(t, s) : X \rightarrow X$ implies the uniform convergence:

$$U(t, s) f_n(s) \longrightarrow U(t, s) f(s), \quad (n \rightarrow \infty)$$

on $[t_1, t]$ so that it holds via Lebesgue’s Theorem:

$$\begin{aligned}
 v(t) &= \int_{t_1}^t U(t, s) f(s) ds = \int_{t_1}^t \lim_{n \rightarrow \infty} U(t, s) f_n(s) ds \\
 &= \lim_{n \rightarrow \infty} \int_{t_1}^t U(t, s) f_n(s) ds = \lim_{n \rightarrow \infty} v_n(t).
 \end{aligned}$$

For $x \in D$, $A(s)x$ is continuously differentiable in s . Moreover the inverses $A^{-1}(s)$ exist and are bounded. Using the identity: $(A^{-1}(s)x)' = -A^{-1}(s)A'(s)A^{-1}(s)x$, $x \in X$, and applying (e) from Preliminaries yields via integration by parts on an arbitrary integral from (3.3):

$$\begin{aligned}
 I_i &:= \int_{s_i}^{s_{i+1}} U(t, s) c_i ds \\
 &= \int_{s_i}^{s_{i+1}} \left\{ \frac{\partial}{\partial s} U(t, s) \right\} A^{-1}(s) c_i ds \\
 &= \{U(t, s_{i+1})A^{-1}(s_{i+1}) - U(t, s_i)A^{-1}(s_i)\} c_i \\
 &\quad + \int_{s_i}^{s_{i+1}} U(t, s) A^{-1}(s) A'(s) A^{-1}(s) c_i ds.
 \end{aligned}$$

That means via (g) and (h) from Preliminaries:

$$\begin{aligned}
 v_n &= \sum_{i=0}^m I_i \\
 &= \sum_{i=0}^m \{U(t, s_{i+1})A^{-1}(s_{i+1}) - U(t, s_i)A^{-1}(s_i)\}c_i \\
 &\quad + \sum_{i=0}^m \int_{s_i}^{s_{i+1}} U(t, s)A^{-1}(s)A'(s)A^{-1}(s)c_i ds \\
 &= A^{-1}(t) \left(\sum_{i=0}^m \{A(t)U(t, s_{i+1})A^{-1}(s_{i+1}) - A(t)U(t, s_i)A^{-1}(s_i)\}c_i \right. \\
 &\quad \left. + \sum_{i=0}^m \int_{s_i}^{s_{i+1}} \{A(t)U(t, s)A^{-1}(s)\} \{A'(s)A^{-1}(s)\}c_i ds \right) \in D.
 \end{aligned}$$

So, we have by reordering of the terms of the finite sum:

$$\begin{aligned}
 A(t)v_n(t) &= c_m - A(t)U(t, t_1)A^{-1}(t_1)c_0 \\
 &\quad - A(t)U(t, s_1)A^{-1}(s_1)(c_1 - c_0) - A(t)U(t, s_2)A^{-1}(s_2)(c_2 - c_1) \\
 &\quad - A(t)U(t, s_3)A^{-1}(s_3)(c_3 - c_2) - \dots - A(t)U(t, s_m)A^{-1}(s_m)(c_m - c_{m-1}) \\
 &\quad + \sum_{i=0}^m \int_{s_i}^{s_{i+1}} \{A(t)U(t, s)A^{-1}(s)\} \{A'(s)A^{-1}(s)\}c_i ds,
 \end{aligned}$$

i.e. with $\|c_i\| \leq \sup_{s \in [t_1, t_2]} \|f(s)\|$ and $s_i \leq s_i^* \leq s_{i+1}$:

$$\begin{aligned}
 \|A(t)v_n(t)\| &\leq \|f(s_m^*) - A(t)U(t, t_1)A^{-1}(t_1)f(s_0^*)\| \\
 &\quad + k_1 \sum_{i=0}^{m-1} \|c_{i+1} - c_i\| + k_1 k_2 \left(\sup_{s \in [t_1, t_2]} \|f(s)\| \right) (t - t_1) \\
 &= \|f(s_m^*) - f(t_1) + f(t_1) - A(t)U(t, t_1)A^{-1}(t_1)\{f(s_0^*) - f(t_1) + f(t_1)\}\| \\
 &\quad + \dots \\
 &\leq \|f(s_m^*) - f(t_1)\| + k_1 \|f(s_0^*) - f(t_1)\| \\
 &\quad + \|\{I - A(t)U(t, t_1)A^{-1}(t_1)\}f(t_1)\| \\
 &\quad + k_1 \int_{t_1}^t \|df(s)\| + k_1 k_2 (\sup_{s \in [t_1, t_2]} \|f(s)\|) (t - t_1) \\
 &\leq \|\{I - A(t)U(t, t_1)A^{-1}(t_1)\}f(t_1)\| + (1 + 2k_1) \int_{t_1}^t \|df(s)\| \\
 &\quad + k_1 k_2 \left(\sup_{s \in [t_1, t_2]} \|f(s)\| \right) (t - t_1).
 \end{aligned}$$

This boundedness of the sequence $\{A(t)v_n(t)\}_{n=0}^\infty$ in reflexive Banach space X implies by reflexivity of X the existence of a subsequence $\{A(t)v_{n_k}(t)\}_{k=0}^\infty$ which converges weakly in X . So, altogether we have:

$$v_{n_k}(t) \longrightarrow v(t) \quad , \quad A(t)v_{n_k}(t) \rightharpoonup g(t) \quad (k \rightarrow \infty).$$

The reflexivity of X yields finally that $v(t) \in D$ and $g(t) = A(t)v(t)$ as well as:

$$\begin{aligned} \|A(t)v(t)\| &\leq \liminf_{k \rightarrow \infty} \|A(t)v_{n_k}(t)\| \\ &\leq \|\{I - A(t)U(t, t_1)A^{-1}(t_1)\}f(t_1)\| + (1 + 2k_1) \int_{t_1}^t \|df(s)\| \\ &\quad + k_1k_2(\sup_{s \in [t_1, t_2]} \|f(s)\|)(t - t_1), \end{aligned}$$

i.e. the desired estimate (3.2).

(II) We prove now the Lipschitz continuity of $v(t)$. By the properties of the evolution family (see Sect. 2):

$$\begin{aligned} v(t + \varepsilon) - v(t) &= \int_{t_1}^{t+\varepsilon} U(t + \varepsilon, s)f(s)ds - \int_{t_1}^t U(t, s)f(s)ds \\ &= U(t + \varepsilon, t) \int_{t_1}^t U(t, s)f(s)ds + \int_t^{t+\varepsilon} U(t + \varepsilon, s)f(s)ds \\ &\quad - \int_{t_1}^t U(t, s)f(s)ds \\ &= \{U(t + \varepsilon, t) - I\}v(t) + \int_t^{t+\varepsilon} U(t + \varepsilon, s)f(s)ds. \end{aligned}$$

Here Lipschitz continuity of the integral term is obvious. By the just proved boundedness of $\|A(t)v(t)\|$ we obtain:

$$\begin{aligned} \|\{U(t + \varepsilon, t) - I\}v(t)\| &= \left\| \int_0^1 \frac{\partial}{\partial \tau} U(t + \tau\varepsilon, t)v(t)d\tau \right\| \\ &= \varepsilon \left\| \int_0^1 A(t + \tau\varepsilon)U(t + \tau\varepsilon, t)v(t)d\tau \right\| \\ &\leq \varepsilon \int_0^1 \|A(t + \tau\varepsilon)U(t + \tau\varepsilon, t)A^{-1}(t)\| \|A(t)v(t)\|d\tau \\ &\leq K_1\varepsilon. \end{aligned}$$

So, we have $v(t) \in C^{0,1}([t_1, t_2], X)$.

(III) Let $f \in C^0([t_1, t_2], X)$. We show in $[t_1, t_2]$:

$$A(t + \varepsilon)v(t + \varepsilon) - A(t)v(t) \longrightarrow 0, \quad (\varepsilon \rightarrow 0).$$

For $\varepsilon > 0$ (for $\varepsilon < 0$ it is similar), it is:

$$\begin{aligned}
 & A(t + \varepsilon) \int_{t_1}^{t+\varepsilon} U(t + \varepsilon, s) f(s) ds - A(t) \int_{t_1}^t U(t, s) f(s) ds \\
 &= A(t + \varepsilon) \int_{t_1}^t U(t + \varepsilon, s) f(s) ds - A(t) \int_{t_1}^t U(t, s) f(s) ds \\
 &\quad + A(t + \varepsilon) \int_t^{t+\varepsilon} U(t + \varepsilon, s) f(s) ds \\
 &= \{A(t + \varepsilon)U(t + \varepsilon, t) - A(t)\} \int_{t_1}^t U(t, s) f(s) ds \\
 &\quad + A(t + \varepsilon) \int_t^{t+\varepsilon} U(t + \varepsilon, s) f(s) ds \\
 &=: T_1(\varepsilon) + T_2(\varepsilon).
 \end{aligned}$$

Part (I) above together with (g) from Preliminaries yields:

$$T_1(\varepsilon) = \{A(t + \varepsilon)U(t + \varepsilon, t)A^{-1}(t) - I\}A(t) \int_{t_1}^t U(t, s) f(s) ds \longrightarrow 0, \quad (\varepsilon \rightarrow 0).$$

Regarding the second term above, it is first according to estimate (3.2) from part (I):

$$\begin{aligned}
 \|T_2(\varepsilon)\| &= \left\| A(t + \varepsilon) \int_t^{t+\varepsilon} U(t + \varepsilon, s) f(s) ds \right\| \\
 &\leq \left\| \{I - A(t + \varepsilon)U(t + \varepsilon, t)A^{-1}(t)\} f(t) \right\| \\
 &\quad + k_1 k_2 (\sup_{s \in [t_1, t_2]} \|f(s)\|) \varepsilon + (2k_1 + 1) \int_t^{t+\varepsilon} \|df(s)\|.
 \end{aligned}$$

With $(\varepsilon \rightarrow 0)$, the first term tends to 0 due to (g) from Preliminaries, the same holds for the last integral term because of the continuity of $f(t)$ (see Jawad [4], Lemma 2.2, holds in Banach space as well).

Summarized, the assertion of (III) follows. □

4 The differentiability of v

In this closing chapter, we prove the differentiability of the integral:

$$v(t) = \int_{t_1}^t U(t, s) f(s) ds.$$

We show then that $u(t) := U(t, t_1)\varphi - v(t)$ represents the unique strong solution of our differential equation (1.1).

Theorem 4.1 (I) (Uniqueness) *Let the function $u : [t_1, t_2] \rightarrow X$ be absolutely continuous and satisfy (1.1) a.e. with $f \in L^1([t_1, t_2], X)$. Then $u(t)$ is uniquely determined by (1.2).*

(II) *Let $f \in BV([t_1, t_2], X)$, then:*

$$\frac{d^\pm v(t)}{dt} = f(t \pm 0) + A(t) \int_{t_1}^t U(t, s)f(s)ds \tag{4.1}$$

$$= f(t \pm 0) - A(t)v(t), \quad t \in [t_1, t_2] \text{ resp. } (t_1, t_2]. \tag{4.2}$$

According to (I), $u(t) := U(t, t_1)\varphi - v(t)$ is the unique strong solution of (1.1).

(III) *If additionally $f \in C^0([t_1, t_2], X)$, then it is moreover:*

$$v(t) \in C^0([t_1, t_2], D) \cap C^1([t_1, t_2], X). \tag{4.3}$$

This means that $u(t)$ is a strict solution.

Proof (I) Multiplying both sides of (1.1) by $U(t, s)$ yields:

$$U(t, s)(u'(s) + A(s)u(s) + f(s)) = 0,$$

i.e.

$$\frac{\partial}{\partial s} U(t, s)u(s) + U(t, s)f(s) = 0.$$

Now integration from $s = t_1$ to t leads to the integral equation (1.2).

(II) Since the function f is of bounded variation, $f(t \pm 0)$ exist, $t \in [t_1, t_2]$ resp. $(t_1, t_2]$. It is for $\varepsilon > 0$:

$$\begin{aligned} v(t + \varepsilon) - v(t) &= \int_{t_1}^{t+\varepsilon} U(t + \varepsilon, s)f(s)ds - \int_{t_1}^t U(t, s)f(s)ds \\ &= \{U(t + \varepsilon, t) - I\} \int_{t_1}^t U(t, s)f(s)ds + \int_t^{t+\varepsilon} U(t + \varepsilon, s)f(s)ds, \end{aligned}$$

where according to Theorem 3.1(I) and (e) from Preliminaries:

$$\frac{1}{\varepsilon} \{U(t + \varepsilon, s) - I\} \int_{t_1}^t U(t, s)f(s)ds \longrightarrow A(t) \int_{t_1}^t U(t, s)f(s)ds, \quad (\varepsilon \rightarrow 0).$$

So, it remains only to show:

$$\frac{1}{\varepsilon} \int_t^{t+\varepsilon} U(t + \varepsilon, s)f(s)ds \longrightarrow f(t + 0), \quad (\varepsilon \rightarrow 0+). \tag{4.4}$$

It is first:

$$\begin{aligned} & \frac{1}{\varepsilon} \int_t^{t+\varepsilon} U(t+\varepsilon, s) f(s) ds \stackrel{(s:=t+\tau)}{=} \frac{1}{\varepsilon} \int_0^\varepsilon U(t+\varepsilon, t+\tau) f(t+\tau) d\tau \\ &= \frac{1}{\varepsilon} \int_0^\varepsilon U(t+\varepsilon, t+\tau) \{f(t+\tau) - f(t+0)\} d\tau \\ & \quad + \frac{1}{\varepsilon} \int_0^\varepsilon U(t+\varepsilon, t+\tau) f(t+0) d\tau \\ &=: I_1(\varepsilon) + I_2(\varepsilon) \end{aligned}$$

with ((b) from Preliminaries):

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \|I_1(\varepsilon)\| &\leq \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_0^\varepsilon M e^{\omega(\varepsilon-\tau)} \|f(t+\tau) - f(t+0)\| d\tau \\ &\leq M \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_0^\varepsilon \|f(t+\tau) - f(t+0)\| d\tau = 0. \end{aligned}$$

So, we just have to show:

$$\lim_{\varepsilon \rightarrow 0} I_2(\varepsilon) = \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_0^\varepsilon U(t+\varepsilon, t+\tau) f(t+0) d\tau = f(t+0), \quad (4.5)$$

which is equivalent to:

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_0^\varepsilon \{U(t+\varepsilon, t+\tau) - I\} f(t+0) d\tau = 0.$$

It is in this context:

$$\begin{aligned} & \frac{1}{\varepsilon} \left\| \int_0^\varepsilon \{U(t+\varepsilon, t+\tau) - I\} f(t+0) d\tau \right\| \\ & \leq \frac{1}{\varepsilon} \int_0^\varepsilon \|\{U(t+\varepsilon, t+\tau) - I\} f(t+0)\| d\tau \\ & \leq \sup_{\tau \in [0, \varepsilon]} \|\{U(t+\varepsilon, t+\tau) - I\} f(t+0)\| \cdot \frac{1}{\varepsilon} \int_0^\varepsilon d\tau \\ & = \|\{U(t+\varepsilon, t+\varepsilon_0) - I\} f(t+0)\|, \quad \varepsilon_0 \in [0, \varepsilon]. \end{aligned}$$

For every $\varepsilon > 0$ there exists at least one $\varepsilon_0 \in [0, \varepsilon]$ so that this relation holds. On the other hand, we have according to (b) from Preliminaries that for every $x \in X$, $U(t, s)x$ is jointly continuous in s and t . It follows:

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \|\{U(t+\varepsilon, t+\varepsilon_0) - I\} f(t+0)\| &= \lim_{\varepsilon_0 \leq \varepsilon \rightarrow 0} \|\{U(t+\varepsilon, t+\varepsilon_0) - I\} f(t+0)\| \\ &= 0, \end{aligned}$$

and (4.5) holds. Summarized, we have:

$$\begin{aligned}
 \frac{d^+v(t)}{dt} &= \lim_{\varepsilon \rightarrow 0} \frac{v(t + \varepsilon) - v(t)}{\varepsilon} \\
 &= f(t + 0) - A(t) \int_{t_1}^t U(t, s) f(s) ds \\
 &= f(t + 0) - A(t)v(t), \quad t \in [t_1, t_2].
 \end{aligned}$$

Arguing similarly for $\varepsilon < 0$ we obtain:

$$\begin{aligned}
 \frac{d^-v(t)}{dt} &= f(t - 0) - A(t) \int_{t_1}^t U(t, s) f(s) ds \\
 &= f(t - 0) - A(t)v(t), \quad t \in (t_1, t_2].
 \end{aligned}$$

Consequently, $u(t) := U(t, t_1)\varphi - v(t)$ is the unique strong solution of (1.1), for obviously $u'(t) \in L^1([t_1, t_2], X)$ (see also (3.2)).

(III) Follows immediately from part (II) and Theorem 3.1(III). \square

References

1. Bárta, T.: A generation theorem for hyperbolic equations with coefficients of bounded variation in time. *Riv. Mat. Univ. Parma* **9**(7), 17–30 (2008)
2. Dieudonné, J.: *Foundations of Modern Analysis*. Academic Press, New York (1960)
3. Dorroh, J.R.: A simplified proof of a theorem of Kato on linear evolution equations. *J. Math. Soc. Jpn.* **27**(3), 274–478 (1975)
4. Jawad, S.: Zur Regularität von drei Integralen im Hilbertraum. *Anal. Int. Math. J. Anal. Appl.* **30**, 261–270 (2010)
5. Jawad, S.: Existenz und Eindeutigkeit starker und klassischer Lösungen für inhomogene hyperbolische Differentialgleichungen im Hilbertraum. *Monatshefte für Mathematik* **180**(4), 765–783 (2016)
6. Kato, T.: Integration of the equation of evolution in Banach space. *J. Math. Soc. Jpn.* **5**, 208–234 (1953)
7. Kato, T.: Linear evolution equations of “hyperbolic” type. *J. Fac. Sci. Univ. Tokyo, Sect. I*, **17**, 241–258 (1970)
8. Pazy, A.: *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Springer, Berlin (1983)