

Alteration in the GABAergic network of the prefrontal cortex in a potential animal model of psychosis

S. Brummelte, J. Neddens, G. Teuchert-Noodt

Department of Neuroanatomy, Faculty of Biology, University of Bielefeld, Bielefeld, Germany

Received: April 4, 2006 / Accepted: November 26, 2006 / Published online: January 3, 2007
© Springer-Verlag 2007

Summary The GABAergic input on cortical pyramidal cells has an important influence on the firing activity of the cortex and thus in regulating the behavioural outcome. The aim of the current study was to investigate the long-term neuroplastic adaptation of the GABAergic innervation pattern after an early severe systemic impact. Therefore 40 Mongolian gerbils (*Meriones unguiculatus*) were either reared under impoverished (IR) or enriched rearing conditions (ER) and received a single early (+)-methamphetamine (MA) challenge (50 mg/kg i.p.) or saline on postnatal day 14. The density of perisomatic immunoreactive GABAergic terminals surrounding layers III and V pyramidal neurons was quantified as well as the overall GABAergic fibre density in layers I/II and V of the medial prefrontal cortex (mPFC) of young adult animals (90 days). We found that IR in combination with an early MA administration led to a significant decrease in GABAergic bouton densities while the overall GABAergic fibre density increased in all investigated layers. The results indicate a shift in inhibition from somatic to dendritic innervation of pyramidal neurons in this potential animal model of psychosis. We conclude that IR combined with early MA trigger changes in the postnatal maturation of the prefrontal cortical GABAergic triggers innervation, which may interfere with proper signal processing within the prefrontal neural network.

Keywords: GABA, immunohistochemistry, schizophrenia, gerbil

Introduction

The interaction of the different transmitter systems plays a decisive role for the functioning of neural circuits throughout the brain. Several transmitters, such as gamma-aminobutyric acid (GABA), serotonin and dopamine, contribute to the modulation of activity of the cortical pyramidal neurons, and thus have an important influence on the behavioural outcome. Every segment of the pyramidal neuron,

from the initial axonal segment and the cell body up to dendritic spines, receives dense GABAergic innervation (Hendry et al., 1983; Houser et al., 1983; Beaulieu et al., 1992). Intriguingly, each of these segments receives its innervation from a distinct subpopulation of GABAergic neurons (Kisvarday et al., 1990).

Somatic GABAergic boutons are mainly build by the basket cell subpopulation, which owe their name to the basket-like arrangement of synapses surrounding pyramidal cell bodies (DeFelipe and Fairen, 1982; Hendry et al., 1983). The majority of cortical basket cells express the calcium-binding protein parvalbumin [PV (Hendry et al., 1989; Kawaguchi and Kubota, 1996)] and they mostly have a fast-spiking firing pattern (Kawaguchi and Kondo, 2002). A second type of cortical GABA cells, the chandelier neurons, are also associated with PV and are known to produce mainly axo-axonic contacts, which form axonal ‘cartridges’ along the initial axonal segment of the pyramidal neurons (Somogyi et al., 1982; Conde et al., 1994; Gabbott and Bacon, 1996). Beside these two populations of powerful interneurons, there are additional groups of GABAergic cells, which usually contain the calcium-binding proteins calbindin (CB) or calretinin (CR) and which are known to innervate primarily the dendritic spines and shafts of the pyramidal neuron (Conde et al., 1994; Gabbott and Bacon, 1996; Radnikow et al., 2002) and are thus less powerful in regulating the firing pattern of pyramidal cells. In summary, the subpopulations of interneurons each participate differently in establishing and maintaining the activity of cortical networks. Disturbances in this inhibitory regulation may result in extensive impairments in cognitive and behav-

Correspondence: Susanne Brummelte, Department of Neuroanatomy, Faculty of Biology, University of Bielefeld, Universitätsstraße 25, 33615 Bielefeld, Germany
e-mail: sbrummelte@uni-bielefeld.de

itorial function, like those seen in schizophrenia (Benes and Berretta, 2001).

Our lab has developed a potential animal model of schizophrenia using a combination of a single early methamphetamine (MA) intoxication on postnatal day 14, which damages monoaminergic fibers (Ricaurte et al., 1980, 1982), and chronically impoverished rearing conditions (IR) of gerbils. Among effects in several areas of the limbic-cortical system, the model impairs the maturation of the prefrontal cortex (PFC) by inducing diminished dopamine innervation (Dawirs et al., 1994; Neddens et al., 2001), increased GABA innervation (Nossoll et al., 1997), altered shape of pyramidal cells (Blaesing et al., 2001), and 'miswiring' of prefrontal efferents (Witte et al., 2006; Bagorda et al., 2006). In effect, the model successfully mimics several characteristics of the schizophrenic human brain (Feinberg, 1982; Weinberger and Lipska, 1995; Akil et al., 1999).

Since numerous alterations concerning the prefrontal GABAergic network have been reported for schizophrenia, as e.g. a particular defect in the parvalbumin-class of interneurons [reviewed in Blum and Mann (2002)], the current study was designed to investigate whether or how the GABAergic system may be affected in our animal model. Therefore, we analyzed different GABAergic structures, namely fibers and somatic terminals in the medial PFC of the developmentally disturbed gerbils.

Material and methods

Animals and rearing conditions

All experimental procedures were approved by the appropriate committee for animal care in accordance with the guidelines of the European Communities Council Directive. Breeding gerbils (*Meriones unguiculatus*) were obtained from Harlan Winkelmann (Borchen, Germany). Gerbils were chosen due to their very small genetic variability (Thiessen and Yahr, 1977), their rich wild-type like behavioural repertoire, and complex social interaction (Rosenzweig and Bennett, 1969).

A total of 40 males (weight 66–90 g) were used in this study. Half of them were bred in standard makrolon cages (type IV) whereas the other half were bred in semi-naturally structured compounds containing branches and hiding opportunities (1 × 1 m; enriched condition). At weaning (30 days), the gerbils that were born in cages were assigned to impoverished conditions (IR, animals kept alone in standard makrolon cages type III without any content except of sawdust), while the other group grew up as a group of siblings under enriched rearing conditions (ER, kept in compounds similar to those they were born in), both for further 60 days. On postnatal day 14 a total of 20 animals received a single injection of (+)-methamphetamine hydrochloride [Sigma (50 mg/kg; i.p.)], whereas the remaining 20 gerbils were sham-treated with saline, resulting in four experimental groups: ER-Sal, ER-MA, IR-Sal, IR-MA; $n = 10$ for each group. All animals had free access to food and water and were kept on natural day/night cycles during summer season.

Immunohistochemistry

On PD 90, animals were transcardially perfused under deep chloralhydrate anesthesia (1.7 g/kg, i.p.). The perfusion was performed with 200 ml 0.05 M

phosphate buffer (pH 6.2), containing 1% sodium metabisulfite, followed by 750 ml 5% glutaraldehyde with 1% sodium metabisulfite in 0.1 M phosphate buffer (pH 7.5). Immediately after perfusion, the brains were removed and postfixed for 30 min. Coronal sections of 50 μ m were cut with a vibratome (Vibratome Series 1000, Technical Products International Inc.) of which every 3rd was collected in wash buffer at 4°C. For immunostaining the sections were rinsed 3 × 10 min in cold wash buffer, followed by a preincubation in 10% normal goat serum and 0.4% Triton X-100 (Sigma) for 30 min. Subsequent the sections were incubated with rabbit anti-GABA (ImmunoStar, Hudson, WI), diluted 1:5000 with 1% normal goat serum and 0.4% Triton X-100 for 48 h.

The following rinses, all three times for 10 min, and dilutions were done in 0.05 M tris-HCL buffered saline pH 7.5 (TBS). The sections were rinsed and incubated for 30 min in biotinylated goat anti-rabbit IgG (Sigma) diluted 1:20 with 1% normal goat serum, rinsed again and incubated with ExtraAvidin-Peroxidase (Sigma) diluted 1:20 for 30 min. After another rinse the sections were stained in 0.05% 3,3-diaminobenzidine (Sigma) with 0.01% H₂O₂ for 4 min. Then the sections were washed, mounted on glass slides, dried overnight, dehydrated with ethanol, cleared with xylene and cover slipped with DePeX (Serva, Heidelberg, Germany). To avoid deviations due to possibly lateralised innervation densities of GABA only right hemispheres were used for quantification.

Quantification of GABAergic profiles

For quantification of bouton and fibre densities, brain sections were chosen in areas of interest (Fig. 1A–E) by means of anatomical characteristics according to brain atlases of the rat (Paxinos and Watson, 1986) and the mouse (Valverde, 1998); identification of the brain regions follows the nomenclature of the atlas of the rat. For the quantification of GABAergic boutons a total number of 3200 cells was analysed, with an average number of 4 analysed sections per animal and an average of 10 clearly identified pyramidal cell somata in standard test fields (0.22 mm²) per section and layer (layers III and V). A cell was chosen if the unstained soma was clearly lying within the range of layer III or V of the cingular cortex area 3 (Cg3) of the mPFC and had a round to slightly oval shape which was clearly surrounded by darkly stained GABAergic boutons (see Fig. 1G). An experimenter blind to the experimental conditions marked the pyramidal cell soma manually. All boutons in a range of 1.66 μ m from this soma were automatically assigned and the density was computed as a percentage of the evaluated test area. The fibre densities were quantified in standard test fields (900 μ m²) in layers V and I/II with an average of 10 test fields per section and layer (see Fig. 1F and H). Layer I/II was chosen due to their high innervation with GABAergic fibres inhibiting distal apical dendrites of pyramidal neurons. All detectable GABAergic boutons and fibres were visualised using a bright field microscope (Olympus BX61, Hamburg, Germany) and a digital camera for microscopy (SIS ColorViewII, Münster, Germany) at 600-fold magnification. Boutons and fibres were quantified by software for image analysis (KS300, Jenoptik, Jena, Germany), which uses a combination of Gauss filter and Gerig operator that depicts differences of grey values of adjacent pixels and transforms the result into binary images. In effect, fibres were depicted as lines of one pixel width, such that different diameters of fibres would not influence the measurement.

Data analysis

The data were computed as arithmetic means by-case and by-group \pm S.E.M. of the respective layers and were analysed for the effects of both rearing conditions and pharmacological treatment. To account for possible interactions between the somatic size of the investigated cells and the area being covered by perisomatic GABAergic boutons, the size of the pyramidal cell bodies was used as a covariate in a 2-way analysis of covariance (ANCOVA) of perisomatic terminals.




Fig. 1. Brightfield photomicrograph of a representative coronar section of the medial prefrontal cortex (A). The rectangle (B) shows the analysed section of the Cg3 region with subsequent rectangles for the analysed layers, which are magnified in (C, D and E). The GABAergic fibre density is generally similar in layers I/II and V (F and H). G shows GABAergic boutons (arrows) around an unstained pyramidal soma. Scale bars: 1 mm (A), 200 μ m (B), 50 μ m (C–E) and 20 μ m (G–H)

Statistical analysis of the overall GABAergic fibre densities was done using a factorial ANOVA. Due to technical problems, sections from 6 animals (two from each group except IR-MA) had to be excluded from

the study. All statistical analysis was computed with Statistica 6 (StatSoft, Tulsa, USA). The levels of significance were set at * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.

Results

Qualitative results

The GABAergic innervation pattern is relatively homogeneous throughout the cortex of gerbils and is similar to rats (Seto-Ohshima et al., 1990). It is characterised by a dense fibre innervation in all cortical layers with the highest density in the molecular layer. We identified immunonegative pyramidal neurons in layers III and V by their round or oval shape, their size and orientation, and the presence of basket-like GABAergic innervation (Fig. 1G).

Quantitative results

GABAergic bouton densities

The 2-way ANCOVA revealed a highly significant effect of rearing conditions on boutons in layer III [$F(1,29) = 28.59$,




Fig. 2. GABAergic bouton (A) and fibre densities (B) in the analysed layers of the PFC of gerbils from enriched (ER) and impoverished rearing (IR) conditions treated with either methamphetamine (MA) or saline given by means + standard error (S.E.M.). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

$p < 0.001$] and layer V [$F(1,29) = 25.58$, $p < 0.001$], and also a significant interaction between rearing and treatment in both layers [L III: $F(1,29) = 6.35$, $p = 0.0175$; L V: $F(1,29) = 5.0806$, $p = 0.0319$]. *Post-hoc* analysis with Newman-Keuls test showed the following results: Isolation rearing (IR) led to a significant decrease in GABAergic bouton density in layer III (-19% , $p = 0.032$), but not in layer V ($p = 0.093$). An early MA intoxication led to a further decrease in the bouton densities of both layers in IR-MA compared to IR-Sal animals (L III: -24% , $p = 0.031$; L V: -22% , $p = 0.032$). However, such an effect was not seen in animals from enriched rearing conditions (ER-MA vs. ER-Sal). Thus, bouton densities were reduced in the IR-MA group (L III: -38% , $p < 0.001$; L V: -33% , $p = 0.001$) compared to ER-Sal animals (cf. Fig. 2A).

GABAergic fibre densities

A factorial ANOVA identified a significant interaction of treatment and rearing conditions in both layer V [$F(1,30) = 13.07$, $p = 0.001$] and layers I/II [$F(1,30) = 9.8844$, $p = 0.004$]. *Post-hoc* Newman-Keuls tests revealed a significant increase in layer I/II fibre density of IR-MA animals ($+15$, $+16$, $+18\%$; all $p < 0.001$) compared to IR-Sal, ER-Sal, and ER-MA animals, respectively. A similar increase in GABAergic fibre density was found in layer V ($+17$ to $+19\%$) of the IR-MA group, compared to IR-Sal, ER-Sal, and ER-MA animals [all $p < 0.001$, except ER-Sal: $p = 0.0012$ (cf. Fig. 2B)].

Discussion

We have demonstrated that a single early MA intoxication combined with impoverished rearing (IR) significantly reduces the density of GABAergic boutons that surround layers III and V pyramidal neurons in the prefrontal cortex of the Mongolian gerbil, whereas the overall GABAergic fibre density in layers I/II and V is increased compared to control animals.

Early MA intoxication and impoverished rearing as a model for schizophrenia

The single early high dose of MA on PD 14 in combination with IR used in the current study is effective to disturb normal postnatal development of the dopaminergic system, by triggering a restraint of the maturation of dopamine fibres in the prefrontal cortex and the nucleus accumbens (Dawirs et al., 1994; Neddens et al., 2001, 2002) as well as a concomitant excessive maturation in several amygdaloid

nuclei and the entorhinal cortex (Busche et al., 2004). A similar pattern of cortical-subcortical dopaminergic imbalance has also been observed in the schizophrenic brain (Laruelle et al., 2003; Abi-Dargham, 2004). Early MA treatment additionally impairs PFC-related abilities and behaviours, such as working memory and spatial learning (Dawirs et al., 1996; Williams et al., 2002). Again, deficits in working memory are well known characteristics of schizophrenic patients (Goldman-Rakic, 1995; Lewis and Anderson, 1995). Furthermore, the early drug challenge in combination with IR leads to a miswiring of prefrontal efferents (Bagorda et al., 2006), in accordance with the dysconnection hypothesis of schizophrenia (Weinberger and Lipska, 1995).

Taken together, our approach using combined early MA intoxication and IR leads to several morphological changes in neuroanatomical brain networks and impairs cognitive functions, resembling some of the changes and deficits seen in schizophrenic individuals, and thus provides a potential animal model of the disease. The present study reveals that an early MA intoxication additionally decreases GABAergic boutons that surround pyramidal cell somata, indicating a loss of somatic synapses (Karube et al., 2004) and a concomitant increase in overall GABAergic fibre density in the medial prefrontal cortex. These findings raise the possibility that the local prefrontal cortical inhibitory network may be functionally disorganised.

The role of somatic inhibition

The distinct classes of GABAergic synapses play differential roles in regulating the activity of pyramidal neurons. The majority of GABAergic synapses terminate on dendrites or spines of the postsynaptic cells (Beaulieu and Somogyi, 1990; Beaulieu et al., 1992; Nitsch and Riesenberger, 1995), thus they are likely to control the efficacy and plasticity of excitatory inputs onto the postsynaptic target (Miles et al., 1996; Tamas et al., 1997, 2003). However, somatic inhibition is thought to be particularly effective in controlling the output of pyramidal neurons and, importantly, has been implicated to synchronize activity patterns of whole pyramidal populations (Miles et al., 1996; Tamas et al., 1997, 2000; Gibson et al., 1999; Freund, 2003; Klausberger et al., 2003). Such oscillatory synchronization is further believed to create the necessary temporal and spatial frame for prefrontal functions such as working memory (Constantinidis et al., 2002; Lewis et al., 2005). In addition, cortical interneurons, in particular 'fast-spiking' neurons, have been shown to play an important role in shaping receptive fields as well as spatial memory fields (Jones, 1993; Rao et al.,

1999, 2000). GABAergic somatic inhibition is thus exceptionally essential for the maintenance of cortical and cognitive functions and one is tempted to suggest that a decrease in this type of GABAergic inhibition and the potential subsequent deficit in synchronization might contribute to reported working memory dysfunction in schizophrenia (Lewis et al., 2005) and our animal model (Dawirs et al., 1996). In fact, post-mortem studies of schizophrenic patients reveal fewer GABAergic synapses on cortical pyramidal cells (Blum and Mann, 2002) and in addition, recent neurophysiological studies have shown, that some cognitive dysfunctions in schizophrenia are associated with an abnormal neural synchronization (Spencer et al., 2003, 2004; Lee et al., 2003; Uhlhaas et al., 2006).

The maturation and shift of GABAergic inhibition

It is well documented that GABA exhibits depolarizing effects at early postnatal stages (Cherubini et al., 1991; Ganguly et al., 2001; Ben-Ari, 2002), due to an inverted electrochemical gradient for Cl^- in neonatal neurons (Ben-Ari, 2002). The shift from an excitatory to an inhibitory transmitter is assumed to coincide with the first expression of PV in cortical interneurons (Berger et al., 1999) and the calcium-binding protein is therefore considered a marker of functional maturity of the neuron (Seto-Ohshima et al., 1990; Solbach and Celio, 1991). In the gerbil mPFC, the first PV-immunoreactive cells appear around PD 14 (unpublished data), that is, at the time of the MA challenge. Interestingly, the maturation of GABAergic synapses in general proceeds until early adulthood (Huang et al., 1999; Morales et al., 2002; Chattopadhyaya et al., 2004; Lewis et al., 2005), and in that, every subpopulation of presynaptic terminals exhibits a particular developmental pattern (Lewis et al., 2005). Therefore, the ability to synchronize pyramidal cell activity is assumed to be in substantial flux until adulthood (Lewis et al., 2005). Although the proliferation and formation of the typical perisomatic basket terminal seems to be a largely stereotypical process, it is additionally also dependent on neuronal activity within cortical circuits (Marty et al., 2000; Chattopadhyaya et al., 2004).

GABAergic interneurons receive direct dopaminergic input (Goldman-Rakic et al., 1989; Verney et al., 1990; Benes et al., 1993), with D1 and D2 receptor types being most abundantly expressed by PV-neurons (Le Moine and Gaspar, 1998). Dopamine modulates cortical GABA cells; both inhibitory (Retaux et al., 1991) and excitatory (Gorelova et al., 2002) effects on fast-spiking interneurons have been reported. The omission of prefrontal dopaminergic afferent fibres by an early MA challenge (Dawirs et al., 1994;

Neddens et al., 2001) might therefore induce significant alterations in the local GABAergic networks.

The dopaminergic afferents to the prefrontal cortex show a prolonged maturation (Kalsbeek et al., 1988; Dawirs et al., 1993; Rosenberg and Lewis, 1995) and continue to form synapses on GABAergic interneurons until early adulthood (Benes et al., 1996b). Pyramidal neurons are also directly innervated by dopaminergic terminals (Jay et al., 1995; Davidoff and Benes, 1998) which demonstrates the rather complex capacity of dopamine to directly and indirectly regulate the firing pattern of pyramidal neurons. By early MA intoxication we induce a restraint of the maturation of prefrontal dopaminergic afferents, which triggers reactive neuroplastic adaptation of the local network. Anatomical data suggest that pyramidal cells may adapt by increasing their dendritic range and their spine density (Blaesing et al., 2001). Our current findings indicate an increase of GABAergic fibre density, which is in line with an earlier study using electron-microscopy that already revealed an increase in non-somatic GABAergic terminals (Nossoll et al., 1997). Therefore, we find it tempting to suggest that an early MA challenge, by acutely reducing the density of monoaminergic innervation of the PFC, might trigger a reactive shift within the GABAergic networks from somatic to dendritic pyramidal inhibition.

GABAergic dysfunction in schizophrenia

GABAergic dysfunction in schizophrenia has first been proposed by Roberts (1972). Since then, several studies have revealed disturbances of GABAergic networks in schizophrenic patients (for review see Benes and Berretta, 2001) or in animal models of schizophrenia (Cochran et al., 2002, 2003; Keilhoff et al., 2004; Reynolds et al., 2004; Penschuck et al., 2006). A decline in PV-immunoreactive structures, particularly in axon cartridges from chandelier neurons, seems to be one of the most prevalent observations in *post-mortem* brains from schizophrenic individuals (Woo et al., 1998; Pierri et al., 1999; Lewis et al., 1999). Furthermore, the GABA_A receptor density was found to be upregulated at the axon initial segment (Volk et al., 2002) as well as at the cell body of pyramidal neurons (Benes et al., 1996a), possibly compensating for a reduction of inhibitory terminals from chandelier and basket cells (Lewis et al., 2005). In contrast to the alterations in PV-containing neurons, only few studies reported on changes in the subpopulation of CB- or CR-immunopositive cells. Iritani and colleagues (1999) found a fibre disarray from CB-containing cells in the PFC, while Daviss and Lewis (1995) described an increase in the density of CB cells but no change

in the CR population in a *post-mortem* study on schizophrenic brains. This would also be in line with our findings, since an increased number of CB cells and an altered fibre pattern are likely to present an elevated GABAergic inhibition of afferent pyramidal parts.

Conclusion

Here we present evidence for a probable dysfunctional reorganization of GABAergic networks in our potential animal model of schizophrenia. GABAergic interneurons critically contribute to the establishment of complex behaviours by controlling and synchronizing the firing patterns of pyramidal neurons. A weakened or altered inhibition may give rise to a broad array of disturbances in cognitive function, like those seen in schizophrenia (Benes and Berretta, 2001). The current study indicates a potential shift from a strong and powerful somatic inhibition to dendritic inhibition, which might attenuate the GABAergic influence on pyramidal activity and thus lead to an uncontrolled firing or abnormal synchronization. Our data coincide with findings of a reduced GABAergic somatic innervation in individuals with schizophrenia. We suggest that, in our animal model, this change in the GABAergic network is secondary, being triggered by the primary impairment of monoaminergic and namely dopaminergic afferents. Further investigations of the separate subpopulations of GABAergic interneurons in the PFC of gerbils are in process to identify the responsible cell classes for the observed alteration in the GABAergic network.

Acknowledgements

The authors wish to thank F. Bagorda, U. Schroeder and S. Graumüller for technical assistance.

References

- Abi-Dargham A (2004) Do we still believe in the dopamine hypothesis? New data bring new evidence. *Int J Neuropsychopharmacol* 7 Suppl 1: S1–S5
- Akil M, Pierri JN, Whitehead RE, Edgar CL, Mohila C, Sampson AR, Lewis DA (1999) Lamina-specific alterations in the dopamine innervation of the prefrontal cortex in schizophrenic subjects. *Am J Psychiatry* 156: 1580–1589
- Bagorda F, Teuchert-Noodt G, Lehmann K (2006) Isolation rearing or methamphetamine traumatization induce a “dysconnection” of prefrontal efferents in gerbils: implications for schizophrenia. *J Neural Transm* 113: 365–379
- Beaulieu C, Somogyi P (1990) Targets and quantitative distribution of GABAergic synapses in the visual cortex of the cat. *Eur J Neurosci* 2: 296–303

- Beaulieu C, Kisvarday Z, Somogyi P, Cynader M, Cowey A (1992) Quantitative distribution of GABA-immunopositive and -immunonegative neurons and synapses in the monkey striate cortex (area 17). *Cereb Cortex* 2: 295–309
- Ben-Ari Y (2002) Excitatory actions of gaba during development: the nature of the nurture. *Nat Rev Neurosci* 3: 728–739
- Benes FM, Berretta S (2001) GABAergic interneurons: implications for understanding schizophrenia and bipolar disorder. *Neuropsychopharmacology* 25: 1–27
- Benes FM, Vincent SL, Molloy R (1993) Dopamine-immunoreactive axon varicosities form nonrandom contacts with GABA-immunoreactive neurons of rat medial prefrontal cortex. *Synapse* 15: 285–295
- Benes FM, Vincent SL, Marie A, Khan Y (1996a) Up-regulation of GABAA receptor binding on neurons of the prefrontal cortex in schizophrenic subjects. *Neuroscience* 75: 1021–1031
- Benes FM, Vincent SL, Molloy R, Khan Y (1996b) Increased interaction of dopamine-immunoreactive varicosities with GABA neurons of rat medial prefrontal cortex occurs during the postweaning period. *Synapse* 23: 237–245
- Berger B, De GN, Alvarez C (1999) Precocious development of parvalbumin-like immunoreactive interneurons in the hippocampal formation and entorhinal cortex of the fetal cynomolgus monkey. *J Comp Neurol* 403: 309–331
- Blaesing B, Nossoll M, Teuchert-Noodt G, Dawirs RR (2001) Postnatal maturation of prefrontal pyramidal neurones is sensitive to a single early dose of methamphetamine in gerbils (*Meriones unguiculatus*). *J Neural Transm* 108: 101–113
- Blum BP, Mann JJ (2002) The GABAergic system in schizophrenia. *Int J Neuropsychopharmacol* 5: 159–179
- Busche A, Polascheck D, Lesting J, Neddens J, Teuchert-Noodt G (2004) Developmentally induced imbalance of dopaminergic fibre densities in limbic brain regions of gerbils (*Meriones unguiculatus*). *J Neural Transm* 111: 451–463
- Chattopadhyaya B, Di CG, Higashiyama H, Knott GW, Kuhlman SJ, Welker E, Huang ZJ (2004) Experience and activity-dependent maturation of perisomatic GABAergic innervation in primary visual cortex during a postnatal critical period. *J Neurosci* 24: 9598–9611
- Cherubini E, Gaiarsa JL, Ben-Ari Y (1991) GABA: an excitatory transmitter in early postnatal life. *Trends Neurosci* 14: 515–519
- Cochran SM, Fujimura M, Morris BJ, Pratt JA (2002) Acute and delayed effects of phencyclidine upon mRNA levels of markers of glutamatergic and GABAergic neurotransmitter function in the rat brain. *Synapse* 46: 206–214
- Cochran SM, Kennedy M, McKerchar CE, Steward LJ, Pratt JA, Morris BJ (2003) Induction of metabolic hypofunction and neurochemical deficits after chronic intermittent exposure to phencyclidine: differential modulation by antipsychotic drugs. *Neuropsychopharmacology* 28: 265–275
- Conde F, Lund JS, Jacobowitz DM, Baimbridge KG, Lewis DA (1994) Local circuit neurons immunoreactive for calretinin, calbindin D-28k or parvalbumin in monkey prefrontal cortex: distribution and morphology. *J Comp Neurol* 341: 95–116
- Constantinidis C, Williams GV, Goldman-Rakic PS (2002) A role for inhibition in shaping the temporal flow of information in prefrontal cortex. *Nat Neurosci* 5: 175–180
- Davidoff SA, Benes FM (1998) High-resolution scatchard analysis shows D1 receptor binding on pyramidal and nonpyramidal neurons. *Synapse* 28: 83–90
- Daviss SR, Lewis DA (1995) Local circuit neurons of the prefrontal cortex in schizophrenia: selective increase in the density of calbindin-immunoreactive neurons. *Psychiatry Res* 59: 81–96
- Dawirs RR, Teuchert-Noodt G, Czaniera R (1993) Maturation of the dopamine innervation during postnatal development of the prefrontal cortex in gerbils (*Meriones unguiculatus*). A quantitative immunocytochemical study. *J Hirnforsch* 34: 281–290
- Dawirs RR, Teuchert-Noodt G, Czaniera R (1994) The postnatal maturation of dopamine innervation in the prefrontal cortex of gerbils (*Meriones unguiculatus*) is sensitive to an early single dose of methamphetamine. A quantitative immunocytochemical study. *J Hirnforsch* 35: 195–204
- Dawirs RR, Teuchert-Noodt G, Czaniera R (1996) Ontogeny of PFC-related behaviours is sensitive to a single non-invasive dose of methamphetamine in neonatal gerbils (*Meriones unguiculatus*). *J Neural Transm* 103: 1235–1245
- DeFelipe J, Fairen A (1982) A type of basket cell in superficial layers of the cat visual cortex. A Golgi-electron microscope study. *Brain Res* 244: 9–16
- Feinberg I (1982) Schizophrenia: caused by a fault in programmed synaptic elimination during adolescence? *J Psychiatr Res* 17: 319–334
- Freund TF (2003) Interneuron diversity series: rhythm and mood in perisomatic inhibition. *Trends Neurosci* 26: 489–495
- Gabbott PL, Bacon SJ (1996) Local circuit neurons in the medial prefrontal cortex (areas 24a, b, c, 25 and 32) in the monkey: I. Cell morphology and morphometrics. *J Comp Neurol* 364: 567–608
- Ganguly K, Schinder AF, Wong ST, Poo M (2001) GABA itself promotes the developmental switch of neuronal GABAergic responses from excitation to inhibition. *Cell* 105: 521–532
- Gibson JR, Beierlein M, Connors BW (1999) Two networks of electrically coupled inhibitory neurons in neocortex. *Nature* 402: 75–79
- Goldman-Rakic PS (1995) Cellular basis of working memory. *Neuron* 14: 477–485
- Goldman-Rakic PS, Leranath C, Williams SM, Mons N, Geffard M (1989) Dopamine synaptic complex with pyramidal neurons in primate cerebral cortex. *Proc Natl Acad Sci USA* 86: 9015–9019
- Gorelova N, Seamans JK, Yang CR (2002) Mechanisms of dopamine activation of fast-spiking interneurons that exert inhibition in rat prefrontal cortex. *J Neurophysiol* 88: 3150–3166
- Hendry SH, Houser CR, Jones EG, Vaughn JE (1983) Synaptic organization of immunocytochemically identified GABA neurons in the monkey sensory-motor cortex. *J Neurocytol* 12: 639–660
- Hendry SH, Jones EG, Emson PC, Lawson DE, Heizmann CW, Streit P (1989) Two classes of cortical GABA neurons defined by differential calcium binding protein immunoreactivities. *Exp Brain Res* 76: 467–472
- Houser CR, Hendry SH, Jones EG, Vaughn JE (1983) Morphological diversity of immunocytochemically identified GABA neurons in the monkey sensory-motor cortex. *J Neurocytol* 12: 617–638
- Huang ZJ, Kirkwood A, Pizzorusso T, Porciatti V, Morales B, Bear MF, Maffei L, Tonegawa S (1999) BDNF regulates the maturation of inhibition and the critical period of plasticity in mouse visual cortex. *Cell* 98: 739–755
- Iritani S, Kuroki N, Ikeda K, Kazamatsuri H (1999) Calbindin immunoreactivity in the hippocampal formation and neocortex of schizophrenics. *Prog Neuropsychopharmacol Biol Psychiatry* 23: 409–421
- Jay TM, Glowinski J, Thierry AM (1995) Inhibition of hippocampoprefrontal cortex excitatory responses by the mesocortical DA system. *Neuroreport* 6: 1845–1848
- Jones EG (1993) GABAergic neurons and their role in cortical plasticity in primates. *Cereb Cortex* 3: 361–372
- Kalsbeek A, Voorn P, Buijs RM, Pool CW, Uylings HB (1988) Development of the dopaminergic innervation in the prefrontal cortex of the rat. *J Comp Neurol* 269: 58–72
- Karube F, Kubota Y, Kawaguchi Y (2004) Axon branching and synaptic bouton phenotypes in GABAergic nonpyramidal cell subtypes. *J Neurosci* 24: 2853–2865
- Kawaguchi Y, Kondo S (2002) Parvalbumin, somatostatin and cholecystokinin as chemical markers for specific GABAergic interneuron types in the rat frontal cortex. *J Neurocytol* 31: 277–287

- Kawaguchi Y, Kubota Y (1996) Physiological and morphological identification of somatostatin- or vasoactive intestinal polypeptide-containing cells among GABAergic cell subtypes in rat frontal cortex. *J Neurosci* 16: 2701–2715
- Keilhoff G, Becker A, Grecksch G, Wolf G, Bernstein HG (2004) Repeated application of ketamine to rats induces changes in the hippocampal expression of parvalbumin, neuronal nitric oxide synthase and cFOS similar to those found in human schizophrenia. *Neuroscience* 126: 591–598
- Kisvarday ZF, Gulyas A, Beroukas D, North JB, Chubb IW, Somogyi P (1990) Synapses, axonal and dendritic patterns of GABA-immunoreactive neurons in human cerebral cortex. *Brain* 113 (Pt 3): 793–812
- Klausberger T, Magill PJ, Marton LF, Roberts JD, Cobden PM, Buzsaki G, Somogyi P (2003) Brain-state- and cell-type-specific firing of hippocampal interneurons in vivo. *Nature* 421: 844–848
- Laruelle M, Kegeles LS, bi-Dargham A (2003) Glutamate, dopamine, and schizophrenia: from pathophysiology to treatment. *Ann Acad Sci NY* 1003: 138–158
- Le Moine C, Gaspar P (1998) Subpopulations of cortical GABAergic interneurons differ by their expression of D1 and D2 dopamine receptor subtypes. *Brain Res Mol Brain Res* 58: 231–236
- Lee KH, Williams LM, Breakspear M, Gordon E (2003) Synchronous gamma activity: a review and contribution to an integrative neuroscience model of schizophrenia. *Brain Res Brain Res Rev* 41: 57–78
- Lewis DA, Anderson SA (1995) The functional architecture of the prefrontal cortex and schizophrenia. *Psychol Med* 25: 887–894
- Lewis DA, Pierri JN, Volk DW, Melchitzky DS, Woo TU (1999) Altered GABA neurotransmission and prefrontal cortical dysfunction in schizophrenia. *Biol Psychiatry* 46: 616–626
- Lewis DA, Hashimoto T, Volk DW (2005) Cortical inhibitory neurons and schizophrenia. *Nat Rev Neurosci* 6: 312–324
- Marty S, Wehrle R, Sotelo C (2000) Neuronal activity and brain-derived neurotrophic factor regulate the density of inhibitory synapses in organotypic slice cultures of postnatal hippocampus. *J Neurosci* 20: 8087–8095
- Miles R, Toth K, Gulyas AI, Hajos N, Freund TF (1996) Differences between somatic and dendritic inhibition in the hippocampus. *Neuron* 16: 815–823
- Morales B, Choi SY, Kirkwood A (2002) Dark rearing alters the development of GABAergic transmission in visual cortex. *J Neurosci* 22: 8084–8090
- Neddens J, Brandenburg K, Teuchert-Noodt G, Dawirs RR (2001) Differential environment alters ontogeny of dopamine innervation of the orbital prefrontal cortex in gerbils. *J Neurosci Res* 63: 209–213
- Neddens J, Lesting J, Dawirs RR, Teuchert-Noodt G (2002) An early methamphetamine challenge suppresses the maturation of dopamine fibres in the nucleus accumbens of gerbils: on the significance of rearing conditions. *J Neural Transm* 109: 141–155
- Nitsch C, Riesenberger R (1995) Synaptic reorganisation in the rat striatum after dopaminergic deafferentation: an ultrastructural study using glutamate decarboxylase immunocytochemistry. *Synapse* 19: 247–263
- Nossoll M, Teuchert-Noodt G, Dawirs RR (1997) A single dose of methamphetamine in neonatal gerbils affects adult prefrontal gamma-aminobutyric acid innervation. *Eur J Pharmacol* 340: R3–R5
- Paxinos G, Watson C (1986) *The rat brain in stereotaxic coordinates*. Academic Press, New York
- Penschuck S, Flagstad P, Didriksen M, Leist M, Michael-Titus AT (2006) Decrease in parvalbumin-expressing neurons in the hippocampus and increased phencyclidine-induced locomotor activity in the rat methylazoxymethanol (MAM) model of schizophrenia. *Eur J Neurosci* 23: 279–284
- Pierri JN, Chaudry AS, Woo TU, Lewis DA (1999) Alterations in chandelier neuron axon terminals in the prefrontal cortex of schizophrenic subjects. *Am J Psychiatry* 156: 1709–1719
- Radnikow G, Feldmeyer D, Lubke J (2002) Axonal projection, input and output synapses, and synaptic physiology of Cajal-Retzius cells in the developing rat neocortex. *J Neurosci* 22: 6908–6919
- Rao SG, Williams GV, Goldman-Rakic PS (1999) Isodirectional tuning of adjacent interneurons and pyramidal cells during working memory: evidence for microcolumnar organization in PFC. *J Neurophysiol* 81: 1903–1916
- Rao SG, Williams GV, Goldman-Rakic PS (2000) Destruction and creation of spatial tuning by disinhibition: GABA(A) blockade of prefrontal cortical neurons engaged by working memory. *J Neurosci* 20: 485–494
- Retaux S, Besson MJ, Penit-Soria J (1991) Synergism between D1 and D2 dopamine receptors in the inhibition of the evoked release of [3H]GABA in the rat prefrontal cortex. *Neuroscience* 43: 323–329
- Reynolds GP, Abdul-Monim Z, Neill JC, Zhang ZJ (2004) Calcium binding protein markers of GABA deficits in schizophrenia postmortem studies and animal models. *Neurotox Res* 6: 57–61
- Ricaurte GA, Schuster CR, Seiden LS (1980) Long-term effects of repeated methylamphetamine administration on dopamine and serotonin neurons in the rat brain: a regional study. *Brain Res* 193: 153–163
- Ricaurte GA, Guillery RW, Seiden LS, Schuster CR, Moore RY (1982) Dopamine nerve terminal degeneration produced by high doses of methylamphetamine in the rat brain. *Brain Res* 235: 93–103
- Roberts E (1972) Prospects for research on schizophrenia. An hypotheses suggesting that there is a defect in the GABA system in schizophrenia. *Neurosci Res Program Bull* 10: 468–482
- Rosenberg DR, Lewis DA (1995) Postnatal maturation of the dopaminergic innervation of monkey prefrontal and motor cortices: a tyrosine hydroxylase immunohistochemical analysis. *J Comp Neurol* 358: 383–400
- Rosenzweig MR, Bennett EL (1969) Effects of differential environments on brain weights and enzyme activities in gerbils, rats, and mice. *Dev Psychobiol* 2: 87–95
- Seto-Ohshima A, Aoki E, Semba R, Emson PC, Heizmann CW (1990) Appearance of parvalbumin-specific immunoreactivity in the cerebral cortex and hippocampus of the developing rat and gerbil brain. *Histochemistry* 94: 579–589
- Solbach S, Celio MR (1991) Ontogeny of the calcium binding protein parvalbumin in the rat nervous system. *Anat Embryol (Berl)* 184: 103–124
- Somogyi P, Freund TF, Cowey A (1982) The axo-axonic interneuron in the cerebral cortex of the rat, cat and monkey. *Neuroscience* 7: 2577–2607
- Spencer KM, Nestor PG, Niznikiewicz MA, Salisbury DF, Shenton ME, McCarley RW (2003) Abnormal neural synchrony in schizophrenia. *J Neurosci* 23: 7407–7411
- Spencer KM, Nestor PG, Perlmuter R, Niznikiewicz MA, Klump MC, Frumin M, Shenton ME, McCarley RW (2004) Neural synchrony indexes disordered perception and cognition in schizophrenia. *Proc Natl Acad Sci USA* 101: 17288–17293
- Tamas G, Buhl EH, Somogyi P (1997) Fast IPSPs elicited via multiple synaptic release sites by different types of GABAergic neurone in the cat visual cortex. *J Physiol* 500 (Pt 3): 715–738
- Tamas G, Buhl EH, Lorincz A, Somogyi P (2000) Proximally targeted GABAergic synapses and gap junctions synchronize cortical interneurons. *Nat Neurosci* 3: 366–371
- Tamas G, Lorincz A, Simon A, Szabadics J (2003) Identified sources and targets of slow inhibition in the neocortex. *Science* 299: 1902–1905
- Thiessen DD, Yahr P (1977) The gerbil in behavioral investigations. Mechanisms of territoriality and olfactory communication. In: University of Texas Press, Austin

- Uhlhaas PJ, Linden DE, Singer W, Haenschel C, Lindner M, Maurer K, Rodriguez E (2006) Dysfunctional long-range coordination of neural activity during Gestalt perception in schizophrenia. *J Neurosci* 26: 8168–8175
- Valverde F (1998) Golgi atlas of the postnatal mouse brain. Springer, Wien, New York
- Verney C, Alvarez C, Geffard M, Berger B (1990) Ultrastructural double-labelling study of dopamine terminals and GABA-containing neurons in rat anteromedial cerebral cortex. *Eur J Neurosci* 2: 960–972
- Volk DW, Pierri JN, Fritschy JM, Auh S, Sampson AR, Lewis DA (2002) Reciprocal alterations in pre- and postsynaptic inhibitory markers at chandelier cell inputs to pyramidal neurons in schizophrenia. *Cereb Cortex* 12: 1063–1070
- Weinberger DR, Lipska BK (1995) Cortical maldevelopment, anti-psychotic drugs, and schizophrenia: a search for common ground. *Schizophr Res* 16: 87–110
- Williams MT, Vorhees CV, Boon F, Saber AJ, Cain DP (2002) Methamphetamine exposure from postnatal day 11 to 20 causes impairments in both behavioral strategies and spatial learning in adult rats. *Brain Res* 958: 312–321
- Witte AV, Bagorda F, Teuchert-Noodt G, Lehmann K (2006) Contralateral prefrontal projections in gerbils mature abnormally after early methamphetamine trauma and isolated rearing. *J Neural Transm*. Epub ahead of print, DOI 10.1007/s00702-006-0506-6
- Woo TU, Whitehead RE, Melchitzky DS, Lewis DA (1998) A subclass of prefrontal gamma-aminobutyric acid axon terminals are selectively altered in schizophrenia. *Proc Natl Acad Sci USA* 95: 5341–5346